

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

Synthesis of Interglycosidically S-Linked 1-Thio-Oligosaccharides Under Microwave Irradiation

El Sayed H. El Ashry^a; Laila F. Awad^a; H. M. Abdel Hamid^a; Atta I. Atta^a

^a Chemistry Department, Faculty of Science, Alexandria University, Alexandria, Egypt

To cite this Article Ashry, El Sayed H. El , Awad, Laila F. , Hamid, H. M. Abdel and Atta, Atta I.(2005) 'Synthesis of Interglycosidically S-Linked 1-Thio-Oligosaccharides Under Microwave Irradiation', *Journal of Carbohydrate Chemistry*, 24: 7, 745 – 753

To link to this Article: DOI: 10.1080/07328300500282540

URL: <http://dx.doi.org/10.1080/07328300500282540>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Synthesis of Interglycosidically S-Linked 1-Thio-Oligosaccharides Under Microwave Irradiation

El Sayed H. El Ashry, Laila F. Awad, H. M. Abdel Hamid, and Atta I. Atta

Chemistry Department, Faculty of Science, Alexandria University, Alexandria, Egypt

Microwave irradiation (MWI) has accelerated the synthesis of *S*-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyl)thiuronium bromide (**2a**), whose reaction with 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl bromide (**1a**) in the presence of Et₃N afforded stereoselectively the acetylated β , β -1-thiotrehalose **4a**. Similarly, the respective D-galactopyranosyl **4b** and 2-acetylamino-2-deoxy-D-glucopyranosyl **4c** analog as well as 4,4'-di-*O*-(2,3,4,6-tetra-*O*-acetyl- β -D-galactopyranosyl) **4d** and 4,4'-di-*O*-(2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl) **4e** derivatives of 2,2',3,3',6,6'-hexa-*O*-acetyl β , β -1-thiotrehalose were prepared.

Keywords Microwave, Thio-oligosaccharides, Thiotrehaloses, Glycosyl isothiuronium salt, Glycosyl halide

INTRODUCTION

Thioglycosides are considered an important class of carbohydrate derivatives.^[1–4] There is increasing interest in using them as glycosyl donors,^[5] and they are promising candidates for carbohydrate mimics, potential thera-

Address correspondence to El Sayed H. El Ashry, Chemistry Department, Faculty of Science, Alexandria University, Alexandria, Egypt. E-mail: eelashry60@hotmail.com; eelashry60@link.net

Atta I. Atta is on leave from Faculty of Medical Sciences, Al-Hodiedah University, Yemen.

peutics, and enzyme inhibitors.^[6] The interest in their enzyme inhibition^[1,2] is due to their stability to the action of glycosidases, which is more promising than that of the corresponding *O*-glycosides. Thus, they can inhibit enzymatic hydrolysis, which may be a versatile approach to map the active site of glycosyl hydrolases. The thioanalog of oligosaccharides have gained a recent continuous development not only devoted to their synthesis but also for biochemical and x-ray studies.^[1,2,7-9] 1,1-Thio- α,α -trehalose and α -D-glucopyranosyl-1-thio- α -D-mannopyranoside have been found to be inhibitors for cockchafer trehalase.^[10] Moreover, thiosucrose inhibits levansucrase from *Bacillus subtilis* and yeast invertase.^[11] Although the β,β -thiotrehalose was synthesized^[12] in 1917, the α,α -thiotrehalose was synthesized^[13] 60 years later, in 1978. Since then a number of their analog have been prepared and various conditions utilized.^[14-23]

RESULTS AND DISCUSSION

Recently there has been a growing interest in employing microwave irradiation (MWI) for the synthesis of organic compounds.^[24-27] This technique is based on the empirical observation that some organic reactions proceed much faster and with higher yields under MWI compared to conventional heating. Continuing our work^[27] on accelerating organic reactions under MWI, we have developed a convenient, efficient, and practical one-pot method for the synthesis of β,β -thio-oligosaccharides.

The β,β -thiotrehaloses have been conventionally prepared from the reaction of 2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl bromide with either potassium sulfide or 2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranose.^[12,16] The former can be catalyzed by the presence of quaternary alkylammonium^[14] or phosphonium^[15] salts.

In the present work, the readily available *S*-(2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyl)-thiuronium bromide^[4] (**2a**) has been used under MWI as a source for generating the respective thiol at the anomeric center to act as a nucleophile that upon reaction with the anomeric electrophile, generated from **1a**, would give the respective thiotrehalose. Thus, the precursor **2a** was prepared by MWI of a solution of **1a** and thiourea in acetone for 3 min. Then the electrophilic thiol was generated, in situ, by MWI of **2a** in a mixture with **1a** in the presence of triethylamine as a catalyst to give 2,3,4,6-tetra-*O*-acetyl- β -D-glucopyranosyl 2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside (acetyl derivative of β,β -1-thiotrehalose) (**4a**) within 3 min in a higher yield than that resulting from the conventional method.^[15] The galacto and 2-acetyl-amino-2-deoxy analog **4b** and **4c** were also obtained from the reaction of **1b** with **2b** and **1c** with **2c**, respectively, under similar conditions (Fig. 1).

Similarly, the reaction was extended to the disaccharides, whereby the isothiuronium derivatives **2d** and **2e** were prepared from the respective

bromides **1d** and **1e** by reaction with thiourea under MWI. Further reaction of **1d** with **2d** and **1e** with **2e** in the presence of Et₃N under MWI gave 4,4'-di-*O*-(2,3,4,6-tetra-*O*-acetyl- β -D-galactopyranosyl) 2,2',3,3',6,6'-hexa-*O*-acetyl- β , β -1-thiotrehalose (**4d**) and 4,4'-di-*O*-(2,3,4,6-tetra-*O*-acetyl- α -D-glucopyranosyl) 2,2',3,3',6,6'-hexa-*O*-acetyl- β , β -1-thiotrehalose (**4e**), respectively. The structures of the products were confirmed by their spectral data. The β -anomeric configuration in the products was readily deduced from their ¹H NMR spectra; H-1 appeared as a doublet with J_{1,2} value 10.2 Hz, and ¹³C-1 appeared at δ_C 80.7 in compound **4a**. Moreover, the DEPT, HMQC, and DQF COSY correlations assisted in assigning the spectra; the symmetrical nature of such oligosaccharides are apparent in their spectra. Similarly, the spectra of **4b–e** showed high coupling constant values between H-1 or H-1a and H-2 or H-2a, respectively, indicating β -orientation at such anomeric positions. Moreover, the orientation of the glycosidic linkage within the lactosyl and maltosyl moieties was confirmed by the coupling of H-1b with H-2b with values 9.0 and 3.8 Hz, respectively for the β - and α -orientations. Their carbons appeared at δ_C 101.1 and 95.6, respectively.

In conclusion, a practical one-pot stereoselective synthesis of the nonreducing thio-oligosaccharides has been achieved by using MWI technology in higher yields than by conventional methods in addition to the quick termination of the reaction. The reactions were clean, and pure products have been readily obtained without chromatography. Moreover, the *S*-glycosyl isothiuronium salts have been also readily prepared under MWI.

EXPERIMENTAL

General Methods

Melting points were determined on a Mel-temp apparatus and are uncorrected. ¹H NMR and ¹³C NMR spectra were recorded on a Bruker DRX 600 MHz, Jeol spectrometer (500 MHz) and a Bruker Advance 300 MHz spectrometer. The chemical shifts are expressed on the δ -scale using Me₄Si as a standard, and coupling-constant values are given in Hz. The assignments of ¹H NMR spectra were based on chemical-shift correlation DQFCOSY spectra, while the assignment of ¹³C NMR spectra were based on heteronuclear multiple quantum coherence (HMQC) experiments. TLC was performed on Merck Silica Gel 60F254 with detection by charring in sulfuric acid and by UV light. Irradiation was achieved using a domestic microwave oven EM-230 M (800-watt output power). Microanalyses were performed in the Microanalysis Unit at the Faculty of Science, Cairo University.

Glycosyl isothiuronium salts (2a–e): General Procedure. A mixture of the glycosyl halide **1a–e** (1.0 mmol) and thiourea (1.0 mmol) in dry acetone




Figure 1: Synthesis of thiotrehaloses.

(5 mL) was irradiated for 3 min in a domestic microwave oven EM-230 M (800-watt output power). The reaction mixture was left to cool at rt to afford products of **2a–e**.

2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyl isothiuronium bromide (2a). mp 203–205°C, lit^[4] mp 205°C; yield 90%.

2,3,4,6-Tetra-O-acetyl- β -D-galactopyranosyl isothiuronium bromide (2b). mp 168–170°C, lit^[28] mp 169.5°C; yield 88%

2-Acetamido-3,4,6-tri-O-acetyl-2-deoxy- β -D-glucopyranosyl isothiuronium chloride (2c). 180–182°C, lit^[29] mp 179–181°C, yield 89%.

2,3,4,6-Tetra-O-acetyl- β -D-galactopyranosyl-(1 \rightarrow 4)-2,3,6-tri-O-acetyl- β -D-glucopyranosyl isothiuronium bromide (2d). Syrup,^[31] yield 83%; ¹H NMR (500 MHz, DMSO-*d*₆): 1.92, 1.95, 1.98, 2.04 (5s, 21H, 7 \times CH₃CO), 3.94–3.99 (m, 2H, H-5a, H-4a), 4.09–4.13 (m, 4H, H-5b, H-6'a, H-6b, H-6'b), 4.41 (dd, 1H, $J_{6a,6'a}$ = 11.5 Hz, H-6a), 4.87 (dd, 1H, $J_{3b,4b}$ = 3.8 Hz, $J_{3b,2b}$ = 9.9 Hz, H-3b), 4.95 (d, 1H, H-1b), 4.98 (dd, 1H, H-2a), 5.19 (t, 1H, $J_{2b,3b}$ = $J_{2b,1b}$ = 9.9 Hz, H-2b), 5.23 (d, 1H, $J_{4b,5b}$ = 3.8 Hz, H-4b), 5.31 (dd, 1H, $J_{3a,4a}$ = 9.2 Hz, $J_{3a,2a}$ = 9.9 Hz, H-3a), 5.63 (d, 1H, $J_{1a,2a}$ = 10.0 Hz, H-1a), 9.25, 9.00 (2s, 2 \times 2H, 2 \times NH₂).

2,3,4,6-Tetra-O-acetyl- α -D-glucopyranosyl-(1 \rightarrow 4)-2,3,6-tri-O-acetyl- β -D-glucopyranosyl isothiuronium bromide (2e). mp 199–201°C; yield 86%; ¹H NMR (500 MHz, DMSO-*d*₆): δ = 1.91, 1.94, 1.95, 1.98, 2.01, 2.03, 2.05 (7s, 21H, 7 \times CH₃CO), 3.94–3.99 (m, 2H, H-5b, H-4a), 4.17–4.06 (m, 4H, H-5a, H-6b, H-6'a, H-6'b), 4.41 (dd, 1H, $J_{6'a,6a}$ = 11.4 Hz, H-6'a), 4.87 (dd, 1H, $J_{2a,1a}$ = 10.7 Hz, $J_{2a,3a}$ = 8.4 Hz, H-2a), 4.94 (dd, 1H, $J_{2b,3b}$ = 10.0 Hz, $J_{2b,1b}$ = 3.1 Hz, H-2b), 4.95 (dd, 1H, $J_{4b,3b}$ = 10.0 Hz, $J_{4b,5b}$ = 6.1 Hz, H-4b), 5.17 (t, 1H, $J_{3b,2b}$ = $J_{3b,4b}$ = 10.0 Hz, H-3b), 5.23 (d, 1H, $J_{1b,2b}$ = 3.1 Hz, H-1b), 5.30 (dd, 1H, $J_{3a,4a}$ = 9.2 Hz, $J_{3a,2a}$ = 8.4 Hz, H-3a), 5.68 (d, 1H, $J_{1a,2a}$ = 10.7 Hz, H-1a), 9.24, 9.06 (2s, 2 \times 2H, 2 \times NH₂).

Anal. Calcd for C₂₇H₃₉BrN₂O₁₇S (775.57): C, 41.81, H, 5.07, N, 3.61. Found: C, 40.12, H, 5.39, N, 3.48.

Acetylated β , β -1-thiotrehaloses (4a–e): General Procedure. A suspension of the glycosyl isothiuronium salts **2a–e** (1.1 mmol) and the respective glycosyl halide (1.0 mmol) in acetonitrile (5 mL) and triethylamine (5 mmol) were placed in a domestic microwave oven and irradiated for 3 min. Methylene chloride (30 mL) was added to the reaction mixture and the solution was washed with water (3 \times 10 mL), dried (Na₂SO₄), and evaporated under reduced pressure. The residue was crystallized from methanol to afford pure products of **4a–e**.

2,3,4,6-Tetra-O-acetyl- β -D-glucopyranosyl 2,3,4,6-tetra-O-acetyl-1-thio- β -D-glucopyranoside (4a). mp 173–175°C, lit^[15] mp 174–175°C; yield 63%; ¹H NMR (600 MHz, CDCl₃): δ = 2.00, 2.01, 2.02, 2.03 (4s, 24H, 8 \times CH₃CO), 3.60 (ddd, 2H, $J_{5,4}$ = 9.6 Hz, $J_{5,6'}$ = 4.9 Hz, $J_{5,6}$ = 2.3 Hz, 2 \times H-5), 4.12 (dd, 2H, $J_{6,5}$ = 2.3 Hz, $J_{6,6'}$ = 12.4 Hz, 2 \times H-6), 4.23 (dd, 2H,

$J_{6',5} = 4.9$ Hz, $J_{6',6} = 12.4$ Hz, $2 \times \text{H-6}'$), 4.81 (d, 2H, $J_{1,2} = 10.2$ Hz, $2 \times \text{H-1}$), 5.03 (dd, 2H, $J_{2,1} = 10.2$ Hz, $J_{2,3} = 9.6$ Hz, $2 \times \text{H-2}$), 5.06 (dd, 2H, $J_{4,3} = 9.6$ Hz, $J_{4,5} = 9.6$ Hz, $2 \times \text{H-4}$), 5.19 (t, 2H, $J_{3,2} = 9.6$ Hz, $J_{3,4} = 9.6$ Hz, $2 \times \text{H-3}$); ^{13}C NMR (150.9 MHz, CDCl_3): $\delta = 20.4, 20.5, 20.6, 20.7$ ($8 \times \text{CH}_3\text{CO}$), 62.1 ($2 \times \text{C-6}$), 68.1 ($2 \times \text{C-4}$), 70.1 ($2 \times \text{C-2}$), 73.8 ($2 \times \text{C-3}$), 76.2 ($2 \times \text{C-5}$), 80.7 ($2 \times \text{C-1}$), 169.2, 169.3, 170.1, 170.5 ($8 \times \text{CO}$).

2,3,4,6-Tetra-O-acetyl- β -D-galactopyranosyl 2,3,4,6-tetra-O-acetyl-1-thio- β -D-galactopyranoside (4b). mp 201–203°C; lit^[14] mp 199–202°C; yield 65%; ^1H NMR (600 MHz, CDCl_3): $\delta = 2.0, 2.01, 2.09, 2.18$ (4s, 24H, $8 \times \text{CH}_3\text{CO}$), 3.89–3.93 (m, 2H, $2 \times \text{H-5}$), 4.12 (dd, 2H, $J_{6,5} = 6.5$ Hz, $J_{6,6'} = 11.3$ Hz, $2 \times \text{H-6}$), 4.18 (dd, 2H, $J_{6',5} = 6.7$ Hz, $J_{6',6} = 11.3$ Hz, $2 \times \text{H-6}'$), 4.80 (d, 2H, $J_{1,2} = 10.0$ Hz, $2 \times \text{H-1}$), 5.05 (dd, 2H, $J_{3,4} = 3.3$ Hz, $J_{3,2} = 10.0$ Hz, $2 \times \text{H-3}$), 5.23 (t, 2H, $J_{2,1} = 10.0$ Hz, $J_{2,3} = 10.0$ Hz, $2 \times \text{H-2}$), 5.45 (dd, 2H, $J_{4,3} = 3.3$ Hz, $J_{4,5} = 2.8$ Hz, $2 \times \text{H-4}$), ^{13}C NMR (150.9 MHz, CDCl_3): $\delta = 19.8, 19.88, 19.9, 20.0$ ($8 \times \text{CH}_3\text{CO}$), 60.6 ($2 \times \text{C-6}$), 66.4 ($2 \times \text{C-4}$), 66.5 ($2 \times \text{C-2}$), 71.0 ($2 \times \text{C-3}$), 73.9 ($2 \times \text{C-5}$), 80.6 ($2 \times \text{C-1}$), 168.7, 169.2, 169.4, 169.6 ($8 \times \text{CO}$).

2-Acetamido-2-deoxy-3,4,6-tri-O-acetyl- β -D-glucopyranosyl 2-acetamido-2-deoxy-3,4,6-tri-O-acetyl-1-thio- β -D-glucopyranoside (4c). mp 312–314°C; lit^[30] mp 314–316°C; yield 62%; ^1H NMR (600 MHz, CDCl_3): $\delta = 1.76$ (s, 6H, $2 \times \text{NCOCH}_3$), 1.90, 1.96, 2.02 (3s, 18H, $6 \times \text{CH}_3\text{CO}$), 3.78 (ddd, 2H, $J_{5,4} = 9.6$ Hz, $J_{5,6'} = 5.1$, $J_{5,6} = 2.1$ Hz, $2 \times \text{H-5}$), 3.92 (dd, 2H, $J_{2,1} = 10.2$ Hz, $J_{2,3} = 9.6$ Hz, $2 \times \text{H-2}$), 4.05 (dd, 2H, $J_{6,5} = 2.1$ Hz, $J_{6,6'} = 12.1$ Hz, $2 \times \text{H-6}$), 4.16 (dd, 2H, $J_{6',5} = 5.1$ Hz, $J_{6',6} = 12.1$ Hz, $2 \times \text{H-6}'$), 4.85 (t, 2H, $J_{4,5} = 9.6$ Hz, $J_{4,3} = 9.6$ Hz, $2 \times \text{H-4}$), 4.92 (d, 2H, $J_{1,2} = 10.2$ Hz, $2 \times \text{H-1}$), 5.08 (t, 2H, $J_{3,2} = 9.6$ Hz, $J_{3,4} = 9.6$ Hz, $2 \times \text{H-3}$); ^{13}C NMR (150.9 MHz, CDCl_3): $\delta = 20.3, 20.4, 20.5$ ($6 \times \text{CH}_3\text{CO}$), 22.6 ($2 \times \text{NCOCH}_3$), 52.0 ($2 \times \text{C-2}$), 62.1 ($2 \times \text{C-6}$), 68.6 ($2 \times \text{C-4}$), 73.7 ($2 \times \text{C-3}$), 75.0 ($2 \times \text{C-5}$), 80.0 ($2 \times \text{C-1}$), 169.3, 169.3, 169.6, 170.1 ($8 \times \text{CO}$).

4,4'-Di-O-(2,3,4,6-tetra-O-acetyl- β -D-galactopyranosyl) 2,2',3,3',6,6'-hexa-O-acetyl- β , β -1-thiotrehalose (4d). mp 143–145°C; yield 58%; ^1H NMR (600 MHz, CDCl_3) $\delta = 1.96$ –2.18 (14s, 42H, $14 \times \text{CH}_3\text{CO}$), 3.58 (ddd, 2H, $J_{5a,4a} = 9.2$ Hz, $J_{5a,6'a} = 5.3$ Hz, $J_{5a,6a} = 3.8$ Hz, $2 \times \text{H-5a}$), 3.78 (dd, 2H, $J_{4a,3a} = 9.9$ Hz, $J_{4a,5a} = 9.2$ Hz, $2 \times \text{H-4a}$), 3.89 (ddd, 2H, $J_{5b,4b} = 7.7$ Hz, $J_{5b,6'b} = 3.1$ Hz, $J_{5b,6b} = 4.6$ Hz, $2 \times \text{H-5b}$), 4.07 (dd, 2H, $J_{6b,5b} = 4.6$ Hz, $J_{6b,6'b} = 11.5$ Hz, $2 \times \text{H-6b}$), 4.13 (dd, 2H, $J_{6'b,5b} = 3.1$ Hz, $J_{6'b,6b} = 11.5$ Hz, $2 \times \text{H-6'b}$), 4.17 (dd, 2H, $J_{6a,5a} = 3.8$ Hz, $J_{6a,6'a} = 12.2$ Hz, $2 \times \text{H-6a}$), 4.46 (d, 2H, $J_{1b,2b} = 9.0$ Hz, $2 \times \text{H-1b}$), 4.57 (dd, 2H, $J_{6'a,5a} = 5.3$ Hz, $J_{6'a,6a} = 12.2$ Hz, $2 \times \text{H-6'a}$), 4.76 (d, 2H, $J_{1a,2a} = 10.0$ Hz, $2 \times \text{H-1a}$), 4.93 (dd, 2H, $J_{2a,1a} = 10.0$ Hz, $J_{2a,3a} = 9.9$ Hz, $2 \times \text{H-2a}$), 4.98 (dd, 2H, $J_{3b,4b} = 3.8$ Hz, $J_{3b,2b} = 9.9$ Hz, $2 \times \text{H-3b}$), 5.11 (dd, 2H, $J_{2b,3b} = 9.9$ Hz, $J_{2b,1b} = 9$ Hz, $2 \times \text{H-2b}$), 5.23 (dd, 2H, $J_{3a,2a} = 9.9$ Hz, $J_{3a,4a} = 9.2$ Hz, $2 \times \text{H-3a}$), 5.34 (dd, 2H, $J_{4b,5b} = 7.7$ Hz, $J_{4b,3b} = 3.8$ Hz, $2 \times \text{H-4b}$). ^{13}C NMR (150.9 MHz,

CDCl₃): δ = 20.4–20.8 (14 \times CH₃CO), 60.7 (2 \times C-6b), 62.1 (2 \times C-6a), 66.5 (2 \times C-4b), 69.0 (2 \times C-2b), 70.4 (2 \times C-2a), 70.5 (2 \times C-5b), 70.9 (2 \times C-3b), 73.8 (2 \times C-3a), 75.7 (2 \times C-4a), 76.1 (2 \times C-5a), 80.4 (2 \times C-1a), 101.1 (2 \times C-1b), 169.0, 169.5, 169.6, 170.0, 170.1, 170.2, 170.3 (14 \times CO).

Anal. Calcd for C₅₂H₇₀O₃₄S (1271.16): C, 49.13, H, 5.55. Found: C, 49.14, H, 5.79.

4,4'-Di-O-(2,3,4,6-tetra-O-acetyl- α -D-glucopyranosyl) 2,2',3,3',6,6'-hexa-O-acetyl- β , β -1-thiotrehalose (4e). mp 110–112°C; yield 57%; ¹H NMR (600 MHz, CDCl₃) δ = 1.99–2.18 (14s, 42H, 14 \times CH₃CO), 3.68–3.73 (m, 2H, 2 \times H-5a), 3.93 (m, 2H, 2 \times H-5b), 4.00 (t, 2H, $J_{4a,5a} = J_{4a,3a} = 9.2$ Hz, 2 \times H-4a), 4.03 (dd, 2H, $J_{6'a,6a} = 12.0$ Hz, 2 \times H-6'a), 4.21 (dd, 2H, $J_{6a,6'a} = 12.0$ Hz, 2 \times H-6a), 4.23 (dd, 2H, $J_{6b,6'b} = 12.2$ Hz, $J_{6b,5b} = 3.8$ Hz, 2 \times H-6b), 4.78 (dd, 2H, $J_{2b,1b} = 3.8$ Hz, $J_{2b,3b} = 9.9$ Hz, 2 \times H-2b), 4.82 (dd, 2H, $J_{2a,1a} = 10.7$ Hz, $J_{2a,3a} = 9.2$ Hz, 2 \times H-2a), 4.83 (d, 2H, $J_{1a,2a} = 10.7$ Hz, 2 \times H-1a), 5.06 (dd, 2H, $J_{4b,3b} = 9.9$ Hz, $J_{4b,5b} = 9.9$ Hz, 2 \times H-4b), 5.28 (t, 2H, $J_{3a,2a} = J_{3a,4a} = 9.2$ Hz, 2 \times H-3a), 5.34 (t, 2H, $J_{3b,2b} = J_{3b,4b} = 9.9$ Hz, 2 \times H-3b), 5.4 (d, 2H, $J_{1b,2b} = 3.8$ Hz, 2 \times H-1b). ¹³C NMR (CDCl₃, 150.9 MHz): δ = 20.5–20.8 (14 \times CH₃CO), 61.5 (2 \times C-6b), 62.9 (2 \times C-6a), 68.0 (2 \times C-4b), 72.2 (2 \times C-4a), 68.5 (2 \times C-3b), 74.9 (2 \times C-3a), 70.1 (2 \times C-2b), 69.4 (2 \times C-5b), 72.4 (2 \times C-5a), 73.8 (2 \times C-2a), 90.0 (2 \times C-1a), 95.6 (2 \times C-1b), 169.3, 169.5, 169.8, 170.0, 170.1, 170.3, 170.4 (14 \times CO).

Anal. Calcd for C₅₂H₇₀O₃₄S (1271.16): C, 49.13, H, 5.55. Found: C, 48.89, H, 5.80.

ACKNOWLEDGMENTS

The continued supports from the AvH and DFG are highly appreciated.

REFERENCES

- [1] Defaye, J.; Gelas, J. Thio-oligosaccharides: their synthesis and reactions with enzymes. In *Studies in Natural Products Chemistry*; Atta-ur-Rahman, Ed.; Elsevier: Amsterdam, 1991; 8E; 315–357.
- [2] Driguez, H. Thiooligosaccharides in glycobiology. *Topics Curr. Chem.* **1997**, *187*, 85–116.
- [3] Gil, V.; Macleod, A.J. Effects of a *Lepidium sativum* enzyme preparation on the degradation of glucosinolates. *Phytochemistry* **1980**, *19*, 2071–2076.
- [4] (a) Horton, D.; Hutson, D.H. Developments in the chemistry of thio sugars. *Adv. Carbohydr. Chem.* **1963**, *18*, 123–199; (b) Witzcak, Z.J. Thiosugars: biological relevance as potential new therapeutics. *Curr. Med. Chem.* **1999**, *6*, 165–178; (c) Wilson, J.C.; Kiefel, M.J.; Angus, D.I.; Itzstein, M.V. Investigation of the stability of thiosialosides toward hydrolysis by sialidases using NMR spectroscopy. *Org. Lett.* **1999**, *1*, 443–446.

- [5] Norberg, T. *Modern Methods in Carbohydrate Synthesis*; Khan, S.H.; O'Neill, R.A., Eds.; Harwood: London, 1995; 82.
- [6] (a) El Ashry, E.S.H.; Rashed, N.; Shobier, A.H. Glycosidase inhibitors and their chemotherapeutic value. Part 1. *Pharmazie* **2000**, *55*, 251–262; (b) El Ashry, E.S.H.; Rashed, N.; Shobier, A.H. Glycosidase inhibitors and their chemotherapeutic value. Part 2. *Pharmazie* **2000**, *55*, 331–348; (c) El Ashry, E.S.H.; Rashed, N.; Shobier, A.H. Glycosidase inhibitors and their chemotherapeutic value. Part 3. *Pharmazie* **2000**, *55*, 403–415; (d) Witczak, Z.J.; Nieforth, K.A. *Carbohydrates in Drug Design*; Marcel Dekker: New York, 1997; (e) Robina, I.; Vogel, P.; Witczak, Z.J. Synthesis and biological properties of monothiosaccharides. *Curr. Org. Chem.* **2001**, *5*, 1177–1214; (f) Witczak, Z.J.; Chabra, R.; Chen, H.; Xie, X.-Q. A novel approach to thiodisaccharides. The synthesis of 3-deoxy-4-thiocellobiose from levoglucosone. *Carbohydr. Res.* **1997**, *301*, 167–175.
- [7] Eisele, T.; Schmidt, R.R. Synthesis of thio-linked analogues of Lewis X and sialyl Lewis X. *Liebigs Ann. Chem.* **1997**, 1303–1313.
- [8] Ding, Y.; Contour-Galceran, M.-O.; Ebel, J.; Ortiz-Mellet, C.; Defaye, J. S-Linked thiomimetics of phytoalexin-elicitor-active, branched oligosaccharides, their synthesis, protein-binding ability and phytoalexin-inducing activity. *Eur. J. Org. Chem.* **1999**, 1143–1152.
- [9] Bock, K.; Defaye, J.; Driguez, H.; Bar-Guilloux, E. Conformations in solution of α , α -trehalose, α -D-glucopyranosyl α -D-mannopyranoside, and their 1-thioglycosyl analogs, and a tentative correlation of their behaviour with respect to the enzyme trehalase. *Eur. J. Biochem.* **1983**, *131*, 595–600.
- [10] Defaye, J.; Driguez, H.; Henrissat, B.; Bar-Guilloux, E. In *Mechanisms of Saccharide Polymerization and Depolymerization*; Marshall, J.J., Ed.; Academic Press: New York, 1980, 331.
- [11] Defaye, J.; Driguez, H.; Poncet, S.; Chombert, R.; Petit-Glatron, M.F. Synthesis of 1-thiosucrose and anomers, and the behaviour of levansucrase and invertase with this substrate analog. *Carbohydr. Res.* **1984**, *130*, 299–315.
- [12] Schneider, W.; Wrede, F. Synthese eines schwefelhaltigen und eines selenhaltigen Disaccharides. *Ber. Dtsch. Chem. Ges.* **1917**, *50*, 793–804.
- [13] Blanc-Muesser, M.; Defaye, J.; Driguez, H. Synthesis stereoselectives de 1-thioglycosides. *Carbohydr. Res.* **1978**, *67*, 305–328.
- [14] Bogusiak, J.; Szeja, W. Synthesis of bis(2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl and -galactopyranosyl)sulfide using phase-transfer catalysis. *Carbohydr. Res.* **1985**, *141*, 165–167.
- [15] Chrétien, F.; Dicesare, P.; Gross, B. Synthesis of thiodisaccharides using phase-transfer catalysis. *J. Chem. Soc., Perkin Trans. 1.* **1988**, 3297–3300.
- [16] Schneider, W.; Beuther, A. Schwefelhaltige Disaccharide aus Galaktose. *Ber. Dtsch. Chem. Ges.* **1919**, *52*, 2135–2149.
- [17] Boss, W.; Schaedel, P.; Wallenfels, K. Untersuchungen zur Induktion der lac-Enzyme. *Eur. J. Biochem.* **1967**, 382–394.
- [18] Gabelle, A.; Defaye, J.; Pedersen, C. A simple preparation of 2,3,4,6-tetra-O-acetyl-1-S-acetyl-1-thio- α -D-glucopyranose. *Carbohydr. Res.* **1990**, *200*, 497–498.
- [19] Blanc-Muesser, M.; Defaye, J.; Driguez, H. Stereoselective synthesis of 1,2-*cis*-thioglycosides. *Tetrahedron Lett.* **1976**, *47*, 4307–4310.

- [20] Ferrier, R.J.; Furneaux, R.H.; Tyler, P.C. Observations on the possible application of glycosyl disulphides, sulphonic esters and sulphones in the synthesis of glycosides. *Carbohydr. Res.* **1977**, *58*, 397–404.
- [21] Korytnyk, W.; Mills, J.A. Preparation and properties of some poly-O-acetylglycosyl chlorides of the “unstable” series. *J. Chem. Soc.* **1959**, 636–649.
- [22] Defaye, J.; Gadelle, A.; Pedersen, C. Hydrogen fluoride-mediated synthesis of 1-thiotrehaloses involving reaction of D-glucose with hydrogen sulfide. *Carbohydr. Res.* **1991**, *217*, 51–58.
- [23] Defaye, J.; Gadelle, A.; Pedersen, C. The behavior of cellulose, amylase, and β -D-xylan towards anhydrous hydrogen fluoride. *Carbohydr. Res.* **1982**, *110*, 217–227.
- [24] Mingos, D.M.P.; Baghurst, D.R. Applications of microwave dielectric heating effects to synthetic problems in chemistry. *Chem. Soc. Rev.* **1991**, *20*, 1–47.
- [25] Abramovitch, R.A. Applications of microwave energy in organic chemistry. *Org. Prep. Proced. Int.* **1991**, *23*, 683–712.
- [26] Bose, A.K.; Manhas, M.S.; Ghosh, M.; Raju, V.S.; Tabei, K.; Urbanczyk-Lipkowska, Z. Highly accelerated reactions in a microwave oven: synthesis of heterocycles. *Heterocycles* **1990**, *30*, 741–744.
- [27] (a) Abdel-Rahman, A.A.-H.; El Ashry, E.S.H. Efficient synthesis of 5-hydroxy methyl pyrimidines and nucleosides using microwave irradiation. *Synlett* **2002**, *12*, 2043–2044; (b) Abdel Hamid, H.M.; Ramadan, E.; Hagar, M.; El Ashry, E.S.H. Synthesis of aryloxyacetic acids, esters and hydrazides assisted by microwave irradiation. *Synth. Commun.* **2004**, *34*, 377–382; (c) El Ashry, E.S.H.; Ramadan, E.; Abdel-Hamid, H.; Hagggar, M. Microwave irradiation for enhancing the regioselective synthesis of 6H-indolo[2,3-b]quinoxaline. *J. Chem. Res.* **2005**, 229–232; (d) El Ashry, E.S.H.; Ramadan, E.; Abdel-Hamid, H.; Hagggar, M. Synthesis of azlactone, phenyl pyruvic acid and 1,2,4-triazine derivatives under microwave irradiation. *Lett. Org. Chem.* **2005**, *2*, 415–418; (e) El Ashry, E.S.H.; Ramadan, E.; Abdel-Hamid, H.; Hagggar, M. Microwave irradiation for enhancing the synthesis of quinoline derivatives from isatin. *Synth. Commun.* **2005**, *in press*; (f) El Ashry, E.S.H.; Ramadan, E.; Kassem, A.; Hagggar, M. Microwave irradiation for accelerating organic reactions. Part 1: three, four and five membered heterocycles. *Adv. Heterocycl. Chem.* **2005**, *88*, 1–114; (g) El Ashry, E.S.H.; Ramadan, E.; Kassem, A. Microwave irradiation for accelerating organic reactions. Part 2: six, seven fused and spiro heterocyclic ring systems. *Adv. Heterocycl. Chem.* **2005**, *accepted*.
- [28] William, A.B.; James, E.K. A study of some S-(polyacetyl- β -D-glycopyranosyl)-thiuronium halides. *J. Am. Chem. Soc.* **1951**, *73*, 2241–2245.
- [29] Horton, D.; Wolfrom, M.L. Thiosugars. I. Synthesis of derivatives of 2-amino-2-deoxy-1-thio-D-glucose. *J. Org. Chem.* **1962**, *27*, 1794–1800.
- [30] Akagi, M.; Tejima, S.; Haga, M. Biochemical studies on thiosugars II. Synthesis of 2-acetamido-2-deoxy-1-thio-glucose and its derivatives. *Chem. Pharm. Bull.* **1961**, *9*, 360–362.
- [31] Xiang-Bao, M.; Lian-Dai, Y.; Hui, L.; Qing, L.; Tie-Ming, C.; Meng-Shen, C.; Zhong-Jun, L. Facile synthesis of 1-thio- β -lactoside clusters scaffolded onto *p*-methoxyphenyl β -D-galactopyranoside, β -D-glucopyranoside, and lactoside. *Carbohydr. Res.* **2002**, *337*, 977–981.