

This article was downloaded by:

On: 22 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713617200>

### Optimized Synthesis of Vinyl Ether Sugars and Vinyl Glycosides through Transfer Vinylation Catalyzed by (4,7-Ph<sub>2</sub>-phen)Pd(OOCCF<sub>3</sub>)<sub>2</sub>

Martin Bosch<sup>a</sup>; Sean Handerson<sup>a</sup>; Marcel Schlaf<sup>a</sup>

<sup>a</sup> The Guelph-Waterloo Centre for Graduate Work in Chemistry (GWC)<sup>2</sup>, Department of Chemistry, University of Guelph, Guelph, Ontario, Canada

**To cite this Article** Bosch, Martin , Handerson, Sean and Schlaf, Marcel(2008) 'Optimized Synthesis of Vinyl Ether Sugars and Vinyl Glycosides through Transfer Vinylation Catalyzed by (4,7-Ph<sub>2</sub>-phen)Pd(OOCCF<sub>3</sub>)<sub>2</sub>', *Journal of Carbohydrate Chemistry*, 27: 2, 103 – 112

**To link to this Article:** DOI: 10.1080/07328300802030852

**URL:** <http://dx.doi.org/10.1080/07328300802030852>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

# Optimized Synthesis of Vinyl Ether Sugars and Vinyl Glycosides through Transfer Vinylation Catalyzed by (4,7-Ph<sub>2</sub>-phen)Pd(OOCCF<sub>3</sub>)<sub>2</sub>

Martin Bosch, Sean Handerson, and Marcel Schlaf

The Guelph-Waterloo Centre for Graduate Work in Chemistry (GWC)<sup>2</sup>,  
Department of Chemistry, University of Guelph, Guelph, Ontario, Canada

Sugar vinyl ethers and vinyl glycosides are conveniently synthesized by catalytic transfer vinylation with butyl vinyl ether, which serves as both the solvent and source of vinyl. The air-stable catalyst (4,7-diphenyl-1,10-phenanthroline)Pd(OOCCF<sub>3</sub>)<sub>2</sub> is prepared in situ from commercially available components.

**Keywords** Vinyl sugars, Catalytic synthesis

## INTRODUCTION

Vinyl glycosides and vinyl ether sugars have been employed as glycosyl donors and acceptors<sup>[1–4]</sup> and also constitute a class of potential components of sugar-based amphiphiles and surfactants or sugar-modified polymers and polymer surfaces obtained by the copolymerization of a vinyl ether sugar with a bulk alkene (e.g., styrene or propylene, methyl methacrylate, vinyl acetate, or acrylonitrile).


Early examples of vinyl sugars are 6-*O*-vinyl-1,2:3,4-di-*O*-isopropylidene-D-galacto-pyranoside,<sup>[5]</sup> obtained in 58% yield by reaction of diacetone-galactose with acetylene in sodium hydroxide melt, and 3-*O*-vinyl-1,2:5:6-di-

---

Received November 30, 2007; accepted February 2, 2008.

Address correspondence to Marcel Schlaf, Department of Chemistry, University of Guelph, Guelph, Ontario N1G 2W1, Canada. E-mail: mschlaf@uoguelph.ca

O-isopropylidene-D-glucufuranoside,<sup>[6]</sup> obtained in 22% yield by transesterification of diacetone-glucose catalyzed by mercuric acetate. More recently sugar vinyl ethers have been synthesized by preparation of the corresponding  $\beta$ -alkoxyvinyl sulfones followed by desulfonylation with Na/Hg amalgam.<sup>[7]</sup> Vinyl glycosides have also been prepared by transformation of vinyl acetates with Tebbe's reagent (yielding the corresponding isopropenyls),<sup>[1]</sup> by Hofmann degradation of *N*-2-trimethyl-ammonium-ethyl-*O*-glycosides,<sup>[8]</sup> through the use of selenium<sup>[9]</sup> or mercury reagents,<sup>[10]</sup> by photochemically induced rearrangement of 4-oxo-pentyl-glycosides,<sup>[11]</sup> and by acid-catalyzed alcohol elimination from mixed acetal glycosides.<sup>[12]</sup> The latter method is now also applicable to the synthesis of sugar vinyl ethers.<sup>[13]</sup> Recently we described a very *atom-efficient* and simple method for the transfer vinylation of a series of sugar substrates catalyzed by the air-stable complex (4,7-diphenyl-1,10-phenanthroline)Pd(OAc)<sub>2</sub> prepared in situ and using butyl vinyl ether as the sole reagent and solvent.<sup>[14–16]</sup> Equation (1) illustrates the reaction principle. The reaction is driven by providing either the sugar alcohol ROH or—as in the present case—the vinyl ether R'OCH=CH<sub>2</sub> in large excess.



We subsequently optimized this catalyst system by exchanging the counterion to the less coordinating trifluoroacetate, which reduces the reaction time required to attain equilibrium by a factor of up to eight, and successfully applied the catalyst to the synthesis of a variety of alkyl and allyl vinyl ethers.<sup>[17]</sup> Here we present the application of this optimized catalyst system to a representative series of protected monosaccharides with one, two, or three free hydroxyl functions resulting in the isolation of the corresponding vinyl glycosides and mono-, di-, or trivinyl ethers in moderate to good and in a few cases excellent yields, which may facilitate their incorporation into new sugar polymer formulations.

## RESULTS AND DISCUSSION

The reaction protocol employed for the transfer vinylation was in all cases 2 mmol of sugar substrate, 0.04 mmol palladium trifluoroacetate (2 mol %), 0.04 mmol 4,7-diphenyl-1,10-phenanthroline ligand, and 0.16 mmol triethyl amine dissolved in 400 mmol (200-fold excess) of butyl vinyl ether (BVE) to give a clear yellow solution, which was heated to 75°C in a sealed flask. After 5 to 12 h qualitative evaluation of TLC samples and quantitative GC of samples drawn in hourly intervals and monitored for the following 24 to 48 h showed that the equilibrium (Eq. (1)) had been established as indicated by

the constant relative concentrations of the *n*-butanol generated by the reactions. The presence of the auxiliary base triethyl amine prevents the formation of mixed ethyl butyl acetals catalyzed by free trifluoro-acetic acid formed in the catalytic cycle.<sup>[14,17]</sup>



















Table 1 summarizes the results obtained with this reaction protocol for the monohydroxyl substrates 2,3,4,6-tetra-*O*-benzyl-*D*-glucopyranose (**1a**), 2,3,4,6-tetra-*O*-acetyl-*D*-glucopyranose (**2a**), 2,3,5-tri-*O*-benzyl-*D*-arabinofuranose (**3a**), 1,2:5,6-di-*O*-isopropylidene- $\alpha$ -*D*-glucofuranoside (**4a**), 1,2:3,4-di-*O*-isopropylidene- $\alpha$ -*D*-galactopyranoside (**5a**), methyl 2,3,6-tri-*O*-benzoyl- $\alpha$ -*D*-galactopyranoside (**6a**), methyl 2,3,6-tri-*O*-benzoyl- $\alpha$ -*D*-mannopyranoside (**7a**), 1,2:4,5-di-*O*-isopropylidene- $\beta$ -*D*-fructopyranoside (**8a**), and 2,3:4,5-di-*O*-isopropylidene- $\beta$ -*D*-fructopyranoside (**9a**). Table 2 lists the results for the dihydroxyl substrates 1,2-*O*-isopropylidene- $\alpha$ -*D*-xylofuranoside (**10a**), 1,2:5,6-di-*O*-isopropylidene-*D*-mannitol (**11a**), methyl 4,6-*O*-isopropylidene- $\alpha$ -*D*-mannopyranoside (**12a**), methyl 4,6-*O*-isopropylidene- $\alpha$ -*D*-glucopyranoside (**13a**), phenyl 4,6-*O*-isopropylidene- $\beta$ -*D*-glucopyranoside (**14a**), methyl 4,6-*O*-benzylidene- $\alpha$ -*D*-glucopyranoside (**15a**), phenyl 4,6-*O*-benzylidene- $\beta$ -*D*-glucopyranoside (**16a**), and the trihydroxyl substrate levoglucosan (**17a**).

The first five entries in Table 1 also compare the reaction times and yields to those achieved earlier with the nonoptimized catalyst.<sup>[14]</sup> Slightly higher yields were realized with the optimized protocol described above that employs a 200:1 rather than the 100:1 BVE-to-sugar ratio used earlier.<sup>[14]</sup>

With the exception of entry 8, 1,2:4,5-di-*O*-isopropylidene- $\beta$ -*D*-fructopyranoside (**8a**), good to excellent yields (68%–95%) of mono vinyl ethers or vinyl glycosides are achieved with sugar substrates protected by acetals or benzyl ethers, while ester protection groups (acetate, benzoyl) result in moderate yields of 45% (**2b**, entry 2, Table 1) and 42% or 43% (**6b** and **7b**, entries 6 and 7, Table 1). As noted previously,<sup>[14]</sup> the presence of ester functionalities (**2a**, **6a**, **7a**, entries 2, 6, and 7) leads to a darkening of the reaction mixture to a more intense yellow, pointing to a coordinative interaction between either the carbonyl oxygen or possibly an ester enolate and the metal center. This can, however, not account for the lower yields with these substrates, as addition of more catalyst at equilibrium did not result in additional product formation excluding irreversible catalyst poisoning as a possible explanation.













As the vinyl exchange Eq. (1) is an isodesmic reaction, the main factors governing the overall yields obtained with any of the substrates should be steric interactions within the vinylated products. However, the conformational variability and subtle structural complexity of the sugar substrates renders a general and predictive model for these interactions and quantitative rationalization of the resulting observed yields elusive. For the monohydroxyl substrates, the observed yields qualitatively scale with the steric accessibility of the free hydroxyl functions, which also correlates with the conformational flexibility of an attached vinyl group in the corresponding

**Table 1:** Vinylation of protected sugars with one free hydroxyl function.

Entry	Sugar substrate (#a)	Vinyl sugar (#b)	Yield with Pd(OAc) <sub>2</sub> (4-7 d) <sup>a</sup>	Yield with Pd(O <sub>2</sub> CCF <sub>3</sub> ) <sub>2</sub> ≤12 h <sup>a</sup>
1	 (1a)	 (1b)	70% $\alpha/\beta = 9:1^b$	74% $\alpha/\beta = 7:1^b$
2	 (2a)	 (2b)	36% $\alpha/\beta = 6.5:1^a$	45% $\alpha/\beta = 7:1^a$
3	 (3a)	 (3b)	60% $\alpha/\beta = 4.4:1^b$	68% $\alpha/\beta = 6:1^b$
4	 (4a)	 (4b)	72%	81%
5	 (5a)	 (5b)	79%	95%
6	 (6a)	 (6b)	not det.	42%
7	 (7a)	 (7b)	not det.	43%
8	 (8a)	 (8b)	not det.	24%
9	 (9a)	 (9b)	not det.	73%











<sup>a</sup>Isolated yield.<sup>b</sup>Anomeric ratios determined by <sup>1</sup>H NMR.

**Table 2:** Vinylation of protected sugars with two or three free hydroxyl functions.

Entry	Sugar substrate	Vinyl sugar	Yield and reaction time (h) <sup>a</sup>
1	 <b>(10a)</b>	 <b>(10b)</b>	51% (6 h)
		 <b>(10c)</b>	25% (6 h)
		 <b>(10d)</b>	6% (6 h)
		Total yield of vinylated products: 82%	
2	 <b>(11a)</b>	 <b>(11b)</b>	24% (6 h)
		Total yield of vinylated products: 56%	
3	 <b>(12a)</b>	 <b>(12b)</b>	6% (12 h)
		 <b>(12c)</b>	30% (12 h)
		 <b>(12d)</b>	20% (12 h)
		Total yield of vinylated products: 56%	
4	 <b>(13a)</b>	 <b>(13b)</b>	5% (12 h)

(continued)

Table 2: Continued.

Entry	Sugar substrate	Vinyl sugar	Yield and reaction time (h) <sup>a</sup>	
5	 (14a)	 (13c)	35% (12 h)	
		Total yield of vinylated products: 50%	 (13d)	20% (12 h)
		 (14b)	36% (24 h)	
		 (14c)	21% (24 h)	
6	 (15a)	 (14d)	33% (24 h)	
		Total yield of vinylated products: 90%	 (15b)	24% (24 h)
		 (15c)	26% (24 h)	
		 (15d)	16% (24 h)	

(continued)

Table 2: Continued.

Entry	Sugar substrate	Vinyl sugar	Yield and reaction time (h) <sup>a</sup>
7	 (16a)	 (16b)	13% (24 h)
		 (16c)	17% (24 h)
		 (16d)	14% (24 h)
		Total yield of vinylated products: 44%	
8	 (17a)	 (17b)	4% (6 h)
		 (17c)	13% (6 h)
		 (17d)	26% (6 h)
		Total yield of vinylated products isolable as pure compounds: 43%	

<sup>a</sup>Isolated yield.

product vinyl ether. For example, the lowest yield of 24% was obtained with 1,2:4,5-di-*O*-isopropylidene- $\beta$ -D-fructopyranoside (**8a**) (entry 8, Table 1). A simple molecular model of **8b** suggests that the  $\beta$ -oriented conformationally fixed isopropylidene ring at the anomeric center hinders a free rotation of the vinyl group about the two oxygen-carbon bonds. The effect is also present in **9b**, but less pronounced, while the vinyl groups in the other sugar substrates listed in Table 1 can freely rotate. Given that the equilibrium constant in equation (1) is close to unity for all substrates, subtle differences in the entropy of the vinyolated products originating from more or less restricted rotation of the vinyl group



compared to BVE could in fact account for the substantially different yields between the acetal protected vinyl sugars **4b**, **5b**, and **8b**.

Based on the results with mono-hydroxyl substrates where the best yields were achieved with acetal protecting groups (Table 1), the di- and trihydroxyl substrates investigated were limited to isopropylidene and benzylidene sugar derivatives. All possible mono- and di-vinylated products derived from compounds **10a** to **16a** (entries 1–7 in Table 2) could be separated and isolated by preparative flash column chromatography. The vinylation of 1,2:5,6-di-*O*-isopropylidene-*D*-mannitol (**11a**), entry 2, gave the monovinylated product in 24% yield. Only one monovinylated product can be formed due to the symmetry of the starting sugar alcohol and the divinyl sugar product did not form. As with **8b**, a simple molecular model of **11b** suggests a severely hindered rotation of the vinyl group compared to BVE leading to the same explanation for the low yield discussed above. For the trihydroxyl substrate 1,6-anhydro- $\beta$ -*D*-glucopyranoside (**17a**) (entry 8, Table 2, levoglucosan), the potential number of vinylated products is seven (three monovinylated, three divinylated, and one pervinylated) and their separation constitutes a major challenge. Of the seven products formed, we were able to purify three fractions: the trivinylated product **17b** and two separate monovinylated products **17c** and **17d**. A fourth fraction appeared to be a mixture of three divinylated products that in our hands could not be individually separated by chromatographic means (both flash and HPLC methods).

## EXPERIMENTAL

### General Procedure for the Transfer Vinylation of Protected Sugar Substrates Using Diacetone- $\alpha$ -*D*-glucopyranoside as an Example

In an oven-dried round bottom Schlenk flask, 4,7-diphenyl-1,10-phenanthroline (13.3 mg = 0.04 mmol) was added to butyl vinyl ether (40 g = 52 mL = 400 mmol) with stirring. Palladium(II) trifluoroacetate (13.3 mg = 0.04 mmol) was added, and the mixture left to stir at 75°C until all solids were dissolved and a clear bright yellow solution was obtained. To prevent acetylation, triethylamine (0.16 mmol) was added before addition of the sugar substrate. The sugar substrate, diacetone- $\alpha$ -*D*-glucopyranoside (520.5 mg = 2.0 mmol), was added to the mixture and allowed to react at 75°C until TLC and quantitative GC analysis for *n*-butanol revealed that equilibrium had been established (10 h). The reaction mixture was passed through a 10 × 2.5 cm plug of charcoal using ethyl acetate as the eluant (removes most of the catalyst) and concentrated under reduced pressure, and the residue was subjected to flash chromatography with various ratios of TLC pretested eluant combinations

(hexane, pentane, ether, ethyl acetate, acetonitrile, dichloromethane) yielding the pure vinylated sugars. See supporting information for details of the chromatographic separation and NMR data.

## SUPPORTING INFORMATION

Sources of sugar starting materials. Detailed experimental procedures for synthesis and purification of vinylated products. Extensive collection of 1D and 2D  $^1\text{H}/^{13}\text{C}$ -NMR data and spectrum images for all new vinyl sugars synthesized (82 pages).

## ACKNOWLEDGMENTS

The authors thank NSERC Canada, the Canadian Foundation for Innovation (CFI), the Ontario Innovation Trust Fund (OIT), and the University of Guelph for funding.

## REFERENCES

- [1] Marra, A.; Esnault, J.; Veyrieres, A.; Sinay, P. Isopropenyl glycosides and congeners as novel classes of glycosyl donors: theme and variations. *J. Am. Chem. Soc.* **1992**, *114*, 6354–6360.
- [2] Chenault, H.K.; Castro, A. Glycosyl transfer by isopropenyl glycosides: trisaccharide synthesis in one pot by selective coupling of isopropenyl and *n*-pentenyl glycosides. *Tetrahedron Lett.* **1994**, *35*, 9145–9148.
- [3] Chenault, H.K.; Castro, A.; Chafin, L.F.; Yang, J. The chemistry of isopropenyl glycopyranosides. Transglycosilations and other reactions. *J. Org. Chem.* **1996**, *61*, 5024–5031.
- [4] Boons, G.-J.; Isles, S. Vinyl glycosides in oligosaccharide synthesis. 2. The use of allyl and vinyl glycosides in oligosaccharide synthesis. *J. Org. Chem.* **1996**, *61*, 4262–4271.
- [5] Whistler, R.L.; Panzer, H.P.; Goatley, J.L. Preparation and polymerization of 6-*O*-vinyl-1,2:3,4-di-*O*-isopropylidene-*D*-galacto-pyranose. *J. Org. Chem.* **1962**, 2961–2962.
- [6] Black, W.A.P.; Dewar, E.T.; Rutherford, D. Polymerisation of unsaturated derivatives of 1,2:3,4-di-*O*-isopropylidene-*D*-glucofuranose. *J. Chem. Soc.* **1963**, 4433–4439.
- [7] Chéry, F.; Desroses, M.; Tatibouët, A.; Lucchi, O.d.; Rollin, P. Synthesis of sugar-based ethenyl ethers through a vinyl bis-sulfone methodology. *Tetrahedron* **2003**, *59*, 4563–4572.
- [8] Perrine, T.D.; Claudemans, C.P.J.; Ness, R.K.; Kyle, J.; Fletcher, H.G. Syntheses with partially benzylated sugars. VII. The anomeric vinyl *D*-glucopyranosides. *J. Org. Chem.* **1967**, *32*, 664–669.
- [9] Rollin, P.; Bencomo, V.V.; Sinay, P. Use of selenium in carbohydrate chemistry: formation of vinyl-glycosides. *Synthesis* **1984**, 134–135.

- [10] Raadt, A.D.; Ferrier, R.J. A simple, efficient synthesis of vinyl- $\beta$ -glucopyranosides. *J. Chem. Soc. Chem. Commun.* **1987**, 1009–1010.
- [11] Cottier, L.; Remy, G.; Descotes, G. Photochemical synthesis of O-vinyl glucosides and their transformations into C-branched sugars. *Synthesis* **1979**, 711–712.
- [12] Edathil, E.; Nguyen, J.; Hughes, K.; Boyko, W.; Giuliano, R.M. Synthesis of vinyl glucopyranosides from mixed acetal glycosides. *J. Carbohydr. Chem.* **2001**, *20*, 81–85.
- [13] Hughes, K.D.; Nguyen, T.L.N.; Dyckman, D.; Dulay, D.; Boyko, W.J.; Giuliano, R.M. Synthesis of vinyl glycosides and carbohydrate vinyl ethers from mixed acetals: a hetero-Diels-Alder approach to deoxygenated disaccharides. *Tetrahedron Asymm.* **2005**, *16*, 273–282.
- [14] Handerson, S.; Schlaf, M. Palladium(II)-catalyzed transfer vinylation of protected monosaccharides. *Org. Lett.* **2002**, *4*, 407–409.
- [15] McKeon, J.E.; Fitton, P.; Griswold, A.A. The palladium(II) catalyzed vinyl interchange reaction-I. *Tetrahedron* **1972**, *28*, 227–232.
- [16] McKeon, J.E.; Fitton, P. The palladium(II) catalyzed vinyl interchange reaction-II. *Tetrahedron* **1972**, *28*, 233–238.
- [17] Bosch, M.; Schlaf, M. Synthesis of allyl and alkyl vinyl ethers using an in situ prepared air-stable palladium catalyst. Efficient transfer vinylation of primary, secondary, and tertiary alcohols. *J. Org. Chem.* **2003**, *68*, 5225–5227.