

- (4) King, L. A., Seegmiller, D. W., *J. Chem. Eng. Data*, **16**, 23 (1971).  
 (5) Sears, J. E., *Proc. Phys. Soc., London*, **28**, 95 (1913); "International Critical Tables", Vol. II, McGraw-Hill, New York, N.Y., 1927, p 457.  
 (6) Seegmiller, D. W., Fannin, Jr., A. A., Olson, D. S., King, L. A., *J. Chem. Eng. Data*, **17**, 295 (1972).  
 (7) Smits, A., Meijering, J. L., *Z. Phys. Chem., Abt. B*, **41**, 98 (1938).

- (8) Viola, J. T., Seegmiller, D. W., Fannin, Jr., A. A., King, L. A., *J. Chem. Eng. Data*, **22**, 367 (1977).  
 (9) Vrieland, G. E., Stull, D. R., *J. Chem. Eng. Data*, **12**, 532 (1967).

Received for review May 26, 1977. Accepted November 15, 1977.

## Vapor Pressure of Aluminum Chloride Systems. 3. Vapor Pressure of Aluminum Chloride–Sodium Chloride Melts

John T. Viola, Lowell A. King,\* Armand A. Fannin, Jr., and David W. Seegmiller

Frank J. Sailer Research Laboratory (Air Force Systems Command) and Department of Chemistry and Biological Sciences, United States Air Force Academy, Colorado 80840

**The vapor pressures of liquid aluminum chloride–sodium chloride mixtures containing 53–74 mol % aluminum chloride were measured from the condensation temperatures of solid aluminum chloride to 251 °C. Samples were contained in Pyrex isoteniscope which utilized mercury as the manometric fluid. The mercury columns were brought to null by an external pressure which was in turn measured at each experimental point. The experimental pressures were fit to a single equation which serves the entire composition–temperature region.**

We measured the vapor pressure of aluminum chloride–sodium chloride molten mixtures as part of our continuing investigation of low-melting molten salt battery electrolytes. Previously reported vapor pressures of this system (2, 3, 5, 6) are in disagreement by more than an order of magnitude.

### Experimental Section

Preparation of the  $\text{AlCl}_3$ , the glovebox atmosphere, loading of the isoteniscope, the constant temperature bath and its temperature regulation and measurement, and the measurement of vapor pressure were all as described previously (10). A mercury manometer was used as an additional external pressure measurement device.

Powdered mixtures of  $\text{AlCl}_3$  and  $\text{NaCl}$  were loaded into a Pyrex isoteniscope similar to that used for saturated  $\text{AlCl}_3$  vapor pressures (10).  $\text{NaCl}$  was dried by maintaining it molten for several hours (7). The  $\text{AlCl}_3$ – $\text{NaCl}$  mixtures were prepared by grinding and mixing weighed amounts of  $\text{AlCl}_3$  crystals with weighed amounts of dry  $\text{NaCl}$  crystals in a glovebox.

### Results

A total of 227 measurements on 12 different compositions were taken at values of mole fraction and temperature that lay within a polygon on the  $X, t$  plane described approximately by the coordinates  $(X, t) = (0.54, 149), (0.58, 106), (0.61, 103), (0.70, 175), (0.74, 182), (0.74, 251),$  and  $(0.54, 251)$ . Pressure measurements were made at both ascending and descending temperature steps, and equilibrium was assured at each point by maintaining constant temperature for many minutes until pressure remained constant within measurement error. The samples were stirred by vigorous shaking before each mea-

surement to prevent the formation of internal concentration gradients.

The data are given in Table I. Pressures have been corrected for the vapor pressure of mercury within the isoteniscope. Interactions between  $\text{Al}_2\text{Cl}_6$  vapor and  $\text{Hg}$  vapor were assumed to be negligible. The measurement method for each datum is indicated in the table. Several additional measurements were made at temperatures lying below the  $X, t$  polygon for each of the eight compositions richest in  $\text{AlCl}_3$ . These pressures lay on the curve of saturated vapor pressure of solid  $\text{AlCl}_3$  (10), and fell distinctly away from the  $X, p, T$  surface for  $\text{AlCl}_3$ – $\text{NaCl}$  melts.

We assumed the vapor pressure of  $\text{AlCl}_3$ – $\text{NaCl}$  melts could be represented by an equation of the form

$$\log p = (A/T) + B \quad (1)$$

where

$$A = A_0 + A_1X + A_2X^2$$

$$B = B_0 + B_1X + B_2X^2 \quad (2)$$

The vapor pressures of individual compositions were fit by other investigators to equations of the form of eq 1 (3, 5, 6) or eq 1 modified by the addition to the right-hand side of the term  $\Delta C_p/R \log T$  (1, 2).

The data were least-squares fit in such a way as to minimize the perpendicular distance,  $z$ , from the experimental points to the function surface in  $X, p$ , and  $T$  given by eq 1 and 2. This procedure is similar to the fitting technique used earlier (10). The equations were solved for the values of  $A_j$  and  $B_j$  which minimized the sum,  $\sum_{i=1}^N z_i^2$  where

$$z_i = \left[ \left( \frac{\delta p_i}{\Delta p_i} \right)^2 + \left( \frac{\delta T}{\Delta T_i} \right)^2 \right]^{-1/2} \quad (3)$$

In eq 3,  $\delta p_i$  = the greater of 0.1 Torr or  $0.005 p$ ,  $\delta T = 1$  K,  $\Delta p_i = p_{\text{exp},i} - \exp[(A_i/T_{\text{exp},i} + B_i) \ln(10)]$ , and  $\Delta T_i = T_{\text{exp},i} - A_i/(\log p_{\text{exp},i} - B_i)$ . As indicated in eq 2,  $A$  and  $B$  are functions of  $X$ . Since there was an appreciable vapor space in the isoteniscope (ca. 85 cm<sup>3</sup>), the true mole fraction,  $x$ , of the melt will be different for each experimental point, depending upon the nominal mole fraction,  $X$ , and upon temperature. Accordingly, for every experimental point

$$A_i = A_0 + A_1x_i + A_2x_i^2$$

$$B_i = B_0 + B_1x_i + B_2x_i^2 \quad (4)$$

Table I. Experimental Data

$t, ^\circ\text{C}$	$p, ^a\text{Torr}$	$t, ^\circ\text{C}$	$p, ^a\text{Torr}$	$t, ^\circ\text{C}$	$p, ^a\text{Torr}$	$t, ^\circ\text{C}$	$p, ^a\text{Torr}$	$t, ^\circ\text{C}$	$p, ^a\text{Torr}$	$t, ^\circ\text{C}$	$p, ^a\text{Torr}$
$X = 0.53790$		242.26	31.35 H	188.25	33.99	226.27	188.68	196.37	159.86 H	230.96	674.34
158.28	3.67 M	242.26	33.14	189.66	34.93 H	231.94	205.07	196.37	162.39	249.68	914.98
158.28	4.33 H	250.59	30.99 H	189.66	35.53	237.23	224.53	204.16	209.38 H	250.95	931.64
190.33	8.52 M	250.59	32.70	222.84	79.45 H	242.75	238.63	204.16	211.83		
190.33	7.97			222.84	81.05	250.90	265.59	206.99	186.25 H	$X = 0.69277$	
217.83	12.79 H	$X = 0.55513$		244.87	109.91 H			206.99	191.41	166.19	322.59
217.83	12.13	142.21	2.64 M	244.87	113.38	$X = 0.62677$		209.63	201.75 H	171.93	380.53
		142.21	4.03 H	247.65	115.87 H	137.00	21.14 H	209.63	208.17	176.47	422.99
$X = 0.54250$		142.27	2.86 M	247.65	119.29	137.00	22.04 M	210.01	202.17 H	183.69	506.47
144.79	3.23 H	142.27	3.90 H	247.65	114.53 H	137.05	21.16	210.01	209.20	190.66	591.83
144.79	3.30 M	152.07	4.68 H	247.67	115.78 H	148.31	32.59 H	224.30	284.72	201.35	735.97
144.79	2.59	152.26	4.86 H	247.67	117.90	148.31	32.50	234.38	343.94	212.22	907.08
155.58	3.16 M	153.06	4.79 M	247.67	119.21	162.05	50.40 H	256.55	506.30	221.97	1076.42
155.58	3.13 H	153.06	4.96 H	247.70	113.53 H	162.05	51.04			231.04	1255.05
155.58	3.29	163.70	6.41 M	247.70	116.64	166.53	54.35 H	$X = 0.65904$		237.21	1385.44
155.87	4.42 M	163.70	6.43 H			166.53	56.70	155.61	114.77 H	250.21	1675.66
155.87	4.38 H	169.21	6.50 H	$X = 0.60977$		173.06	71.21 H	155.61	117.53		
155.87	3.37	169.21	7.11 M	134.86	12.39 M	173.06	72.61	177.72	199.88 H	$X = 0.70255$	
182.34	6.00 M	179.69	8.86 H	134.86	12.84	185.66	98.60 H	177.72	205.24	174.67	470.55
182.34	6.62 H	179.69	9.14 M	139.73	14.54 M	185.66	98.53	193.58	295.20	183.22	573.93
182.34	7.32 H	186.49	10.64 H	139.73	14.81	198.48	136.68 H	209.87	395.03	192.82	698.04
182.34	7.80 M	186.49	11.34 M	144.91	17.70 H	198.48	140.69	212.21	408.33	201.36	835.45
182.34	6.26	194.50	13.14 H	144.91	18.19 M	210.15	175.69 H	225.92	533.74	209.66	976.29
182.34	7.72	202.75	16.04 H	144.91	18.70	210.15	185.43	226.22	508.17	$X = 0.73929$	
190.41	10.28 H			150.12	21.64 H	224.48	247.37	226.33	529.68	182.02	934.82
190.41	10.68 M	$X = 0.58062$		150.12	22.75 M	236.39	305.04	240.06	669.23	189.20	1122.00
190.41	9.65	105.67	2.26 M	150.12	22.78	250.89	396.51	242.59	675.56	190.31	1146.63
195.93	9.81 H	105.67	2.11	155.74	26.00 H			250.84	799.90	197.83	1336.35
195.93	10.97 M	127.55	5.10 M	155.74	27.56	$X = 0.63046$		250.98	800.52	204.09	1524.16
195.93	9.94	127.55	4.74	163.35	33.40 H	134.17	26.76 H			211.59	1748.73
199.10	12.01 H	128.95	5.13 M	171.69	44.26 H	134.17	26.75	$X = 0.66036$		225.99	2254.49
199.10	12.12	128.95	5.03	179.55	55.89 H	151.03	44.83 H	153.55	102.90 H	226.61	2294.22
209.33	14.86 H	150.05	9.40 H	187.95	71.90 H	151.03	45.38	153.55	104.10	230.85	2478.87
209.33	15.12	150.05	9.56 M	196.40	90.26 H	152.39	44.11 H	157.88	118.12 H	239.46	2825.83
220.61	20.53 H	160.07	13.98 M	204.32	110.78 H	152.39	47.74	157.88	118.89	239.54	2846.17
220.61	20.68	160.07	14.30 H	212.49	134.75 H	166.95	72.58 H	174.16	187.30 H	239.62	2829.08
225.85	21.63 H	160.07	14.17	212.49	137.08	166.95	74.44	174.16	192.13	245.17	3089.51
225.85	22.42	183.10	28.87 H	220.72	163.42 H	179.71	97.30 H	180.13	229.65	246.00	3113.01
237.59	25.93 H	183.10	28.54	220.72	167.36	179.71	98.86	195.82	341.97	250.89	3353.37
237.59	27.09	188.25	33.91 H	226.27	182.89 H	182.58	108.14 H	204.20	401.04	251.03	3358.08
						182.58	109.87	222.71	583.06		

<sup>a</sup> The symbols "H" and "M" indicate the use of a mercury manometer and a McLeod gauge, respectively, for pressure measurement; all other pressure measurements were made with a capacitance manometer.

The van der Waals equation obtained previously (9) was solved for the number of moles of  $\text{Al}_2\text{Cl}_6$  in the vapor phase at each experimental point, and the nominal mole fractions were corrected to obtain  $x_i$ . The equation is cubic in the number of vapor-phase moles and was solved by the Newton-Raphson technique. The solution was iterated until  $x_i$  did not change more than one part in 1000. (Actually, only a single application of the Newton-Raphson technique was necessary for all points except the ten highest temperature data of the sample for which  $X = 0.73929$ .) Mole fraction corrections ranged from  $-0.00001$  to  $-0.009$ . Many of the corrections were very small but were included because all were in the same direction.

The least-squares fit was iterative. After each iteration, any points for which  $z \geq 3\sigma$  were eliminated and the remaining data were fit in the next iteration. The standard deviation,  $\sigma$ , was calculated from

$$\sigma = \left[ \frac{1}{N} \sum_{i=1}^N z_i^2 \right]^{1/2}$$

Iterations were continued until

$$\frac{2(\sigma - \sigma_{\text{previous iteration}})}{\sigma + \sigma_{\text{previous iteration}}} \leq 1.0 \times 10^{-7} \quad (5)$$

The values of  $A_j$  and  $B_j$  for eq 1 and 2 which are given in Table II are those obtained when the criterion of eq 5 was first satisfied. Seven of the original 227 points were more than  $3\sigma$  away from the calculated  $X, p, T$  surface. These do not appear

Table II. Least-Squares Fit Parameters for Equations 1 and 2


$A_0 = 6064.90$	$s_p = 1.4\text{ Torr}$
$A_1 = -29406.3$	$s'_p = 2.5\%$
$A_2 = 25360.7$	$s_T = 3.1\text{ K}$
$B_0 = -26.2772$	$s'_T = 0.65\%$
$B_1 = 100.6062$	
$B_2 = -75.1432$	

in Table I. The root mean square value of the quantity  $|z - \text{true perpendicular}| / (\text{true perpendicular})$  was 0.008. The root mean square percentage errors in pressure and temperature for all points were 2.5 and 0.65%, respectively. The surface generated by eq 1 and 2 is shown in Figure 1.

In addition to the 12 samples described above, two more  $\text{AlCl}_3$ - $\text{NaCl}$  melts were examined, for which the nominal mole fractions were 0.896 and 0.925. We noted the same separation into two phases which Kendall et al. (4) reported for  $X > 0.82$ . The pressures of these two samples were the same as the pressures of saturated liquid  $\text{AlCl}_3$  (10) at the same temperatures over the temperature range 184–238  $^\circ\text{C}$ , with a root mean square pressure difference of 1.5%.

## Discussion

Remarkable discrepancies among reported  $\text{AlCl}_3$ - $\text{NaCl}$  melt vapor pressures exist in the literature. Naryshkin (6) reported vapor pressures smaller than are reported in the present work at the same temperatures and compositions by factors ranging from 2 to 17. Galitskii's vapor pressures (3) are as much as



**Figure 1.**  $X$ ,  $\log p$ ,  $1/T$  surface for vapor pressure over  $\text{AlCl}_3$ - $\text{NaCl}$  liquid mixtures. Points are calculated from eq 1 and 2:  $\cdot$ , density within  $X$ ,  $t$  range experimentally covered;  $+$ , extrapolated values. For clarity, surface is extrapolated to intersect faces of cube.

70-fold smaller! Dewing (2) studied four compositions in the range  $X = 0.54$ – $0.66$ . His apparatus utilized an internal manometer wherein the  $\text{AlCl}_3$ - $\text{NaCl}$  melt itself was the manometric fluid. His results are depicted graphically only, but Davey et al. (1) calculated vapor pressure equations of the form

$$\log p = \Delta C_p/R \log T + A/T + B$$

from Dewing's results. These values agree reasonably well with the present work; the differences range from  $-1$  to  $-4\%$  for one composition and from  $12$  to  $-21\%$  overall. Davey et al. (1) also gave the unpublished results of Rogers (8), who investigated five compositions in the composition range  $X = 0.53$ – $0.69$ . Rogers' pressures match both those obtained in this study and Dewing's but show more scatter and yield both higher and lower pressures than either of the two other studies.

The most complete study previously reported, both in numbers of samples, 7, and in composition range,  $X = 0.54$ – $0.75$ , is that of Narita et al. (5). They used an isotenscope with mercury as the manometric fluid. Their results are lower than those of Dewing (2), Rogers (8), and the present work by a factor of 2–5.

No readily obvious reason has been suggested for the great discrepancies noted among the several studies reported. However,  $\text{AlCl}_3$ -containing melts are notoriously difficult to obtain and keep free from contamination, and most likely contaminants would reduce the vapor pressure. An exception to this is the presence of dissolved gases. Both Dewing (2) and Narita et al. (5) have commented on the presence of a residual gas pressure once a melt had been heated to well above its melting point then cooled to some lower temperature. Narita et al. were able to pump out the residual gas. Dewing corrected for its presence by assuming the residual gas partial pressure was governed by the ideal gas law. We pumped out the residual gas, but first froze the sample and cooled it to room temperature to avoid removing any  $\text{AlCl}_3$  at the same time. Initial residual pressures varied from a few Torr to around 100 Torr. We would repeatedly remelt and refreeze each sample until the residual pressure was undetectably small. The residual gas from two samples was analyzed mass spectrometrically. One sample was essentially all  $\text{HCl}$  and the other contained in addition a small amount of  $\text{N}_2$ .

The volatility correction referred to earlier was not mentioned by the other investigators. The correction may be rather substantial at high  $T$  and  $X$ . For example, the vapor pressure of the last point in Table I is 12% lower than the calculated vapor pressure of a sample whose composition is actually at the nominal value,  $X = 0.73929$ . It is not possible to tell from the information furnished whether volatilization errors were large in the previously published studies, or even if they contributed to the much smaller pressures observed by Galitskii (3), Narita et al. (5), and Naryshkin (6).

The study presented here is the most extensive reported on this system, both in regard to the number of measurements made in the temperature–composition range investigated and in the numerical evaluation. A detailed thermodynamic analysis of the data presented in this paper will be made in a forthcoming publication in conjunction with another study on the  $\text{AlCl}_3$ - $\text{NaCl}$  system.

### Glossary

$A$	fitted constant
$B$	fitted constant
$C_p$	heat capacity at constant pressure, $\text{cal mol}^{-1} \text{K}^{-1}$
$i$	index
$j$	index
$\ln$	natural logarithm
$\log$	logarithm to the base 10
$N$	the total number of experimental points used in fitting
$p$	pressure, Torr
$R$	gas constant, $1.987 \text{ cal mol}^{-1} \text{K}^{-1}$
$s_p$	root mean square of the differences between experimental pressure and pressure at the corresponding intersection of the perpendicular to the chord, Torr
$s'_p$	root mean square of $100 \times$ the difference between experimental pressure and pressure at the corresponding intersection of the perpendicular to the chord, divided by experimental pressure, %
$s_T$	root mean square of the differences between experimental temperature and temperature at the corresponding intersection of the perpendicular to the chord, K
$s'_T$	root mean square of $100 \times$ the difference between experimental temperature and temperature at the corresponding intersection of the perpendicular to the chord, divided by experimental temperature, %
$t$	temperature, $^{\circ}\text{C}$
$T$	temperature, K
$x$	apparent mole fraction of $\text{AlCl}_3$ on $\text{AlCl}_3$ - $\text{NaCl}$ scale after volatility correction
$X$	nominal apparent mole fraction of $\text{AlCl}_3$ on $\text{AlCl}_3$ - $\text{NaCl}$ scale
$z$	perpendicular distance to chord along equation; function to be treated by least-squares fitting
$\delta p$	the greater of 0.1 Torr or $0.5\% p$ ; estimated uncertainty in pressure measurement
$\delta T$	1 K; estimated uncertainty in temperatures of sample
$\sigma$	standard deviation

### Literature Cited

- (1) Davey, T. R. A., Floyd, J. M., Marcantonio, P. J., "The  $\text{AlCl}_3$ - $\text{NaCl}$  System-Vapor Pressures and Liquidus", presented before the Meeting of American Institute of Metallurgical Engineers, New York, N.Y., March 1971.
- (2) Dewing, E. W., *J. Am. Chem. Soc.*, **77**, 2639 (1955).
- (3) Galitskii, N. V., *Russ. J. Inorg. Chem.*, **13**, 1607 (1968).
- (4) Kendall, J., Crittenden, E. D., Miller, H. K., *J. Am. Chem. Soc.*, **45**, 963 (1923).
- (5) Narita, T., Ishikawa, T., Midorikawa, R., *Denki Kagaku*, **36**, 300 (1968).

- (6) Naryshkin, I. I., *Zh. Fiz. Khim.*, **13**, 690 (1939).  
 (7) Pinching, G. D., Bates, R. G., *J. Res. Natl. Bur. Stand.*, **37**, 311 (1946).  
 (8) Rogers, L., Division of Mineral Chemistry, Commonwealth Scientific and Industrial Research Organization, Melbourne, Australia, 1962, unpublished results given in ref 1.  
 (9) Viola, J. T., Fannin, Jr., A. A., King, L. A., Seegmiller, D. W., *J. Chem.*

- Eng. Data*, preceding paper in this issue.  
 (10) Viola, J. T., Seegmiller, D. W., Fannin, Jr., A. A., King, L. A., *J. Chem. Eng. Data*, **22**, 367 (1977).

Received for review June 23, 1977. Accepted November 21, 1977.

# Enthalpies of Combustion, Vaporization, and Formation of Phenylbenzene, Cyclohexylbenzene, and Cyclohexylcyclohexane; Enthalpy of Hydrogenation of Certain Aromatic Systems

Robert L. Montgomery and Frederick D. Rossini\*

Department of Chemistry, Rice University, Houston, Texas 77001

Margret Månsson

Chemical Center, University of Lund, S-220 07, Lund, Sweden

The enthalpies of combustion and vaporization were measured. Standard enthalpies of formation at 298.15 K were calculated as follows, in kcal<sub>th</sub> mol<sup>-1</sup>: for phenylbenzene, crystal 23.12 ± 0.62, hypothetical liquid 27.10 ± 0.63, gas 42.66 ± 0.62; for phenylcyclohexane, liquid -18.9 ± 1.34, gas -4.60 ± 1.35; for cyclohexylcyclohexane, liquid -66.05 ± 0.74, gas -52.19 ± 0.74. Analysis of these and earlier data shows that, within the assigned limits of uncertainty, the standard enthalpy of complete hydrogenation, in the gaseous state at 298.15 K, may be taken as (-15.88 ± 0.13) kcal<sub>th</sub>/mol of H<sub>2</sub>, for the following aromatic systems: naphthalene, phenylbenzene, phenylcyclohexane, and *n*-butylbenzene. This quantity may be useful in helping to estimate the enthalpy of hydrogenation of coal.

## 1. Introduction

This investigation is an extension of the experimental work on the relation between energy and molecular structure of hydrocarbons in which the senior author (F.D.R.) has been engaged for many years (20). With the availability of highly purified samples of phenylbenzene (diphenyl), cyclohexylbenzene (phenylcyclohexane), and cyclohexylcyclohexane (bicyclohexyl), it seemed desirable to confirm previously reported values of their standard enthalpies of combustion and obtain new values of their enthalpies of vaporization. Also, by having measurements on all three compounds made at the same time, in the same laboratory, by one investigator, and thus eliminating unknown systematic errors between them, one can obtain reliable values

of the relevant enthalpies of hydrogenation for both the gaseous and liquid states.

The combustion experiments on the three compounds were performed in the Thermochemical Laboratory at Rice University and the vaporization experiments at the University of Lund.

The values of the atomic weights used were as follows (6): C, 12.011; H, 1.0079. These yield the following molecular weights: phenylbenzene, C<sub>12</sub>H<sub>10</sub>, 154.2110; cyclohexylbenzene, C<sub>12</sub>H<sub>18</sub>, 160.2584; cyclohexylcyclohexane, C<sub>12</sub>H<sub>22</sub>, 166.3058.

## 2. Experimental Section

**Materials.** The materials used were American Petroleum Institute standard samples (purchased from the American Petroleum Institute Office of Standard Reference Materials, Carnegie-Mellon University, Pittsburgh, Pa.) having certified mole percentage impurities as follows: phenylbenzene, 0.02 ± 0.02; cyclohexylbenzene, 0.07 ± 0.05; cyclohexylcyclohexane, 0.023 ± 0.008.

**Enthalpy of Combustion.** The apparatus and procedures used in these experiments were essentially as previously reported (7, 8), with the bomb combustion apparatus having been assembled by Margrave and associates (10). The symbols used are the same as in the preceding reports (7, 8).

Fuse wire, Chromel C, was used as in the preceding investigation (8). The values used for determining the energy of ignition and the energy of combustion of the polyester (Mylar) were the same as before (8). The value used for the polyester (Mylar) was checked, satisfactorily, by a single experiment, in which enough polyester (Mylar) was burned to produce a temperature rise near 2 K.

Table I. Results of Five Experiments to Determine  $\epsilon_s$ , the Energy Equivalent of the Standard Calorimeter System at 298.15 K

$m_{BA}/g$	$10^3 k/s^{-1}$	$10^4 u/K$ $min^{-1}$	$\Delta t/K$	$\Delta t_c/K$	$\Delta \epsilon/J K^{-1}$	$q_n/J$	$q_i/J$	$\epsilon_s/J K^{-1}$
1.14659-1.16788	3.32-3.59	0.93-4.47	1.9993-2.0372	1.99105-2.02800	19.37-19.40	3.36-4.07	59.49-76.78	15238.36-15243.26
1.15669 <sup>a</sup>			2.0176 <sup>a</sup>	2.00857 <sup>a</sup>	19.38 <sup>a</sup>	3.64 <sup>a</sup>	72.12 <sup>a</sup>	15241.43 <sup>a,b</sup>

<sup>a</sup> Mean value. <sup>b</sup> Standard deviation of the mean: ±0.93.