

A Lanczos Approach to the Inverse Square Root of a Large and Sparse Matrix

Artan Boriçi

Paul Scherrer Institute, CH-5232 Villigen PSI, Switzerland

E-mail: Artan.Borici@psi.ch

Received November 18, 1999; revised March 23, 2000

I construct a Lanczos process on a large and sparse matrix and use the results of this iteration to compute the inverse square root of the same matrix. The algorithm is a stable version of an earlier proposal by the author. It can be used for problems related to the matrix sign and polar decomposition. The application here comes from the theory of chiral fermions on the lattice. © 2000 Academic Press

1. INTRODUCTION

The computation of the inverse square root of a matrix is a special problem in scientific computing. It is related to the matrix sign and polar decomposition [1].

One may define the *matrix sign* by

$$\text{sign}(A) = A(A^2)^{-1/2}, \quad (1)$$

where A is a complex matrix with no pure imaginary eigenvalues.

In polar coordinates, a complex number $z = x + iy$, is represented by

$$z = |z|e^{i\phi}, \quad \phi = \arctan \frac{y}{x}. \quad (2)$$

In analogy, the polar decomposition of a matrix A is defined by

$$A = V(A^\dagger A)^{1/2}, \quad V^{-1} = V^\dagger, \quad (3)$$

where V is the polar part and the second factor corresponds to the absolute value of A .

The mathematical literature involving the matrix sign function traces back to 1971 when it was used to solve the Lyapunov and algebraic Riccati equations [1].

In computational physics one may face a similar problem when dealing with Monte Carlo simulations of fermion systems, the so-called *sign problem* [2]. In this case the integration

measure is proportional to the determinant of a matrix and the polar decomposition may be helpful to monitor the sign of the determinant.

The example brought in this paper comes from recent progress in formulating quantum chromodynamics (QCD) on a lattice with exact chiral symmetry [3].

In continuum, the massless Dirac propagator D_{cont} is chirally symmetric; i.e.,

$$\gamma_5 D_{\text{cont}} + D_{\text{cont}} \gamma_5 = 0. \quad (4)$$

On a regular lattice with spacing a the symmetry is suppressed according to the Ginsparg–Wilson relation [4],

$$\gamma_5 D + D \gamma_5 = a D \gamma_5 D, \quad (5)$$

where D is the lattice Dirac operator.

An explicit example of a Dirac operator obeying this relation is the so-called overlap operator [5]

$$aD = 1 - A(A^\dagger A)^{-1/2}, \quad A = M - aD_W, \quad (6)$$

where M is a shift parameter in the range $(0, 2)$, which I have fixed at one.

D_W is the Wilson operator,

$$D_W = \sum_{\mu=1}^4 \gamma_\mu \nabla_\mu - \frac{a}{2} \sum_{\mu=1}^4 \Delta_\mu, \quad (7)$$

which is a nearest-neighbors discretization of the continuum Dirac operator (it violates the Ginsparg–Wilson relation). ∇_μ and Δ_μ are first and second order covariant differences given by

$$\begin{aligned} (\nabla_\mu \psi)_i &= \frac{1}{2a} (U_{\mu,i} \psi_{i+\hat{\mu}} - U_{\mu,i-\hat{\mu}}^\dagger \psi_{i-\hat{\mu}}) \\ (\Delta_\mu \psi)_i &= \frac{1}{a^2} (U_{\mu,i} \psi_{i+\hat{\mu}} + U_{\mu,i-\hat{\mu}}^\dagger \psi_{i-\hat{\mu}} - 2\psi_i), \end{aligned}$$

where ψ_i is a fermion field at the lattice site i and $U_{\mu,i}$ an $SU(3)$ lattice gauge field associated with the oriented link $(i, i + \hat{\mu})$. These are unitary 3×3 complex matrices with determinant one. A set of such matrices forms a lattice gauge “configuration.”

γ_μ , $\mu = 1, \dots, 5$ are 4×4 Dirac matrices which anticommute with each-other.

Therefore, if there are N lattice points in four dimensions, the matrix A is of order $12N$. A restive symmetry of the matrix A that comes from the continuum is the so-called γ_5 —*symmetry*, which is the hermiticity of the $\gamma_5 A$ operator.

By definition, computation of D involves the inverse square root of a matrix. This is a non-trivial task for large matrices. Therefore several algorithms have been proposed by lattice QCD physicists [6–10].

All these methods rely on matrix–vector multiplications with the sparse Wilson matrix D_W , being therefore feasible for large simulations.

In fact, methods that approximate the inverse square root by Legendre [6] and Chebyshev polynomials [7] need to know *a priori* the extreme eigenvalues of $A^\dagger A$ to be effective. This requires computational resources for at least one conjugate gradients (CG) inversion.

In [8] the inverse square root is approximated by a rational approximation, which allows an efficient computation via a multi-shift CG iteration. Storage here may be an obstacle, which is remedied by a second CG step [11].

The Pade approximation used by [9] needs the knowledge of the smallest eigenvalue of $A^\dagger A$. Therefore the method becomes effective only in connection with the D inversion [12].

The method presented earlier by the author [10] relies on taking exactly the inverse square root from the Ritz values. These are the roots of the Lanczos polynomial approximating the inverse of $A^\dagger A$.

In that work the Lanczos polynomial was constructed by applying the Hermitian operator $\gamma_5 A$. The latter is indefinite, thereby responsible for observed oscillations in the residual vector norm [10].

Here I use a Lanczos polynomial on the positive definite matrix $A^\dagger A$. In this case the residual vector norm decreases monotonically and leads to a stable method. This is a crucial property that allows a reliable stopping criterion that I will present here.

The paper is self-contained: in the next section I will briefly present the Lanczos algorithm and set the notations. In Section 3, I use the algorithm to solve linear systems, and in Section 4, give the computation of the inverse square root. The method is tested in Section 5 and conclusions are drawn in the end.

2. THE LANCZOS ALGORITHM

The Lanczos iteration is known to approximate the spectrum of the underlying matrix in an optimal way and, in particular, it can be used to solve linear systems [13].

Let $Q_n = [q_1, \dots, q_n]$ be the set of orthonormal vectors, such that

$$A^\dagger A Q_n = Q_n T_n + \beta_n q_{n+1} (e_n^{(n)})^T, \quad q_1 = \rho_1 b, \quad \rho_1 = 1/\|b\|_2, \quad (8)$$

where T_n is a tridiagonal and symmetric matrix, b is an arbitrary vector, and β_n is a real and positive constant. $e_m^{(n)}$ denotes the unit vector with n elements in the direction m .

By writing down the above decomposition in terms of the vectors q_i , $i = 1, \dots, n$ and the matrix elements of T_n , I arrive at a three-term recurrence that allows these vectors to be computed in increasing order, starting from the vector q_1 . This is the *Lanczos Algorithm*,

$$\begin{aligned} \beta_0 &= 0, \quad \rho_1 = 1/\|b\|_2, \quad q_0 = o, \quad q_1 = \rho_1 b \\ \text{for } i &= 1, \dots \\ v &= A^\dagger A q_i \\ \alpha_i &= q_i^\dagger v \\ v &:= v - q_i \alpha_i - q_{i-1} \beta_{i-1} \\ \beta_i &= \|v\|_2 \\ \text{if } \beta_i &< \text{tol}, \quad n = i, \quad \text{end for} \\ q_{i+1} &= v/\beta_i \end{aligned} \quad (9)$$

where tol is a tolerance which serves as a stopping condition.

The Lanczos algorithm constructs a basis for the Krylov subspace [13]:

$$\text{span}\{b, A^\dagger Ab, \dots, (A^\dagger A)^{n-1}b\}. \quad (10)$$

If the Algorithm stops after n steps, one says that the associated Krylov subspace is invariant.

In floating point arithmetic, there is the danger that once the Lanczos algorithm (polynomial) has approximated well some part of the spectrum, the iteration reproduces vectors which are rich in that direction [13]. As a consequence, the orthogonality of the Lanczos vectors is spoiled with an immediate impact on the history of the iteration: if the algorithm stopped after n steps in exact arithmetic, in the presence of round-off errors the loss of orthogonality would keep the algorithm going on.

3. THE LANCZOS ALGORITHM FOR SOLVING $A^\dagger Ax = b$

Here I will use this algorithm to solve linear systems, where the loss of orthogonality will not play a role in the sense that I will use a different stopping condition.

I ask for the solution in the form

$$x = Q_n y_n. \quad (11)$$

By projecting the original system onto the Krylov subspace I get

$$Q_n^\dagger A^\dagger Ax = Q_n^\dagger b. \quad (12)$$

By construction, I have

$$b = Q_n e_1^{(n)} / \rho_1. \quad (13)$$

Substituting $x = Q_n y_n$ and using (8), my task is now to solve the system

$$T_n y_n = e_1^{(n)} / \rho_1. \quad (14)$$

Therefore the solution is given by

$$x = Q_n T_n^{-1} e_1^{(n)} / \rho_1. \quad (15)$$

This way, using the Lanczos iteration, one reduces the size of the matrix to be inverted. Moreover, since T_n is tridiagonal, one can compute y_n by short recurrences.

If I define

$$r_i = b - A^\dagger Ax_i, \quad q_i = \rho_i r_i, \quad \tilde{x}_i = \rho_i x_i, \quad (16)$$

where $i = 1, \dots$, it is easy to show that

$$\begin{aligned} \rho_{i+1} \beta_i + \rho_i \alpha_i + \rho_{i-1} \beta_{i-1} &= 0 \\ q_i + \tilde{x}_{i+1} \beta_i + \tilde{x}_i \alpha_i + \tilde{x}_{i-1} \beta_{i-1} &= 0. \end{aligned} \quad (17)$$

Therefore the solution can be updated recursively and I have the following *Algorithm1* for solving the system $A^\dagger Ax = b$:

$$\begin{aligned}
& \beta_0 = 0, \quad \rho_1 = 1/\|b\|_2, \quad q_0 = 0, \quad q_1 = \rho_1 b \\
& \text{for } i = 1, \dots \\
& \quad v = A^\dagger A q_i \\
& \quad \alpha_i = q_i^\dagger v \\
& \quad v := v - q_i \alpha_i - q_{i-1} \beta_{i-1} \\
& \quad \beta_i = \|v\|_2 \\
& \quad q_{i+1} = v/\beta_i \\
& \quad \tilde{x}_{i+1} = -\frac{q_i + \tilde{x}_i \alpha_i + \tilde{x}_{i-1} \beta_{i-1}}{\beta_i} \\
& \quad \rho_{i+1} = -\frac{\rho_i \alpha_i + \rho_{i-1} \beta_{i-1}}{\beta_i} \\
& \quad r_{i+1} := q_{i+1}/\rho_{i+1} \\
& \quad x_{i+1} := y_{i+1}/\rho_{i+1} \\
& \quad \text{if } \frac{1}{|\rho_{i+1}|} < \text{tol}, \quad n = i, \quad \text{end for}
\end{aligned} \tag{18}$$

4. THE LANCZOS ALGORITHM FOR SOLVING $(A^\dagger A)^{1/2} x = b$

Now I would like to compute $x = (A^\dagger A)^{-1/2} b$ and still use the Lanczos algorithm. In order to do so I make the following observations:

Let $(A^\dagger A)^{-1/2}$ be expressed by a matrix-valued function, for example the integral formula [1]:

$$(A^\dagger A)^{-1/2} = \frac{2}{\pi} \int_0^\infty dt (t^2 + A^\dagger A)^{-1} \tag{19}$$

From the previous section, I use the Lanczos algorithm to compute

$$(A^\dagger A)^{-1} b = Q_n T_n^{-1} e_1^{(n)} / \rho_1. \tag{20}$$

It is easy to show that the Lanczos algorithm is shift-invariant; i.e., if the matrix $A^\dagger A$ is shifted by a constant, say t^2 , the Lanczos vectors remain invariant. Moreover, the corresponding Lanczos matrix is shifted by the same amount.

This property allows one to solve the system $(t^2 + A^\dagger A)x = b$ by using the same Lanczos iteration as before. Since the matrix $(t^2 + A^\dagger A)$ is better conditioned than $A^\dagger A$, it can be concluded that once the original system is solved, the shifted one is solved too. Therefore I have

$$(t^2 + A^\dagger A)^{-1} b = Q_n (t^2 + T_n)^{-1} e_1^{(n)} / \rho_1. \tag{21}$$

Using the above integral formula and putting everything together, I get

$$x = (A^\dagger A)^{-1/2} b = Q_n T_n^{-1/2} e_1^{(n)} / \rho_1. \tag{22}$$

There are some remarks to be made here:

(a) As before, by the application of the Lanczos iteration on $A^\dagger A$, the problem of computing $(A^\dagger A)^{-1/2}b$ reduces to the problem of computing $y_n = T_n^{-1/2}e_1^{(n)}/\rho_1$ which is typically a much smaller problem than the original one. But since $T_n^{1/2}$ is full, y_n cannot be computed by short recurrences. It can be computed, for example, by using the full decomposition of T_n in its eigenvalues and eigenvectors; in fact this is the method I have employed too, for its compactness and the small overhead for moderate n .

(b) The method is not optimal, as it would have been if one had applied it directly to the matrix $(A^\dagger A)^{1/2}$. By using $A^\dagger A$ the condition is squared, and one loses a factor of 2 compared to the theoretical case!

(c) From the derivation above, it can be concluded that the system $(A^\dagger A)^{1/2}x = b$ is solved at the same time as the system $A^\dagger Ax = b$.

(d) To implement the result (22), I first construct the Lanczos matrix and then compute

$$y_n = T_n^{-1/2}e_1^{(n)}/\rho_1 \quad (23)$$

To compute $x = Q_n y_n$, I repeat the Lanczos iteration. I save the scalar products, though it is not necessary.

Therefore I have the following

Algorithm2 for solving the system $(A^\dagger A)^{1/2}x = b$:

$$\beta_0 = 0, \quad \rho_1 = 1/\|b\|_2, \quad q_0 = o, \quad q_1 = \rho_1 b$$

for $i = 1, \dots$

$$v = A^\dagger A q_i$$

$$\alpha_i = q_i^\dagger v$$

$$v := v - q_i \alpha_i - q_{i-1} \beta_{i-1}$$

$$\beta_i = \|v\|_2$$

$$q_{i+1} = v/\beta_i$$

$$\rho_{i+1} = -\frac{\rho_i \alpha_i + \rho_{i-1} \beta_{i-1}}{\beta_i}$$

$$\text{if } \frac{1}{|\rho_{i+1}|} < \text{tol}, \quad n = i, \text{ end for} \quad (24)$$

Set $(T_n)_{i,i} = \alpha_i$, $(T_n)_{i+1,i} = (T_n)_{i,i+1} = \beta_i$, otherwise $(T_n)_{i,j} = 0$

$$y_n = T_n^{-1/2}e_1^{(n)}/\rho_1 = U_n \Lambda_n^{-1/2} U_n^T e_1^{(n)}/\rho_1$$

$$q_0 = o, \quad q_1 = \rho_1 b, \quad x_0 = o$$

for $i = 1, \dots, n$

$$x_i = x_{i-1} + q_i y_n^{(i)}$$

$$v = A^\dagger A q_i$$

$$v := v - q_i \alpha_i - q_{i-1} \beta_{i-1}$$

$$q_{i+1} = v/\beta_i$$

By o I denote a vector with zero entries and by U_n, Λ_n the matrices of the eigenvectors and eigenvalues of T_n . Note that there are only four large vectors necessary to store: q_{i-1}, q_i, v, x_i .

5. TESTING THE METHOD

I propose a simple test: I solve the system $A^\dagger A x = b$ by applying *Algorithm2* twice; i.e., I solve the linear systems

$$(A^\dagger A)^{1/2} z = b, \quad (A^\dagger A)^{1/2} x = z \quad (25)$$

in the above order. For each approximation x_i , I compute the residual vector

$$r_i = b - A^\dagger A x_i. \quad (26)$$

The method is tested for an SU(3) configuration at $\beta = 6.0$ on an $8^3 16$ lattice, corresponding to an order 98,304 complex matrix A .

In Fig. 1 I show the norm of the residual vector decreasing monotonically. The stagnation of $\|r_i\|_2$ for small values of *tol* may come from the accumulation of round-off error in the 64-bit precision arithmetic used here.

This example shows that the tolerance line is above the residual norm line, which confirms the expectation that *tol* is a good stopping condition for *Algorithm2*.




FIG. 1. Test of the Lanczos algorithm for the inverse square root $(A^\dagger A)^{-1/2} b$.

6. CONCLUSIONS

I have presented a Lanczos method to compute the inverse square root of a large and sparse positive definite matrix.

The method is characterized by a residual vector norm that decreases monotonically and a consistent stopping condition. This stability should be compared with that of a similar method presented earlier by the author [10], where the underlying Hermitian but indefinite matrix $\gamma_5 A$ led to appreciable instabilities in the norm of the residual vector.

In terms of complexity this algorithm requires less operations for the same accuracy than its indefinite matrix counterpart. This property is guaranteed by the monotonicity of the residual vector norm. Nonetheless, the bulk of the work remains the same.

With the improvement in store the method is complete.

It shares with methods presented in [8, 9] the same underlying Lanczos polynomial. As is well known [13], CG and Lanczos methods for solving a linear system produce the same results in exact arithmetic. In fact, CG derives from the Lanczos algorithm by solving the coupled two-term recurrences of CG for a single three-term recurrence of Lanczos. However, the coupled two-term recurrences of CG accumulate less round-off. This makes CG preferable for ill-conditioned problems.

There are two main differences between the method presented here and those in [8, 9]:

(a) Since CG and Lanczos are equivalent, they produce the same Lanczos matrix. Therefore, any function of $A^\dagger A$ translates for both algorithms into a function of T_n (given the basis of Lanczos vectors). The latter function translates into a function of the Ritz values, the eigenvalues of T_n . That is, whenever the methods of papers [8, 9] try to approximate the inverse square root of $A^\dagger A$, the underlying CG algorithm shifts this function to the Ritz values. It is clear now that if I take the inverse square root from the Ritz values exactly, I have no approximation error. This is done in *Algorithm2*.

(b) *Algorithm2* sets no limits on the amount of memory required, whereas the multi-shift CG needs to store as many vectors as the number of shifts. For high accuracy approximations the multi-shift CG is not practical. However, one may lift this limit at the expense of a second CG iteration (two-step CG) [11]. Therefore *Algorithm2* and the two-step CG have the same iteration workload, with *Algorithm2* computing exactly the inverse square root.

Additionally, *Algorithm2* requires the calculation of Ritz eigenpairs of T_n , which makes for an overhead proportional to $\sim n^2$ when the QR algorithm is used for the eigenvalues and the inverse iteration for the eigenvectors [13]. Since the complexity of the Lanczos algorithm is $\sim nN$, the relative overhead is proportional to $\sim n/N$. For moderate gauge couplings and lattice sizes this is a small percentage.

I conclude that the algorithms of [8, 9] may be used in situations where high accuracy is not required and/or A is well conditioned.

Experience with overlap fermions shows that high accuracy is often essential [7, 10]. *Algorithm2* is best suited in such situations.

REFERENCES

1. For a starting point and a thorough review of the problem see N. J. Higham, *Proceedings of "Pure and Applied Linear Algebra: The New Generation," Pensacola, FL, March 1993.*

2. For an example in simulating finite density QCD, see P. Hasenfratz and F. Karsch, *Phys. Lett. B* **125**, 308 (1983). For the Hubbard model, see D. J. Scalapino and R. L. Sugar, *Phys. Rev. Lett.* **46**, 519 (1981).
3. For a recent review see H. Neuberger, hep-lat/9909042.
4. P. H. Ginsparg and K. G. Wilson, *Phys. Rev. D* **25**, 2649 (1982).
5. H. Neuberger, *Phys. Lett. B* **417**, 141 (1998); *Phys. Rev. D* **57**, 5417 (1998).
6. B. Bunk, *Nucl. Phys. Proc. Suppl. B* **63**, 952 (1998).
7. P. Hernandez, K. Jansen, and M. Lüscher, *Nucl. Phys. B* **552**, 363 (1999).
8. H. Neuberger, *Phys. Rev. Lett.* **81**, 4060 (1998).
9. R. G. Edwards, U. M. Heller, and R. Narayanan, *Nucl. Phys. B* **540**, 457 (1999).
10. A. Boriçi, *Phys. Lett. B* **453**, 46 (1999).
11. H. Neuberger, *Int. J. Mod. Phys. C* **10**, 1051 (1999).
12. R. G. Edwards, U. M. Heller, and R. Narayanan, *Phys. Rev. D* **59**, 094510 (1999).
13. G. H. Golub and C. F. Van Loan, *Matrix Computations*, The Johns Hopkins University Press, Baltimore, 1989.
[This is meant as a general reference with original references included therein.]