

Continued from outside back cover

- 1116 **Gas-kinetic numerical studies of three-dimensional complex flows on spacecraft re-entry**
Z.-H. Li and H.-X. Zhang
- 1139 **A lattice Boltzmann method for immiscible multiphase flow simulations using the level set method**
G. Thömmes, J. Becker, M. Junk, A.K. Vaikuntam, D. Kehrwald, A. Klar, K. Steiner and
A. Wiegmann
- 1157 **Triple-decker: Interfacing atomistic–mesoscopic–continuum flow regimes**
D.A. Fedosov and G.E. Karniadakis
- 1172 **Efficient computation of the spectrum of viscoelastic flows**
J.V. Valério, M.S. Carvalho and C. Tomei
- 1188 **A spectral Galerkin method for the coupled Orr–Sommerfeld and induction equations for free-surface MHD**
D. Giannakis, P.F. Fischer and R. Rosner
- 1234 **The MAST FV/FE scheme for the simulation of two-dimensional thermohaline processes in variable-density saturated porous media**
C. Aricò and T. Tucciarelli
- 1275 **Micromixing models for turbulent flows**
D.W. Meyer and P. Jenny
- 1294 **Vertically nested nonhydrostatic model for multiscale resolution of flows in the upper troposphere and lower stratosphere**
A. Mahalov and M. Moustaoi
- 1312 **HyPAM: A hybrid continuum-particle model for incompressible free-surface flows**
Q. Zhang and P.L.-F. Liu