

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Coordination Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713455674>

X-RAY CRYSTAL STRUCTURE OF AN ORGANOINDIUM ALKOXIDE DIMER, $(\text{Ph}_2\text{InOSiMe}_3)_2$

Mark F. Self^a; Andrew T. McPhail^a; Richard L. Wells^a

^a Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University, Durham, North Carolina, USA

To cite this Article Self, Mark F. , McPhail, Andrew T. and Wells, Richard L.(1993) 'X-RAY CRYSTAL STRUCTURE OF AN ORGANOINDIUM ALKOXIDE DIMER, $(\text{Ph}_2\text{InOSiMe}_3)_2$ ', *Journal of Coordination Chemistry*, 29: 1, 27 – 32

To link to this Article: DOI: 10.1080/00958979308037122

URL: <http://dx.doi.org/10.1080/00958979308037122>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

NOTE

X-RAY CRYSTAL STRUCTURE OF AN ORGANOINDIUM ALKOXIDE DIMER, (Ph₂InOSiMe₃)₂

MARK F. SELF, ANDREW T. McPHAIL and RICHARD L. WELLS*

*Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University,
Durham, North Carolina 27708, USA*

(Received July 2, 1992; in final form October 6, 1992)

The solid-state structure of (Ph₂InOSiMe₃)₂, **1**, has been established by single-crystal X-ray analysis. Monoclinic crystals of **1** belong to space group *P*2₁/*c*, with *a* = 9.136(1), *b* = 15.503(2), *c* = 11.836(1) Å, *β* = 102.46(1)°, *Z* = 2. Refinement of atomic parameters converged at *R* = 0.050 (*R*_w = 0.063) over 1718 observed reflections with *I* > 3.0σ(*I*). The dimeric molecule lies on a crystallographic centre of symmetry with In–O = 2.159(7) and 2.149(8) Å, O–In–O = 78.9(3)°, and In–O–In = 101.1(3)°. Compound **1** is only the second example of a simple dimeric organoindium alkoxide to be structurally characterized in this manner.

KEY WORDS: Indium, alkoxide, siloxide, X-ray structure.

INTRODUCTION

Chemical investigations involving alkoxides of indium have recently undergone a resurgence^{1,2} due in part to their potential as precursors to III–VI materials, some of which have found utility in applications such as conducting and transparent oxide films for displays and solar cell windows.³ Although a variety of organoindium alkoxides have been fabricated for use in relation to these material science studies, determinations of their detailed structures have thus far been lacking.

During the period 1956 to 1974, there appeared numerous literature reports of simple organoindium alkoxides (e.g., R₂InOR': R = Me; R' = SiMe₃,⁴ SiPh₃,⁵ CMe₃,⁶ CPH₃,⁷ Me,⁸ CD₃,⁹ *t*-Bu;¹⁰ R = CD₃; R' = Me, CD₃;⁹ R = Et; R' = Me, Et⁹ R = Bu; R' = Bu, Ph).¹¹ The nature of these compounds was established by a combination of NMR spectroscopy, cryoscopic molecular weight determination, and infrared techniques. It is surprising that no X-ray crystallographic analyses of these compounds have appeared to date. In fact, the first and only report of the determination of the crystal structure of a simple dimeric organoindium alkoxide, (*t*-Bu₂InOEt)₂, was published only recently.¹² The solid-state structure of (Ph₂InOSiMe₃)₂ described herein represents only the second complete characterization of a member of this class.

* Author for correspondence.

EXPERIMENTAL

Materials and Measurements

All manipulations were performed using general Schlenk techniques in a Vacuum/Atmospheres HE-493 Dri-Lab containing an argon atmosphere. Toluene and benzene- d_6 were distilled from sodium/benzophenone ketyl under dry dinitrogen. $\text{P}(\text{SiMe}_3)_3$ ¹³ and Ph_2InCl ¹⁴ were prepared by literature procedures. The ^1H NMR spectrum was recorded on a Varian XL-300 spectrometer using a flame-sealed 5 mm tube and referenced to TMS by use of the residual protons of deuterated benzene at δ 7.15 ppm. The melting point (Thomas-Hoover Uni-melt, sealed capillary) is uncorrected. Elemental analysis was performed by E + R Microanalytical Laboratory, Inc., Corona, New York.

Preparation of $(\text{Ph}_2\text{InOSiMe}_3)_2$ 1

The title compound was prepared *via* the adventitious hydrolysis of a reaction mixture containing $\text{P}(\text{SiMe}_3)_3$ and Ph_2InCl in 100 cm³ of toluene. This hydrolysis occurred through the probable contamination of the solvent by a trace amount of water. The procedure employed was as follows. In the dry box, Ph_2InCl (0.250 g, 0.82 mmol) in 25 cm³ of toluene and $\text{P}(\text{SiMe}_3)_3$ (0.103 g, 0.41 mmol) in 75 cm³ of toluene were combined in a 150 cm³ one-necked round-bottomed flask equipped with a Teflon valve and a small magnetic follower. The resulting *milky* suspension was taken out of the box and stirred at room temperature for 24 h. Removal of the volatiles *in vacuo* yielded an oily, off-white residue that was dissolved in approximately 15 cm³ of hot toluene and cooled to -15°C for 1 day. Compound **1** was isolated in 71% yield (0.416 g, based on Ph_2InCl) from an intractable yellow oil as clear colourless platelets. $\text{Mp} = 230^\circ\text{C}$ (dec), Anal.: calc (found) for $\text{C}_{30}\text{H}_{38}\text{In}_2\text{O}_2\text{Si}_2$: C, 50.30 (49.93); H, 5.35 (5.46)%. ^1H NMR; δ 0.02 (s, 18 H, OSiMe_3), 7.02 (m, 20 H, Ph).

X-Ray Crystal Structure Analysis of 1

All measurements were made on a sample mounted inside a flame-sealed 0.6 mm thin-walled glass capillary under an inert argon atmosphere. Oscillation and Weissenberg photographs provided preliminary unit cell parameters and space group information. Crystallographic data are summarized in Table 1. The monoclinic space group $P2_1/c$ was established unequivocally from the Laue symmetry and systematic absences. Intensity data ($+h$, $+k$, $\pm l$, 3590 reflections, $\theta_{\text{max}} = 75^\circ$), recorded on an Enraf-Nonius CAD-4 diffractometer using $\text{Cu-K}\alpha$ radiation ($\lambda = 1.5418 \text{ \AA}$, graphite monochromator), were corrected for the usual Lorentz and polarization effects. An empirical absorption correction, based on the ϕ -dependency of the intensities of two reflections with ψ *ca* 90° , was applied, and equivalent reflections were averaged [$R_{\text{merge}} = 0.038$ on (I)] to yield 3378 reflections out of which those 1718 with $I > 3.0\sigma(I)$ were retained for the analysis.

The crystal structure was solved by the heavy-atom approach. Approximate indium atom coordinates were derived from a Patterson map. The remaining non-hydrogen atoms were located in a weighted F_o Fourier synthesis phased by the indium atom. Several rounds of full-matrix least-squares adjustment of non-hydrogen atom positional and thermal parameters (at first isotropic, then anisotropic) followed. Hydrogen atoms

Table 1 Crystallographic data for $(\text{Ph}_2\text{InOSiMe}_3)_2$, **1**

Molecular formula	$\text{C}_{30}\text{H}_{38}\text{In}_2\text{O}_2\text{Si}_2$
Formula weight	716.45
Crystal system	monoclinic
Space group	$P2_1/c(C_{2h})$ —No. 14
$a/\text{\AA}$	9.136(1)
$b/\text{\AA}$	15.503(2)
$c/\text{\AA}$	11.826(1)
$\beta(^{\circ})$	102.46(1)
No. of orientation refls.; $\theta(^{\circ})$ range	25; 36–40
$U/\text{\AA}^3$	1635.5(6)
Z	2
$D_c/\text{g cm}^{-3}$	1.455
μ/cm^{-1} (Cu-K α radiation, λ 1.5418 \AA)	124
T/K	298
Crystal size/mm	$0.14 \times 0.18 \times 0.22$
$T_{\text{max}}:T_{\text{min}}$	1.00:0.37
Scan type	ω -2 θ
Scanwidth/ $^{\circ}$	$1.00 + 0.14 \tan \theta$
$\theta_{\text{max}}/^{\circ}$	75
Intensity control refls.	132, 113, 123, 231;
Overall variation in standards (%);	<2
Repeat time/hr	2
Total no. of refls. ($+h, +k, \pm l$) recorded	3590
No. of non-equiv. refls. recorded	3378
R_{merge} (on I)	0.038
No. of refls. retained [$I > 3.0\sigma(I)$]	1718
No. of parameters refined	163
$R = \sum F_o - F_c / \sum F_o $	0.050
$R_w = [\sum w(F_o - F_c)^2 / \sum w F_o ^2]^{1/2}$	0.063
$\Sigma = [\sum w\Delta^2 / (N_{\text{obs.}} - N_{\text{para.}})]^{1/2}$	1.39
Max. shift/esd in final least-squares cycle	0.02
Final $\Delta\rho(\text{e}/\text{\AA}^3)$ max.;min.	0.91; -0.84

were incorporated at their calculated positions in the later iterations which converged (max. shift/esd = 0.02) at $R = 0.050$ ($R_w = 0.063$). A final difference Fourier synthesis revealed no unusual features (max. 0.91, min. $-0.84 \text{ e}/\text{\AA}^3$; both in the vicinity of the In atom). Crystallographic calculations were performed using the Enraf-Nonius Structure Determination Package.¹⁵ For all structure-factor calculations, neutral atom scattering factors and their anomalous dispersion corrections were taken from ref. 16. In the least-squares iterations, $\Sigma w\Delta^2$ [$w = 1/\sigma^2(|F_o|)$, $\Delta = (|F_o| - |F_c|)$] was minimized. Fractional atomic coordinates are listed in Table 2; selected distances and angles are provided in Table 3. A view of the solid-state structure of **1**, with the atom numbering scheme, is presented in Figure 1.

RESULTS AND DISCUSSION

From solution studies, the complexes $\text{R}_2\text{InOR}'$ where $\text{R} = \text{Me}$; $\text{R}' = \text{SiMe}_3$,⁴ SiPh_3 ,⁵ CMe_3 ,⁶ CPh_3 ,⁷ or $t\text{-Bu}$:¹⁰ $\text{R} = \text{Bu}$; $\text{R}' = \text{Bu}$ or Ph ¹¹ are postulated to exist primarily as dimers whereas when $\text{R} = \text{Me}$, $\text{R}' = \text{Me}$,⁸ $\text{R} = \text{CD}_3$, $\text{R}' = \text{Me}$ or CD_3 ,⁹ $\text{R} = \text{Et}$, $\text{R}' = \text{Me}$ or Et ,⁹ the complexes are said to be trimeric. It is unfortunate that the limited solubility of **1** in organic solvents precludes a solution cryoscopic molecular weight

Table 2 Non-hydrogen atom fractional coordinates and equivalent isotropic thermal parameters ($\times 10^3$) for $(\text{Ph}_2\text{InOSiMe}_3)_2$, **1**, with estimated standard deviations in parentheses

Atom	<i>x/a</i>	<i>y/b</i>	<i>z/c</i>	$b_{\text{eq}}(\text{\AA}^2)$
In	0.03169(8)	0.8616(5)	-0.07494(6)	3.77(1)
Si	-0.2315(3)	0.0991(2)	0.0930(2)	4.26(6)
O	-0.1169(7)	0.0416(5)	0.0320(5)	3.7(1)
C(1)	-0.3412(13)	0.1719(9)	-0.0152(10)	6.6(3)
C(2)	-0.3531(13)	0.0203(10)	0.1476(10)	6.3(4)
C(3)	-0.1190(16)	0.1622(10)	0.2162(11)	7.1(4)
C(11)	-0.0888(12)	0.0809(7)	-0.2525(6)	4.2(2)
C(12)	-0.2364(12)	0.0555(8)	-0.2769(9)	4.7(3)
C(13)	-0.3160(15)	0.0498(9)	-0.3938(11)	6.3(3)
C(14)	-0.2451(16)	0.0688(9)	-0.4801(11)	7.2(4)
C(15)	-0.0961(19)	0.0954(11)	-0.4547(9)	9.1(4)
C(16)	-0.0218(13)	0.1017(8)	-0.3413(10)	5.4(3)
C(21)	0.1857(10)	0.1767(7)	0.0214(9)	3.9(2)
C(22)	0.2155(14)	0.2539(8)	-0.0280(9)	5.5(3)
C(23)	0.3158(15)	0.3139(8)	0.0392(13)	6.9(4)
C(24)	0.3860(13)	0.2958(9)	0.1477(11)	6.3(3)
C(25)	0.3569(14)	0.2201(11)	0.1967(11)	6.8(4)
C(26)	0.2568(14)	0.1612(8)	0.1327(10)	5.7(3)

Table 3 Interatomic distances (\AA) and angles (deg.) for $(\text{Ph}_2\text{InOSiMe}_3)_2$, **1**, with estimated standard deviations in parentheses

In-O	2.159(7)	Si-O	1.654(8)
In-O'	2.149(8)	Si-C(1)	1.834(12)
In-C(11)	2.152(7)	Si-C(2)	1.858(14)
In-C(21)	2.135(10)	Si-C(3)	1.868(13)
O-In-O'	78.9(3)	C(1)-Si-C(3)	110.4(6)
O-In-C(11)	107.8(3)	C(2)-Si-C(3)	110.1(6)
O-In-C(21)	108.9(3)	In-O-In'	101.1(3)
O'-In-C(11)	106.2(3)	In-O-Si	128.3(4)
O'-In-C(21)	108.5(3)	In'-O-Si	128.7(4)
C(11)-In-C(21)	133.3(4)	In-C(11)-C(12)	119.2(7)
O-Si-C(1)	109.2(5)	In-C(11)-C(16)	121.5(8)
O-Si-C(2)	106.2(5)	In-C(21)-C(22)	120.7(7)
O-Si-C(3)	109.2(5)	In-C(21)-C(26)	121.6(9)
C(1)-Si-C(2)	111.7(6)		

determination. As is evident from the Figure, **1** exists in the solid state as a siloxy-bridged dimer. This degree of oligomerization is undoubtedly due to size effects of the ligands bonded to the indium and oxygen atoms as the aforementioned trimeric alkoxides all involve relatively small substituents.

Though the formation of **1** was accomplished by a rather serendipitous event, this does not diminish the fundamental value that the elucidation of its structure and bonding geometry contributes to the general knowledge of alkoxide chemistry. As was stated earlier, the only previously reported structure of a simple organoindium alkoxide involved the dimer (*t*-Bu₂InOEt)₂, **2**.¹² Interestingly, **2** was also prepared by a non-conventional method involving the reaction of *n*-BuLi, NHCMe₂(CH₂)₃CMe₂,




Figure 1 ORTEP diagram (40% probability ellipsoids) showing the atom numbering scheme and solid-state conformation of $(\text{Ph}_2\text{InOSiMe}_3)_2$, **1**; primed atoms are related to the unprimed atoms by a crystallographic centre of symmetry. Small circles represent hydrogen atoms.

and $t\text{-Bu}_2\text{InCl}$ in Et_2O solvent. The shortest intermolecular separations between the discrete dimeric units in crystals of **1** and **2** correspond to normal van der Waals distances. Compounds **1** and **2** both lie on crystallographic centres of symmetry and thus involve planar four-membered In_2O_2 cores. The mean In-O bond distances [2.154 Å **1**; 2.156 Å **2**] are essentially identical in both compounds, and the Si-O distance of 1.654(8) Å in **1** is in accord with the expected value of 1.63(2) Å.¹⁷ One of the oxygen atoms in **1** lies close to one of the phenyl ring planes [O-In-C(11)-C(12) torsion angle = $3(1)^\circ$] whereas both are nearly equidistant from the other phenyl ring plane [torsion angles: O-In-C(21)-C(26) = $46(1)^\circ$, O'-In-C(21)-C(26) = $-39(1)^\circ$] which, accordingly, approximately bisects the O-In-O' angle. Ring aryl substituents are similarly oriented in crystals of $[\text{In}(\text{2,4,6-trimethylphenyl})_2\text{I}]_2$, **3**,¹⁸ where the mean In-C distance of 2.169 Å associated with the bulkier ligands is slightly longer than that of 2.149 Å in **2**. In both **1** and **2**, the geometry about the oxygen atoms is distorted trigonal planar while that at the indium atoms is distorted tetrahedral. The O-In-O' and In-O-In' angles in **1** at $78.9(3)^\circ$ and $101.1(3)^\circ$, respectively, are similar to corresponding values in **2** [$75.2(3)^\circ$ and $104.8(2)^\circ$]. The considerable enlargement of the C-In-C angles [$133.3(4)^\circ$ in **1**, $127.2(4)^\circ$ in **2**, $127.7(1)^\circ$ in **3**] over the ideal tetrahedral value is probably due principally to electronic factors with intramolecular non-bonded steric interactions between ring substituents also making some contribution.

SUPPLEMENTARY DATA

Additional material consisting of H-atom coordinates, thermal parameters, observed and calculated structure factors and a complete list of bond distances and angles are available from R.L.W.

Acknowledgements

We thank the Office of Naval Research for financial support.

References

1. R. Nomura, S. Inazawa, H. Matsuda and S. Saeki, *Polyhedron*, **6**, 507 (1987).
2. R. Nomura, S. Fujii, K. Kanaya and H. Matsuda, *Polyhedron*, **9**, 361 (1990).
3. K.L. Chopra, S. Major and D.K. Pandya, *Thin Solid Films*, **102**, 1 (1983).
4. B. Armer and H. Schmidbaur, *Chem. Ber.*, **100**, 1521 (1967).
5. H. Schmidbaur, G. Kuhr and U. Kruger, *Angew. Chem. Int. Ed.*, **4**, 877 (1965).
6. H. Schmidbaur, *Angew. Chem. Int. Ed.*, **4**, 152 (1965).
7. H. Schmidbaur and F. Schindler, *Chem. Ber.*, **99**, 2178 (1966).
8. G.E. Coates and R.A. Whitcombe, *J. Chem. Soc.*, 3351 (1956).
9. V.G. Man, H. Olapinski, R. Ott and J. Weidlein, *Z. Anorg. Allg. Chem.*, **410**, 195 (1974).
10. H. Schmidbaur, *Angew. Chem.*, **77**, 169 (1965).
11. H.C. Clark and A.L. Pickard, *J. Organomet. Chem.*, **8**, 427 (1967).
12. D.C. Bradley, D.M. Frigo, M.B. Hursthouse and B. Hussain, *Organometallics*, **7**, 1112 (1988).
13. L. Chao and R.D. Rieke, *Syn. React. Inorg. Metal-Org. Chem.*, **4**, 373 (1974).
14. G. Becker and W. Holderich, *Chem. Ber.*, **108**, 2484 (1975).
15. Enraf-Nonius Structure Determination Package (SDP 3.0), Enraf-Nonius, Delft, The Netherlands (1985).
16. *International Tables for X-ray Crystallography*; The Kynoch Press, Birmingham, England (1974), Vol IV.
17. F.H. Allen, O. Kennard, D.G. Watson, L. Brammer, A.G. Orpen and R. Taylor, *J. Chem. Soc., Perkin Trans.*, **2**, S2 (1987).
18. J.T. Leman, J.W. Ziller and A.R. Barron, *Organometallics*, **10**, 1766 (1991).