

This article was downloaded by:

On: 16 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Immunoassay and Immunochemistry

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597271>

Erratum

To cite this Article (1991) 'Erratum', *Journal of Immunoassay and Immunochemistry*, 12: 4, 611 – 612

To link to this Article: DOI: 10.1080/01971529108053283

URL: <http://dx.doi.org/10.1080/01971529108053283>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ERRATUM

Due to a printing error, the following figures were inadvertently transposed on pp. 436-437 in the article "Diagnostic Use of Polyclonal Antibodies Raised in Mouse Ascitic Fluid in Bancroftian Filariasis" by K. Cheirmaraj, M.V.R. Reddy, and B.C. Harinath. The article appeared in the Vol. 11, No. 4, 1990 issue of JOURNAL OF IMMUNOASSAY, on pp. 429-444. The order in which the figures should have appeared is as follows.




FIGURE 1 : Scattergram of the filarial antigen levels against anti *Bm*A (PBS) SAg-antibody in sera samples of (A) microfilaraemia, (B) acute filariasis (C) subacute filariasis (D) chronic filariasis, (E) endemic normals, (F) non-endemic normals, (G) Onchocerciasis, (H) Strongyloides (I) *Toxocara canis*, (J) *Echinococcus* and (K) *Brugia malayi*. The sera dilution at 1:300 was taken as the threshold level for positivity.




FIGURE 2 : Scattergram of the filarial antigen levels against anti BmA (SDS) SAg-antibody in sera samples of (A) microfilaraemia, (B) acute filariasis, (C) subacute filariasis, (D) chronic filariasis, (E) endemic normals, (F) non-endemic normals, (G) Onchocerciasis, (H) Strongyloides, (I) Toxocara canis, (J) Echinococcus and (K) Brugia malayi. The sera dilution at 1:300 was taken as the threshold level for positivity.