

This article was downloaded by:

On: 23 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Liquid Chromatography & Related Technologies

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597273>

Subject Index to Volume 26

Online publication date: 12 February 2003

To cite this Article (2003) 'Subject Index to Volume 26', *Journal of Liquid Chromatography & Related Technologies*, 26: 20, 3545 – 3581

To link to this Article: DOI: 10.1081/JLC-120028188

URL: <http://dx.doi.org/10.1081/JLC-120028188>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Subject Index to Volume 26

A

- Abscisic acid enantiomers, chiral separation by capillary electrophoresis with cyclodextrin additives, 1709–1717
- Accelerated solvent extraction (ASE) conditions, optimization for the HPLC determination of rutin and isoquercitrin in *Sambucus nigra* L., 2381–2397
- Acetaminophen
determination in porcine microsomal samples, 925–935
HPLC determination of dextropropoxyphene related substances in combination with in tablet and capsule formulations, 1977–1990
HPTLC with densitometry for assay in multicomponent extra strength analgesic tablets, 3453–3462
- Achyranthes bidentata*, CCC with an aqueous two-phase system using a cross-axis coil planet centrifuge for purification of polysaccharides from, 1895–1903
- Acrylamide, determination in fried foods by HPLC–electrospray mass spectrometry with SEC, 1877–1884
- Active pharmaceutical ingredient, high-low chromatography for quantitation of propyl gallate in, 1743–1753
- Acylglycerols, identification and quantification in castor oil by HPLC with ELSD, 773–780
- Adamantyl modified silica, via olefin hydrosilation on a hydride intermediate, 1169–1195
- S-Adenosyl-L-methionine, determination in dietary supplement preparations by capillary electrophoresis, 2481–2490
- Adsorption separation, for the extracts from *Ginkgo biloba* leaves using intermediate polarity resins, 2941–2960

3545

DOI: 10.1081/JLC-120028188
Copyright © 2003 by Marcel Dekker, Inc.

1082-6076 (Print); 1520-572X (Online)
www.dekker.com

MARCEL DEKKER, INC.
270 Madison Avenue, New York, New York 10016



- Alkylamide phase, application to the separation of inorganic anions by RP-HPLC, 1197–1206
- Aluminum, extraction with toluene and HPLC determination in the form of an 8-hydroxyquinoline derivative, 273–283
- Amaryllidaceae plants, HPLC determination of galanthamine in, 3217–3233
- Amines, cresyl violet for indirect fluorescence in capillary electrophoresis, 3387–3400
- Amino acids
 free pool, TLC analysis in cercariae, rediae, encysted metacercariae, and excysted metacercarie of *Echinostoma caproni*, 2697–2702
 phenyl isothiocyanated, HPLC enantiomeric resolution on teicoplanin-bonded phase using an acetonitrile-based mobile phase, 3475–3495
 underivatized, capillary zone electrophoresis techniques combined with indirect UV, direct UV, and mass spectrometric detection for determination in infusion solutions, 2783–2795
- β -Amino acids, effect of the tethering groups of chiral stationary phases based on (+)-(18-crown-6)-2,3,11,12-tetracarboxylic acid on the liquid chromatographic resolution of, 1083–1094
- Amino stationary phase, for study of the vancomycin dimerization dependence on solute enantioselectivity, 1027–1039
- trans*-2-Aminocyclohexanol, enantiomeric separation on a crown ether stationary phase using ELSD, 425–442
- Amiodarone hydrochloride, RP-HPLC estimation of, 1315–1326
- Amoxicillin, simultaneous determination with sulbactam pivoxil in a combination formulation, 1927–1936
- Anilines, positional isomeric mono- and di-substituted, RP-HPLC separation and determination on an *R,S*-hydroxypropyl ether β -cyclodextrin column, 1–15
- Anions, new electrolyte composition for determination by capillary electrophoresis with indirect UV detection, 709–722
- Antibiotics, ionophore, HPTLC detection in feeds and animal tissues, 147–156
- Apple catechin oligomers, separation by CCC, 1609–1621
- Aristolochic acid and aristolactam analogues, RP-HPLC resolution and quantitative determination in traditional Chinese medicine, 3057–3067
- Aromatic amine isomers, ion-pair RP-HPLC determination of, 85–98
- Aromatic diamidines, optimized analysis by HPLC, 3511–3525
- ASE. *See* Accelerated solvent extraction



- Aspartate neurotransmitters, HPLC analysis following *o*-phthalaldehyde-mercaptoethanol derivatization, 1797–1808
- Aspirin
coupled flow injection–capillary electrophoresis system for kinetic study of hydrolysis in aqueous solution, 1695–1707
HPLC determination of dextro-propoxyphene related substances in combination with in tablet and capsule formulations, 1977–1990
- Aspirin–butalbital, analysis in serum using a monolithic C₁₈ column, 2567–2579
- Association constants, chromatographic determination between psoralen derivatives and modified β -cyclodextrin, 871–882
- Atrazine, HPLC analysis in freshwater bivalves, 1885–1894
- B**
- Balsamic vinegar, HPLC determination of organic acids in, 2177–2187
- Barbiturates, TLC determination of lipophilicity of, 3277–3287
- Barium (II), HPCPC separation with organophosphorus compounds, 629–640
- Basic compounds, mixed mode of hydrophilic and ionic interaction pressurized capillary electrochromatography separation of, 2119–2131
- Benzoyl chloride derivatization, determination of propylene glycol by HPLC with, 3413–3431
- Benzoyllecgonine, HPLC determination in human hair, 2003–2012
- Benzoylphenylurea insecticides, correction of the matrix effect in determination in vegetables, 297–316
- Betamethasone, HPTLC determination in tablets, 2709–2717
- Bile acids
separation by TLC, 1095–1108
TLC determination of lipophilicity for, 2741–2750
- Bile salts, RP-HPLC estimation in complex tanning brines, 1041–1050
- Biomimetic synthesis, HPLC with UV detection for quantitative analysis of kielcorins in, 29–41
- 1-1'-Bi-2-naphthol enantiomers, continuous separation by an eight-column simulated moving bed system, 1853–1875
- Biopolymer retention process, thermodynamics in reversed-phase liquid chromatography, 3199–3215
- Biotin, determination in pharmaceutical preparations by HPLC and/or MEKC, 195–205
- Bivalves, HPLC analysis of atrazines in, 1885–1894
- Bronopol, as an ingredient of a new test mixture for evaluation of HPLC columns, 737–750



Bucillamine, HPLC analysis by derivatization with *N*-(1-pyrenyl)maleimide in human blood, 3265–3275

C

Caffeine

HPLC determination of dextro-propoxyphene related substances in combination with in tablet and capsule formulations, 1977–1990

HPTLC with densitometry for assay in multicomponent extra strength analgesic tablets, 3453–3462

Calcium (II), HPCPC separation with organophosphorus compounds, 629–640

Candesartan cilexetil, simultaneous analysis with hydrochlorothiazide in human plasma and dosage forms using HPLC with a photodiode array detector, 2581–2591

Capillary electrochromatography mixed mode hydrophilic and ionic interaction pressurized, for separation of basic compounds, 2119–2131

ultrashort open-tubular capillary column with modified silica-gel thin layer for, 697–707

Capillary electrophoresis for analysis of pharmacologically active xanthone compounds from *Swertia przewalskii pissjauk* extract, 1219–1229

for chiral separation of fluoxetine and its analogs with charged cyclodextrins, 2351–2367

co-electroosmotic flow, influence of organic modifiers on the separation of carboxylic acids using, 455–468

comparison with HPLC for the separation of xanthenes, 993–1003

coupled with flow injection for kinetic study of aspirin hydrolysis in aqueous solution, 1695–1707

cresyl violet for indirect fluorescence of amines in, 3387–3400

with cyclodextrin additives, for chiral separation of enantiomers of abscisic acid, 1709–1717

for determination of *S*-adenosyl-*L*-methionine in dietary supplement preparations, 2481–2490

evaluation of cyclodextrins as chiral selectors in the separation of monoterpenes by, 505–516

with indirect UV detection, new electrolyte composition for determination of anions by, 709–722

of polyamines with universal indirect chemiluminescence detection using cobalt (II) as a probe ion, 355–367

for quantitative analysis of biological and chemical



- insecticide mixture (erratum), 1161
for separation and quantitation of ractopamine stereoisomers using cyclodextrins as chiral additives, 1677–1693
for simultaneous determination of methylxanthines in human plasma, 1913–1925
for studies of molecular association of glutenins and maltodextrin, 2465–2479
See also Chiral capillary electrophoresis; High-performance capillary electrophoresis
- Capillary liquid chromatography, evaluation of cyclodextrins as chiral selectors in the separation of monoterpenes by, 505–516
- Capillary zone electrophoresis techniques, combined with indirect UV, direct UV, and mass spectrometric detection for determination of underivatized amino acids and vitamin B₆ in infusion solutions, 2783–2795
- Carbamazepine, LC–MS/MS determination in human serum, 1937–1957
- Carboxylic acids, influence of organic modifiers on separation using co-electroosmotic flow capillary electrophoresis, 455–468
- Carotenoid colorings, TLC simultaneous analysis in foods, 819–832
- Caseinomacropptide, determination by an RP-HPLC method and monitoring of the addition of rennet whey to powdered milk, 99–107
- Castor oil, HPLC with ELSD for identification and quantification of molecular species of acylglycerols in, 773–780
- Catechin oligomers, apple, separation by CCC, 1609–1621
- Catecholamine, quantification by RP-HPLC with CoulArray multi-electrode chemical detector system, 3433–3451
- Cation exchange packings, strong, based on monodisperse poly(glycidyl methacrylate-co-ethylenedimethacrylate) particles, 963–976
- CCC. *See* Countercurrent chromatography
- Cefaclor, HPLC determination in human plasma, 3315–3323
- Cellulose *tris*-(3,5-dimethylphenyl-carbamate) chiral stationary phase, for HPLC enantio-separation of chiral tetrahedral heterometal clusters, 2505–2516
- Cellulose *tris*-(phenylcarbamate) (Chiralcel[®] OC) stationary phase, for preparative chiral chromatography of trans-stilbene oxide, 3235–3248
- Cephalosporins, zwitterionic, effect of ion pairing agents in the retention profile of, 937–952
- Ceramide III, quantitative analysis in *Saccharomyces cerevisiae* by normal phase HPLC, 617–627



- Chiral additives, capillary electrophoretic separation and quantitation of ractopamine stereoisomers using cyclodextrins as, 1677–1693
- Chiral analysis, of *R*- and *S*-1-phenyl-1-butanol in biphasic *Saccharomyces cerevisiae* mediated culture, 3249–3264
- Chiral capillary electrophoresis, for determination of 2',3'-dideoxy-5-fluoro-3'-thiacytidine in rat plasma, 3037–3044
- Chiral HPLC, for resolution of the Wieland–Miescher ketone and derivatives, 409–424
- Chiral liquid chromatography, for direct separation and quantitative determination of flurbiprofen enantiomers in pharmaceuticals, 517–527
- Chiral resolution system, for development of preparative chiral separations, 529–543
- Chiral tetrahedral heterometal clusters, HPLC enantio-separation on a cellulose *tris*-(3,5-dimethylphenylcarbamate) chiral stationary phase, 2505–2516
- Chiralcel[®] OC. See Cellulose *tris*(phenylcarbamate)
- Chloramphenicol, analysis and stability of residues in food samples during storage, 2535–2549
- Chlorinated species, trace, enrichment in a complex matrix of fatty acids using HPLC in conjunction with gas chromatography–halogen specific detection, 1809–1826
- 4-Chloro-3,5-dinitrobenzoic acid amides of α,β -aminoalcohols and α -arylethylamines, enantiomer separation and molecular recognition with chiral stationary phases on, 63–83
- Chlorophyll degradation products, HPLC-UV-APCI-MS separation and characterization in silkworm, 3183–3197
- Chromatographic models, as tools for scale-up of isolation of natural products by semi-preparative HPLC, 177–193
- Chromatography theory (book review), 2023–2024
- Chromium (III), silica gel phases–immobilized-8-hydroxyquinoline for separation and pre-concentration from water samples, 3045–3056
- Cleavage position, HPSEC determination for a protein in enzymatic hydrolysis, 1787–1796
- Cobalt (II), as a probe ion for capillary electrophoresis with universal indirect chemiluminescence of polyamines, 355–367
- Cocaine, HPLC determination in human hair, 2003–2012
- Cocoa powder, HPLC determination of ochratoxin A in, 585–598
- Collagen and collagen-like proteins, roles within the proteome, 2225–2254



- Colloidal particles, high-speed analysis by micro-channel thermal field-flow fractionation, 849–869
- Column switching, for analysis of *R*- and *S*-1-phenyl-1-butanol in biphasic *Saccharomyces cerevisiae* mediated culture, 3249–3264
- Columns
- based on polysaccharide, effects on enantioselectivity by use for multimodal elution, 2083–2101
 - HPLC, bronopol as an ingredient of a new test mixture for evaluation of, 737–750
 - R,S*-hydroxypropyl ether β -cyclodextrin, for RP-HPLC separation and determination of positional isomeric mono- and di-substituted anilines and phenols, 1–15
 - monolithic C₁₈, for analysis of aspirin-butalbital in serum, 2567–2579
 - monolithic silica HPLC, for determination of sildenafil citrate in pharmaceutical preparations, 2897–2908
 - porous graphitic carbon, for HPLC determination of haloperidol and its degradation products in pharmaceuticals, 261–271
 - rotating coiled, retention of solids in, 1649–1657
 - Sephadex LH-20, chromatographic separation of phenolic compounds from rapeseed with ethanol as the mobile phase, 2157–2165
 - series coupled ion exclusion, for HPLC determination of low molecular weight organic acids in honey, 1231–1253
 - silica-based and polymer-based ion exchange, retention of glucosinolates and inorganic ions on, 751–761
 - ultrashort open-tubular capillary, with modified silica-gel thin layer for capillary electrochromatography, 697–707
- Computational chemical simulation, of chromatographic retention of phenolic compounds, 2031–2039
- Corn protein zein
- TLC study of the binding of nonionic surfactants to, 2751–2761
 - TLC study of the effects of pH and salts on binding of ring-substituted phenol derivatives to, 2303–2313
- Corrinoid compounds, TLC analysis in fish sauce, 2703–2707
- CoulArray multi-electrode chemical detector system, catecholamine quantification by RP-HPLC with, 3433–3451
- Coumarin based fluorogenic derivatization reagent, for labelling free carboxyl groups, 385–400



Countercurrent chromatography (CCC)
 with an aqueous two-phase system using a cross-axis coil planet centrifuge for purification of polysaccharides from *Achyranthes bidentata*, 1895–1903
 characterization of new phase systems in, 1551–1564
 effect of operating conditions and phase system properties on scale-up on resolution in, 1475–1491
 ICP-MS determination of plutonium in soil after pre-separation by, 1593–1607
 modelling using an eluting countercurrent distribution model, 1449–1474
 a new class of solvents for, 1493–1508
 organic phase retention in, 1537–1550
 for preparation of national certified reference materials of active compounds from natural products, 1565–1577
 property calculation and prediction for selecting solvent systems in, 1397–1415
 selection of aqueous two-phase solvent systems in, 1509–1520
 for separation of apple catechin oligomers, 1609–1621
 for separation and identification of flavonoids in an extract

from the seeds of *Oroxylum indicum*, 1623–1636
 for the separation of lutein from a crude extract of marigold flower petals, 1659–1669
 for separation of soybean isoflavones by different types of coil planet centrifuges with two-phase solvent systems, 3497–3509
 slow rotary
 large convoluted tubing for scale-up of, 1991–2002
 parameter study, 1827–1838
 small coil-volume instrument for direct interfacing with mass spectrometry, 1345–1354
 stationary phase retention in, 1373–1396
 UNIFAC estimation of separation factor for analogues in, 1431–1447
See also High-speed countercurrent chromatography; Toroidal-coil countercurrent chromatography
 Creatinine, simultaneous measurement with purine derivatives in ruminant's urine using ion-pair HPLC, 2961–2968
 Cresyl violet, for indirect fluorescence of amines in capillary electrophoresis, 3387–3400
 Crude oil spills, comparison of modern extraction techniques for analysis of soil contaminated with, 247–259
 Curcuminoids, validation of liquid chromatography/electrospray-mass spectro-



- metric determination in foods, 53–62
- β -Cyclodextrin, modified, determination of association constants with psoralen derivatives, 871–882
- Cyclodextrins
- as additives for chiral separation of abscisic acid by capillary electrophoresis, 1709–1717
 - charged, for chiral separation of fluoxetine and its analogs by capillary electrophoresis, 2351–2367
 - as chiral additives for capillary electrophoretic separation and quantitation of ractopamine stereoisomers, 1677–1693
 - as chiral selectors in the separation of monoterpenes by capillary liquid chromatography and capillary electrophoresis, 505–516
- Cyclohexanediol fatty acid diesters, as model compounds for mechanistic studies in silver ion HPLC, 1905–1912
- D**
- Dammarane-saponins, separation from notoginseng by HSCCC coupled with ELSD, 1579–1591
- Deflazacort, RP-HPLC drug dissolution studies and determination in pharmaceutical formulations and human serum, 2141–2156
- Densitometry
- assay of caffeine and acetaminophen in multicomponent extra strength analgesic tablets by HPTLC with, 3453–3462
 - determination of mometasone furoate in topical preparations by TLC with, 109–117
- Dextro-propoxyphene related substances, HPLC determination in combination with aspirin, acetaminophen, and caffeine in tablet and capsule formulations, 1977–1990
- Diamide-type chiral stationary phase, for chiral resolution by normal and reversed-phase HPLC, 2103–2117
- Diazinon, HPLC determination in environmental water and soil samples, 483–497
- 2',3'-Dideoxy-5-fluoro-3'-thiacytidine, chiral capillary electrophoretic determination in rat plasma, 3037–3044
- Dietary supplement preparations, capillary electrophoresis determination of *S*-adenosyl-*L*-methionine in, 2481–2490
- Di-(2-ethylhexyl) phosphoric acid zirconium salt, HPLC/mass spectrometry determination of radiolysis products in, 2593–2604
- Di-(2-ethylhexyl)phthalate metabolites, high throughput analysis in urine for exposure assessment, 2167–2176



- Digitalis ambigua* × *Digitalis lanata* leaves, HPLC quantitative determination of lanatosides in, 2013–2021
- Dimethoate, SPE for determination in environmental water and soil samples by MEKC, 545–557
- Diode array detection, study of stability of solutions containing hydromorphone, ketorolac, haloperidol, midazolam, famotidine, metoclopramide, dimenhydrinate, and scopolamine by HPLC with, 2909–2929
- N*-(Dithiocarboxy)sarcosine, determination of mercury(II) ion by solvent extraction with, 2797–2808
- DNA, removal from a protein solution with cross-linked poly(ethyleneimine) spherical particles, 231–246
- DNA stretching, slalom chromatography of viscosity-temperature dependence on, 883–893
- Drug control analysis, optimization of an RP-HPLC method for, 3401–3412
- Dual parallel liquid chromatography/dual mass spectrometry, of bovine brain total lipid extract, 3147–3181

E

- Echinostoma caproni*, TLC analysis of the free pool amino acids in cercariae, rediae, encysted metacercariae, and excysted metacercariae of, 2697–2702

- Electrochemical detection, determination of scutellarin in *Scutellaria barbata* extract by liquid chromatography with, 2133–2140
- Electrospray ionization mass spectrometry (ESI-MS)/photodiode array detection, analysis of stibonic acids by ion exchange chromatography with, 2041–2051
- Electrospray mass spectrometric detection
for analysis of safflower yellow in food, 1207–1217
HPLC with SEC for determination of acrylamide in fried foods, 1877–1884
- ELSD. *See* Evaporative light scattering detection
- Eluting countercurrent distribution model, for modelling CCC, 1449–1474
- Empirical peak shape equations, for separate determination of rate constants in a chromatographic column and eluent, 285–296
- Enalaprilat, HPLC determination in human plasma, 987–991
- Enantioselectivity, of solute, use of an amino stationary phase for study of vancomycin dimerization dependence on, 1027–1039
- Enantioseparation
of derivatives of phenylcarbamic acid using π -complex stationary phase in HPLC, 2331–2350
of organophosphonate derivatives on amylose



- tris*(3,5-dimethylphenyl-carbamate) chiral stationary phase by HPLC, 3075–3084
- Endosulfan and its major metabolites, RP-HPLC analysis of, 1069–1082
- Environmental analysis (book review), 833–834
- Enzymatic hydrolysis, HPSEC determination of the cleavage position of a protein in, 1787–1796
- ESI-MS. *See* Electrospray ionization mass spectrometry
- Etoricoxib, HPLC for impurity and quantitative analysis of, 2551–2566
- Evaporative light scattering detection (ELSD) coupled with HSCCC for separation of dammaranesaponins from notoginseng, 1579–1591
- for determination of piperazine in pharmaceutical drug substances, 3093–3104
- for enantiomeric separation of *trans*-2-aminocyclohexanol on a crown ether stationary phase, 425–442
- identification and quantification of molecular species of acylglycerols in castor oil by HPLC with, 773–780
- inorganic anion analysis by ion exchange chromatography with, 3117–3128
- Experimental design, for evaluation of HPLC assay robustness, 2975–2985
- Experimental parameters, effect on the HPLC separation of peptides and proteins, 2255–2283
- Exposure assessment, of di-(2-ethylhexyl)phthalate metabolites in urine by high throughput analysis, 2167–2176
- F**
- Famotidine, HPTLC determination in acid reduction tablets and videodensitometry of fluorescence quenched zones, 2719–2727
- Fat soluble vitamins, determination by RP-HPLC with UV detection, 641–663
- Fatty acids
- complex matrix, HPLC combined with gas chromatography–halogen specific detection for enrichment of trace chlorinated species in, 1809–1826
 - TLC investigation of a homologous series of, 2663–2671
- Fenitrothion, HPLC determination in environmental water and soil samples, 483–497
- Ficus* leaves, medium pressure solid–liquid extraction compared with rotation planar extraction of, 3289–3305
- Field-flow fractionation, frit inlet asymmetrical flow, effect of inlet frit lengths on the hydrodynamic relaxation efficiency in, 2369–2379



- Fish sauce, TLC analysis of
corrinoid compounds in,
2703–2707
- Flavonoids, CCC separation and
identification in an extract
from the seeds of *Oroxylum*
indicum, 1623–1636
- Flow field-flow fractionation
and MALLS, for characteriza-
tion of polymeric wheat
proteins, 2771–2781
retention behavior of polystyrene
latex particles and gold
colloids on different
channel walls in,
3003–3035
- Flow injection
coupled with capillary electro-
phoresis for kinetic study
of aspirin hydrolysis in
aqueous solution,
1695–1707
using UV detection, for
determination of naproxen
sodium in pharmaceutical
formulations, 401–408
- Flowing eluent wall, for operating
segmentation for single
and multi-channel OPLC
separation, 2611–2627
- Flunarizine dihydrochloride, HPLC
determination in tablets,
1059–1067
- Flunixin, determination in meat
using liquid chromatography/
mass spectrometry, 791–801
- Fluorescence, indirect, of amines
in capillary electrophoresis
using cresyl violet,
3387–3400
- Fluoride, determination in
biological samples, 763–772
- Fluorogenic derivatization reagent,
coumarin based, for labelling
free carboxyl groups,
385–400
- Fluorogenic reagent, for labelling
carboxylic acids in HPLC
(erratum), 1327
- Fluoroquinolone derivatives, SPE
from a biological matrix for
biodegradation by liquid
chromatography, 1731–1741
- Fluoroquinolones, TLC separation
of, 2673–2685
- Fluoxetine and its analogs, chiral
separation with charged
cyclodextrins by capillary
electrophoresis, 2351–2367
- Flurbiprofen enantiomers, direct
separation and quantitative
determination in pharmaceu-
ticals, 517–527
- Fly ash, microwave-assisted
extraction of polychlorinated
biphenyls and polychlorinated
dibenzo-*p*-dioxins from,
803–818
- Fourier transform infrared (FTIR),
combined with HPTLC for
identification of phthalazine
derivatives, 2687–2696
- Free carboxyl groups, coumarin
based fluorogenic derivati-
zation reagent for labelling of,
385–400
- FTIR. *See* Fourier transform
infrared
- Fuel spills, comparison of modern
extraction techniques for
analysis of soil contaminated
with, 247–259
- Furosine, determination in honey,
317–326



G

- Galanthamine, HPLC determination in Amaryllidaceae plants, 3217–3233
- Gas chromatography, controlling the structure of trifluoroacetylated hydrazino pharmaceutical intermediate for improved analyses by, 3371–3386
- Gas chromatography–halogen specific detection, combined with HPLC for enrichment of trace chlorinated species in a complex matrix of fatty acids, 1809–1826
- Gel permeation chromatography (GPC), with viscometry for characterization of hyperbranched aliphatic polyesters and their trimethylsilylated derivatives, 207–230
- Ginkgo biloba* leaf extracts, adsorption separation using intermediate polarity resins, 2941–2960
- Glipizide and gliclazide, micellar liquid chromatographic simultaneous determination of metformin in its multi-component dosage forms with, 1117–1133
- Glucosinolates, retention on silica-based and polymer-based ion exchange columns, 751–761
- Glutamate neurotransmitters, HPLC analysis following *o*-phthalaldehyde-mercaptoethanol derivatization, 1797–1808
- Glutenins, capillary electrophoresis of molecular association with maltodextrin, 2465–2479
- Glycolipids, separation with TC-CCC, 1135–1147
- Glycyrrhetic acid, HPCE quantitative determination in compound liquorice tablets, 43–51
- Glycyrrhizic acid, HPCE quantitative determination in compound liquorice tablets, 43–51
- Gold colloids, retention behavior on different channel walls in flow field-flow fractionation, 3003–3035
- GPC. *See* Gel permeation chromatography
- Gravity field, influence on the interface of two immiscible liquids, 1521–1535
- Guanxin II, HPLC simultaneous separation and determination of bioactive components in, 469–482

H

- Haloperidol and its degradation products, HPLC determination in pharmaceuticals using a porous graphitic carbon column, 261–271
- Herbicides
acidic, TC-CCC analysis in water, 1637–1648
micro-HPLC with temperature changes investigation of retention behavior of, 3357–3369



- High performance capillary electrophoresis (HPCE), for quantitative determination of glycyrrhizic acid, glycyrrhetic acid, morphine, and sodium benzoate in compound liquorice tablets, 43–51
- High performance centrifugal partition chromatography (HPCPC)
for isolation and purification of kava lactones, 3069–3074
for separation of calcium (II), magnesium (II), strontium (II), and barium (II) with organophosphorus compounds, 629–640
- High performance frontal analysis, of protein binding for perillyl alcohol enantiomers, 2861–2871
- High performance hydrophobic interaction chromatography, renaturation with simultaneous purification of recombinant human proinsulin from *E. coli* with, 683–695
- High performance liquid chromatography (HPLC)
for analysis of aspartate and glutamate neurotransmitters following o-phthalaldehyde-mercaptoethanol derivatization, 1797–1808
for analysis of atrazine in freshwater bivalves, 1885–1894
for analysis of bucillamine by derivatization with *N*-(1-pyrenyl)maleimide in human blood, 3265–3275
- for analysis of propolis extract, 2399–2409
- bronopol as an ingredient of a new test mixture for evaluation of columns for, 737–750
- chiral, for resolution of the Wieland–Miescher ketone and derivatives, 409–424
- comparison with capillary electrophoresis for the separation of xanthenes, 993–1003
- in conjunction with gas chromatography–halogen specific detection, for enrichment of trace chlorinated species in a complex matrix of fatty acids, 1809–1826
- controlling the structure of trifluoroacylated hydrazino pharmaceutical intermediate for improved analyses by, 3371–3386
- for determination of acetaminophen and phenacetin in microsomal porcine amples, 925–935
- for determination of aluminum in the form of an 8-hydroxyquinoline derivative, 273–283
- for determination of biotin in pharmaceutical preparations, 195–205
- for determination of cefaclor in human plasma, 3315–3323
- for the determination of cocaine and benzoylecgonine in human hair, 2003–2012
- for determination of dextropropoxyphene related



- substances in combination with aspirin, acetaminophen, and caffeine in tablet and capsule formulations, 1977–1990
- for determination of diazinon and fenitrothion in environmental water and soil samples, 483–497
- for determination of enalaprilat in human plasma, 987–991
- for determination of flunarizine dihydrochloride in tablets, 1059–1067
- for determination of galanthamine in Amaryllidaceae plants, 3217–3233
- for determination of haloperidol and its degradation products in pharmaceuticals using a porous graphitic carbon column, 261–271
- for determination of impurities in 1-(2-thienylethyl)-4-imidazoline-2-thione, 573–583
- determination of low molecular weight organic acids in honey, 1231–1253
- for determination of ochratoxin A
- in cocoa powder, 585–598
 - in wine samples, 119–133
- for determination of organic acids in traditional balsamic vinegar, 2177–2187
- for determination of phenolic acids in *Lamiaceae* family plants, 443–453
- for determination of sulfathiazole in French honeys, 953–961
- with diode array detection, for studying the stability of solutions containing hydromorphone, ketorolac, haloperidol, midazolam, famotidine, metoclopramide, dimenhydrinate, and scopolamine, 2909–2929
- of dissolution of tamsulosin from pharmaceutical dosage forms, 1109–1116
- effect of experimental parameters on separation of peptides and proteins by, 2255–2283
- and electrospray mass spectrometry with SEC for determination of acrylamide in fried foods, 1877–1884
- with ELSD, for identification and quantification of molecular species of acylglycerols in castor oil, 773–780
- for enantiomeric resolution of phenyl isothiocyanated amino acids on teicoplanin-bonded phase using an acetonitrile-based mobile phase, 3475–3495
- for enantioseparation
- of chiral tetrahedral heterometal clusters on a cellulose *tris*-(3,5-dimethylphenylcarbamate) chiral stationary phase, 2505–2516
 - of derivatives of phenylcarbamic acid using π -complex stationary phase in, 2331–2350



- [High performance liquid chromatography (HPLC)]
of organophosphonate
derivatives on amylose

phenylcarbamate) chiral
stationary phase,
3075–3084
- for impurity and quantitative
analysis of etoricoxib,
2551–2566
- ion pair, for simultaneous
measurement of creatinine
and purine derivatives
in ruminant's urine,
2961–2968
- with mass spectrometry, for
determination of radiolysis
products in zirconium
salt of di-(2-ethylhexyl)
phosphoric acid,
2593–2604
- micro-, for investigation of
retention behavior of her-
bicides, 3357–3369
- normal phase, for quantitative
analysis of ceramide III of
Saccharomyces cerevisiae,
617–627
- normal and reversed phase, dia-
midate-type chiral stationary
phase for chiral resolution
by, 2103–2117
- on-line detection for analysis of
water-soluble vitamins in
multivitamin tablets,
1769–1786
- optimization of ASE conditions
for determination of
rutin and isoquercitrin in
Sambucus nigra L.,
2381–2397
- optimized analysis of aromatic
diamidines by, 3511–3525
- with a photodiode array detector,
for simultaneous analysis
of candesartan cilexetil and
hydrochlorothiazide in
human plasma and dosage
forms, 2581–2591
- with photodiode array and
mass spectrometry, for
differential metabolic
profiling of wild type
and transgenic tobacco
plants constitutively
producing salicylic acid,
369–383
- preparation conditions of silica
monoliths for, 2881–2986
- for quantitation of phytochelatins
in plant tissue extracts,
3463–3473
- for quantitative determination of
lanatosides in the hybrid
Digitalis ambigua ×
Digitalis lanata leaves,
2013–2021
- retention time, prediction using
multiple linear regression,
2987–3002
- robotic sample preparation and
determination of the major
component of ivermectin
in human plasma by,
895–910
- robustness, evaluation with
experimental design,
2975–2985
- semi-preparative, chromato-
graphic models as tools for
scale-up of isolation of
natural products by,
177–193



- silver ion, cyclohexanediol fatty acid diesters as model compounds for mechanistic studies in, 1905–1912
- for simultaneous determination of piperacillin sodium and sulbactam sodium, 665–676
- for simultaneous separation and determination of bioactive components in traditional Chinese medicinal formula guanxin II, 469–482
- with SPE, for determination of lomefloxacin in seminal plasma, 2053–2063
- with tandem mass spectrometry and benzoyl chloride derivatization, for determination of propylene glycol, 3413–3431
- toxic/harmful solvents-free technique for determination of sulfonamides in meat, 2931–2939
- with UV detection
- for the determination of venlafaxine and O-desmethylvenlafaxine in human plasma, 1297–1313
 - for quantitative analysis of kielcorins in biomimetic synthesis, 29–41
 - for simultaneous determination of tyrosol and oleuropein in plasma, 599–616
- High performance size exclusion chromatography (HPSEC), for determination of the cleavage position of a protein in enzymatic hydrolysis, 1787–1796
- High performance thin-layer chromatography (HPTLC)
- with densitometry, for assay of caffeine and acetaminophen in multicomponent extra strength analgesic tablets, 3453–3462
 - for detection of ionophore antibiotics in feeds and animal tissues, 147–156
 - for determination of beta-methasone in tablets, 2709–2717
 - for determination of famotidine in acid reduction tablets, 2719–2727
 - for determination of rosiglitazone in pharmaceutical formulations, 3307–3314
 - and FTIR, for identification of phthalazine derivatives, 2687–2696
 - for quantification of the sweetener Sucralose, 2729–2739
- High throughput organic synthesis (HTOS) libraries, purification by supercritical fluid chromatography, 333–354
- High-low chromatography, for quantitation of propyl gallate in an active pharmaceutical ingredient, 1743–1753
- High-speed countercurrent chromatography (HSCCC) coupled with ELSD for separation of dammarane-saponins from notoginseng, 1579–1591



- [High-speed countercurrent chromatography (HSCCC)]
for preparation of mevinolinic acid from *Monascus purpurces*, 3085–3092
for separation of high purity przewaquinone A, 1267–1274
spiral disk assembly for, 1355–1372
- HIV protease inhibitor, RP-HPLC separation from its piperazine diastereomer, 3343–3355
- Honey
determination of furosine in, 317–326
French, HPLC determination of sulfathiazole in, 953–961
HPLC determination of low molecular weight organic acids in, 1231–1253
- HPCE. *See* High performance capillary electrophoresis
- HPCCP. *See* High performance centrifugal partition chromatography
- HPLC. *See* High performance liquid chromatography
- HPLC-UV-APCI-MS, for separation and characterization of chlorophyll degradation products in silkworm, 3183–3197
- HPLC with a photodiode array detector, 2581–2591
- HPSEC. *See* High performance size exclusion chromatography
- HPTLC. *See* High performance thin-layer chromatography
- HSCCC. *See* High-speed countercurrent chromatography
- HTOS. *See* High throughput organic synthesis
- Hydrochlorothiazide, simultaneous analysis with candesartan cilexetil in human plasma and dosage forms using HPLC with a photodiode array detector, 2581–2591
- Hydrocortisone and associated compounds, MEKC determination in pharmaceutical preparations, 1011–1025
- Hydrodynamic relaxation efficiency, in frit inlet asymmetrical flow field-flow fractionation, effect of inlet frit lengths on, 2369–2379
- Hydrophilic interaction chromatography, for determination of piperazine in pharmaceutical drug substances, 3093–3104
- R,S*-Hydroxypropyl ether β -cyclodextrin column, for RP-HPLC separation and determination of positional isomeric mono- and di-substituted anilines and phenols, 1–15
- 8-Hydroxyquinoline, immobilized on silica gel phase for separation and pre-concentration of chromium (III) from water samples, 3045–3056
- 8-Hydroxyquinoline derivative, HPLC determination of aluminum as, 273–283
- Hyperbranched aliphatic polyesters, and their trimethylsilylated derivatives, characterization by GPC-viscometry, 207–230
- Hyperici* herba, supercritical fluid extraction of quercetin and rutin from, 2517–2533



I

- ICP-MS. *See* Inductively coupled plasma mass spectrometry
- Immiscible liquids, influence of gravity field on the interface of, 1521–1535
- Inductively coupled plasma mass spectrometry (ICP-MS), for determination of plutonium in soil after CCC preseparation, 1593–1607
- Inlet frit lengths, effect on the hydrodynamic relaxation efficiency in frit inlet asymmetrical flow field-flow fractionation, 2369–2379
- Inorganic anions, application of alkylamide phase to the separation by RP-HPLC, 1197–1206
- Inorganic ions, retention on silica-based and polymer-based ion exchange columns, 751–761
- Intermediate polarity resins, for adsorption separation for the extracts from *Ginkgo biloba* leaves, 2941–2960
- Intermolecular interaction, thermodynamic approaches in liquid chromatography, 2809–2838
- International Conference on Counter-current Chromatography, trends in research and applications, 1335–1344
- Ion exchange chromatography with ELSD, for inorganic anion analysis, 3117–3128
- with ESI-MS/photodiode array detection for analysis of stibonic acids, 2041–2051
- Ion exchange columns, silica based and polymer based, comparative study of retention of glucosinolates and inorganic ions on, 751–761
- Ion pairing agents, effect in the retention profile of zwitterionic cephalosporins, 937–952
- Ion-exclusion columns, series coupled, for HPLC determination of low molecular weight organic acids in honey, 1231–1253
- Ionic liquids, room temperature, for a new class of solvents for CCC, 1493–1508
- Ionophore antibiotics, HPTLC detection in feeds and animal tissues, 147–156
- Ion-pair HPLC, for simultaneous measurement of creatinine and purine derivatives in ruminant's urine, 2961–2968
- Ion-pair RP-HPLC, for determination of aromatic amine isomers, 85–98
- Isoflavones, soybean, CCC separation by different types of coil planet centrifuges with two-phase solvent systems, 3497–3509
- Isoquercitrin, optimization of ASE conditions for HPLC determination in *Sambucus nigra* L., 2381–2397
- Isoquinoline alkaloids, optimized separation in *Thalictrum* herbal medicine with microemulsion electrokinetic chromatography, 1719–1730



Ivermectin, robotic sample preparation for HPLC determination of the major component in human plasma, 895–910

J

J-type centrifuge extra-coil volume, determination using stationary phase retentions at differing flow rates, 1417–1430

K

Kava lactones, isolation and purification by high performance centrifugal partition chromatography, 3069–3074

Kielcorins, quantitative analysis in biomimetic synthesis by HPLC with UV detection, 29–41

L

Lamiaceae family plants, HPLC determination of phenolic acids in, 443–453

Lanatosides, HPLC quantitative determination in the hybrid *Digitalis ambigua* × *Digitalis lanata* leaves, 2013–2021

LC-MS/MS. *See* Liquid chromatography-mass spectrometry/mass spectrometry

Ligand exchange, for determination of mercury(II) ion, 2797–2808

Linear solvation energy relationships, for evaluation of octa-

decene-polybutadiene-coated zirconia stationary phase, 723–736

Lipophilicity, TLC determination for barbiturates, 3277–3287 for bile acids, 2741–2750

Liquid chromatography/atmospheric pressure chemical ionization-ion trap mass spectrometry, for analysis of urushiols, 17–28

Liquid chromatography/atmospheric pressure chemical ionization/mass spectrometry, of stability of ¹³C-urea/PEG capsules, 1275–1286

Liquid chromatography/electrospray-mass spectrometry, for determination of major curcuminoids in foods, 53–62

Liquid chromatography/mass spectrometry application of a polymeric SPE for analysis of sulfonamides in milk by, 1149–1159 for determination of flunixin and tiamulin hydrogen fumarate in meat and toltrazuril and the metabolite toltrazurilsufon in meat and eggs, 791–801

open access, critical considerations for high reliability, 2839–2859

for routine proteome analyses, 3331–3342

Liquid chromatography-mass spectrometry/mass spectrometry



- (LC-MS/MS), for determination of carbamazepine, pindolol, and theophylline in human serum, 1937-1957
- Liquid chromatography-nuclear magnetic resonance (book review), 835-837
- Liquorice tablets, HPCE quantitative determination of glycyrrhizic acid, glycyrrhetic acid, morphine, and sodium benzoate in, 43-51
- Lomefloxacin, determination in seminal plasma by HPLC with SPE, 2053-2063
- Lutein, CCC separation from a crude extract of marigold flower petals, 1659-1669
- M**
- Magnesium (II), HPCPC separation with organophosphorus compounds, 629-640
- Malic acid, influence on the mobility of metal ions, 1969-1976
- MALLS. *See* Multiangle laser light scattering
- Maltodextrin, capillary electrophoresis of molecular association with glutenins, 2465-2479
- Marigold flower petals, CCC separation of lutein from a crude extract of, 1659-1669
- Mass spectrometry
determination of radiolysis products in zirconium salt of di-(2-ethylhexyl) phosphoric acid by HPLC with, 2593-2604
- a small coil-volume CCC instrument for direct interfacing with, 1345-1354
- Matrix effect, correction in the determination of benzoyl-phenylurea insecticides in vegetables, 297-316
- Medium pressure solid-liquid extraction, compared with rotation planar extraction of *Ficus* leaves, 3289-3305
- MEKC. *See* Micellar electrokinetic capillary chromatography
- Mercury(II), selective determination by solvent extraction with *N*-(dithiocarboxy)sarcosine, diammonium salt, followed by ligand exchange and RP-HPLC with photometric detection, 2797-2808
- Metabolic profiling, differential, by HPLC with photodiode array and mass spectrometry of wild type and transgenic tobacco plants constitutively producing salicylic acid, 369-383
- Metal ions, influence of malic acid, phosphate ion, and urea on the mobility of, 1969-1976
- Metformin, micellar liquid chromatographic simultaneous determination in its multi-component dosage forms with glipizide and gliclazide, 1117-1133
- Methylxanthines, simultaneous determination in human plasma by capillary electrophoresis, 1913-1925



- Mevinolinic acid, preparation from *Monascus purpurces* using HSCCC, 3085–3092
- Mexiletine hydrochloride, modified RP-HPLC analysis of, 1287–1296
- Micellar electrokinetic capillary chromatography (MEKC) for determination of biotin in pharmaceutical preparations, 195–205
- for determination of hydrocortisone and associated compounds in pharmaceutical preparations, 1011–1025
- SPE for determination of dimethoate in environmental water and soil samples by, 545–557
- Micellar liquid chromatography, for simultaneous determination of metformin in its multicomponent dosage forms with glipizide and gliclazide, 1117–1133
- Micro-channel thermal field-flow fractionation, high-speed analysis of colloidal particles by, 849–869
- Microemulsion electrokinetic chromatography, for optimized separation of isoquinoline alkaloids in *Thalictrum* herbal medicine, 1719–1730
- Microsomal samples, porcine, determination of acetaminophen and phenacetin in, 925–935
- Microwave-assisted extraction, of polychlorinated biphenyls and polychlorinated dibenzo-*p*-dioxins from fly ash and sea sediments, 803–818
- Milk
 application of a polymeric SPE for liquid chromatography/mass spectrometric analysis of sulfonamides in, 1149–1159
 powdered, monitoring of the addition of rennet whey to, 99–107
- Mobile phases, automatic selection of, 1255–1266
- Modern extraction techniques, comparison for analysis of soil contaminated with fuel and crude oil spills, 247–259
- Mometasone furoate, TLC densitometric determination in topical preparations, 109–117
- Monascus purpurces*, HSCCC preparation of mevinolinic acid from, 3085–3092
- Monosulfides, investigation with complexation TLC, 2651–2661
- Monoterpenes, evaluation of cyclodextrins as chiral selectors in separation by capillary liquid chromatography and capillary electrophoresis, 505–516
- Morphine, HPCE quantitative determination in compound liquorice tablets, 43–51
- Moving bed system, eight column, continuous separation of 1-1'-bi-2-naphthol enantiomers by, 1853–1875
- Multiangle laser light scattering (MALLS), combined with flow field-flow fractionation for characterization of



polymeric wheat proteins,
2771–2781

Multimodal elution, effects on
enantioselectivity using
columns based on polysac-
charide, 2083–2101

Multiple linear regression, for
prediction of HPLC retention
time, 2987–3002

N

Naproxen sodium, flow injection
analysis using UV detection
determination in pharmaceu-
tical formulations, 401–408

Nateglinide, reversed-phase liquid
chromatographic determina-
tion of the *L*-enantiomer
in a bulk drug substance,
1839–1845

National certified reference materi-
als, of active compounds,
preparation from natural
products by CCC,
1565–1577

Natural products, chromatographic
models as tools for scale-up
of isolation by semi-
preparative HPLC, 177–193

Nonionic surfactants, TLC study of
binding to the corn protein
zein, 2751–2761

Non-segmented sorbent bed, for
operating segmentation for
single and multi-channel
OPLC separation, 2611–2627

Notoginseng, HSCCC coupled
with ELSD for separation of
dammarane-saponins from,
1579–1591

O

Ochratoxin A, HPLC determination
in cocoa powder, 585–598
in wine samples, 119–133

Octadecene–polybutadiene-coated
zirconia stationary phase,
evaluation based on linear
solvation energy relation-
ships, 723–736

Octanol-water partition coefficients,
RP-HPLC estimation for
polycyclic aromatic
hydrocarbons, 135–146

Olefin hydrosilation, on a hydride
intermediate for adamantyl
modified silica, 1169–1195

Oleuropein, simultaneous determi-
nation with tyrosol in plasma
by HPLC with UV detection,
599–616

Oligodeoxynucleotide–doxorubicin
conjugate, RP-HPLC study of
stability in vitro and pharma-
cokinetics in vivo, 3105–3115

On-line matrix elimination, for
analysis of *R*- and *S*-1-phenyl-
1-butanol in biphasic
Saccharomyces cerevisiae
mediated culture, 3249–3264

Open access liquid chromatogra-
phy/mass spectrometry,
critical considerations for
high reliability, 2839–2859

Open-tubular capillary column,
ultrashort, with modified
silica-gel thin layer for
capillary electrochromato-
graphy, 697–707

Operating conditions, on scale-up,
effect on resolution in CCC,
1475–1491



- OPLC. *See* Overpressurized layer chromatography
- Organic acids
 HPLC determination in traditional balsamic vinegar, 2177–2187
 low molecular weight, HPLC determination in honey, 1231–1253
- Organic extractables, from packaging materials, 2449–2464
- Organic modifiers, influence on the separation of carboxylic acids using co-electroosmotic flow capillary electrophoresis, 455–468
- Organic phase
 poly(4-vinylpyridine) as for RP-HPLC, unique selectivity for polycyclic aromatic hydrocarbons, 2491–2503
 retention in CCC, 1537–1550
- Organic polymer additives, chromatographic methods used in the identification of, 2417–2447
- Organophosphonate derivatives, enantioseparation on amylose *tris*(3,5-dimethylphenylcarbamate) chiral stationary phase by HPLC, 3075–3084
- Organophosphorus compounds, HPCPC separation of calcium (II), magnesium (II), strontium (II), and barium (II) with, 629–640
- Oroxylum indicum* seeds, CCC separation and identification of flavonoids in an extract from, 1623–1636
- Ovarian cancer cell lines, surface-enhanced laser desorption ionization time-of-flight mass spectrometry profiling of secreted proteins from, 2315–2328
- Overpressurized layer chromatography (OPLC) separation, single and multi-channel, on non-segmented sorbent bed using flowing eluent wall for operating segmentation, 2611–2627

P

- Packaging materials, organic extractables from, 2449–2464
- Peptide mass modification, integration with chromatography for quantitative proteomics, 2285–2301
- Peptides
 chromatographic fractionation techniques in the characterization of, 2195–2224
 effect of experimental parameters on the HPLC separation of, 2255–2283
- Perillyl alcohol enantiomers, high performance frontal analysis of protein binding for, 2861–2871
- Pesticides, chiral and related intermediates, direct separation on polysaccharide-based chiral stationary phases, 2873–2880
- pH, TLC study of effect on the binding of ring-substituted phenol derivatives to the corn protein zein, 2303–2313



- Pharmaceutical process impurity, selective removal using a reactive resin, 1959–1968
- Phase system properties, on scale-up, effect on resolution in CCC, 1475–1491
- Phase systems, characterization in CCC, 1551–1564
- Phenacetin, determination in porcine microsomal samples, 925–935
- Phenol derivatives, ring substituted, TLC study of the effects of pH and salts on binding to the corn protein zein, 2303–2313
- Phenolic acids, HPLC determination in *Lamiaceae* family plants, 443–453
- Phenolic compounds
chromatographic separation from rapeseed by a Sephadex LH-20 column with ethanol as the mobile phase, 2157–2165
computational chemical simulation of chromatographic retention of, 2031–2039
- Phenols, positional isomeric mono- and di-substituted, RP-HPLC separation and determination on an R,S-hydroxypropyl ether β -cyclodextrin column, 1–15
- Phenyl isothiocyanated amino acids, HPLC enantiomeric resolution on teicoplanin-bonded phase using an acetonitrile-based mobile phase, 3475–3495
- Phenylacrylates, QSPR prediction of partition properties for, 2065–2082
- 1-Phenyl-1-butanol, *R*- and *S*-, column switching technique, assisted on-line matrix elimination, and chiral analysis in biphasic *Saccharomyces cerevisiae* mediated culture, 3249–3264
- Phenylcarbamic acid derivatives, HPLC enantioseparation using π -complex stationary phase in HPLC, 2331–2350
- Phosphate ion, influence on the mobility of metal ions, 1969–1976
- Phospholipids, separation with TC-CCC, 1135–1147
- Photodiode array detection
for analysis of safflower yellow in food, 1207–1217
HPLC with for simultaneous analysis of candesartan cilexetil and hydrochlorothiazide in human plasma and dosage forms, 2581–2591
- Photometric detection
determination of mercury(II) ion by RP-HPLC with, 2797–2808
for selective determination of tin (II) ion in water, 3129–3139
- Phthalazine derivatives, identification by HPTLC-FTIR, 2687–2696
- o*-Phthaldialdehyde-mercaptoethanol derivatization, validation of HPLC analysis of aspartate and glutamate neurotransmitters following, 1797–1808



- Phytochelatins, HPLC quantitation in plant tissue extracts, 3463–3473
- Phytoecdysteroids, TLC of, 2629–2649
- Pindolol, LC-MS/MS determination in human serum, 1937–1957
- Piperacillin sodium, HPLC simultaneous determination with sulbactam sodium, 665–676
- Piperazine
determination in pharmaceutical drug substances using hydrophilic interaction chromatography and ELSD, 3093–3104
diastereomer, RP-HPLC separation of HIV protease inhibitor from, 3343–3355
- Plutonium, determination in soil by ICP-MS after CCC pre-separation, 1593–1607
- Polyamines, capillary electrophoresis with universal indirect chemiluminescence detection using cobalt (II) as a probe ion, 355–367
- Polychlorinated biphenyls and polychlorinated dibenzo-*p*-dioxins, microwave-assisted extraction from fly ash and sea sediments, 803–818
- Polycyclic aromatic hydrocarbons
RP-HPLC estimation of octanol-water partition coefficients for, 135–146
unique selectivity of poly(4-vinylpyridine) as an organic phase for RP-HPLC for, 2491–2503
- Poly(ethyleneimine) spherical particles, cross-linked, for removal of DNA from a protein solution, 231–246
- Poly(glycidyl methacrylate-co-ethylenedimethacrylate) particles, monodisperse, for preparation of strong cation exchange packings, 963–976
- Polysaccharides
direct separation of chiral pesticides and related intermediates on chiral stationary phases based on, 2873–2880
multimodal elution effects on enantioselectivity using columns based on, 2083–2101
purification from *Achyranthes bidentata* by CCC with an aqueous two-phase system using a cross-axis coil planet centrifuge, 1895–1903
- Polystyrene latex particles, retention behavior on different channel walls in flow field-flow fractionation, 3003–3035
- Poly(4-vinylpyridine), as an organic phase for RP-HPLC, unique selectivity for polycyclic aromatic hydrocarbons, 2491–2503
- Polyvinylpyridines, salt effect on separation by thermal field-flow fractionation, 165–175
- Porous graphitic carbon column, for HPLC determination of haloperidol and its degradation products in pharmaceuticals, 261–271



- Porphyrin, RP-HPLC speciation of complexes with thulium, 911–923
- Preparative chiral chromatography, of *trans*-stilbene oxide using cellulose *tris*(phenyl-carbamate) (Chiralcel[®] OC) stationary phase, 3235–3248
- Preparative chiral separations, development using an intelligent chiral resolution system, 529–543
- Proinsulin, recombinant human, renaturation with simultaneous purification from *E. coli* with high performance hydrophobic interaction chromatography, 683–695
- Propolis extract, HPLC analysis of, 2399–2409
- Propyl gallate, quantitation in an active pharmaceutical ingredient using high-low chromatography, 1743–1753
- Propylene glycol, HPLC with tandem mass spectrometry and benzoyl chloride derivatization for determination of, 3413–3431
- Protein binding, high performance frontal analysis for perillyl alcohol enantiomers, 2861–2871
- Proteins
chromatographic fractionation techniques in the characterization of, 2195–2224
collagen-like, roles within the proteome, 2225–2254
effect of experimental parameters on the HPLC separation of, 2255–2283
secreted, profiling from human ovarian cancer cell lines by surface-enhanced laser desorption ionization time-of-flight mass spectrometry, 2315–2328
- Proteome
roles of collagen and collagen-like proteins in, 2225–2254
routine analyses by liquid chromatography/mass spectrometry, 3331–3342
- Proteomics
(book review), 839–841
quantitative, integration of chromatography and peptide mass modification for, 2285–2301
- Przewaquinone A, high purity, separation by HSCCC, 1267–1274
- Psoralen derivatives, determination of the association constants with modified β -cyclodextrin, 871–882
- Purine derivatives, simultaneous measurement with creatinine in ruminant's urine using ion-pair HPLC, 2961–2968
- N*-(1-Pyrenyl)maleimide, HPLC analysis of buccillamine in human blood by derivatization with, 3265–3275
- Q**
- QSPR, for prediction of partition properties for phenylacrylates, 2065–2082



Quercetin, supercritical fluid
extraction from *Hyperici*
herba, 2517–2533

R

- Ractopamine stereoisomers,
capillary electrophoretic
separation and quantitation
using cyclodextrins as chiral
additives, 1677–1693
- Radiolysis products, determination
in zirconium salt of di-(2-
ethylhexyl) phosphoric acid
by HPLC/mass spectrometry,
2593–2604
- Rapeseed, chromatographic separa-
tion of phenolic compounds
by a Sephadex LH-20 column
with ethanol as the mobile
phase, 2157–2165
- Rate constants, separate determina-
tion in a chromatographic
column and eluent using
empirical peak shape
equations, 285–296
- Reactive resin, for selective removal
of pharmaceutical process
impurity, 1959–1968
- Relative retention times, of mole-
cular species of acylglycerols,
phosphatidylcholines, and
phosphatidylethanolamines
containing ricinoleate in
RP-HPLC, 1051–1058
- Renaturation, with simultaneous
purification of recombinant
human proinsulin from
E. coli with high
performance hydrophobic
interaction chromatography,
683–695
- Retention behavior
of herbicides in micro-HPLC
with temperature changes,
3357–3369
of polystyrene latex particles and
gold colloids on different
channel walls in flow field-
flow fractionation,
3003–3035
thermodynamic approaches in
liquid chromatography,
2809–2838
- Retinoic acid, simultaneous
determination with retinol
and retinyl palmitate in ram
plasma, 559–572
- Reversed-phase high performance
liquid chromatography
(RP-HPLC)
for analysis of endosulfan and
its major metabolites,
1069–1082
for analysis of mexiletine
hydrochloride,
1287–1296
application of alkylamide
phase to the separation
of inorganic anions by,
1197–1206
of aristolochic acid and aristo-
lactam analogues in
traditional Chinese
medicine, 3057–3067
with CoulArray multi-electrode
chemical detector system,
for catecholamine
quantification, 3433–3451
for determination of caseinoma-
cropeptide and monitoring
of the addition of rennet
whey to powdered milk,
99–107



- for determination of sibutramine hydrochloride in the presence of its oxidatively induced degradation products, 977–986
- for determination of valacyclovir in tablets and human serum, 1755–1767
- for drug dissolution studies and determination of deflazacort in pharmaceutical formulations and human serum, 2141–2156
- for estimation of amiodarone hydrochloride, 1315–1326
- for estimation of bile salts in complex tanning brines, 1041–1050
- for estimation of octanol-water partition coefficients for polycyclic aromatic hydrocarbons, 135–146
- ion pair, for determination of aromatic amine isomers, 85–98
- optimization for drug control analysis, 3401–3412
- with photometric detection
- determination of mercury(II) ion by, 2797–2808
 - for selective determination of tin (II) ion in water, 3129–3139
- poly(4-vinylpyridine) as an organic phase for, unique selectivity for polycyclic aromatic hydrocarbons, 2491–2503
- relative retention times of molecular species of acylglycerols, phosphatidylcho-
- lines, and phosphatidylethanolamines containing ricinoleate in, 1051–1058
- for separation and determination of positional isomeric mono- and di-substituted anilines and phenols on an *R,S*-hydroxypropyl ether β -cyclodextrin column, 1–15 for separation of HIV protease inhibitor from its piperazine diastereomer, 3343–3355
- for speciation of thulium porphyrin complexes, 911–923
- for study of the stability of oligodeoxynucleotide–doxorubicin conjugate in vitro and pharmacokinetics in vivo, 3105–3115
- with UV detection, for determination of fat soluble vitamins, 641–663
- Reversed-phase liquid chromatography
- chiral, determination of the *L*-enantiomer of nateglinide in a bulk drug substance by, 1839–1845
 - thermodynamics during the biopolymer retention process in, 3199–3215
- Ricinoleate, relative retention times of molecular species of acylglycerols, phosphatidylcholines, and phosphatidylethanolamines containing in RP-HPLC, 1051–1058



- Robotic sample preparation, for HPLC determination of the major component of ivermectin in human plasma, 895–910
- Rosiglitazone, rapid HPTLC determination in pharmaceutical formulations, 3307–3314
- Rotation planar extraction, compared with medium pressure solid-liquid extraction of *Ficus* leaves, 3289–3305
- RP-HPLC. *See* Reversed-phase high performance liquid chromatography
- Rutin
 optimization of ASE conditions for HPLC determination in *Sambucus nigra* L., 2381–2397
 supercritical fluid extraction from *Hyperici* herba, 2517–2533
- S**
- Saccharomyces cerevisiae*, normal phase HPLC quantitative analysis of ceramide III of, 617–627
- Saccharomyces cerevisiae* mediated culture, biphasic, column switching technique, assisted on-line matrix elimination, and chiral analysis of *R*- and *S*-1-phenyl-1-butanol in, 3249–3264
- Safflower yellow, analysis in food by liquid chromatography with photodiode array and electrospray mass spectrometric detection, 1207–1217
- Salicylidenamino-2-thiophenol, selective determination of tin (II) ion in water by solvent extraction with, 3129–3139
- Salt effect, on separation of polyvinylpyridines by thermal field-flow fractionation, 165–175
- Salts, TLC study of effect on the binding of ring-substituted phenol derivatives to the corn protein zein, 2303–2313
- Sambucus nigra* L., optimization of ASE conditions for HPLC determination of rutin and isoquercitrin in, 2381–2397
- Scutellarin, determination in *Scutellaria barbata* extract by liquid chromatography with electrochemical detection, 2133–2140
- Sea sediments, microwave-assisted extraction of polychlorinated biphenyls and polychlorinated dibenzo-*p*-dioxins from, 803–818
- SEC. *See* Size exclusion chromatography
- Separation factor, UNIFAC estimation for analogues in CCC, 1431–1447
- Sephadex LH-20, for chromatographic separation of phenolic compounds from rapeseed with ethanol as the mobile phase, 2157–2165
- Sibutramine hydrochloride, RP-HPLC determination in the presence of its oxidatively induced degradation products, 977–986



- Sildenafil citrate, monolithic HPLC column for determination in pharmaceutical preparations, 2897–2908
- Silica
- adamantyl modified, via olefin hydrosilation on a hydride intermediate, 1169–1195
 - monoliths, preparation conditions for HPLC, 2881–2986
 - TLC of 2,3,4-trisubstituted tetrahydroisoquinolinones on, 1255–1266
- Silica gel phases-immobilized-8-hydroxyquinoline, for separation and pre-concentration of chromium (III) from water samples, 3045–3056
- Silica-gel thin layer, capillary electrochromatography with ultrashort open-tubular capillary column with, 697–707
- Silver ion HPLC, cyclohexanediol fatty acid diesters as model compounds for mechanistic studies in, 1905–1912
- Single molecule detection (book review), 837–839
- Size exclusion chromatography (SEC), HPLC–electrospray mass spectrometry with for determination of acrylamide in fried foods, 1877–1884
- Slalom chromatography, of viscosity-temperature dependence on DNA stretching, 883–893
- Sodium benzoate, HPCE quantitative determination in compound liquorice tablets, 43–51
- Soil, contaminated with fuel and crude oil spills, comparison of modern extraction techniques for analysis of, 247–259
- Soil environmental samples
- HPLC determination of diazinon and fenitrothion in, 483–497
 - SPE for the determination of dimethoate by MEKC, 545–557
- Solid-phase extraction (SPE)
- for the determination of dimethoate in environmental water and soil samples by MEKC, 545–557
 - determination of lomefloxacin in seminal plasma by HPLC with, 2053–2063
 - of fluoroquinolone derivatives from a biological matrix for biodetermination by liquid chromatography, 1731–1741
 - polymeric, for the analysis of sulfonamides in milk by liquid chromatography/mass spectrometry, 1149–1159
 - for triazines, 781–790
- Solids, retention in rotating coiled columns, 1649–1657
- Solvent extraction
- with *N*-(dithiocarboxy)sarcosine for selective determination of mercury(II), 2797–2808
 - with salicylidenamino-2-thiophenol for selective determination of tin (II) ion in water, 3129–3139



- Solvent systems
 aqueous two phase, selection in CCC, 1509–1520
 property calculation and prediction for selection in CCC, 1397–1415
- Solvents, for CCC, 1493–1508
- Soybean isoflavones, CCC
 separation by different types of coil planet centrifuges with two-phase solvent systems, 3497–3509
- SPE. *See* Solid-phase extraction
- Spiral disk assembly, for HSCCC, 1355–1372
- Stability
 of chloramphenicol residues in food samples during storage, 2535–2549
 of solutions containing hydromorphone, ketorolac, haloperidol, midazolam, famotidine, metoclopramide, dimenhydrinate, and scopolamine, HPLC with diode array detection for study of, 2909–2929
 of ¹³C-urea/PEG capsules, liquid chromatography/atmospheric pressure chemical ionization/mass spectrometry of, 1275–1286
- Stationary phases
 amino, for study of the vancomycin dimerization dependence on solute enantioselectivity, 1027–1039
 cellulose *tris*(phenylcarbamate) (Chiralcel[®] OC), for

- preparative chiral chromatography of *trans*-stilbene oxide, 3235–3248
- chiral
 amylose *tris*(3,5-dimethylphenylcarbamate), for HPLC enantioseparation of organophosphonate derivatives, 3075–3084
 based on (+)-(18-crown-6)-2,3,11,12-tetracarboxylic acid, effect of the tethering groups on the liquid chromatographic resolution of β -amino acids, 1083–1094
 cellulose *tris*-(3,5-dimethylphenylcarbamate), for HPLC enantioseparation of chiral tetrahedral heterometal clusters, 2505–2516
 on 4-chloro-3,5-dinitrobenzoic acid amides of α,β -aminoalcohols and α -arylethylamines for enantiomer separation and molecular recognition, 63–83
 diamide type, for chiral resolution by normal and reversed-phase HPLC, 2103–2117
 polysaccharide based, for preparation and direct separation of chiral pesticides and related intermediates, 2873–2880
 crown ether, enantiomeric separation of *trans*-2-amino-



- cyclohexanol using ELSD, 425–442
- octadecene–polybutadiene-coated zirconia, evaluation based on linear solvation energy relationships, 723–736
- π -complex, use in HPLC for enantioseparation of derivatives of phenylcarbamic acid, 2331–2350
- retention in CCC, 1373–1396
- retentions at differing flow rates for determination of J-type centrifuge extra-coil volume, 1417–1430
- Stibonic acids, analysis by ion exchange chromatography with ESI-MS/photodiode array detection, 2041–2051
- trans*-Stilbene oxide, preparative chiral chromatography using cellulose *tris*(phenylcarbamate) (Chiralcel[®] OC) stationary phase, 3235–3248
- Strontium (II), HPCPC separation with organophosphorus compounds, 629–640
- Subproteomics, in analytical chemistry, chromatographic fractionation techniques in the characterization of proteins and peptides, 2195–2224
- Sucralose[®] sweetener, HPTLC quantification of, 2729–2739
- Sucrose, effect as a co-enhancer association agent, 871–882
- Sulbactam pivoxil, simultaneous determination with amoxicillin in a combination formulation, 1927–1936
- Sulbactam sodium, HPLC simultaneous determination with piperacillin sodium, 665–676
- Sulfathiazole, HPLC determination in French honeys, 953–961
- Sulfonamides
- application of a polymeric SPE for analysis in milk by liquid chromatography/mass spectrometry, 1149–1159
 - toxic/harmful solvents-free technique for HPLC determination in meat, 2931–2939
- Supercritical fluid chromatography, for purification of HTOS libraries, 333–354
- Supercritical fluid extraction, of quercetin and rutin from *Hyperici* herba, 2517–2533
- Surface-enhanced laser desorption ionization time-of-flight mass spectrometry, for profiling of secreted proteins from human ovarian cancer cell lines, 2315–2328
- Swertia przewalskii pissjauk* extract, capillary electrophoretic analysis of pharmacologically active xanthone compounds from, 1219–1229

T

- Tamsulosin, automated HPLC of dissolution from pharmaceutical dosage forms, 1109–1116
- Tandem mass spectrometry, determination of propylene glycol by HPLC with, 3413–3431



- Tanning brines, complex, RP-HPLC estimation of bile salts in, 1041–1050
- TC-CCC. *See* Toroidal-coil counter-current chromatography
- Teicoplanin-bonded phase, for HPLC enantiomeric resolution of phenyl isothiocyanated amino acids using an acetonitrile-based mobile phase, 3475–3495
- Tetrahydroisoquinolinones, 2,3,4-trisubstituted, TLC on silica of, 1255–1266
- Thalictrum* herbal medicine, microemulsion electrokinetic chromatography for optimized separation of isoquinoline alkaloids in, 1719–1730
- Theophylline, LC-MS/MS determination in human serum, 1937–1957
- Thermal field-flow fractionation micro-channel, high-speed analysis of colloidal particles by, 849–869
- salt effect on separation of polyvinylpyridines by, 165–175
- Thermodynamics
- during the biopolymer retention process in reversed-phase liquid chromatography, 3199–3215
 - of intermolecular interaction and retention behavior in liquid chromatography, 2809–2838
- 1-(2-Thienylethyl)-4-imidazoline-2-thione, HPLC test for impurities in, 573–583
- Thin-layer chromatography (TLC)
- for analysis of corrinoid compounds in fish sauce, 2703–2707
 - for analysis of free pool amino acids in cercariae, rediae, encysted metacercariae, and excysted metacercariae of *Echinostoma caproni*, 2697–2702
 - of the binding of nonionic surfactants to the corn protein zein, 2751–2761
 - complexation, for investigation of monosulfides, 2651–2661
 - densitometry, for determination of mometasone furoate in topical preparations, 109–117
 - for determination of lipophilicity of barbiturates, 3277–3287
 - of bile acids, 2741–2750
 - for investigation of a homologous series of fatty acids, 2663–2671
 - of phytoecdysteroids, 2629–2649
 - separation of bile acids by, 1095–1108
 - for separation of fluoroquinolones, 2673–2685
 - on silica of 2,3,4-trisubstituted tetrahydroisoquinolinones, 1255–1266
 - for simultaneous analysis of carotenoid colorings in foods, 819–832
 - for study of effects of pH and salts on the binding of ring-substituted phenol



- derivatives to the corn protein zein, 2303–2313
- Thulium porphyrin complexes, RP-HPLC speciation of, 911–923
- Tiamulin hydrogen fumarate, determination in meat using liquid chromatography/mass spectrometry, 791–801
- Tin (II) ion, selective determination in water by solvent extraction with salicylidenamino-2-thiophenol followed by RP-HPLC with photometric detection, 3129–3139
- TLC. *See* Thin-layer chromatography
- Tobacco plants, constitutively producing salicylic acid, differential metabolic profiling by HPLC with photodiode array and mass spectrometry of 369–383
- Toltrazuril and metabolite toltrazurilsulfon, determination in meat and eggs using liquid chromatography/mass spectrometry, 791–801
- Topical preparations, TLC densitometric determination of mometasone furoate in, 109–117
- Toroidal-coil countercurrent chromatography (TC-CCC) for analysis of acidic herbicides in water, 1637–1648
for separation of phospholipids and glycolipids, 1135–1147
- Total lipid extract, of bovine brain, dual parallel liquid chromatography/dual mass spectrometry of, 3147–3181
- Toxic/harmful solvents-free technique, for HPLC determination of sulfonamides in meat, 2931–2939
- Traditional Chinese medicine guanxin II, HPLC simultaneous separation and determination of bioactive components in, 469–482
RP-HPLC resolution and quantitative determination of aristolochic acid and aristolactam analogues in, 3057–3067
- Triazines, SPE of, 781–790
- Trifluoroacetylated hydrazino pharmaceutical intermediate, controlling the structure for improved HPLC and gas chromatography analyses, 3371–3386
- Tyrosol, simultaneous determination with oleuropein in plasma by HPLC with UV detection, 599–616
- U**
- Ultraviolet (UV) detection determination of venlafaxine and O-desmethylvenlafaxine in human plasma by HPLC with, 1297–1313
indirect, new electrolyte composition for determination of anions by capillary electrophoresis with, 709–722



- [Ultraviolet (UV) detection]
 quantitative analysis of kielcor-
 ins in biomimetic synthesis
 by HPLC with, 29–41
 RP-HPLC with for determina-
 tion of fat soluble
 vitamins, 641–663
 simultaneous determination of
 tyrosol and oleuropein in
 plasma by HPLC with,
 599–616
 use in flow injection analysis for
 determination of naproxen
 sodium in pharmaceutical
 formulations, 401–408
- UNIFAC, for estimation of separa-
 tion factor for analogues in
 CCC, 1431–1447
- Universal indirect chemilumines-
 cence detection, using cobalt
 (II) as a probe ion for
 capillary electrophoresis of
 polyamines, 355–367
- Urea, influence on the mobility of
 metal ions, 1969–1976
- ¹³C-Urea/PEG capsules, liquid
 chromatography/atmospheric
 pressure chemical ioniza-
 tion/mass spectrometry of
 stability of, 1275–1286
- Urushiols, analysis by liquid
 chromatography/atmospheric
 pressure chemical ionization–
 ion trap mass spectrometry,
 17–28
- UV. *See* Ultraviolet
- V**
- Valacyclovir, RP-HPLC determina-
 tion in tablets and human
 serum, 1755–1767
- Vancomycin, use of an amino
 stationary phase for study of
 dimerization dependence on
 solute enantioselectivity,
 1027–1039
- Vegetables, correction of the matrix
 effect in the determination
 of benzoylphenylurea
 insecticides in, 297–316
- Venlafaxine and O-desmethylvenla-
 faxine, HPLC with UV
 detection for determination in
 human plasma, 1297–1313
- Videodensitometry, of fluorescence
 quenched zones of famoti-
 dine, 2719–2727
- Viscometry, characterization of
 hyperbranched aliphatic
 polyesters and their
 trimethylsilylated derivatives
 by GPC with, 207–230
- Viscosity behavior, abnormal SEC
 and rheology study in dilute
 oligomer solutions (erratum),
 157–158
- Viscosity-temperature dependence,
 on DNA stretching, 883–893
- Vitamin B₆, capillary zone
 electrophoresis techniques
 combined with indirect UV,
 direct UV, and mass
 spectrometric detection for
 determination in infusion
 solutions, 2783–2795
- Vitamins
 fat soluble, determination by RP-
 HPLC with UV detection,
 641–663
 water soluble, on-line detection
 for HPLC analysis in
 multivitamin tablets,
 1769–1786



W

- Water environmental samples
HPLC determination of diazinon
and fenitrothion in,
483–497
SPE for the determination of
dimethoate by MEKC,
545–557
- Wheat proteins, polymeric,
characterization by
flow field-flow
fractionation/MALLS,
2771–2781
- Wieland–Miescher ketone
and derivatives, chiral
HPLC for resolution of,
409–424
- Wine samples, HPLC deter-
mination of ochratoxin A in,
119–133

X

- Xanthone compounds, from *Swertia
przewalskii pissjauk* extract,
capillary electrophoretic
analysis of, 1219–1229
- Xanthenes, comparison of capillary
electrophoresis modes and
HPLC for the separation of,
993–1003

Z

- Zirconium salt of di-(2-ethylhexyl)
phosphoric acid, HPLC/mass
spectrometry determination
of radiolysis products in,
2593–2604
- Zwitterionic cephalosporins, effect
of ion pairing agents in the
retention profile of, 937–952

