

exactly as described in A above, with the only difference that cyclohexane was used in place of benzene. The crude oxidation product (2.3 g.), showing no absorption maximum in the ultraviolet, was dissolved in benzene and adsorbed on a column of neutral alumina (200 g.). Elution with benzene-ethyl acetate (1:1, 300 ml.) and two crystallizations of the residue (1.1 g.) from ethyl acetate gave 17 α -methylandrostan-17 β -ol-3-one (0.33 g.), m.p. and m.m.p. 190–193°. Identity was also confirmed by their infrared spectra. The column was eluted further with benzene-ethyl acetate (1:1), and the eluates were collected in 200-ml. portions. Evaporation gave a colorless oil (0.35 g.) and colorless solids (0.4 g., 0.25 g., and 0.1 g.) in that order. The last three solids were mixtures of diols containing some 17 α -methylandrostan-3 β ,17 β -diol, detected by its characteristic triple peaks at 3480, 3400, and 3210 cm.⁻¹ in the O-H stretching region of the infrared spectrum.

Benzo[*b*]thiophene Derivatives. V. Mannich Bases With Antimicrobial Activity¹

E. CAMPAIGNE, E. D. WEINBERG, G. CARLSON,^{2a}
AND E. S. NEISS^{2b}

Departments of Chemistry and Bacteriology,
Indiana University, Bloomington, Indiana 47405

Received September 11, 1964

In a continuation of our studies³ of benzo[*b*]thiophene as a source of compounds of potential biological activity, a series of β -amino ketone hydrochlorides of this heterocycle have been prepared and tested for antimicrobial activity. A wide variety of ketonic bases are reported in the literature and many of these compounds have been found to exhibit analgesic, local anesthetic, antispasmodic, and chemotherapeutic activity.^{4–10}

The Mannich reaction has been reviewed by Blicke¹¹ and more recently by Hellmann and Opitz.¹² Only one Mannich base of 3-acetylbenzo[*b*]thiophene (I) has been reported,¹³ but no biological activity evaluation was included in this report. This compound, 3-(β -dimethylaminopropionyl)benzo[*b*]thiophene hydrochloride (II), has been included in this work for comparison of its antimicrobial properties with the other members of the series described in Table I. Com-

TABLE I
MANNICH BASE HYDROCHLORIDES OF 3-ACETYL-BENZO[*b*]THIOPHENE^a

No.	Amine	M.p., °C.	Purified yield, %	Formula	Carbon, % Calc.	Found	Hydrogen, % Calc.	Found	Nitrogen, % Calc.	Found	S. aureus	MIC ^b <i>E. coli</i>	S. cerevisiae
II	Dimethyl-	173–174	65	C ₁₃ H ₁₆ ClN ₂ OS	57.85	57.56	5.99	6.20	5.19	5.55	10–35	>100	>100
III	Diethyl-	135–137	55	C ₁₅ H ₂₀ ClN ₂ OS	60.05	60.24	6.77	6.91	4.70	4.87	10–33	>100	>100
IV	Dibenzyl-	213–214	44	C ₂₅ H ₂₄ ClN ₂ OS	71.25	71.48	5.73	6.07	3.32	3.64	>100	>100	>100
V	Pyrrrolidine	198–199	67	C ₁₃ H ₁₅ ClN ₂ OS	61.40	61.03	6.14	6.11	4.74	4.75	33–100	33–100	>100
VI	Piperidine	228–229	71	C ₁₄ H ₁₈ ClN ₂ OS	61.95	61.86	6.46	6.62	4.53	4.82	33–100	>100	>100
VII	Morpholine	205–206	57	C ₁₃ H ₁₇ ClN ₂ OS	57.75	58.10	5.82	5.93	4.49	4.51	3.3–10	10–35	10–35
VIII	Hexamethylenimine	194–195	70	C ₁₅ H ₂₂ ClN ₂ OS	65.10	65.10	6.81	7.04	4.32	4.21	10–35	33–100	10–35
IX	3-Azabicyclo[3,2,2]nonane	180–190	54	C ₁₄ H ₂₁ ClN ₂ OS	65.25	64.93	6.92	7.14	4.05	4.27	10–35	>100	10–35
	Penicillin G										0.35–1.0	35–100	>100
	Tetracycline										0.55–1.0	1.0–3.3	>100
	Streptomycin										1.0–3.3	1.0–3.3	>100
	Chloramphenicol										3.3–10	3.3–10	>100

^a Compounds are all white solids. Melting points are corrected. Microanalyses were performed by Midwest Microchem, Inc., Indianapolis, Ind. ^b Expressed in μ mol. of agent for 100% inhibitory concentration. ^c Reported previously by E. F. Blicke and D. G. Sheets, *J. Am. Chem. Soc.*, **71**, 2856 (1949).

(1) Contribution No. 1248. This investigation was supported by the United States Public Health Service by Research Grants AI-04184-03 and GM-10366-02, and Fellowship No. MFM-14,696.

(2) (a) National Science Foundation High School Summer Science Institute Participant, 1964, under Grant GE-3706 to Indiana University. (b) Taken in part from the Ph.D. thesis to be submitted by E. S. Neiss, National Institutes of Health Predoctoral Fellow, 1961–1964.

(3) E. Campaigne and W. E. Kreighbaum, *J. Org. Chem.*, **26**, 359, 363, 1326, 1327 (1961).

(4) C. Mannich and U. Lammering, *Ber.*, **55**, 3510 (1922).

(5) F. F. Blicke and E. S. Blake, *J. Am. Chem. Soc.*, **52**, 235 (1930).

(6) J. J. Denton, R. J. Turner, W. B. Neier, V. A. Lawson, and H. P. Schedl, *ibid.*, **71**, 2048 (1949).

(7) G. A. Levy and H. B. Nisbet, *J. Chem. Soc.*, 1053 (1938).

(8) J. H. Burekhalter and S. H. Johnson, *J. Am. Chem. Soc.*, **73**, 4835 (1951).

(9) W. L. Nobles, *et al.*, *J. Pharm. Sci.*, **43**, 641 (1954); **43**, 644 (1954); **44**, 273 (1955); **44**, 717 (1955); **47**, 77 (1958); **51**, 878 (1962).

(10) P. Da Re, L. Verlicchi, and I. Setnikar, *J. Org. Chem.*, **25**, 1097 (1960).

(11) F. F. Blicke, *Org. Reactions*, **1**, 303 (1942).

(12) H. Hellmann and G. Opitz, "α-Aminoalkylisierung," Verlag Chemie, Weinheim, Germany, 1960.

(13) F. F. Blicke and D. G. Sheets, *J. Am. Chem. Soc.*, **71**, 2856 (1949).

pond I was prepared by the method of Farrar and Levine.¹⁴ The product, which was an azeotrope, was found by gas phase chromatography to consist of 70% of I and 30% of the 2-isomer, in close agreement with the authors. Only chromatographically pure I was used to avoid contamination of the products by the Mannich bases of the 2-isomer. Except for II and III, which were prepared in isoamyl alcohol, each of the β -amino ketones was prepared from formaldehyde and the base in absolute ethanol by the usual procedure.¹¹

Method of Testing for Antimicrobial Activity.—

The test microorganisms consisted of stock laboratory strains of *Staphylococcus aureus*, *Escherichia coli*, and *Saccharomyces cerevisiae*. The minimal inhibitory concentration (MIC) of each compound, as well as of each of four antibiotics, for the test organisms was determined by the agar dilution method. Appropriate concentrations of each compound were incorporated in 15-ml. portions of liquified nutrient agar, the medium was then poured into Petri plates, and 0.05 ml. of 24-hr. nutrient broth cultures of the microbial species were spread on the solidified agar surfaces. For the yeast species, glucose yeast infusion agar and broth, instead of nutrient agar and broth, were employed.

The lowest concentrations of Mannich bases or of antibiotics that prevented the development of visible growth are listed in Table I. The range of values represents maximum fluctuation in a series of assays. The data were obtained from tests with solutions sterilized by filtration; autoclaving destroyed as much as 90% of the activity of the various bases. Additional tests carried out with the most active of the Mannich bases (VII) demonstrated that the substance is not germicidal at low concentrations nor is its antimicrobial potency inactivated by lecithin. Inasmuch as the compound suppresses the growth of gram-positive and gram-negative bacteria, and yeast cells at similar concentrations, the mechanism of action is probably not associated with suppression of cell wall synthesis. Further speculation concerning the mode of action must await additional observations of microorganisms exposed to the compound.

Experimental

3-Acetylbenzo[b]thiophene (I).—To a rapidly stirred mixture of 134.2 g. (1.0 mole) of benzo[b]thiophene and 112 g. (1.1 moles) of acetic anhydride was added 10 g. of ferric chloride in one portion. The temperature immediately rose to 96°, and the reaction mixture became very dark. After stirring for 1.5 hr., 400 ml. of cold water was added, and the dark mixture was extracted with three 200-ml. portions of ether. The combined ether phase was washed with 10% sodium carbonate solution and dried with MgSO₄. After removal of the solvent and fractionation, 105 g. (70%) of a clear colorless liquid boiling at 129–131° (1 mm.) was obtained. Upon cooling and seeding, a white crystalline mass formed which was recrystallized three times from ethanol to yield a product melting at 64–65°, which showed only one peak in the gas chromatograph. The reported melting point is 64–65°,¹³ $\lambda_{\text{max}}^{\text{KB}}$ 6.05 (C=O) and 7.35 μ (CH₂—C=O). The oxime was prepared as white platelets which melted at 123–124°.

Anal. Calcd. for C₁₀H₉NOS: C, 62.85; H, 4.75; N, 7.32. Found: C, 62.91; H, 4.69; N, 7.24.

Mannich Bases From I (II–IX).—In a 50-ml. flask containing 25 ml. of absolute ethanol (dry isoamyl alcohol for II and III) was added 0.05 mole of the respective amine, and the pH was adjusted to 3–4 with concentrated HCl. To this was added 8.8 g. (0.05 mole) of the ketone and 2.3 g. of paraformaldehyde. The reaction mixture was allowed to reflux for 4 hr. and was then poured into 100 ml. of dry acetone. After cooling in the refrigerator overnight, the white precipitate was collected and recrystallized from absolute ethanol.

4-(Alkoxystryl)quinolines¹

C. T. BAHNER, H. KINDER, AND T. RIGDON

Carson-Newman College, Jefferson City, Tennessee

Received August 26, 1964

The activity of 4-(4-aminostyryl)quinolines against Walker 256 tumors in rats and against KB tumor cells in cell culture² led us to prepare the series of 4-(methoxystyryl)quinolines listed in Table I for similar testing. In each case a mixture of lepidine hydrochloride and an equimolar quantity of the appropriate aldehyde was heated until reaction had taken place, the mixture was dissolved in hot methanol and neutralized by adding an excess of concentrated ammonium hydroxide, and the oil which separated upon addition of water was recrystallized from isohexane, isooctane, or ethanol until acceptably pure. Samples were submitted for antitumor tests *in vivo* and *in vitro*. None of the compounds tested showed clear-cut activity against the Walker tumor at a single 200 mg./kg. dose level, but several of them produced 50% reduction in growth rate of KB cells in tissue culture at concentrations below 4 γ /ml., which placed them in the same range with 4-(4-dimethylaminostyryl)quinoline.

It has been suggested that there might be a correlation between the ultraviolet absorption maxima of such compounds and their cytotoxicity. Absorption spectra were examined with a Beckman DU spectrophotometer. Most of the peaks were not sharp but there were observable differences between the compounds. One interesting observation was that the wave length and intensity of absorption both diminished when a methanol solution of the compound was allowed to stand 1 day, but the acetic acid solutions did not undergo this change. On the other hand, the acetic acid solutions had an additional peak at a longer wave length than the methanol solutions. The shift in the methanol solutions was least for the dimethoxy and trimethoxy compounds containing a methoxy group at the 2-position, but 4-(2-methoxystyryl)quinoline itself exhibited a decided shift. The greatest difference between log ϵ for the acetic acid solution peak and the peak in methanol was exhibited by the three monomethoxy compounds and the 2,4-dimethoxy compound. The presence of a 2-methoxy group seemed to increase cytotoxicity except in the 2,3-dimethoxy compound.

(1) This work was supported by Public Health Service Research Grants CA-03717-05-7 from the National Cancer Institute.

(2) C. T. Bahner, L. M. Rives, and C. Breder, *J. Med. Chem.*, **7**, 818 (1964); C. T. Bahner, D. Brotherton, W. H. Chapman, Jr., W. Longuire, H. B. Orr, L. M. Rives, E. B. Senter, and W. Yee, *ibid.*, submitted for publication.

(14) M. W. Farrar and R. Levine, *J. Am. Chem. Soc.*, **72**, 4433 (1950).