

Acknowledgments.—The authors wish to express appreciation to the Upjohn Company for financial assistance and Wyeth Laboratories for sodium γ -hydroxybutyrate.

Some 3-Arylaminoquinoxaline-2-carboxylic Acids

JOSEPH G. LOMBARDINO

Medical Research Laboratories, Chas. Pfizer & Co., Inc.,
Groton, Connecticut

Received May 14, 1966

A hypothetical receptor site has been proposed by Scherrer, *et al.*,¹ in an attempt to correlate activity in some N-arylanthranilic acids with other known anti-inflammatory compounds. Important features of the

receptor site include a cavity to accommodate an out-of-plane aryl ring and a cationic site 5.2 Å from a good hydrogen-bonding site. The present paper describes some compounds which might possibly fit such a receptor.

The readily available² ethyl 3-chloroquinoxaline-2-carboxylate provided a direct route to 3-arylaminoquinoxaline-2-carboxylic acids including some with out-of-plane aryl rings. Displacement of the relatively reactive chloride³ by substituted anilines followed by hydrolysis produced the compounds shown in Table I. Table II presents ultraviolet absorption data for some of these compounds. As found by Scherrer, *et al.*,¹ for N-arylanthranilic acids, an absorption at $285 \pm 5 \text{ m}\mu$ is present in those structures possessing an in-plane arylamino function. A hypsochromic shift is observed in those structures with out-of-plane arylamino functions (7, 11) or in a 3-alkylamino (17) derivative.

When examined in the carrageenin rat foot edema-

TABLE I
SODIUM SALTS OF 3-ARYLAMINOQUINOXALINE-2-CARBOXYLIC ACIDS

No.	Ar	Method of prep ^a	Yield, ^b %	Formula	Calcd			Found		
					C	H	N	C	H	N
1	Phenyl	A	97	C ₁₅ H ₁₀ N ₂ O ₂ Na	62.72	3.51	14.63	62.70	3.69	14.63
2	3-Trifluoromethylphenyl	A	83	C ₁₆ H ₉ F ₃ N ₂ O ₂ Na	54.09	2.55	11.83	53.88	2.62	11.66
3	2,4-Dichlorophenyl	A	94	C ₁₅ H ₈ Cl ₂ N ₂ O ₂ Na	50.58	2.26	11.80	50.71	2.50	12.07
4	2,3-Dimethylphenyl	A	59	C ₁₇ H ₁₄ N ₂ O ₂ Na	64.80	4.48	13.35	64.97	4.78	13.35
5	3-Chloro-4,6-dimethoxyphenyl	A	95	C ₁₇ H ₁₃ ClN ₂ O ₄ Na	53.48	3.43	11.00	53.19	3.57	11.01
6	3-Chloro-2-methylphenyl	A	18	C ₁₆ H ₁₁ ClN ₂ O ₂ Na	57.30	3.30	12.50	56.99	3.35	12.51
7	2,6-Dimethylphenyl	B	2	C ₁₇ H ₁₄ N ₂ O ₂ Na·H ₂ O	61.25	4.88	12.60	60.95	4.75	12.58
8	2,3-Dichlorophenyl	C	52	C ₁₅ H ₈ Cl ₂ N ₂ O ₂ Na·H ₂ O	48.14	2.67	11.23	48.40	2.71	11.36
9	2-Methylphenyl	B	42	C ₁₆ H ₁₁ N ₂ O ₂ Na·0.5H ₂ O	62.10	3.89	13.59	61.92	4.20	13.84
10	2-Methoxyphenyl	A	91	C ₁₆ H ₁₂ N ₂ O ₃ Na	60.56	3.81	13.25	60.34	3.81	13.19
11	1-Naphthyl	C	39	C ₁₈ H ₁₂ N ₂ O ₂ Na	67.69	3.56	12.49	67.42	3.44	12.29
12	3-Nitrophenyl	A	96	C ₁₅ H ₉ N ₃ O ₄ Na·H ₂ O	51.50	3.16	15.99	51.69	3.24	15.97
13	2-Bromophenyl	C	82	C ₁₅ H ₉ O ₂ BrN ₂ Na	49.20	2.48	11.48	49.45	2.51	11.52
14	4-Chloro-2-methylphenyl	A	16	C ₁₆ H ₁₁ ClN ₂ O ₂ Na·0.5H ₂ O	55.70	3.48	12.19	55.55	3.53	12.30
15	3-Methoxyphenyl	A	96	C ₁₆ H ₁₂ N ₂ O ₃ Na	60.56	3.81	13.24	60.30	3.80	13.27
16	4-Chlorophenyl	A	76	C ₁₅ H ₁₀ ClN ₂ O ₂ Na	54.51	3.03	12.72	54.79	3.16	13.03
17	2-Morpholinoethyl	B	45 ^c	C ₁₅ H ₁₇ N ₄ O ₃ Na	55.55	5.28	17.28	55.44	5.44	17.44

^a See Experimental Section. ^b None of these compounds melted below 340°. ^c Recrystallized from acetone, mp 298-300° dec.

TABLE II
ULTRAVIOLET SPECTRA OF 3-ARYLAMINOQUINOXALINE-2-CARBOXYLIC ACID SODIUM SALTS

No.	$\lambda_{\text{max}}^{\text{EtOH}}$, m μ	ϵ , respectively
2	290, 390	30,700, 6870
3	294, 390	30,400, 7050
4	283, 400	24,000, 5600
5	288, 321, 408	26,800, 19,000, 6050
6	286, 393	25,200, 6150
7	262, 385	25,300, 5250
9	285, 400	25,700, 5830
11	248, 260, 328, 405	21,100, 19,100, 17,000, 6400
12	221, 288, 387	36,000, 31,400, 7050
13	288, 390	27,000, 6500
16	290, 390	27,000, 6060
17	256, 298, 384	30,800, 3300, 5600

test of Winter, *et al.*,⁴ at an oral dose of 100 mg/kg only **2** and **6** exhibited very weak antiedema activity. The relative lack of solubility of this class of compounds may possibly reduce oral absorption, although this was not specifically investigated.

Experimental Section

Compounds in Table I were synthesized by one of three methods.

A.—A combination of 0.015 mole of ethyl 3-chloroquinoxaline-2-carboxylate and 0.045 mole of a substituted aniline was heated in an oil bath at 125° in a nitrogen atmosphere for 2 hr. After cooling, 25 ml of ethanol and 25 ml of 10% NaOH were added and

(1) R. A. Scherrer, C. V. Winder, and F. W. Short, 9th National Medicinal Chemistry Symposium of the American Chemical Society, June 1964, Abstracts, pp 11f, 11g.

(2) A. H. Gowenlock, G. T. Newbold, and F. S. Spring, *J. Chem. Soc.*, 622 (1945).

(3) R. G. Shepiner, J. L. Fedrick, *Advan. Heterocyclic Chem.*, **4**, 374 (1965).

(4) C. A. Winter, E. A. Risley, and G. W. Nuss, *J. Pharmacol. Exptl. Ther.*, **141**, 369 (1963).

the mixture refluxed 2 hr. Cooling, evaporation to half volume, and addition of 200 ml of water produced a solid. Filtration and thorough washing with water and ether yielded analytically pure material which did not melt below 340°.

Many of these compounds form tenacious hydrates and were dried for analysis at 132° under vacuum over P₂O₅ for 48 hr. Certain compounds (especially 2-methyl analogs) retained water of hydration even under these drying conditions.

B.—After attempting method A and detecting incomplete reaction by thin layer chromatography, the reaction mixture was heated at 125° for 17–20 hr and then hydrolyzed and purified as above.

C.—This method was similar to method A except that the reactants were combined in a pressure bottle and heated for 18 hr at 125°.

Acknowledgment.—The author is grateful to Dr. Stuart Paulson for carrying out the biological testing.

Total Synthesis of Unsaturated Phosphatidylserine Using the N-4-Chlorobutyryl Protecting Group¹

D. L. TURNER, M. J. SILVER, AND E. BACZYNSKI

Carleza Foundation, Jefferson Medical College,
Philadelphia, Pennsylvania 19107

Received May 18, 1966

Large amounts of unsaturated phosphatidylserines with defined fatty acid composition are needed for the systematic study of the effect of phospholipids on hemostasis, and these phosphatidylserines can only be obtained by total synthesis. The available syntheses have suffered in yield from the relatively energetic methods used for the removal of protecting groups.

Recently, Eschenmoser's group² has introduced an easily removable protecting group, the 4-chlorobutyryl group, and we decided to explore the applicability of this group in phosphatide synthesis.

Experimental Section³

N-(4-Chlorobutyryl)-O-benzyl-(±)-serine methyl ester was prepared exactly as described for the corresponding tryptophan derivative,² replacing the tryptophan methyl ester hydrochloride with O-benzyl-(±)-serine methyl ester hydrochloride.⁴ The product was crystallized from CCl₄-hexane, mp 42°, yield 84%.

Anal. Calcd for C₁₅H₂₀ClNO₄: C, 57.41; H, 6.43; N, 4.47. Found: C, 57.39; H, 6.54; N, 4.60.

N-(4-Chlorobutyryl)-O-benzyl-(±)-serine. A. By Hydrolysis of Methyl Ester.—The methyl ester (20 g) was dissolved in 60 ml of acetone and to this was added 1 N NaOH dropwise as the alkali was consumed. The hydrolysis was slow at first but speeded up as part of the alkali was consumed. When close to the theoretical amount of alkali was used up, 100 ml of water was added and crystals of O-benzyl-(±)-serine were filtered out. The

mother liquor was cooled and acidified with ice-cold 1 N phosphoric acid, giving an oil which solidified to 8.2 g (43%) of crystals, mp 117°, identical by infrared spectrum and mixture melting point with material obtained by method B below.

B. By Ronwin's Acylation of Acid.—Dry O-benzyl-(±)-serine⁴ (48.5 g, 0.25 mole) was refluxed and stirred with 28 ml (0.25 mole) of 4-chlorobutyryl chloride in 2 l. of dry ethyl acetate for 6 hr. The solution was filtered and evaporated *in vacuo* to a small volume. Addition of CCl₄ gave 42 g of the acylated material (57%). Recrystallized from ethyl acetate-petroleum ether, it had mp 120°.

Anal. Calcd for C₁₄H₁₈ClNO₄: C, 56.09; H, 6.05. Found: C, 56.27; H, 6.04.

Phthalimidomethyl Ester of O-Benzyl-N-(4-chlorobutyryl)-(±)-serine.—A solution of 3 g (0.01 mole) of the preceding acylated O-benzylserine in 20 ml of dry DMF was treated with 1.97 ml (0.01 mole) of dicyclohexylamine and a solution of 1.95 g (0.01 mole) of phthalimidomethyl chloride in 20 ml of dry DMF. The mixture was kept at 37° overnight. It was diluted with water and allowed to crystallize. Recrystallized from ethanol the collected crystals had mp 125°, yield 2.8 g.

Anal. Calcd for C₂₃H₂₈ClN₂O₈: C, 60.19; H, 5.05; N, 6.11. Found: C, 60.44; H, 5.24; N, 5.89.

Phthalimidomethyl Ester of N-(4-Chlorobutyryl)-(±)-serine.—The benzyl ester (10 g) was suspended in peroxide-free tetrahydrofuran (THF) and hydrogenated at 3 atm over the Pd catalyst of Tausz and Von Putnokoy.⁶ After the addition of boiling acetone to dissolve precipitated product, the catalyst was filtered, and the solution was concentrated in part and cooled to give crystals, mp 145°, yield 5 g. This hydrogenation did not always proceed as smoothly as described and the product gave a test with ninhydrin indicating an N → O shift. Such material was purified by chromatography on silica gel (British Drug Houses, Inc.) by eluting the desired product with acetone. The purity of the material was conveniently investigated by the using the solvent system *n*-butyl ether-propionic acid-water (120:90:11) with silica gel H (E. Merck, Darmstadt) and the staining procedure of Pataki.⁷

Anal. Calcd for C₁₈H₁₇ClN₂O₈: C, 52.11; H, 4.65; N, 7.60. Found: C, 51.84; H, 4.53; N, 7.49.

DL-Phosphatidyl-(oleoyl stearoyl)-DL-serine.⁸—1-Oleoyl-2-stearoylglycerol¹⁰ (9 g) was brought into reaction with equivalent quantities of POCl₃ and the protected serine in the presence of pyridine and quinoline under standard conditions.^{9,11} The crude product was extracted with anhydrous ether and with hexane, and the material insoluble in these two solvents was rejected. The phosphatide appeared to be in the soluble portion by infrared spectrum. This portion weighed 7 g. The material was dissolved in 75 ml of peroxide-free THF (distilled over excess triphenylphosphine) and to the solution was added a mixture of 48 ml of 0.5 N aqueous silver perchlorate and 75 ml of water. The mixture was shaken mechanically under nitrogen in the dark for 40 hr. The THF was removed by distillation and the aqueous layer was extracted (CHCl₃). The CHCl₃ solution was washed with dilute KCl solution, dried, and distilled finally in a high vacuum. The dry residue was dissolved in 90 ml of ethanol and cooled in an ice bath. To the solution was added 8.4 mM hydrazine in 90 ml of ethanol. The reaction mixture was stored at room temperature for 30 min and then evaporated.¹² The crude product was purified by the usual techniques of chromatography on silicic acid and DEAE-cellulose (acetate form).⁹ The yield was 727 mg of crude phosphatidylserine (one chromatogram) of which half could be recovered as completely homogeneous material after a second chromatograph. The product showed only one spot moving opposite to natural phosphatidylserine when chromatographed in amounts of 250 μg on silicic acid impregnated paper; it had the infrared spectrum of phos-

(1) Supported by Grant No. A-533 of the U. S. Public Health Service.

(2) H. Peter, M. Brugger, J. Schreiber, and A. Eschenmoser, *Helv. Chim. Acta*, **46**, 577 (1963).

(3) Solvents were dried over Molecular Sieve 4A of Linde Products Division, Union Carbide Corp. 4-Chlorobutyryl chloride was purchased from Aldrich Chemical Co., Inc. and was used as received. Oleic acid was 99% purchased from Applied Science Laboratories, Inc., State College, Pa. Stearic acid 99% was purchased from Stearinerie Dubois Fils, Usine de Scoury (Indre), France. Melting points were determined on the Kofler bar and infrared spectra on the Perkin-Elmer Infracord spectrophotometer. Evaporations were done *in vacuo* in a rotating evaporator. All operations with unsaturated material were under oxygen-free nitrogen.

(4) W. Grassmann, E. Wünsch, P. Deufel, and A. Zwick, *Chem. Ber.*, **91**, 538 (1958).

(5) Cf. (a) E. Ronwin, *J. Org. Chem.*, **18**, 127 (1953); (b) *ibid.*, **18**, 1546 (1953); (c) *ibid.*, **22**, 1180 (1957); (d) E. Ronwin and C. B. Warren, *ibid.*, **29**, 2276 (1964).

(6) J. Tausz and N. Von Putnokoy, *Chem. Ber.*, **52**, 1573 (1919).

(7) G. Pataki, *J. Chromatog.*, **12**, 541 (1963).

(8) For nomenclature see ref 9, footnote 1.

(9) D. L. Turner, M. J. Silver, E. Baczynski, N. Giordano, and I. Rodalewicz, *J. Lipid Res.*, **5**, 616 (1964).

(10) N. A. Bogoslovskii, G. I. Samoshvalov, and N. A. Preobrazhenskii, *Zh. Obshch. Khim.*, **31**, 1143 (1961).

(11) E. Baer and D. Buchnea, *J. Am. Chem. Soc.*, **81**, 1758 (1959).

(12) The reaction period was reduced because the more usual conditions^{9,11} gave poorer yields, possibly connected with the presence of colloidal silver.