

Forschungsinstitut für Naturarzneimittel, Prague, for supplies of material in this investigation. He is also grateful to Doctors Francis G. Henderson, Alena

Kováříková, and Ken Ito, and Messrs. Delbert Campbell, Harold E. Roeder, and Terry Davis for their assistance in these experiments.

## Small Ring Analogs of Acetylcholine. Synthesis and Absolute Configurations of Cyclopropane Derivatives<sup>1a</sup>

P. DOUGLAS ARMSTRONG AND JOSEPH G. CANNON<sup>1b</sup>

Division of Medicinal Chemistry, College of Pharmacy, The University of Iowa, Iowa City, Iowa 52240


Received June 13, 1970

Conformational rigidity has been conferred upon the OCCN portion of acetylcholine by incorporation of the C atoms into a cyclopropane ring. *trans*-2-Acetoxypropyltrimethylammonium iodide has been prepared from an olefinic starting material, 2-vinyloxytetrahydropyran; assignment of the *trans* configuration to the 1,2-disubstituted cyclopropane systems was based upon literature precedent and upon nmr data, and was confirmed by X-ray crystallographic analysis. Resolution of one of the racemic intermediates in the reaction sequence was achieved, which permitted preparation of both enantiomers of the final product. X-Ray crystallographic analysis has demonstrated that the muscarinically active (+)-*trans*-2-acetoxypropyltrimethylammonium iodide possesses the same absolute configuration as the muscarinically active enantiomers of acetyl- $\beta$ -methylcholine and muscarine.

Earlier communications have presented preliminary reports of the synthesis<sup>2</sup> of (+)- and (-)-*trans*-2-acetoxypropyltrimethylammonium iodide **9** and the details of the cholinergic effects and enzymatic hydrolysis rates<sup>3</sup> of the two isomers. Herein are presented the details of synthesis of (+)- and (-)-**9** (outlined in Scheme I), and assignments of absolute configurations to the enantiomers.

The 2-tetrahydropyranyl moiety was utilized to mask the cyclopropanol group which is highly susceptible to ring-opening and other undesirable reactions.<sup>4</sup> This protecting group is stable in neutral and basic media, and is easily cleaved under very mild acidic conditions.<sup>5</sup> 2-Vinyloxytetrahydropyran (**2**) reacted with ethyl diazoacetate to give a mixture of the *cis*- and *trans*-cyclopropane isomers (**10** and **3**) which were separable by distillation. The configurational assignments at this stage were tentative and were based upon the expected predominance of the sterically favored *trans* isomer in reactions of this type.<sup>6</sup> Vpc analysis of the crude reaction mixture indicated a ratio of *trans*-**3** to *cis*-**10** of 45:1. Attempts to modify this ring-closure reaction to attain larger proportions of the *cis* isomer failed, as did efforts to photoisomerize the *trans* isomer to *cis*.<sup>7</sup>

The *trans* isomer **3** could be converted into the amide **4** either with anhyd NH<sub>3</sub> in ethylene glycol or by use of *n*-BuLi and liq NH<sub>3</sub>,<sup>8</sup> while the *n*-BuLi method afforded lower yields than the ethylene glycol method, it



(1) (a) This investigation was supported in part by Grant NS-06100, National Institute of Neurological Diseases and Stroke. Abstracted in part from a thesis submitted by P. D. A. in partial fulfillment of the requirements for the degree of Doctor of Philosophy, University of Iowa, 1968.

(b) To whom all correspondence should be addressed.

(2) P. D. Armstrong, J. G. Cannon, and J. P. Long, *Nature (London)*, **220**, 65 (1968).

(3) C. Y. Chiou, J. P. Long, J. G. Cannon, and P. D. Armstrong, *J. Pharmacol. Exp. Ther.*, **166**, 243 (1969).

(4) C. H. DePuy, *Accounts Chem. Res.*, **1**, 33 (1968).

(5) D. N. Robertson, *J. Org. Chem.*, **25**, 931 (1960).

(6) P. S. Skell and R. M. Etter, *Proc. Chem. Soc.*, 443 (1961).

(7) G. W. Griffin, J. Covell, R. C. Petterson, R. M. Dodson, and G. Klose, *J. Amer. Chem. Soc.*, **87**, 1410 (1965); D. W. Setser, B. S. Rabinovitch, and E. G. Spittler, *J. Chem. Phys.*, **35**, 1840 (1961).

(8) K. W. Yang, J. G. Cannon, and J. G. Rose, *Tetrahedron Lett.*, 1791 (1970).

permitted a less tedious, less time-consuming work-up. Ammonolysis of the *cis* ester **10** by the ethylene glycol method (Chart I) gave rise to an amide which could not be obtained analytically pure. Both the *trans* amide **4** and its *cis* isomer **11** underwent the Hofmann hypohalite reaction to form the *trans* and *cis* primary amines **5** and **12**. The *cis* amine **12** did not yield a satisfactory analysis for all elements, although spectral data supported its structure. The use of the reaction sequence in Scheme I as a route to the *cis* isomer of structure **9** was concluded to be unpromising and was abandoned.




Figure 1.—Absolute configurations of muscarinically active compounds. Structural representation for (+)-*trans*-**9** adapted from Chothia and Pauling.<sup>11</sup>

The nmr spectrum of the *cis* primary amine **12** showed a sharp singlet at  $\delta$  1.7, which integrated for 2 protons and was assigned to the  $\text{NH}_2$  group. The *trans* isomer **5** showed a sharp singlet at  $\delta$  1.4 which integrated for 2 protons and was assigned to the  $\text{NH}_2$  group. The  $\text{NH}_2$  group of cyclopropylamine exhibits a sharp singlet at  $\delta$  1.83.<sup>9</sup> The positions of the chemical shifts of the  $\text{NH}_2$  protons of the *cis* and *trans* isomers **12** and **5** support the configurational assignments, since intramolecular H bonding involving the  $\text{NH}_2$  group, which may be expected to be stronger in the *cis* isomer than in the *trans*, has been reported<sup>10</sup> to produce a downfield shift. Crystallographic studies<sup>11</sup> have proved the correctness of these assignments of geometry to the *cis*-*trans* isomers.

The tetrahydropyranyl group was removed from **6** under mild conditions to afford a quantitative yield of ( $\pm$ )-*trans*-2-dimethylamino-cyclopropanol (**7**) which was resolved with *l*-tartaric acid. Each enantiomer of **7** was converted into the final structure **9**.

The (+) enantiomer of **9**, which exhibits powerful muscarinic effects,<sup>3</sup> has been shown by X-ray crystallographic analysis to have the S-S absolute configuration<sup>11</sup> (Figure 1). Models indicate that conformations are possible in which the train of atoms in (+)-*trans*-**9**, C1, C2, C3; N; C4; C5; O1; C6; C7 superimpose perfectly upon C1, C2, C3; N; C4; C5; O1; C6; C7 of L-(+)-muscarnine (Figure 1), the muscarinically active stereoisomer.<sup>12</sup> The absolute configuration of (+)-*trans*-**9** is the same as that of the corresponding asymmetric center of the muscarinically potent acetyl

(+)-(*S*)- $\beta$ -methylcholine.<sup>13</sup> The conformation suggested<sup>11</sup> for (+)-*trans*-**9** "relevant to muscarinic receptors" differs from a conformation for "receptor-bound" acetylcholine proposed by Belleau and Puranen.<sup>14</sup> Data are not yet available which would permit selection one of these conformations over the other.

## Experimental Section<sup>15</sup>

**2-(2-Tetrahydropyranyloxy)ethyl chloride (1)** was prepared in 80% yield by the method of Frearson, *et al.*:<sup>16</sup> bp 103–104° (25 mm);  $n_D^{25}$  1.4528 [lit.<sup>16</sup> bp 93° (20 mm);  $n_D^{20}$  1.4580].

**2-Vinyloxytetrahydropyran (2)**.—A 1-l. flask was charged with 509 g (9.1 moles) of KOH, and the system was connected to a source of reduced pressure ( $\text{H}_2\text{O}$  aspirator). The flask was maintained at 175–180° while 279 g (1.60 moles) of **1** was added dropwise over 2 hr, while maintaining a gentle flow of air through the apparatus. After addition was complete, stirring was continued at 170–180° for 1.5 hr, then at 195–200° for 1 hr, and finally at 215–220° for 0.5 hr. A total of 195 g of colorless liquid distilled. The residue in the reaction vessel was taken up in  $\text{H}_2\text{O}$  and was extracted with  $\text{Et}_2\text{O}$ . The extract was added to the distillate, the combined organic phases were dried ( $\text{K}_2\text{CO}_3$ ) and concentrated at atm pressure, and the residue was distilled through a column packed with Berl saddles to yield 122.7 g of a colorless liquid, bp 55–56° (25 mm). The pot residue was distilled to afford of 37.6 g of starting material **1**; yield of **2** (based upon amount of **1** recovered), 66%. Distillation of **2** through a spinning band column gave a center fraction: bp 60° (21 mm);  $n_D^{25}$  1.4413. *Anal.* ( $\text{C}_7\text{H}_{12}\text{O}_2$ ) C, H.

**Ethyl *cis*- and *trans*-2-(2-Tetrahydropyranyloxy)cyclopropanecarboxylate (10 and 3)**.—Ethyl diazoacetate (185 g, 1.62 moles, Aldrich Chemical Co.) in 180 g (1.40 moles) of **2** was added dropwise to a stirred mixture of 3.5 g of Cu powder (Matheson, Coleman, and Bell, purified) and 58 g (0.45 mole) of **2** at 125–130° over 5.5 hr. After addition was complete, stirring was continued at 150–160° for 0.5 hr. The reaction mixture was cooled to room temp and was filtered. The combined filtrates from two such reactions were distilled through a Vigreux column; the fraction bp 54–114° (3 mm), amounting to 507 g, was distilled through a spinning band column. The fraction bp 73–74° (1 mm), consisting of the *cis* isomer **10** was homogeneous by vpc: yield, 9.0 g (1.3%);  $n_D^{20}$  1.4572; ir (film) 1720  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ). *Anal.* ( $\text{C}_{11}\text{H}_{18}\text{O}_4$ ) C, H.

The fraction bp 92–94° (1 mm) consisted of 433 g (62%) of *trans* isomer **3**:  $n_D^{20}$  1.4531; ir (film) 1725  $\text{cm}^{-1}$  ( $\text{C}=\text{O}$ ). *Anal.* ( $\text{C}_{11}\text{H}_{18}\text{O}_4$ ) C, H. This material demonstrated a different vpc retention time than the bp 73–74° fraction, and was homogeneous.

***trans*-2-(2-Tetrahydropyranyloxy)cyclopropanecarboxamide (4)**. **Method A**.—Anhydrous  $\text{NH}_3$  was passed into a stirred mixture of 543 g (2.5 moles) of **3** and 3.8 l. of ethylene glycol at room temp for 62 hr. Removal of the  $\text{NH}_3$  and ethylene glycol under reduced pressure at a temp not exceeding 75° gave 476 g of a light yellow-orange solid which was treated with 500 ml of  $\text{C}_6\text{H}_6$  on a steam bath. The resulting soln was filtered, to the filtrate was added 2.5 l. of anhyd  $\text{Et}_2\text{O}$ , which resulted in separation of a light yellow orange crystalline solid which was collected on a filter. Dilution of the filtrate with 500 ml of heptane caused a light yellow oil to separate, which was discarded. The organic soln, upon standing overnight, deposited crystals which were collected on a filter. This filtrate was again diluted with 500 ml of heptane; the oil which sepd was removed; and the organic layer deposited an additional crop of crystals upon standing; total yield, 258 g (55%). This material was sufficiently pure for

(13) A. B. Beckett, *Ann. N. Y. Acad. Sci.*, **144**, (75) (1967).

(14) B. Belleau and J. Puranen, *J. Med. Chem.*, **6**, 325 (1963).

(15) Melting points were determined in open capillaries on a Thomas-Houwer Uni-Melt apparatus and are corrected. Boiling points are uncorrected. Ir spectra were recorded on Beckman IR5-A and IR-10 instruments, and nmr spectra on a Varian Associates A-60 instrument (Me<sub>2</sub>SO). Optical rotations were determined on a Zeiss circular scale polarimeter. Gas chromatograms were obtained using an F & M Model 500 instrument with a Model 1600 flame ionization detector, using He as the carrier. Where analyses are indicated only by symbols of the elements, the analytical results for those elements were within  $\pm 0.4\%$  of the theoretical value.

(16) P. M. Frearson, D. G. Hardy, and E. S. Stern, *J. Chem. Soc.*, 2193 (1960).

(9) N. S. Blacca, L. F. Johnson, and J. N. Shoolery, "High Resolution NMR Spectra Catalog," Varian Associates, Palo Alto, Calif., 1962, Spectrum no. 37.

(10) J. A. Hyde, W. G. Schneider, and H. J. Bernstein, "High-Resolution Nuclear Magnetic Resonance," McGraw-Hill Book Co., New York, N. Y., 1958, Chapter 15.

(11) C. Chothia and P. Pauling, *Nature (London)*, **226**, 541 (1970).

(12) P. Waser, *Experientia*, **17**, 300 (1961).

subsequent steps. A small portion of it was recryst repeatedly from EtOAc: mp 92.5–93.5°; ir (KBr) 1620 ("amide II"), 1655 ("amide I"), 3190 (NH<sub>2</sub>), and 3360 cm<sup>-1</sup> (NH<sub>2</sub>). *Anal.* (C<sub>8</sub>H<sub>15</sub>NO<sub>2</sub>) C, H, N.

**Method B.**—To a stirred mixture of 75 ml of liquid NH<sub>3</sub> and 25 ml of purified THF was added in a slow stream 0.1 mole of *n*-BuLi in hexane (Alpha Inorganics, Inc.); a copious white ppt formed. Compound **3** (20.8 g 0.097 mole) in an equal vol of THF was added dropwise with stirring; the white ppt dissolved. After stirring 4 hr, the excess NH<sub>3</sub> was removed at room temp, the reaction mixture was poured into an ice–H<sub>2</sub>O slush, and the resulting soln was extracted repeatedly with Et<sub>2</sub>O. Evaporation of the Et<sub>2</sub>O left a light yellow gum which was chromatographed in C<sub>6</sub>H<sub>6</sub> on neutral alumina and eluted with Et<sub>2</sub>O–MeOH (9:1). The solid material from the eluate was recryst from EtOAc to yield 4.6 g (26%) of white crystals: mp 118–120°; ir (CHCl<sub>3</sub>) of the product of method B was superimposable upon a similar spectrum of the product of method A.

**cis-2-(2-Tetrahydropyranloxy)cyclopropanecarboxamide (11).**—Method A, described for the trans isomer **4**, was employed, using 10.8 g (0.05 mole) of **10** and 900 ml of ethylene glycol. Removal of the ethylene glycol from the reaction mixture left a yellow solid residue which was treated with 275 ml of EtOAc on a steam bath. On filtration and cooling overnight, this soln deposited 5.7 g of colorless crystals, the mother liquor yielded additional crystals; total yield, 6.8 g (72%). Repeated successive recrystallizations from EtOAc, C<sub>6</sub>H<sub>6</sub>, and Me<sub>2</sub>CO and column chromatography on silica gel afforded colorless crystals: mp 180–181°; ir (KBr) 1615 ("amide II"), 1660 ("amide I"), 3185 (NH<sub>2</sub>), and 3350 cm<sup>-1</sup> (NH<sub>2</sub>). *Anal.* (C<sub>9</sub>H<sub>15</sub>NO<sub>3</sub>) C, H, N; calcd, 7.56; found, 8.07.

**trans-2-(2-Tetrahydropyranloxy)cyclopropylamine (5).**—A mixture of 250 g (1.35 moles) of **4** and a soln of 101 g (1.4 moles) of Cl<sub>2</sub> and 285 g (7.1 moles) of NaOH in 3.24 l. of H<sub>2</sub>O was stirred at room temp for 0.5 hr, then at 50–60° for 19 hr. The reaction mixture was saturated with K<sub>2</sub>CO<sub>3</sub> and extracted several times with Et<sub>2</sub>O. An orange–yellow liquid which separated from the Et<sub>2</sub>O extracts was removed, and the Et<sub>2</sub>O phase was dried (K<sub>2</sub>CO<sub>3</sub>) and concentrated under reduced pressure. The residual liquid was distilled through a Vigreux column, bp 69–72° (0.8 mm), to yield 140 g (67%) of a colorless liquid. For characterization, a small portion of this material was distilled through a spinning band column, bp 67° (1.3 mm). Gc indicated that this distillate was homogeneous: *n*<sub>D</sub><sup>25</sup> 1.4698; ir (film) 3300 (NH<sub>2</sub>) and 3375 cm<sup>-1</sup> (NH<sub>2</sub>); nmr (CCl<sub>4</sub>) δ 1.4 (s, 2H).

A phenylthiourea derivative was recrystd repeatedly from EtOH, mp 160–161°. *Anal.* (C<sub>15</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub>S) C, H, N, S.

**cis-2-(2-Tetrahydropyranloxy)cyclopropylamine (12).**—A soln of 2.3 g (0.03 mole) of Cl<sub>2</sub>, 6.3 g (0.16 mole) of NaOH, and 4.5 g (0.02 mole) of **11** in 58 ml of H<sub>2</sub>O was stirred at room temp for 0.5 hr, then at 60–65° for 21 hr. The reaction mixture was saturated with K<sub>2</sub>CO<sub>3</sub> and extracted repeatedly with Et<sub>2</sub>O, the combined extracts were dried (Na<sub>2</sub>SO<sub>4</sub>) and filtered, and the Et<sub>2</sub>O was removed. The light yellow residue was distilled; bp 58–59° (4.2 mm); yield, 2.8 g (73%) of a colorless liquid; *n*<sub>D</sub><sup>20</sup> 1.4779; ir (film) 3300 (NH<sub>2</sub>) and 3370 cm<sup>-1</sup> (NH<sub>2</sub>); nmr (CCl<sub>4</sub>) δ 1.7 (s, 2H). *Anal.* (C<sub>8</sub>H<sub>15</sub>NO<sub>2</sub>) H, N; C: calcd, 61.12; found, 63.45.

A phenylthiourea derivative was recrystd from anhyd EtOH, mp 167.5–168°. *Anal.* (C<sub>15</sub>H<sub>20</sub>N<sub>2</sub>O<sub>2</sub>S) C, H, N; calcd, 9.58; found, 10.82; S, calcd, 10.97; found, 12.85.

A benzamide derivative was recryst repeatedly from C<sub>6</sub>H<sub>6</sub>, mp 162–163°. *Anal.* (C<sub>11</sub>H<sub>15</sub>NO<sub>3</sub>) H; C: calcd, 68.94; found, 71.18.

**trans-2-(2-Tetrahydropyranloxy)cyclopropyldimethylamine (6).**—A mixture of 190 ml of anhyd MeOH, 31.8 g of anhyd Na<sub>2</sub>CO<sub>3</sub>, and 19.1 g (0.12 mole) of **5** was stirred under reflux for 10 min, then 45.2 g (0.24 mole) of methyl *p*-toluenesulfonate in 190 ml of anhyd MeOH was added dropwise with stirring over 21 hr. After addition was complete, stirring and refluxing were continued for 7.25 hr, the reaction mixture was allowed to cool to room temp and was filtered. The solid which collected on the filter was washed with 3 portions of anhyd Et<sub>2</sub>O; the washings were added to the filtrate, which caused sepn of additional solid which was removed by filtration. The combined filtrate and

Et<sub>2</sub>O washings were concentrated under reduced pressure to give a light yellow liquid which was distd through a Vigreux column: bp 54–61° (0.6 mm); yield, 7.0 g (31%). A MeI salt of this product was recryst from Me<sub>2</sub>CO–EtOAc, mp 133.0–134.5° dec, when introduced into the melting point bath at 128°. *Anal.* (C<sub>11</sub>H<sub>22</sub>INO<sub>2</sub>) C, H, I.

**(±)-trans-2-Dimethylaminocyclopropanol (7).**—A soln of 19.5 g (0.105 mole) of **6** in 230 ml of 3% HCl was stirred at room temp for 9 hr. The reaction mixture was washed several times with Et<sub>2</sub>O, then was satd with K<sub>2</sub>CO<sub>3</sub> while immersed in an ice bath. The resulting mixture was extracted repeatedly with Et<sub>2</sub>O, the combined extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), and the solvent was removed under reduced pressure to leave a light yellow liquid which was distd through a "short path" apparatus, bp 58–60° (3 mm). This distillate was redistd through a Vigreux column, collecting the fraction at bp 54° (2 mm): yield, 10.4 g (98%); *n*<sub>D</sub><sup>20</sup> 1.4428.

A bifumarate salt of **7** was recrystd twice from EtOAc, mp 114–117.5°. *Anal.* (C<sub>9</sub>H<sub>15</sub>NO<sub>5</sub>) C, H, N.

**Resolution of (±)-trans-2-Dimethylaminocyclopropanol (7).**—To 14.7 g (0.1 mole) of *l*-tartaric acid in 1.18 l. of anhyd EtOH and 1.78 l. of EtOAc was added 9.9 g (0.1 mole) of (±)-**7** in 600 ml of anhyd EtOH. After cooling for 36 hr, the crystals which separated were collected on a filter, the mother liquor was reserved as solution A. The crystals were recrystd 4 times from EtOH–EtOAc, then once from MeCN–*i*-PrOH, mp 170–171° dec. *Anal.* (C<sub>9</sub>H<sub>17</sub>NO<sub>7</sub>) C, H, N. A soln of this material (2.1 g) in 25 ml of H<sub>2</sub>O was saturated with K<sub>2</sub>CO<sub>3</sub> and was extracted repeatedly with Et<sub>2</sub>O. The combined extracts were dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated under reduced pressure, and the residual liquid was distd through a "short path" apparatus: bp 54–55° (3.0 mm); yield, 0.6 g [*α*]<sub>D</sub><sup>25</sup> –70.1° (*c* 3.9, H<sub>2</sub>O). An ir spectrum of this material (film) was superimposable upon one of (±)-**7**.

Solution A was concentrated under reduced pressure to 1.1 l. Anhyd EtOH (350 ml) and 100 ml of EtOAc were added and the resulting soln was permitted to stand at room temp for 1 day, resulting in separation of crystals. Additional crops of crystals were obtained by successive additions to the filtrate of 100-ml portions of EtOAc. The combined crops were recryst from MeCN and from anhyd EtOH, mp 123–124° dec. *Anal.* (C<sub>9</sub>H<sub>17</sub>NO<sub>7</sub>) C, H, N. The free base was liberated as described for the (–) isomer: bp 53° (2.4 mm); yield, 0.6 g; [*α*]<sub>D</sub><sup>25</sup> +59.2° (*c* 3.6, H<sub>2</sub>O). An ir spectrum (film) of this material was superimposable upon one of (±)-**7**.

**(–)-trans-2-Trimethylammoniumcyclopropanol Iodide (8).**—A soln of 0.5 g (0.005 mole) of (–)-**7** and 2.0 g (0.014 mole) of MeI in 15 ml of anhyd MeOH was stirred at room temp for 40 min and at reflux for 20 min. The solid which separated was collected on a filter, diln of the filtrate with EtOAc and Et<sub>2</sub>O resulted in separation of additional solid. The combined solids were recrystd from anhyd MeOH to give 0.6 g (54%) of crystals: mp 250–253° dec; [*α*]<sub>D</sub><sup>25</sup> –13.5° (*c* 2.1, H<sub>2</sub>O). *Anal.* (C<sub>8</sub>H<sub>14</sub>INO) C, H, I, N.

**(+)-trans-2-Trimethylammoniumcyclopropanol Iodide (8).**—Using the procedure described for (–)-**8**, 0.4 g (0.004 mole) of (+)-**7** yielded a solid which was recryst from anhyd MeOH to afford 0.5 g (51%) of crystals: mp 254–255° dec; [*α*]<sub>D</sub><sup>25</sup> +14.2° (*c* 1.9, H<sub>2</sub>O). *Anal.* (C<sub>8</sub>H<sub>14</sub>INO) C, H, I, N.

**(–)-trans-2-Acetoxypropyltrimethylammonium Iodide (9).**—A mixture of 0.564 g (0.002 mole) of (–)-**8** and 6.0 g (0.06 mole) of Ac<sub>2</sub>O was stirred on a steam bath for 13 hr. The cooled reaction mixture was diluted with 400 ml of anhyd Et<sub>2</sub>O, the solid which sepd was collected on a filter, washed with anhyd Et<sub>2</sub>O, and recrystd repeatedly from EtOAc–MeOH to yield 0.25 g (43%) of a light tan solid: mp 173–175.5°; [*α*]<sub>D</sub><sup>20</sup> –89° (*c* 1.01 H<sub>2</sub>O). *Anal.* (C<sub>8</sub>H<sub>16</sub>INO<sub>2</sub>) C, H, I, N.

**(+)-trans-2-Acetoxypropyltrimethylammonium Iodide (9).**—The procedure described for (–)-**9** was carried out on 0.495 g (0.002 mole) of (+)-**8**, to yield 0.196 g (34%) of an almost colorless solid: mp 172–174°; [*α*]<sub>D</sub><sup>20</sup> +93.2° (*c* 1.13, H<sub>2</sub>O). *Anal.* (C<sub>8</sub>H<sub>16</sub>INO<sub>2</sub>) C, H, I, N.

**Acknowledgment.**—We thank Dr. Peter Pauling for his interest in this research and for determining the absolute configurations of the final compounds.