

Carbamate Ester Latentiation of Physiologically Active Amines

ANTHONY J. VERBISCAR

Institute of Drug Design, Sierra Madre, California 91024

AND LEO G. ABOOD*


*Center for Brain Research and Department of Biochemistry,
University of Rochester School of Medicine, Rochester, New York 14625*

Carbamate ester latentiation of phenethylamine, *dl*- and *d*-amphetamine, *l*-ephedrine, and *dl*-*p*-hydroxyamphetamine is described, particularly in regard to the central activity of these drugs. By increasing lipophilicity of the amine and protecting it against enzymatic deactivation, carbamoylation facilitates transport into the brain. The lack of a carbamate-specific enzyme in mammals requires that a potency toward biological hydrolysis be intrinsic in the drug in order for amine release to occur in a useful period of time. Eight different alcoholic leaving groups were incorporated into the carbamate esters and studied for their ability to facilitate such hydrolysis. Functional groups in the alcoholic moiety which can act on the carbamate ester linkage by electron withdrawal and neighboring group participation appear to facilitate amine release. The most active carbamate esters in this series have a biphasic spectrum of activity characterized by mild sedation initially followed by moderate stimulation. Anorexigenic activity of the carbamates occurs with considerably less central stimulation than with amphetamine, and a degree of selectivity in modifying such drug action is apparent. Preliminary studies with the *o*-nitrophenyl and *o*-carbomethoxyphenyl carbamate esters of α -[¹⁴C]amphetamine demonstrate that these carbamates rapidly enter the mouse brain where they are readily hydrolyzed.¹

Since Ehrlich's original observation about restricted penetration of drugs into the brain² the blood-brain barrier has come to be recognized as a complex biochemical phenomenon with no single hypothesis explaining all of the events.³⁻⁶ Nevertheless, in the case of amines, transport seems to depend primarily on two aspects of metabolism, lipophilicity and biotransformation. For compounds which ionize, the rate of transport appears to be proportional to the concentration of undissociated molecules in the blood and its degree of lipid solubility. Since carbamoylated amines do not ionize, they are more soluble in organic and lipoidal systems. This lipophilicity theory is one that has been broadly useful in correlating the action of centrally active drugs.⁷⁻⁹ In the case of amines, however, an enzymatic barrier must be considered in addition to the lipid barrier. Metabolizing enzymes present in the brain may deactivate amines before they reach their proper site of activity. For example, presence of monoamine oxidase (MAO) in brain capillaries destroys dopamine and other phenethylamines before their entry into the brain.⁹ By protecting the amine from deactivation by MAO, carbamoylation facilitates its transport.

Success with carbamate ester latentiation of amines requires that the drug be hydrolyzed to a carbamic acid and alcohol moiety after penetration into the

brain. At physiological pH values carbamic acids are unstable and decompose to the parent amine and CO₂.¹⁰ The release of CO₂ is physiologically comparable to the decarboxylation of indigenous amino acids to form their amine analogs, such as the decarboxylation of 5-hydroxytryptophan in brain tissue to form the biogenic amine, serotonin.



The alcoholic leaving group can act like a "carrier" and contains a potential for variation whereby amine drug transport and release characteristics can be modified for specific uses. Several ethyl carbamate latentiated drugs, such as normeperidine¹¹ and chlorphentermine,¹² have been reported and the low toxicity of EtOH favors its use as a leaving group. However, the use of simple alcohols as leaving groups in the carbamate esters has not often met with success, probably because the esters are relatively stable to biological hydrolysis. In this regard, there does not appear to be a carbamate ester specific hydrolytic enzyme in mammals, although one has been found in soil organisms.¹³ Cholinesterase will hydrolyze carbamates and become reversibly inhibited during the process.¹⁴ The hydrolysis proceeds *via* carbamoylation of the enzyme with release of the alcohol moiety, followed by decarbamoylation with release of amine. Since both rates are relatively slow, cholinesterase cannot be enticed upon to split carbamate esters. Although chymotrypsin can undergo carbamoylation,¹⁵ this reaction does not generally occur with carbamate esters.

* To whom correspondence should be addressed.

(1) N. J. Harper, *Fortschr. Arzneimittelforsch.*, **4**, 221 (1962); *J. Mol. Pharm. Chem.*, **1**, 467 (1959).

(2) P. Ehrlich, "Das Sauerstoff-Bedürfnis des Organismus. Eine Farbenanalytische Studie," Berlin (1885), from "The Collected Papers of Paul Ehrlich," F. Himmelweit, Ed., Vol. 1, Pergamon Press, New York, N. Y.

(3) B. B. Brodie in "Absorption and Distribution of Drugs," T. B. Bins, Ed., The Williams and Wilkins Company, Baltimore, Md., 1964, pp. 16-18, 199-251.

(4) J. Dobbins, *Physiol. Rev.*, **41**, 130 (1961).

(5) C. F. Barlow, *Annu. Rev. Med.*, **15**, 187 (1964).

(6) E. J. Ariens, Ed., "Molecular Pharmacology—The Mode of Action of Biologically Active Compounds," Vol. 1, Academic Press, Inc., New York, N. Y., particularly sections 1A and 1B, 1964.

(7) S. Mayer, R. P. Maickel, and B. B. Brodie, *J. Pharmacol. Exp. Ther.*, **127**, 205 (1959); B. B. Brodie, H. Kurz, and L. S. Schanker, *ibid.*, **130**, 20 (1960).

(8) C. Hansch, A. R. Steward, S. M. Anderson, and D. Bentley, *J. Mol. Chem.*, **11**, 1 (1968).

(9) A. Benler, B. Falck, Ch. Owman, and E. Rosengren, *Pharmacol. Rev.*, **18**, 369 (1966).

(10) C. Faurholt, *J. Chim. Phys.*, **22**, 1 (1925); J. Olesen, K. Veilby, and C. Faurholt, *Acta Chem. Scand.*, **6**, 398 (1952).

(11) S. M. Kupelian and A. C. Isenberg, *J. Med. Chem.*, **10**, 960 (1967).

(12) P. Bjurulf, S. Carlstrom, and G. Rorsman, *Acta Med. Scand.*, **182**, 273 (1967).

(13) P. C. Kerney and D. D. Kaufman, *Science*, **147**, 740 (1965).

(14) R. D. O'Brien, B. D. Hilton, and L. Gilmour, *Mol. Pharmacol.*, **2**, 593 (1966); R. L. Metcalf, and T. R. Fukuto, *J. Agr. Food Chem.*, **13**, 220 (1965); I. B. Wilson, M. A. Harrison, and S. Ginsburg, *J. Biol. Chem.*, **236**, 1498 (1961); (15) J. E. Cassida, K. R. Augustinsson, and G. Jonsson, *J. Econ. Entomol.*, **53**, 205 (1960).

(16) R. F. Erlanger and F. Edel, *Biochemistry*, **3**, 346 (1964).

TABLE I
CARBAMATE ESTERS

No.	Compound	Alkyl moiety	Purification solvent	Mp, °C	$[\alpha]^{25}_D$, deg (CHCl ₃ , c 1)	Formula ^a
1	I	<i>o</i> -MeO ₂ CC ₆ H ₄	CCl ₄ -hexane	96-98		C ₁₇ H ₁₇ NO ₄
2	<i>dl</i> -II	Ph	Hexane	66-67		C ₁₆ H ₁₇ NO ₂
3	<i>dl</i> -II	<i>o</i> -MeO ₂ CC ₆ H ₄	<i>i</i> -Pr ₂ O-hexane	71-77 ^b		C ₁₈ H ₁₉ NO ₄
4	<i>dl</i> -II	<i>p</i> -MeOC ₆ H ₄	CHCl ₃ -hexane	87-88		C ₁₇ H ₁₉ NO ₃
5	<i>dl</i> -II	<i>o</i> -OCHC ₆ H ₄	CHCl ₃ -hexane	79-81		C ₁₇ H ₁₇ NO ₃
6	<i>dl</i> -II	<i>o</i> -O ₂ NC ₆ H ₄	<i>i</i> -Pr ₂ O	86-90		C ₁₆ H ₁₆ N ₂ O ₄
7	<i>dl</i> -II	α -EtO ₂ CC ₂ H ₄	Hexane	Oil ^c		C ₁₅ H ₂₁ NO ₄
8	<i>d</i> -II	Ph	Hexane	64-65	-24	C ₁₆ H ₁₇ NO ₂
9	<i>d</i> -II	<i>o</i> -MeO ₂ CC ₆ H ₄	<i>i</i> -Pr ₂ O-hexane	87-88	-20	C ₁₈ H ₁₉ NO ₄
10	<i>d</i> -II	<i>o</i> -O ₂ NC ₆ H ₄	<i>i</i> -Pr ₂ O	92.5-94	-41	C ₁₆ H ₁₆ N ₂ O ₄
11	<i>d</i> -II	<i>m</i> -O ₂ NC ₆ H ₄	C ₆ H ₆ - <i>i</i> -Pr ₂ O	123-123.5	-11	C ₁₆ H ₁₆ N ₂ O ₃
12	<i>dl</i> -II	<i>p</i> -O ₂ NC ₆ H ₄	C ₆ H ₆ - <i>i</i> -Pr ₂ O	124-124.5	-29	C ₁₆ H ₁₆ N ₂ O ₄
13	<i>l</i> -III	Ph	Pet ether	72-74	-13	C ₁₇ H ₁₉ NO ₃
14	<i>l</i> -III	<i>o</i> -MeO ₂ CC ₆ H ₄	CHCl ₃ -hexane	95-96	-39	C ₁₉ H ₂₁ NO ₃
15	<i>l</i> -III	<i>p</i> -MeOC ₆ H ₄	CHCl ₃ -hexane	Oil	-29	C ₁₈ H ₂₁ NO ₄ ^d
16	<i>l</i> -III	<i>o</i> -OCHC ₆ H ₄	CHCl ₃ -hexane	Oil	-24	C ₁₈ H ₁₉ NO ₄ ^d
17	<i>l</i> -III	<i>o</i> -O ₂ NC ₆ H ₄	CHCl ₃ -hexane	Oil	-21	C ₁₇ H ₁₈ N ₂ O ₃ ^e
18	<i>l</i> -III	α -EtO ₂ CC ₂ H ₄	Hexane	Oil	-27	C ₁₆ H ₂₃ NO ₅
19	<i>dl</i> -IV	Ph	C ₆ H ₆	115-116 ^f		C ₃₆ H ₄₄ N ₃ O ₃
20	<i>dl</i> -IV	<i>o</i> -OCHC ₆ H ₄	C ₆ H ₆ -hexane	114-116		C ₁₇ H ₁₇ NO ₄

^a All compounds were analyzed for C, H, N. ^b After several crystallizations the product was chromatographically pure on tlc. ^c Compounds 7 and 15-18 were purified by pptg and evacuating the oil. They were chromatographically pure on tlc. ^d C: calcd, 69.00; found, 68.21. ^e Calcd: C, 61.81; H, 5.49; N, 8.48. ^f Bispiperazine salt complex [J. D. Benigni and A. J. Verbiscar, *J. Med. Chem.*, **6**, 607 (1963)]. ^g C: calcd, 68.55; found, 67.42.

In searching for leaving alkyl groups that would provide some control of hydrolytic stability in a physiological environment, consideration was given to structural features which influence chemical hydrolysis.¹⁶ The carbamate esters were readily prepared by standard methods¹⁷ from the amine and an appropriate ester of chloroformic acid.¹⁸ Most of the carbamates were crystalline although several of the *l*-ephedrine carbamates were resistant to crystallization and were obtained as oils. Four different physiologically active amines were carbamoylated including phenethylamine, *dl*- and *d*-amphetamine, *l*-ephedrine, and *dl*-*p*-hydroxyamphetamine (Table I). The pharmacology of these amines is relatively well known while the release of the amine could be easily confirmed by its known pharmacological action.

Pharmacological Results.—In Table II the pharmacological effects of 8 of these carbamate esters are compared with those of *d*-amphetamine. The most active compounds in this group appeared to have a biphasic spectrum of activity, characterized by mild sedation initially, followed by moderate stimulation. By comparison with amphetamine the most active carbamate derivatives were no more than 0.1 as effective in producing central stimulation (hyperactivity) and peripheral sympathetic effects. As for anorexigenic potency, the nitrophenyl derivatives of the amphetamine carbamates compared favorably with amphetamine it-

TABLE II
PHARMACOLOGICAL EFFECTS OF THE CARBAMATE ESTERS

Compd	Hyperactivity ^a	Initial tranquilization ^b	Anorexigenic ^c
<i>d</i> -Amphetamine	1	0	1
5	0	+	1/5
6	1/10	++	1/2
10	1/10	++	1/2
11	1/15	+	1/3
12	1/15	+	1/3
16	0	0	1/15
17	1/30	+	1/3
19	1/100	+	1/4

^a Hyperactivity is expressed relative to amphetamine (10 mg/kg) and as for the 1-2 hr period following administration of 25 mg/kg ip the "initial tranquilization" period referred to the first 40 min. ^b Initial tranquilization observed for 0-40 min. ^c Anorexigenic action was measured for a period of 5 days, following administration of 10 mg/kg of the drug twice daily, 12 hr apart. The results are expressed relative to a 1 mg/kg dose of amphetamine, also administered twice daily.

self. Although less active than the NO₂, the formyl derivatives exhibited both stimulating and anorexigenic activity. This series of compounds possess anorexigenic activity at doses somewhat less than those required to produce moderate central stimulation. It would appear that the anorexigenic action of the carbamates occurs with considerably less central stimulation than with amphetamine.

Gross behavioral effects in mice were observed for all of the carbamate esters. Sedative action and central stimulation appeared in various degrees depending upon the particular amine and structure of the leaving alkyl group. There was always a delayed sedative action prior to the onset of central stimulation although

(16) M. J. Kolbezen, R. L. Metcalf, and T. R. Fukuto, *J. Agr. Food Chem.*, **2**, 864 (1954); L. W. Dittert and T. Higuchi, *J. Pharm. Sci.*, **52**, 852 (1963); I. Christenson, *Acta Chem. Scand.*, **18**, 904 (1964); M. L. Bender and R. B. Homer, *J. Org. Chem.*, **30**, 3975 (1965).

(17) P. Adams and F. A. Baron, *Chem. Rev.*, **65**, 567 (1965).

(18) M. Matzner, R. P. Kurkijy, and R. J. Cotter, *ibid.*, **64**, 645 (1964).

the degree of hyperactivity caused by certain drugs such as **1**, **4**, **5**, **13**, **16**, and **19** was slight. It is not known for certain whether the initial depressant action is centrally mediated and attributable to the carbamate as such.

The 5 carbamate esters of amphetamine and ephedrine with nitrophenyl leaving groups showed the highest degree of activity. A contributory factor to the higher activity of the nitrophenolic carbamates is their ease of hydrolysis in aqueous media, which is attributable to the electron-withdrawing character of the NO₂ substituent. It also appears that the *o*-NO₂ participates in this hydrolysis to some degree because of the higher overall activity of *o*-nitrophenyl *d*-amphetaminecarbamate compared with the *m*- and *p*-NO₂ isomers. The *o*-carbomethoxyphenyl carbamates **3**, **9**, and **14** also show somewhat higher activities as compared with the phenyl, *p*-methoxyphenyl, and lactyl carbamates. Although the *o*-carbomethoxyphenyl carbamates are not as active as the nitrophenyl carbamates, the toxicity of salicylate as a leaving group is somewhat lower.

Direct evidence that the carbamate esters were hydrolyzed after entering the brain was obtained from radiotracer studies. Two carbamate esters of *d*-amphetamine labeled with ¹⁴C in the amine side chain α to the ring were found to enter the brain (Table III).

TABLE III

DISTRIBUTION OF ¹⁴C-LABELED COMPOUNDS IN MOUSE BRAIN

Drug administered	Dose		Time of sacrifice, hr	Amount in brain, μ g	
	μ Ci	μ g		Carbamate	<i>d</i> -Amphetamine
¹⁴ C- <i>d</i> -Amphetamine ^a	1.20	200	0.5		12.0
	21 ^b	0.70	200	0.5	3.0
22 ^c	0.74	200	1.0	2.0	4.0
			0.5	3.3	2.8
			1.0	3.8	

^a Sulfate salt. ^b *o*-Carbomethoxyphenyl [¹⁴C]-*d*-amphetaminecarbamate. ^c *o*-Nitrophenyl [¹⁴C]-*d*-amphetaminecarbamate.

Although it is not known whether the hydrolyzed products were formed entirely in the brain, it is reasonable to assume that the carbamates can be hydrolyzed in the brain, particularly since they can be shown to do so *in vitro*. Both the delayed onset and the greater duration of action of the carbamates support the notion that the hydrolytic product is required for their stimulating action.

The present studies suggest that latentiation of centrally active amines appears feasible *via* carbamoylation. A degree of selectivity in modifying the drug action is possible through the leaving alkyl group, which temporarily protects the amine against deactivation by MAO and increases lipophilicity. Further manipulation is possible through leaving groups containing substituents which can act on the carbamate ester linkage by electron withdrawal or by neighboring group participation to facilitate hydrolysis. The concepts developed in this study are being extended to the latentiation of biogenic and other amines.

Experimental Section

Melting points were determined using a Thomas-Hoover melting point apparatus and are corrected. Analyses were per-

formed by Midwest Microlab, Inc., Indianapolis, Ind. 46226. Where analyses are indicated only by symbols of the elements analytical results obtained for those elements were within \pm 0.4% of the theoretical values. Specific rotations were measured on a Franz Schmidt and Haensch, Berlin, S., polarimeter. Radioscanning of the Hc sheets was done with an atomic scanner, Atomic Accessories, Inc., New York, N. Y.

Pharmacological Measurements.—The measurement of motor activity was carried out as described elsewhere.¹⁹ For the results described in Table II a dose of 25 mg/kg (in propylene glycol) administered ip was employed, although similar effects were obtained in the 10–50 mg/kg dose range. Usually the animals were sedated the first 30–50 min prior to the onset of increased motor activity. During this initial period the animals also exhibited decreased skeletal muscle tonus and were generally less responsive to noxious stimuli. The increase in motor activity was observable 1–2 hr following drug administration. With **6**, **10**, and **11** some pupillary dilation and piloerection were observed during the 1–2 hr period. The anorexigenic action was measured over a period of 5 days utilizing a dose of 10 mg/kg administered ip twice daily at intervals of 12 hr. Amphetamine (1 mg/kg twice daily) was used as a reference standard. The food consumption of both the experimental and control groups was measured.

Radiochemical Drug Study.—The drugs in Table III were given to mice in a single injection at 10 mg/kg ip in propylene glycol. One-half and one hour later the mice were sacrificed and their brains removed and analyzed. An average of 4 animals were used to obtain each value and agreement was good. The original drug penetrating into the brain and the hydrolyzed products were separated and identified using tic and autoradiography. The values given in Table III are for micrograms present in the whole brain. All mice were extremely hyperactive at time of sacrifice. Onset of activity was within 3–5 min with amphetamine and 8–10 min with **21** and **22**.

Chloroformic Acid Esters. *o*-Formylphenyl Chloroformate.—A soln of 97.6 g (0.8 mole) of salicylaldehyde, 96.7 g (0.8 mole) of PhNMe₂, and 8.1 g (0.08 mole) of Et₃N in 100 ml of solvent A (10% PhMe and 90% C₆H₆) was added slowly with cooling to 90 g (0.9 mole) of COCl₂ in 400 ml of solvent A. After stirring for 6 hr, H₂O was added cautiously. The organic phase was washed with 5% HCl and H₂O, and dried (CaCl₂). The solvent was removed and the residue crystallized nicely from hexane as white needles, 97.5 g (66%), mp 80.5–81.5°. The product is unstable and slowly decomposes while standing in a closed container, apparently eliminating HCl. It can be stored for only a short time. *Anal.* (C₈H₇ClO₂) C, H, Cl.

The other chloroformic acid esters were prepared similarly from COCl₂, an alcohol, and with an organic base as a scavenger for HCl and as a catalyst. The various products, their yields and a brief description are given here: *p*-methoxyphenyl chloroformate,²⁰ 89% colorless oil, bp 85–88° (3 mm), n_D^{20} 1.558; α -carboethoxyethyl chloroformate,²¹ 66%, colorless lachrymatory oil, bp 55–57° (4 mm), n_D^{20} 1.558, 1.566; *o*-carbomethoxyphenyl chloroformate,^{22,23} 65%, colorless oil, bp 106–108° (3 mm), n_D^{20} 1.558, 1.577; *p*-nitrophenyl chloroformate,^{26,27} 42%, white crystals (from *l*-Pr₃O), mp 65–72° (reported mp 80–81°). *m*-Nitrophenyl chloroformate²⁸ and *o*-nitrophenyl chloroformate^{26,27} were prepared by standard methods and were used immediately without isolation. Phenyl chloroformate was a commercial product.

Carbamate Esters.—The carbamate esters in Table I were prepared by methods similar to the examples given below, with minor variations.

Phenyl *l*-Ephedrinecarbamate.—A mixture of 6.6 g (0.04 mole) of *l*-ephedrine in 40 ml of CHCl₃ and 2.33 g (0.022 mole) of Na₂CO₃ in 12 ml of H₂O was cooled to 10°. A soln of 6.66 g (0.044 mole) of phenyl chloroformate in 10 ml of CHCl₃ was added

119) L. G. Abood, in "Drugs Affecting the Central Nervous System," A. Burger, Ed., Vol. 2, M. Dekker, New York, N. Y., 1968, p 127.

(20) M. J. Zabik and R. D. Schuetz, *J. Org. Chem.*, **32**, 300 (1967).

(21) J. Thiele and F. Dent, *Justus Liebigs Ann. Chem.*, **302**, 245 (1898).

(22) A. Einhorn and A. von Bagh, *Chem. Ber.*, **43**, 322 (1910).

(23) N. Sueiu, *Rev. Roum. Chim.*, **11**, (6), 745 (1966).

(24) Y. Iwakura and A. Nabeya, *J. Org. Chem.*, **25**, 1118 (1960).

(25) R. E. Oesber, W. Broker, and W. A. Cook, *J. Amer. Chem. Soc.*, **47**, 2609 (1925).

(26) L. C. Ralford and G. O. Luman, *ibid.*, **56**, 1586 (1934).

(27) Y. Iwakura, Y. Isidzuka, and T. Saito, *Nippon Kagaku Zasshi*, **76**, 1408 (1955); *Chem. Abstr.*, **51**, 17,750 (1957).

over 10 min with good stirring, keeping the temp below 20°. The cooling bath was removed and stirring was continued for 1 hr, after which time the evolution of CO₂ stopped. The CHCl₃ layer was washed with 20 ml of 5% HCl, then H₂O and dried (Na₂SO₄) and the solvent evaporated. The oily product was crystallized from CCl₄-hexane, decolorizing with charcoal, to give 6.7 g (59%) of small white needles, mp 68–70°. Recrystallization from pet ether (bp 60–110°) gave an analytical sample: mp 72–74°; $\lambda_{\text{max}}^{\text{CHCl}_3}$ 3.0 (OH), 5.87 (C=O).

***o*-Carbomethoxyphenyl *l*-Ephedrinecarbamate.**—A mixture of 4.95 g (0.03 mole) of *l*-ephedrine in 50 ml of CHCl₃ and 1.7 g (0.016 mole) of Na₂CO₃ in 20 ml of H₂O was cooled to 5°. A soln of 6.66 g (0.031 mole) of *o*-carbomethoxyphenyl chloroformate in 20 ml of CHCl₃ was added over 10 min with good stirring, keeping the temp below 15°. The cooling bath was removed and the reaction mixture was stirred for 1.5 hr. The organic layer was washed with 20 ml of 5% HCl and 20 ml of H₂O and dried (Na₂SO₄) and the solvent evaporated. The solid product was recrystallized from CHCl₃-hexane, decolorizing with charcoal, to give 7.66 g (74%) of fine long white needles, mp 95–96°.

Bis(phenyl *dl*-*p*-hydroxy- α -methylphenethylcarbamate)piperazine Salt.—A stirred mixture of 6.96 g (0.03 mole) of *dl*-*p*-hydroxyamphetamine-HBr, 3.4 g (0.032 mole) of Na₂CO₃, 30 ml of H₂O and 60 ml of CHCl₃ was cooled in an ice bath. A soln of 4.85 g (0.031 mole) of phenyl chloroformate in 20 ml of CHCl₃ was added over 5 min. After stirring at room temp for 2 hr, the partially soluble free phenolic amine gradually reacted and dissolved in the CHCl₃ giving 2 clear layers. The CHCl₃ layer was washed with 2 × 50 ml of 5% HCl, dried (Na₂SO₄), and the solvent evaporated to give 8 g of a colorless oil. This was dissolved in 150 ml of C₆H₆ and 4 g of piperazine was added to the warm soln, followed by 80 ml of hexane. The resulting ppt was recrystallized from C₆H₆ giving 4.9 g (52%) of white crystalline product, mp 115–116°, that analyzed correctly for the bis salt.

***o*-Nitrophenyl *dl*- α -Methylphenethylcarbamate.**—A soln of 13.9 g (0.1 mole) of *o*-nitrophenol in 30 ml of solvent A was added to a solution of 17 g (0.17 mole) of COCl₂ in 150 ml of solvent A at 0° with no noticeable rise in temperature. However, an exothermic reaction occurred during the gradual addition of 10.1 g (0.1 mole) of Et₃N in 30 ml of solvent A and a solid formed. After stirring out of the cooling bath for 1 hr, 100 ml of H₂O was added cautiously to dissolve the ppt salt. The organic phase

was washed with H₂O, dried (CaCl₂), and evapd to about 50 ml of yellow soln containing the *o*-nitrophenyl chloroformate, from which the carbamate was prepared.

A mixture of 9.02 g (0.245 mole) of *dl*-amphetamine sulfate and 5.3 g (0.05 mole) of Na₂CO₃ in 70 ml of H₂O with 50 ml of CHCl₃ was cooled to 0°. The *o*-nitrophenyl chloroformate solution was added with stirring and cooling over 10 min. The cooling bath removed and the mixture was stirred for 2 hr whereupon both layers became clear yellow at pH 7–7.5. The organic phase was washed with 50 ml of H₂O, two 50-ml portions of 5% HCl, H₂O again and then dried (Na₂SO₄). Most of the solvent was evaporated and upon the addition of hexane a solid formed. Recrystallization from CCl₄ gave 12.3 g (84%) of crude product, mp 80–86°. Further recrystn from dry *i*-Pr₂O gave analytically pure product, mp 89–90°, as very light yellow needles.

***o*-Carbomethoxyphenyl [¹⁴C]-*d*- α -Methylphenethylcarbamate (21).**—A mixture of 20 mg (0.055 μ mole) of [¹⁴C]-*d*-amphetamine sulfate (6.0 μ Ci/mg), 35 mg (0.33 μ mole) of Na₂CO₃, 40 mg (0.2 μ mole) of *o*-carbomethoxyphenyl chloroformate, 1 ml of H₂O, and 3 ml of CHCl₃ was shaken for 0.5 hr at room temperature. The mixture was dild with 10 ml of CHCl₃ and 4 ml of H₂O. The organic phase was washed with dil HCl and H₂O, dried (Na₂SO₄), and evaporated to an oil that crystallized. After trituration with pet ether the solid was recrystallized from CCl₄ and pet ether, eventually giving 30 mg (87%) of white needles with a specific activity of 3.5 μ Ci/mg. Tlc on an Eastman Kodak 6060 chromatogram silica gel sheet using C₆H₆ developer indicated that the product was homogeneous with an *R*_f 0.21 corresponding exactly to that of the unlabeled material. A radioscan of the strip showed a single peak corresponding to the visual spot, indicating radioactive homogeneity as well.

***o*-Nitrophenyl [¹⁴C]-*d*- α -methylphenethylcarbamate (22)** was prepd from 25 mg (0.068 μ mole) of [¹⁴C]-*d*-amphetamine sulfate (6.0 μ Ci/mg) by procedures similar to those in the preceding experiment and those given here for the prepn of the unlabeled product. A 30-mg yield of long, yellow-tinted needles was eventually obtained with a specific activity of 3.7 μ Ci/mg. Tlc on Eastman Kodak 6060 Chromagram silica gel sheet using C₆H₆ as the developer showed the product to be homogeneous with an *R*_f 0.54 corresponding exactly to that of the unlabeled product. A radioscan indicated that the product was radiochemically pure and corresponded to the visual spot.

Peripheral Inhibition of Thyroxine by Thiohydantoins Derived from Amino Acids^{1a}

JOSEPH V. MARX, DAN A. RICHERT, AND W. W. WESTERFELD*

Department of Biochemistry, State University of New York, Upstate Medical Center, Syracuse, New York 13210

Received November 29, 1969

A number of 3-phenyl-2-thiohydantoins with nonpolar substituents at the 5 position inhibited the peripheral effect of T₄ as measured by the liver GPD response. Like TU and PTU, they also increased the PBI in rats given exogenous T₄. These PTH derivatives of valine, leucine, norleucine, isoleucine, etc., were very weak goiterogens which had only little or no effect on thyroid weight or radioiodine uptake.

Barker, *et al.*^{2a} originally showed that a given dose of T₄ did not restore the metabolic rate as well in rats made hypothyroid with TU as it did in surgically thyroidectomized rats. In a recent summary by de Escobar and del Rey^{2b} the minimal daily T₄ require-

ment was found to be increased twofold in rats receiving TU, PTU, or MTU when a variety of biological endpoints was used (metabolic rate, liver GPD, goiter prevention, pituitary basophilia, normal plasma TSH, and suppression of thyroid ¹³¹I release). These thiouracils, in addition to preventing the biosynthesis of thyroid hormones, inhibited the peripheral action of T₄.^{3,4}

This paper reports a study of the inhibition of the GPD response to exogenous T₄ by a series of 5-alkyl-3-phenyl-2-thiohydantoins derived from amino acids by the Edman reaction^{5,6} in an attempt to establish the

* To whom correspondence should be addressed.

(1) (a) This study was aided by Grant No. 5-RO1-AM-09106 from the National Institute of Arthritis and Metabolic Diseases of the National Institutes of Health, and Grant No. 5-RO1-CA-01852-17 from the National Cancer Institute, U. S. Public Health Service. (b) Abbreviations used are: GPD, mitochondrial α -glycerophosphate dehydrogenase (EC 1.1.99.5); MTU, 6-methyl-2-thiouracil; PBI, protein-bound iodine; PTH-AA, 5-alkyl-3-phenyl-2-thiohydantoin derived from amino acid; PTU, 6-*n*-propyl-2-thiouracil; T₄, 3,5,3',5'-tetraiodo-L-thyronine, L-thyroxine; TSH, thyroid stimulating hormone; TU, 2-thiouracil; RAIU, radioiodine uptake.

(2) (a) S. Barker, C. Kiely, and H. Lipner, *Endocrinology*, **45**, 624 (1949); (b) C. Morreale de Escobar and F. Escobar del Rey, *Recent Progr. Horm. Res.*, **23**, 87 (1967).

(3) W. Ruegamer, W. W. Westerfeld, and D. A. Richert, *Endocrinology*, **75**, 908 (1964).

(4) W. Hoffman, D. Richert, and W. W. Westerfeld, *ibid.*, **78**, 1189 (1966).

(5) P. Edman, *Acta Chem. Scand.*, **4**, 277 (1950).

(6) W. Westerfeld, R. Doisy, and D. Richert, *J. Nutr.*, **78**, 393 (1962).