

Table VIII. 2-Carboxymethylbenzenesulfonamides 8

No.	R	Mp, °C	Yield, %	Recrystn solvent	Formula	Analyses
8a ^a	CH ₃	174 ^b	78	H ₂ O	C ₉ H ₁₁ NO ₄ S	C, H, N, S
8b	<i>i</i> -C ₂ H ₅	115–117	45	H ₂ O	C ₁₁ H ₁₃ NO ₄ S	C, H, N, S
8c ^a	CH ₂ C ₆ H ₅	124 ^c	86	H ₂ O	C ₁₅ H ₁₅ NO ₄ S	C, H, N
8d	<i>o</i> -C ₆ H ₄ Cl	146–148	95	EtOH–H ₂ O	C ₁₄ H ₁₂ ClNO ₄ S	C, H, N
8e	<i>m</i> -C ₆ H ₄ Cl	159–163	86	H ₂ O	C ₁₄ H ₁₂ ClNO ₄ S	Cl, N, S
8f	<i>p</i> -C ₆ H ₄ Cl	128–130	94	EtOH–H ₂ O	C ₁₄ H ₁₂ ClNO ₄ S	Cl, N, S
8g	<i>p</i> -C ₆ H ₄ SO ₂ NH ₂	202–204	84	H ₂ O	C ₁₄ H ₁₄ N ₂ O ₆ S ₂	C, H, S

^a See ref 9. ^b Lit.⁹ 158–164°, uncrystallized. ^c Lit.⁹ 107–109°.

hr, evaporation of this solution and treatment of the residue as above gave finally 0.27 g (2.8%) of the crystalline isomer 1e (mixture melting point, tlc, and ir).

(b) 1a (5.91 g, 0.03 mol) and PCl₅ (6.25 g, 0.03 mol) were mixed and heated at 160° for 30 min, then POCl₃ was removed *in vacuo*, and the residue was triturated with a little CHCl₃, collected, and crystallized (C₆H₆–CCl₄) to give 4.55 g (70%) of 4-chloro-1*H*-2,3-benzothiazine 2,2-dioxide (11), mp 172–174°. *Anal.* (C₈H₆ClNO₂S) C, H, Cl, N, S.

A suspension of 11 (3.23 g, 0.015 mol) in 2-PrOH (20 ml) was refluxed for 4 hr and the obtained solution let stand in the cold to complete separation of the product, that was collected, washed, dried, and recrystallized (2-PrOH) to afford 2.6 g (72%) of 10, mp 182–183°, identical (analysis, tlc, mixture melting point, and ir) with the product described under (a).

Supplementary Material Available. A listing of the pharmacological data for weakly active or inactive compounds will appear following these pages in the microfilm edition of this volume of the journal. Photocopies of the supplementary material from this paper only or microfiche (105 × 148 mm, 20× reduction, negatives) containing all of the supplementary material for the papers in this issue may be obtained from the Journals Department, American Chemical Society, 1155 16th St., N.W., Washington, D. C. 20036. Remit check or money order for \$3.00 for photocopy or \$2.00 for microfiche, referring to code number JMED-73-1133.

References

- (1) E. Sianesi, G. Bonola, R. Pozzi, and P. Da Re, *Chem. Ber.*, 104, 1880 (1971) (paper 3).
- (2) A. A. Rubin, F. E. Roth, and M. M. Winbury, *Nature (London)*, 192, 176 (1961).
- (3) J. K. Wales, S. V. Krees, A. M. Grant, J. K. Viktora, and F. W. Wolff, *J. Pharmacol. Exp. Ther.*, 164, 421 (1968).
- (4) W. Kunz, *Fortschr. Arzneimittelforsch.*, 3, 383 (1961), and references cited therein.
- (5) (a) M. Bauer, *Arzneim.-Forsch.*, 15, 806 (1965); (b) S. Hayao, W. G. Stricker, B. M. Phillips, and H. Fujimori, *J. Med. Chem.*, 11, 1246 (1968).
- (6) A. Burger in "Medicinal Chemistry," 3rd ed, A. Burger, Ed., Wiley, New York, N. Y., 1970, p 77.
- (7) C. Elison, E. J. Lien, A. P. Zinger, M. Hussain, G. L. Tong, and M. Golden, *J. Pharm. Sci.*, 60, 1058 (1971).
- (8) G. Satzinger, U. S. Patent 3,408,346 (1968); *Chem. Abstr.*, 70, 87826 (1969).
- (9) J. G. Lombardino and E. H. Wiseman, *J. Med. Chem.*, 14, 973 (1971).
- (10) E. Sianesi, R. Redaelli, M. Bertani, and P. Da Re, *Chem. Ber.*, 103, 1992 (1970).
- (11) S. Irwin in "Animal and Clinical Pharmacologic Technique in Drug Evaluation," J. H. Nodine and P. E. Siegler, Ed., Year Book Publishing, Chicago, Ill., 1964, p 46.
- (12) J. E. P. Toman, E. A. Swinyard, and L. S. Goodman, *J. Neurophysiol.*, 9, 231 (1946).
- (13) P. Ruggli, *Helv. Chim. Acta*, 14, 544 (1931).

Synthesis of the Thyrotropin-Releasing Hormone Enantiomer and Some Diastereoisomers and *in Vitro* Studies of Their Biological Activity

G. Flouret,*† R. Morgan, R. Gendrich,

Organic Chemistry and Biochemistry Departments, Research Division, Abbott Laboratories, North Chicago, Illinois 60064

J. Wilber, and M. Seibel

Northwestern University, School of Medicine, Chicago, Illinois 60611. Received March 15, 1973

The isomers of TRH with DDD, DLL, LDL, and LLD configurations were prepared unequivocally by coupling pentachlorophenyl pyroglutamate of the required configuration with the appropriate histidylprolinamide diastereoisomer. The dipeptide amide was made by condensation of *tert*-butyloxycarbonylhistidine with prolinamide mediated by dicyclohexylcarbodiimide, removal of the blocking group in trifluoroacetic acid, and treatment of the peptide amide hydrochloride with a basic ion exchanger. Protected peptide intermediates and the final isomers were purified by chromatography on silica gel. Formation of diastereoisomers of the protected dipeptide was detected by nmr and tlc techniques. The LDL isomer was approximately 2–3% as active as TRH for TSH release in rat hemipituitaries *in vitro*, the DLL and LLD isomers were about 0.1% active, and the DDD isomer was inactive. None of these analogs inhibited TSH release *in vitro*.

The recent availability of synthetic thyrotropin-releasing hormone (TRH) and analogs has made possible for the first time studies of TRH structure-activity relationships. Burgus and collaborators¹ have demonstrated the critical

importance of the N-terminal pyrrolidone group and the C-terminal amide for full biological expression of TRH, and Bowers and associates² have shown that of several C-terminal analogs only those containing proline were inactivated by serum. Experiments with labeled TRH suggested that the C-terminal carboxamide group hydrolysis by serum is a first metabolic step^{2,3} in the inactivation

*Address correspondence to this author at Northwestern University, The Medical School, Department of Physiology, Chicago, Ill. 60611.

mechanism which probably plays an important physiologic role.

In the present studies, the DDD, LLD, LDL, and DLL stereoisomers of TRH were synthesized and their biological activity was investigated in an *in vitro* system, where the variable of serum inactivation is not operative. These isomers were prepared with the expectation that they would be substantially resistant to enzymatic cleavage and that they might preserve TRH activity but with longer duration of action. Furthermore, the importance of TRH as a synthetic hormone of potential commercial interest made it desirable to ascertain whether epimers that might arise during synthesis would have inhibitory or other undesirable properties.

The TRH stereoisomers were made by a method similar to the one employed previously for a synthesis of TRH,⁴ with the selection of a D- or an L-amino acid derivative as required. The desired TRH isomer was obtained by coupling pentachlorophenyl pyroglutamate of the required configuration with the appropriate His-Pro-NH₂ isomer. The free dipeptide amide was made by the condensation of *N*^α-*tert*-butyloxycarbonylhistidine⁵ with prolinamide in the presence of DCC,⁶ removal of the blocking group in acid, and treatment of the peptide hydrochloride with a basic ion-exchange resin. Peptide intermediates were purified by chromatography on silica gel.⁴

For the synthesis of *tert*-butyloxycarbonyl-D-histidine, *tert*-butyloxycarbonylhistidine (Boc) was refluxed in MeOH solution for 24 hr. Under these conditions, the imidazolyl-*tert*-butyloxycarbonyl group which is more susceptible to alcoholysis was cleanly removed. The *tert*-butyloxycarbonylhistidine (Boc) was made by treating D-histidine with excess *tert*-butyloxycarbonyl azide⁷ in TEA, DMF, and H₂O for 24 hr. The dipeptide Boc-His-Pro-NH₂ was conveniently analyzed for optical purity by thin-layer chromatography (tlc) with 15% MeOH-CHCl₃ (solvent A) as the eluting solvent and Cl₂-toluidine, ninhydrin, or Pauly's spray as the detection method. Under these conditions the LL isomer and its enantiomer, the DD isomer, had an *R*_f of 0.30 while the DL and LD isomers had an *R*_f of 0.25. A combination of LL and DL isomers, for example, was usually resolved as two spots resembling a figure 8.

Similarly, in the nuclear magnetic resonance (nmr) spectrum (C₅D₅N) of this dipeptide, the chemical shift for the *tert*-butyloxycarbonyl methyl proton was identical with the LL and DD isomers (δ 1.50) and with the DL and LD isomers (δ 1.40). A combination of LL and DL isomers was easily detectable by nmr analysis. All synthetic preparations of Boc-His-Pro-NH₂ showed after column chromatography the presence of tlc fractions suggesting that diastereoisomers had formed during coupling. An nmr of such fractions revealed the CH₃-proton absorption due to the "incorrect" isomer. Appropriate pooling of clean chromatographic fractions led to optically pure dipeptide isomers according to the tlc and nmr criteria. In one purification, where 33% MeOH-CHCl₃ was employed to elute dipeptide material, indiscriminate elution of isomers occurred. The nmr of the "mixed" Boc-His-Pro-NH₂ revealed the presence of 5.3% of the "wrong" isomer; this value was obtained by dividing the peak height for diastereoisomer by the sum of the peak height for diastereoisomer plus the peak height for the expected isomer. One possible explanation for the formation of the "wrong" isomer is that the presence of the free imidazole group in *tert*-butyloxycarbonylhistidine leads to some epimerization of the α position of this amino acid during carbodiimide activation of the carboxyl group due to the formation of a base-sensitive acylimidazole.⁸ Recently, Windridge re-

ported a study on racemization during synthesis of histidine dipeptides and he too concluded that *tert*-butyloxycarbonylhistidine can undergo racemization to the extent of forming in one case approximately 5% of D isomer in DMF with DCC as the coupling agent.⁹ Our results thus seem consistent with his findings. A reexamination of Boc-His-Pro-NH₂ (LL), previously made by the solid-phase technique,⁴ revealed that 5.9% wrong diastereoisomer (DL?) was also obtained as detected by nmr analysis. The free dipeptide His-Pro-NH₂ was analyzed in the Beckman amino acid analyzer with both the short and long columns, but it was always eluted unresolved as one peak when mixtures of LL and DL dipeptide were applied; thus, this technique which is useful for dipeptides¹⁰ could not be successfully applied to the resolution of these dipeptide amides. The final TRH isomers (DDD, LLL, LLD, LDL, and DLL) could not be differentiated by tlc in several solvent systems. The nmr spectra of these isomers were identical except for a small difference in the splitting of imidazolyl protons, which was very weak and thus could not be used to detect the presence of incorrect isomers. The high-resolution mass spectra of the TRH isomers showed the correct molecular ion as calculated for TRH (*m/e* 362.1702) and were consistent with the structure of the tripeptide amide. The intensity of several peaks showed subtle differences, however, which can only be ascribed to conformational effects but were of no value for the quantitative estimation of isomers. Therefore, no convenient physical method was found to check directly the optical purity of these analogs.

The thyrotropin (TSH)-releasing activity of the four TRH analogs was investigated *in vitro* with rat hemipituitaries. TSH release was determined by a specific radioimmunoassay for rat TSH.¹¹ The observed potency of "LDL" TRH was 2-3% of native TRH activity. In contrast, the LLD and DLL isomers possessed only approximately 0.1% of native TRH activity and the DDD isomer was inactive. None of the four analogs inhibited TSH release by TRH.

It is of interest that the biological potencies of these analogs have corresponded quantitatively with their binding affinities for a specific anterior pituitary receptor.¹²

Experimental Section†

D-Prolinamide Hydrochloride. To a suspension of D-proline (5.75 g, 50 mmol) in MeOH (50 ml) was added SOCl₂ (6.5 ml) dropwise with the temperature being kept at 0° with external cooling. The reaction mixture was stirred at room temperature for 2 hr and then heated at reflux for 2 hr. The resulting solution was evaporated *in vacuo* to an oil which was dissolved in MeOH and the solution evaporated again to an oil, the process being repeated three times. The final oil was dried over P₂O₅ and KOH pel-

†A Thomas-Hoover apparatus was used for melting point determinations in capillary tubes. Optical rotations were measured in 1-dm polarimeter tubes with Hilger-Watts polarimeter reading with a precision of $\pm 0.01^\circ$. Where analyses are indicated only by symbols of the elements, analytical results were obtained for the elements within $\pm 0.4\%$ of the theoretical value. For tlc or electrophoretic analysis several samples ranging between 20 and 100 μ g of peptide were usually employed to reveal impurities. Where tlc was used to determine purity of intermediates and products, silica gel G plates were used and were developed with one of the following solvent combinations: A, 15% MeOH-CHCl₃; B, MeOH-CHCl₃ (1:2); C, MeOH-CHCl₃ (2:1); with Pauly and/or Cl₂-toluidine color sprays. Where the ninhydrin color reaction was the detection method of choice, the Boc intermediate was exposed to HCl vapors in a chromatographic chamber for 10 min prior to spraying. For electrophoresis, Brinkmann precoated thin-layer cellulose sheets were employed with Py-AcOH buffer, pH 6.3 (25-45 min). The nmr spectrum of the dipeptides was obtained at 100 MHz using a Varian Associates HA-100 spectrometer. Chemical shifts are reported in parts per million from internal TMS. Mass spectra were recorded on an AEI MS-902 mass spectrometer. The following abbreviations were used throughout the text: DCC = dicyclohexylcarbodiimide; DCU = dicyclohexylurea; TFA = trifluoroacetic acid; TEA = triethylamine; Py = pyridine.

lets *in vacuo*. The dried oil (8.4 g) was dissolved in MeOH (50 ml) and saturated with NH₃. The insoluble crystals which formed (mostly NH₄Cl) were removed by filtration. The clear solution was evaporated *in vacuo* to an oil which was redissolved in MeOH and the resulting solution was evaporated to an oil (three times). The residual oil was dissolved in MeOH (50 ml) and the solution was acidified with concentrated HCl until it turned Hydrion paper to a color corresponding to a pH of 3. The solution was evaporated to an oil which was dissolved in EtOH (150 ml) and the solvent was removed *in vacuo*, the whole process being repeated three times. The resulting oil was dissolved in EtOH (75 ml) and the clear solution was allowed to cool slowly at room temperature. The crystals were collected and washed with EtOH-Et₂O (1:1) and finally with Et₂O; the first crop weighed 4.95 g, mp 169–172°; the second crop weighed 1.25 g, mp 165–168°. Both crops were combined and recrystallized from EtOH (35 ml) yielding 4.97 g; mp 178–180°; $[\alpha]^{23D} + 69.5^\circ$ (c 2, EtOH) [lit.¹³ L isomer mp 182°; $[\alpha]^{25D} - 70.7^\circ$ (c 2, EtOH)]. Anal. (C₅H₁₁ClN₂O) C, H, N.

tert-Butyloxycarbonyl-D-histidine. To a solution of D-histidine (20.9 g) in H₂O (75 ml) was added TEA (42 ml), tert-butyloxycarbonyl azide (43 ml), and DMF (150 ml). The reaction mixture was stirred overnight at room temperature. The progress of the reaction was followed by tlc with solvent C and ninhydrin color reaction. As some histidine and tert-butyloxycarbonylhistidine were still detectable, the reaction mixture was treated with additional tert-butyloxycarbonyl azide (15 ml) and TEA (14 ml). After 2 hr saturated citric acid solution was added to a pH of 7 and the reaction mixture was evaporated *in vacuo* to an oil. This oil was dissolved in a small volume of H₂O and the solution was acidified with Et₂O (three times). The aqueous phase was acidified with saturated citric acid solution to pH 2–3 and extracted with CHCl₃ (six times). The extracts were combined and washed twice with a small volume of H₂O and saturated NaCl solution (three times) and dried (Na₂SO₄). Evaporation of the CHCl₃ solution *in vacuo* led to an oil. Analysis of this oil by tlc as described above showed a slow-moving minor component with R_f 0.3 equivalent to that of tert-butyloxycarbonyl-L-histidine and a major fast-moving component (R_f 0.6). The crude Boc-His (Boc) was refluxed in MeOH solution (1000 ml). After 24 hr tlc analysis indicated almost complete disappearance of the fast-moving component. The reaction mixture was concentrated to an oil which was dissolved in EtOH (100 ml). The EtOH solution was concentrated to a lower volume and allowed to cool slowly. The crystals which formed were collected, washed with EtOH, and dried over P₂O₅ in a desiccator; the yield was 12.1 g; mp 197–199°; $[\alpha]^{25D} - 14.1^\circ$ (c 2, H₂O). Anal. (C₁₁H₁₇N₃O₄) C, H, N.

tert-Butyloxycarbonyl-L-histidine. This material was made by the method described above for the D isomer. The yields and properties were similar: mp 198° dec; $[\alpha]^{25D} + 14^\circ$ (c 2, H₂O). A commercial sample obtained from Schwarz BioResearch, Inc., had a rotation of $[\alpha]^{25D} + 14^\circ$ (c 2, H₂O).

D-Pyroglutamic Acid. A solution of D-glutamic acid (10 g) was refluxed in H₂O (250 ml) for 2 days. Analysis by tlc solvent C and Cl₂-toluidine color reaction shows two components, a minor component at the origin with R_f equivalent to glutamic acid and a faster component with R_f equivalent to that of L-pyroglutamic acid. The reaction mixture was evaporated to a residue which was treated with EtOH and the solvent was removed *in vacuo*. The latter treatment was repeated three times and the product obtained was treated with boiling EtOH (100 ml) and the resulting suspension filtered. The EtOH-insoluble material was mostly D-glutamic acid (1.47 g) as shown by tlc. The EtOH solution was concentrated to approximately 40 ml and allowed to crystallize overnight. The crystals formed were collected and washed with EtOH to yield a first crop of 4.50 g, mp 156–159°, $[\alpha]^{23D} + 11.5^\circ$ (c 2, H₂O) and from the mother liquor a second crop, 1.15 g, mp 154–157°, $[\alpha]^{23D} + 11.5^\circ$ (c 2, H₂O), for a combined yield of 64%, or 75% based on consumed D-glutamic acid [lit.¹⁴ L isomer mp 160–161°; $[\alpha]^{24D} - 11.8^\circ$ (c 1.4, H₂O)]. Anal. (C₅H₇NO₃) C, H, N.

Pentachlorophenyl D-Pyroglutamate. This derivative was prepared essentially as described for the L isomer:⁴ mp 195–198°; $[\alpha]^{23D} - 21.2^\circ$ (c 2, DMF) [lit.⁴ L isomer mp 196–199°; $[\alpha]^{26D} + 21^\circ$ (c 2, DMF)]. Anal. (C₁₁H₆Cl₅N₁O₃) C, H, N.

Boc-L-His-D-Pro-NH₂. To a solution of tert-butyloxycarbonyl-L-histidine (2.55 g, 10 mmol) in 20 ml of DMF was added D-prolinamide hydrochloride (1.65 g, 11 mmol). The resulting clear solution was cooled to 0° and treated with TEA (1.5 ml) and DCC (2.06 g, 10 mmol). The reaction mixture was then stirred for 40 hr

at room temperature, the DCU which precipitated removed by filtration, and the filtrate evaporated to an oil. For purification this product was chromatographed on silica gel with 5% MeOH-CHCl₃ as the eluent to remove by-products and 12% MeOH-CHCl₃ to elute the desired dipeptide. All effluents were monitored by tlc with solvent A (Cl₂-toluidine or ninhydrin). All desired fractions (R_f 0.25) containing none of the slightly faster moving component (R_f 0.30, corresponding probably to DD isomer) were combined. The yield of crude product was 1.91 g. A methanolic solution of this product was treated with Rexyn 201 (OH⁻), filtered, and evaporated to a residue which crystallized from MeOH-EtOH. The crystals formed were collected and recrystallized from MeOH yielding the analytical sample: mp 133–137°; $[\alpha]^{23D} + 53.2^\circ$ (c 1, H₂O), +58.0° (c 1, DMF). Anal. (C₁₆H₂₅N₅O₄) C, H, N. An nmr spectrum (C₅D₅N) was consistent with the structure and the (CH₃)₃C-proton peak (δ 1.40) was a sharp singlet showing no evidence of the presence of DD isomer.

DL Isomer. This isomer crystallized from MeOH: mp 129–133°; $[\alpha]^{23D} - 53.5^\circ$ (c 1, H₂O), -57.4° (c 1, DMF). The nmr was identical with that of the LD isomer with the CH₃ protons showing as a singlet (δ 1.40): R_f^A 0.25. Anal. (C₁₆H₂₅N₅O₄) C, H, N.

DD Isomer. This isomer was obtained by lyophilization of chromatographically pure fractions: $[\alpha]^{23D} + 50^\circ$ (c 1, H₂O); $[\alpha]^{24D} - 26.1^\circ$ (c 1, DMF). The nmr spectrum was consistent with the structure and the CH₃ protons showed as a singlet (δ 1.50): R_f^A 0.30. Anal. (C₁₆H₂₅N₅O₄) C, H, N.

LL Isomer. This isomer was prepared as a fluffy powder obtained by lyophilization of chromatographically pure fractions: $[\alpha]^{24D} - 49^\circ$ (c 1, H₂O), +26.4° (c 1, DMF). The nmr spectrum was consistent with the structure, the CH₃ protons showing as a singlet (δ 1.50): R_f^A 0.30. Anal. (C₁₆H₂₅N₅O₄) C, H, N.

D-<Glu-L-His-L-Pro-NH₂. A solution of Boc-L-His-L-Pro-NH₂ (LL) (703 mg) in 10 ml of TFA was allowed to stand for 15 min at room temperature. At this time tlc with solvent B revealed disappearance of the protected dipeptide amide spot (Cl₂-toluidine) and the appearance of His-Pro-NH₂ as a single spot, R_f^A 0.1 (ninhydrin +). The TFA was removed from the reaction mixture by evaporation to dryness followed by the addition of Et₂O and evaporation to dryness three additional times. The residue was dried over P₂O₅ and KOH *in vacuo* and then dissolved in MeOH, and the solution was treated with Rexyn 201 (OH⁻) and filtered. The filtrate was evaporated under reduced pressure and the residue dried over P₂O₅ *in vacuo*. The dried amino dipeptide was dissolved in 2 ml of DMF and treated with pentachlorophenyl pyroglutamate (755 mg). The resulting clear solution was allowed to stand overnight at room temperature. The reaction mixture was chromatographed on silica gel employing 15% MeOH-CHCl₃ to elute by-products, the desired product being eluted with 33% MeOH-CHCl₃. Progress of the reaction and chromatographic separation was followed by tlc with solvent B (with Cl₂-toluidine and ninhydrin color reactions); fractions containing the TRH (DLL) isomer were combined and evaporated to an oil which crystallized slowly. Recrystallization from MeOH-EtOH gave 480 mg; mp 231–233°; $[\alpha]^{24D} - 66.4^\circ$ (c 1, H₂O). Thin-layer electrophoresis on Brinkman precoated thin-layer cellulose sheets with pyridine-AcOH buffer, pH 6.3 (25–45 min), showed this isomer to have only one component (Pauly color reaction), R_f^C 0.30. Anal. (C₁₆H₂₂N₆O₄) C, H, N.

LDL Isomer. This isomer was obtained by the method described above, except for lyophilization of the final product which yielded an amorphous powder: $[\alpha]^{24D} - 41.2^\circ$. The nmr (C₅D₅N) was consistent with the structure. The tlc (R_f^C 0.30) and tlc patterns showed one component. Anal. (C₁₆H₂₂N₆O₄) C, H, N.

LLD Isomer. This isomer was similarly obtained as an amorphous powder after lyophilization: $[\alpha]^{24D} + 43.8^\circ$ (c 1, H₂O). The nmr (C₅D₅N) was consistent with the structure. The tlc (R_f^C 0.30) and tlc patterns showed one component. Anal. (C₁₆H₂₂N₆O₄·0.4H₂O) C, H, N.

DDD Isomer. This isomer was obtained as an amorphous powder after lyophilization: $[\alpha]^{24D} + 66.4^\circ$ (c 1, H₂O) [lit. L isomer $[\alpha]^{25D} - 65.5^\circ$ (c 1, H₂O)];⁴ $[\alpha]^{25D} - 64.8^\circ$ (c 1, H₂O)];¹⁵ also $[\alpha]^{25D} - 50^\circ$ (c 1.5, H₂O)]¹⁶. The nmr (C₅D₅N) was consistent with the structure and identical with that for the LLL isomer. The tlc (R_f^C 0.30) and tlc patterns showed one component with mobility identical with TRH. Anal. (C₁₆H₂₂N₆O₄·0.7H₂O) C, H, N.

Biologic Activity of TRH and Diastereoisomeric Analogs of TRH. Thyrotropin (TSH)-releasing activity of TRH and analogs was investigated *in vitro*. Single rat hemipituitaries were preincubated for 4 hr in 2 ml of Krebs-Ringer bicarbonate (KRB)




Figure 1.

medium to permit TSH release to equilibrate at a basal level. Six tissues were used per test substance. The medium was then removed and discarded and replaced by 1 ml of KRB for the succeeding 1-hr control period. Medium was then replaced again with media containing TRH or analogs at concentrations indicated in Figure 1 (abscissa). Medium TSH concentrations during control and experimental periods were determined by a specific radioimmunoassay for rat TSH.¹¹ Incremental changes in medium TSH are expressed as per cent of control [TSH (hr 2)/TSH (hr 1) \times 100].

Native TRH yielded a sigmoidal dose-response function between 0.020 and 0.2 μ g/ml (Figure 1). LDL-TRH gave a parallel dose-response function shifted to the right. The observed potency of LDL-TRH was about 2-3% of native TRH activity. In contrast, the analogs LLD- and DLL-TRH possessed only about 0.1% of native TRH activity. DDD-TRH failed to exhibit any TSH-releasing activity with medium concentrations as high as 1 μ g/ml.

In order to test for inhibitory properties, hemipituitaries were incubated with a submaximal TRH stimulus (100 μ g/ml) and substimulating concentrations of the four analogs.

The LDL analog did not inhibit thyrotropin release at a concen-

tration 20 times that of the hormone nor did the LLD, DLL, or DDD isomers at concentrations 1000 times that of TRH.

Acknowledgment. The authors wish to thank Victor Rauschel and associates for elemental analyses, Richard Egan for nmr spectra, and Dr. W. Cole for discussions concerning this work.

References

- (1) R. Burgus, T. F. Dunn, D. Desiderio, W. Vale, and R. Guillemin, *C. R. Acad. Sci., Paris*, **269**, 226 (1969).
- (2) C. Y. Bowers, A. Weil, J. K. Chang, H. Sievertsson, F. Enzmann, and K. Folkers, *Biochem. Biophys. Res. Commun.*, **40**, 683 (1970).
- (3) T. W. Redding and A. V. Schally, *Endocrinology*, **89**, 1075 (1971).
- (4) G. Flouret, *J. Med. Chem.*, **13**, 843 (1970).
- (5) A. Loffet, *Experientia*, **23**, 406 (1967).
- (6) J. C. Sheehan and G. P. Hess, *J. Amer. Chem. Soc.*, **77**, 1067 (1955).
- (7) R. Schwyzer, P. Sieber, and H. Kappeler, *Helv. Chim. Acta*, **42**, 2622 (1959).
- (8) H. C. Beyerman, W. Maassen Van Den Brink, F. Weygand, A. Prox, W. König, L. Schmidhammer, and E. Nintz, *Recl. Trav. Chim. Pays-Bas*, **84**, 213 (1965).
- (9) G. C. Windridge and E. C. Jorgensen, *Intra-Sci. Chem. Rep.*, **5** (4), 375 (1971).
- (10) J. M. Manning and S. Moore, *J. Biol. Chem.*, **243**, 5591 (1968).
- (11) J. F. Wilber and R. D. Utiger, *Endocrinology*, **81**, 145 (1967).
- (12) J. F. Wilber and M. J. Seibel, *ibid.*, in press.
- (13) J. P. Greenstein and M. Winitz in "Chemistry of the Amino Acids," Vol. 3, Wiley, New York, N. Y., 1961, p 2198.
- (14) W. E. Hanby, S. G. Waley, and J. Watson, *J. Chem. Soc.*, 3239 (1950).
- (15) G. Flouret, A. Alter, and R. Gendrich, *J. Label. Compounds*, **8**, 53 (1972).
- (16) H. C. Beyerman, P. Kranenberg, and J. L. M. Syrier, *Recl. Trav. Chim. Pays-Bas*, **90**, 791 (1971).

Effect of Simple Amino Acid Replacements on the Biological Activity of Luteinizing Hormone-Releasing Hormone

David H. Coy,* Esther J. Coy, and Andrew V. Schally

Veterans Administration Hospital and Department of Medicine, Tulane University School of Medicine, New Orleans, Louisiana 70146.
Received April 27, 1973

Six decapeptides having structures based on the LH-RH sequence, <Glu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH₂>, have been synthesized by modifications of the solid-phase technique *via* protected peptide intermediates. Thus, pyroglutamic acid, tryptophan, serine, glycine, proline, and glycine amide residues in the LH-RH sequence were replaced by leucine, tyrosine, threonine, isoleucine, alanine, and glycine dimethylamide, respectively. The compounds were assayed *in vivo* and found to possess the following LH-releasing potencies relative to pure, natural LH-RH: [Leu¹]-LH-RH, 0.003%; [Tyr³]-LH-RH, 0.13%; [Thr⁴]-LH-RH, 19%; [Ile⁶]-LH-RH, 0.034%; [Ala⁹]-LH-RH, 0.8%; and [(Gly-dimethylamide)¹⁰]-LH-RH, 14%.

Recent studies^{1,2} on the effects of replacement of tyrosine in position 5 of LH-RH by closely related amino acids with aromatic side chains established that the phenolic hydroxyl group does not contribute significantly toward biological activity. This paper reports methods of synthesis and biological characteristics of decapeptides containing replacements for many of the other residues.

Of the amino acids with functional side chains in LH-RH, cyclic pyroglutamic acid was replaced by leucine, tryptophan by tyrosine, and serine by the closely related threonine. While this work was in progress, data on the threonyl peptide, synthesized by fragment condensation, were reported by Fujino, *et al.*³ The glycine residue in the sterically crowded center of the molecule and the confor-

mationally important proline at position 9 were particularly intriguing and were replaced by isoleucine and alanine, respectively. Finally, the importance of the amide part of C-terminal glycine was examined by replacing it with a dimethylamide group.

Synthesis. All protected peptides were prepared on a resin support by versions of the Merrifield solid-phase method⁴ which have been described in part previously.^{1,2,5} Protected peptide amides were cleaved from the resin by treatment with ammonia in methanol or, in the case of [(glycine-dimethylamide)¹⁰]-peptide, with dimethylamine.

In all the peptides the following groups were used for the protection of sensitive side chains during synthesis: arginine, *N*^C-tosyl; serine, threonine; and tyrosine, *O*-ben-