

- (2) N. C. Brown, *J. Mol. Biol.*, **59**, 1 (1971).
- (3) M. M. Neville and N. C. Brown, *Nature (London), New Biol.*, **240**, 80 (1972).
- (4) G. W. Bazill and J. D. Gross, *Nature (London), New Biol.*, **240**, 82 (1972).
- (5) K. B. Gass, R. L. Low, and N. R. Cozzarelli, *Proc. Nat. Acad. Sci. U. S.*, **70**, 103 (1973).
- (6) J. M. Mackenzie, M. M. Neville, G. E. Wright, and N. C. Brown, *Proc. Nat. Acad. Sci. U. S.*, **70**, 512 (1973).
- (7) N. R. Cozzarelli and R. L. Low, *Biochem. Biophys. Res. Commun.*, **51**, 151 (1973).
- (8) E. F. Mooney, "An Introduction of ^{19}F NMR Spectroscopy," Heydon and Son, Ltd., London, 1970, p 39.
- (9) E. F. Mooney, ref 8, p 45.
- (10) J. Clements, J. d'Ambrosio, and N. C. Brown, *J. Biol. Chem.*, submitted for publication.
- (11) N. C. Brown and J. E. Clements, *Fed. Proc., Fed. Amer. Soc. Exp. Biol.*, **33**, 1420 (1974).
- (12) R. L. Low, S. A. Rashbaun, and N. R. Cozzarelli, *Fed. Proc., Fed. Amer. Soc. Exp. Biol.*, **33**, 1420 (1974).
- (13) A. I. Vogel, "A Textbook of Practical Organic Chemistry," 3rd ed, Longmans, Green and Co., Ltd., London, 1956, p 636.
- (14) S. Levy and G. Schultz, *Justus Liebigs Ann. Chem.*, **210**, 133 (1881).
- (15) E. Sarauw, *Justus Liebigs Ann. Chem.*, **209**, 93 (1881).
- (16) A. I. Vogel, ref 13, p 745.
- (17) R. B. Wickner, B. Ginsberg, I. Berkower, and J. Hurwitz, *J. Biol. Chem.*, **247**, 489 (1972).

Synthesis and Biological Activity of Selected 2-Substituted 6-(β -D-Ribofuranosyl)oxazolo[5,4-*d*]pyrimidin-7-ones

Vemanna D. Patil, Dean S. Wise, Leroy B. Townsend,*

Department of Biopharmaceutical Sciences and Department of Chemistry, University of Utah, Salt Lake City, Utah 84112

and Alexander Bloch

Department of Experimental Therapeutics, Roswell Park Memorial Institute, Buffalo, New York 14203. Received June 7, 1974

Several 2-substituted 6-(β -D-ribofuranosyl)oxazolo[5,4-*d*]pyrimidin-7-ones have been prepared by condensation of the appropriate silylated heterocycles with 2,3,5-tri-*O*-acetyl- β -D-ribofuranosyl bromide and the subsequent removal of blocking groups from the carbohydrate moiety with methanolic ammonia. The site of ribosylation was established by comparison of the uv spectra of the nucleosides with that of an appropriate model compound. The anomeric configuration was determined by pmr spectroscopy. These nucleosides have been evaluated for their inhibitory activity against leukemia L1210 and *Escherichia coli* cells *in vitro* and for their effects on leukemia L1210 growth *in vivo*. Among these compounds the 2-methyl, 2-ethyl, and 2-propyl derivatives markedly inhibited the *in vitro* growth of both cell types, the inhibitory concentrations ranging from 5×10^{-7} to 8×10^{-4} M. Only the 2-methyl derivative was significantly active against leukemia L1210 *in vivo* with a dose of 200 mg/kg/day \times 5, causing a 31% increase in the life span of the tumor-bearing mice.

The characterization¹ of uric acid ribonucleoside isolated from bovine erythrocytes as 3-(D-ribofuranosyl)uric acid has generated interest in the chemical synthesis^{2,4} of bicyclic heterocyclic nucleosides with the glycosidic linkage on a nitrogen atom in the pyrimidine ring rather than in the five-membered ring portion of the molecule. 3-(β -D-Ribofuranosyl)adenosine (isoadenosine) was prepared and found⁵ to inhibit the growth of various tumor cell lines *in vitro* and *in vivo*, as well as exhibiting some activity against adeno III virus in culture. This interest has been further stimulated by the recent isolation⁶ of 7-(D-ribose)pyrazolo[3,4-*d*]pyrimidine-4,6-dione (oxoallopurinol ribonucleoside) from the urine of patients treated with allopurinol and the report⁷ that oxoallopurinol ribonucleoside, presumably as the corresponding 5'-phosphate derivative, inhibits pyrimidine biosynthesis *de novo*. These findings have prompted us to synthesize a number of 2-substituted 6-(β -D-ribofuranosyl)oxazolo[5,4-*d*]pyrimidin-7-ones and to study their biological effects.^{8,9}


Results and Discussion

Chemistry. The trimethylsilylation of 2-substituted oxazolo[5,4-*d*]pyrimidin-7-ones¹⁰ with hexamethyldisilazane in the presence of a catalytic amount of ammonium sulfate was accomplished in excellent yield. These silyl derivatives (1a-f) were subsequently condensed¹¹ with 2,3,5-tri-*O*-acetyl-D-ribofuranosyl bromide in benzene at reflux temperature in the presence of mercuric oxide-mercuric bromide to furnish good yields of the blocked nucleosides 2a-f as syrups. Thin-layer chromatography of these syrups revealed the presence of only one nucleoside in each reaction mixture.

Deacetylation of 2a-f was accomplished with methanol-

ic ammonia to furnish the nucleosides 3a-f (Table I). That complete deblocking had occurred without ring opening of the labile oxazole ring[†] was established by uv and pmr spectroscopy. The site of ribosylation was established as N₆ by a comparison of the uv spectral data (Table II) obtained for 3a and the data reported¹² for 2,6-dimethyloxazolo[5,4-*d*]pyrimidin-7-one. The anomeric configuration of the deblocked nucleosides could not be assigned on the basis of pmr spectral data (Table II) since the peaks assigned to the anomeric proton (H_{1'}) of 3b and 3f revealed a coupling constant ($J_{1',2'}$) of a magnitude which precluded an unequivocal anomeric assignment.¹³ Therefore, the isopropylidene derivatives 4a and 4b were prepared from 3b and 3f using a standard procedure,¹⁴ and the 5'-*O*-tosyl-2',3'-*O*-isopropylidene derivative 6 was prepared by tosylation of 4a (Table I). The pmr spectra of 4a, 4b, and 6 revealed singlets for their respective anomeric protons, which now allowed an unequivocal assignment of the β configuration to these nucleosides (Table III). This anomeric assignment is further substantiated by utilizing a recently reported¹⁵ criterion for determining anomeric configuration for β -D-ribofuranosides. The difference between the chemical shifts of the two methyl signals for the 2',3'-*O*-isopropylidene group is between 0.18 and 0.22 ($\Delta\delta$) for β -D-ribofuranosides and 0.0 and 0.10 ($\Delta\delta$) for α -D-ribofuranosides. The pmr spectral data in Table III corroborate the β assignment for the nucleosides reported herein.

[†]The oxazole ring is, however, susceptible to ring opening. This was established by the frequent pmr spectroscopic monitoring of a reaction mixture of the 2-ethyl nucleoside derivative 3b dissolved in NaOD; at least two additional compounds can be formed.

Table I. Chemical-Physical Data for Some Oxazolo[5,4-d]pyrimidin-7-one 6-Ribonucleosides


| Compd no. | R | R' | R'' | Mp, °C | % yield (purified) | Formula ^a |
|-----------|---|-------------------------------------|-------------------------------------|---------|--------------------|--|
| 3a | -CH ₃ | H | H | 181.5 | 42 | C ₁₁ H ₁₃ N ₃ O ₆ |
| 3b | -CH ₂ CH ₃ | H | H | 172 | 55 | C ₁₂ H ₁₅ N ₃ O ₆ |
| 3c | -CH ₂ CH ₂ CH ₃ | H | H | 191-192 | 32 | C ₁₃ H ₁₇ N ₃ O ₆ ·H ₂ O |
| 3d | -CH(CH ₃)CH ₃ | H | H | 197-198 | 16 | C ₁₃ H ₁₇ N ₃ O ₆ |
| 3e | -CH ₂ CH ₂ CH ₂ CH ₃ ^b | H | H | 164-165 | 37 | C ₁₄ H ₁₉ N ₃ O ₆ |
| 3f | -C ₆ H ₅ | H | H | 258-259 | 43 | C ₁₆ H ₁₅ N ₃ O ₆ |
| 4a | -CH ₂ CH ₃ | Isopropylidene | H | 165-166 | 41 | C ₁₅ H ₁₉ N ₃ O ₆ |
| 4b | -C ₆ H ₅ | Isopropylidene | H | 248 | 45 | C ₁₉ H ₁₉ N ₃ O ₆ |
| 5 | -CH ₂ CH ₃ | -C(=O)C ₆ H ₅ | -C(=O)C ₆ H ₅ | 110 | 73 | C ₃₃ H ₂₇ N ₃ O ₉ ·0.5H ₂ O |
| 6 | -C ₂ H ₅ | Isopropylidene | Tosyl | 75-80 | 25 | C ₂₂ H ₂₅ N ₃ O ₈ S |

^a All compounds were analyzed for C, H, and N and the results were within $\pm 0.4\%$ of theoretical values. ^b The base was prepared by the procedure outlined in ref 10; mp 168-169°.

Table II. Spectral Data for Some 2-Substituted 6-(β -D-Ribofuranosyl)oxazolo[5,4-d]pyrimidin-7-ones^{a,b}


| Compd | Uv, λ max, nm ($\epsilon \times 10^3$) | | | Pmr | | |
|-------|--|--|--|--------------------------------------|---------------------|-------------------------|
| | MeOH | pH 1 | pH 11 | δ H ₁ ^c | J_{H_1, H_2} , Hz | δ H ₆ |
| 3a | 236 (17.5), 246.5 (sh, 14.3), 274 (8.9) | 240 (16.4), 246 (sh, 14.7), 270 (8.7) | 240 (15.2), 246 (sh, 14.1), 295 (11.9) | 6.18 | 2.6 | 8.98 |
| 3b | 240 (25.6), 246 (sh, 21.1), 275 (12.8) | 241 (24.4), 247 (sh, 22.0), 271 (12.2) | 241 (21.7), 246 (sh, 20.5), 290 (14.5) | 6.16 | 3.0 | 8.96 |
| 3c | 240 (14.9), 247 (sh, 13.4), 275 (8.4) | 241 (14.0), 247 (sh, 13.1), 272 (7.5) | 241 (14.0), 246 (sh, 13.1), 290 (10.6) | 6.22 | 3.4 | 8.84 |
| 3d | 239 (17.8), 247 (sh, 15.1), 274 (9.2) | 241 (16.1), 247 (sh, 14.9), 271 (8.7) | 241 (16.5), 247 (sh, 15.5), 290 (11.5) | 6.18 | 2.6 | 9.01 |
| 3e | 240 (15.8), 246 (sh, 13.9), 274 (8.8) | 240 (15.0), 246 (sh, 13.8), 270 (8.6) | 241 (15.1), 246 (sh, 14.0), 291 (10.8) | 6.20 | 2.8 | 9.01 |
| 3f | 286 (sh, 26.9), 295 (29.6), 303 (sh, 28.4) | 286 (sh, 29.4), 293 (32.8), 304 (sh, 30.4) | 305 (36.2) | 6.10 | 2.7 | 8.80 |
| 6 | 275 (10.0) | 238 (29.6), 246 (31.8), 280 (20.7) | 238 (36.8), 243 (sh, 36.1), 278 (20.1) | | | |

^a Ultraviolet spectra were obtained on a Beckman DK-2 spectrophotometer. ^b Pmr spectra were recorded in DMSO-*d*₆ on a Varian A-56/60 spectrometer using TMS as an internal standard. ^c In all cases H₁ peak shape was a doublet.


Biological. The effect of some of the newly prepared nucleosides on the *in vitro* growth of leukemia L1210 and *Escherichia coli* K₁₂ cells is shown in Table IV. Substitution at the 2 position with a methyl, ethyl, or propyl group gives rise to a growth inhibitory activity in both the L1210 and *E. coli* systems, whereas an isopropyl or phenyl group at the 2 position results in compounds which are inactive against L1210 cells but, in the case of the phenyl

derivative, shows some activity against *E. coli*. The corresponding 2-substituted heterocycles are inactive in these test systems, except for the isopropyl derivative which inhibits the growth of *E. coli* by 50% at 3×10^{-5} M.

Preliminary evaluation of the activity of these compounds against leukemia L1210 *in vivo* showed that, at 200 mg/kg/day \times 5, the 2-methyl-substituted nucleoside 3a increased the life span of the tumor-bearing mice by



- 1a. R = CH₃
 b. R = CH₂CH₃
 c. R = CH₂CH₂CH₃
 d. R = CH(CH₃)₂
 e. R = CH₂CH₂CH₂CH₃
 f. R = C₆H₅



31%. At this dose, none of the other analogs were significantly active.

Since these nucleosides can be viewed either as derivatives of a pyrimidine nucleus substituted with a cyclic moiety at positions 5 and 6, or as purine analogs with the carbohydrate moiety attached to the pyrimidine moiety, it was of interest to determine whether, in the cells, these compounds act as purine or pyrimidine antagonists. Con-

Table III. Pmr Spectral Data of 2',3'-*O*-Isopropylidenes of Some Oxazolo[5,4-*d*]pyrimidin-7-one 6-Ribonucleosides

| Compd | δ CH ₃ | $\Delta\delta^a$ | δ H _{1'} | $J_{H_1', H_2'}$, Hz |
|-------|--------------------------|------------------|--------------------------|-----------------------|
| 4a | 1.36, 1.54 | 0.18 | 6.18 | <1 |
| 4b | 1.37, 1.58 | 0.21 | 6.30 | s ^b |
| 6 | 1.30, 1.50 | 0.20 | 6.12 | s |

^a $\Delta\delta$ is the difference between the chemical shifts observed for the methyl protons of the isopropylidene group. ^b s = singlet.

sequently, an inhibition analysis¹⁶ was carried out, which showed that the inhibition of growth exerted, for example, by the 2-ethyl nucleoside **3b**, is prevented by pyrimidines but not by purines. Thus, the inhibition of the growth of L1210 cells by 1×10^{-4} M **3b** was completely prevented by uridine or cytidine at 1×10^{-4} M. Smaller concentrations of these metabolites decreased the inhibition to a proportionately smaller extent. 2'-Deoxyuridine or 2'-deoxycytidine was approximately ten times less effective in preventing the inhibition than were the corresponding ribonucleosides and thymidine (at 1×10^{-5} M) which appeared to act like a product of the inhibited path, reversing the inhibition to the same extent (80%) at all concentrations of inhibitor. The reversal pattern obtained in *E. coli* was quite similar to that seen in the leukemic cells, thymidine again providing a product effect, whereas the extent of reversal provided by the other pyrimidine nucleosides was approximately proportional to their concentration. Thus, it appears that the compounds act as pyrimidine antagonists, exerting their inhibitory activity along the metabolic path followed by the pyrimidines, rather than as purine analogs which could mimic the feedback effect exerted by purine nucleosides and nucleotides in enzymes concerned with *de novo* purine synthesis.¹⁷

Experimental Section

General Procedure for the Preparation of 2-Substituted 6-(β -D-Ribofuranosyl)oxazolo[5,4-*d*]pyrimidin-7-ones (3a-f). The silylated heterocycle 1a-f (0.04 mol) [prepared by heating the heterocycle¹⁰ in hexamethyldisilane in the presence of a catalytic amount of (NH₄)₂SO₄ at reflux temperature] was dissolved in 50 ml of anhydrous benzene and then added to a suspension of HgBr₂ (7.5 g) and HgO (7.5 g) in 250 ml of azeotropically dried benzene. 2,3,5-Tri-*O*-acetyl-D-ribofuranosyl bromide (prepared from 10.5 g of 1,2,3,5-tetra-*O*-acetyl-D-ribofuranose) dissolved in 50 ml of anhydrous benzene was added to the reaction mixture. The suspension was heated at reflux temperature for 20 hr, protected from moisture. The reaction mixture was cooled, the mercury salts were removed by filtration through a Celite pad, and the pad was washed with chloroform (3 \times 50 ml). The filtrate and washings were combined and evaporated *in vacuo* to afford a syrup. This syrup was dissolved in chloroform (350 ml); the solution was washed successively with a 30% aqueous solution of KI (3 \times 50 ml), a cold saturated sodium carbonate solution (4 \times 50 ml), and distilled water (4 \times 50 ml) and was then dried over anhydrous sodium sulfate. The drying agent was removed by filtration and the chloroform mixture evaporated under reduced pressure to afford a syrup. The syrup was treated with methanol, which had been previously saturated with ammonia at 0° (400 ml) and the solution was allowed to stand at room temperature for 20 hr with occasional shaking. The solution was filtered and the filtrate evaporated *in vacuo* to yield a solid material which was collected by filtration. Recrystallization from methanol yielded pure 2-substituted 6-(β -D-ribofuranosyl)oxazolo[5,4-*d*]pyrimidin-7-ones.

2-Ethyl-6-(2,3,5-tri-*O*-benzoyl- β -D-ribofuranosyl)oxazolo[5,4-*d*]pyrimidin-7-one (5). The same ribosylation procedure was followed as outlined above, with the following reactants being used. The trimethylsilyl derivative formed from 3.3 g of 2-ethyl-oxazolo[5,4-*d*]pyrimidin-7-one was dissolved in 40 ml of benzene and added to a suspension of mercuric oxide (5 g) and mercuric

Table IV. Effect of Some 2-Substituted 6-(β -D-Ribofuranosyl)oxazolo[5,4-d]pyrimidin-7-ones on the *in Vitro* Growth of Leukemia L1210 and *E. coli* Cells

| Compd | 2-Substituent | Concn (M) for 50% growth inhibition of | |
|-------|--|--|--------------------------------------|
| | | Leukemia L1210 cells | <i>E. coli</i> K ₁₂ cells |
| 3a | -CH ₃ | 5×10^{-5} | 8×10^{-4} |
| 3b | -CH ₂ CH ₃ | 7×10^{-6} | 6×10^{-6} |
| 3c | -CH ₂ CH ₂ CH ₃ | 3×10^{-5} | 5×10^{-7} |
| 3d | -CH(CH ₃) ₂ | $>10^{-4}$ | $>10^{-3}$ |
| 3f | -C ₆ H ₅ | $>10^{-4}$ | 4×10^{-4} |

bromide (5 g) in 250 ml of benzene and this was then added to 2,3,5-tri-*O*-benzoyl-D-ribofuranosyl bromide (prepared from 10 g of 1-*O*-acetyl-2,3,5-tri-*O*-benzoyl-D-ribofuranose) dissolved in 40 ml of anhydrous benzene. After work-up of the reaction mixture the chloroform was evaporated to a hard foam residue which was recrystallized from benzene to yield 5 (9 g, 72.8%).

2-Ethyl-6-(2,3-*O*-isopropylidene- β -D-ribofuranosyl)oxazolo[5,4-d]pyrimidin-7-one (4a). To 450 ml of dry acetone was added 1.6 ml of 2,2-dimethoxypropane and 1.5 ml of 70% perchloric acid. The mixture, protected from moisture, was stirred at room temperature for 5 min and then 1.3 g of 3b was added in one portion. Pyridine (1.6 ml) was added after the mixture had been stirred for 45 min. The volume was reduced to 50 ml *in vacuo* and 23 ml of 10% aqueous sodium carbonate was added before the remaining acetone was removed. Cold water (40 ml) was then added and the solution was allowed to stand at 5° for 12 hr. The white crystalline solid which separated from the solution was collected by filtration and was washed with a small amount of cold water. Recrystallization from methanol afforded 0.60 g (40.7%) of 4a.

2-Phenyl-6-(2,3-*O*-isopropylidene- β -D-ribofuranosyl)oxazolo[5,4-d]pyrimidin-7-one (4b). Compound 4b was prepared in the same manner as 4a by treatment of 3f (1.4 g) with 1.6 ml of 2,2-dimethoxypropane and 1.5 ml of 70% perchloric acid, followed by 1.6 ml of pyridine. This procedure furnished 4b (0.7 g, 44.8%) as white needles.

2-Ethyl-6-(2,3-*O*-isopropylidene-5-*O*-*p*-toluenesulfonyl- β -D-ribofuranosyl)oxazolo[5,4-d]pyrimidin-7-one (6). 2-Ethyl-6-(2,3-*O*-isopropylidene- β -D-ribofuranosyl)oxazolo[5,4-d]pyrimidin-7-one (4b, 0.405 g) was dissolved in pyridine (8 ml) and *p*-toluenesulfonyl chloride (0.286 g) was added to this solution. The reaction mixture was stored in the dark at 5° for 36 hr and was then poured onto ice water (100 ml). The aqueous mixture was extracted with chloroform (2 \times 5 ml); the organic layer was washed with 1 M H₂SO₄ (2 \times 25 ml) and then with water until the aqueous layer was neutral. The chloroform solution was dried over sodium sulfate and its volume was reduced to 10 ml. Methanol (20 ml) was added and the solution was then evaporated to dryness and the residue dried in a vacuum desiccator to afford 0.150 g (25%) of 6.

Biological. The procedures used for the evaluation of the biological effects of the nucleosides have been described previously.¹⁸

Acknowledgments. This work was supported by Research Grants CA-11147 and CA-12585 from the National Cancer Institute, National Institutes of Health, U. S.

Public Health Service. One of the authors (V. D. Patil) wishes to thank Dr. G. R. Revankar for his encouragement.

References

- (1) A. R. Davis, E. B. Newton, and S. R. Benedict, *J. Biol. Chem.*, **54**, 595 (1922); R. Falconner and J. M. Gulland, *J. Chem. Soc.*, 1369 (1939); M. M. Jeyewska, B. Gorykowski, and T. Sawicka, *Acta Biochim. Pol.*, **14**, 71 (1967).
- (2) H. S. Forrest, D. Hatfield, and J. M. Lagowski, *J. Chem. Soc.*, 963 (1961); R. Lohrmann, J. M. Lagowski, and H. S. Forrest, *ibid.*, 451 (1964).
- (3) C. L. Schmidt, W. J. Rusho, and L. B. Townsend, *Chem. Commun.*, 1515 (1971).
- (4) C. L. Schmidt and L. B. Townsend, *J. Heterocycl. Chem.*, **10**, 687 (1973).
- (5) K. Gerzon, I. S. Johnson, G. B. Boder, J. C. Cline, P. J. Simpson, C. Speth, N. J. Leonard, and R. A. Laursen, *Biochim. Biophys. Acta*, **119**, 445 (1966).
- (6) T. A. Krenitsky, G. B. Elion, R. A. Strelity, and G. H. Hitchings, *J. Biol. Chem.*, **212**, 2675 (1967).
- (7) T. D. Beardmore and W. W. Kelley, *J. Lab. Clin. Med.*, **78**, 696 (1971).
- (8) L. B. Townsend, V. D. Patil, and A. Bloch, *Proc. Amer. Ass. Cancer Res.*, **12**, 74 (1971).
- (9) V. D. Patil, D. S. Wise, and L. B. Townsend, *J. Heterocycl. Chem.*, **10**, 277 (1973).
- (10) V. D. Patil and L. B. Townsend, *J. Heterocycl. Chem.*, **8**, 503 (1971).
- (11) E. Wittenberg, *Chem. Ber.*, **101**, 1095 (1968).
- (12) Y. Ohtsuka, *Chem. Pharm. Bull.*, **18**, 2242 (1970).
- (13) L. B. Townsend in "Synthetic Procedures in Nucleic Acid Chemistry," Vol. II, W. W. Zorbach and R. S. Tipson, Ed., Interscience, New York, N.Y., 1973, Chapter 7.
- (14) G. R. Revankar and L. B. Townsend, *J. Heterocycl. Chem.*, **7**, 117 (1970), and references cited therein.
- (15) J.-L. Imbach, J.-L. Barascut, B. Kam, B. Rayner, and C. Tapiero, *J. Heterocycl. Chem.*, **10**, 1069 (1973).
- (16) W. Shive, *Amer. N. Y. Acad. Sci.*, **52**, 1212 (1950).
- (17) E. Besnick and K. Blatchford, *Biochim. Biophys. Acta*, **81**, 150 (1964).
- (18) A. Bloch, G. Dutschman, B. L. Currie, R. K. Robins, and M. J. Robins, *J. Med. Chem.*, **16**, 294 (1973).