

- (6) J. A. Marshall and S. F. Brady, *J. Org. Chem.*, **35**, 4068 (1970).  
 (7) E. A. Steck, R. P. Brundage, and L. T. Fletcher, *J. Amer. Chem. Soc.*, **73**, 1117 (1953).  
 (8) G. Bachmann, British Patent 1,168,291 (Oct 22, 1969).  
 (9) J. R. Cummings, A. M. Welter, J. L. Grace, Jr., and L. M. Lipchuck, *J. Pharmacol. Exp. Ther.*, **161**, 88 (1968).  
 (10) (a) G. Bachmann and A. Amann, U. S. Patent 3,475,431 (Oct 28, 1969); (b) G. Bachmann and A. Amann, British Patent 1,164,139 (Sept 17, 1969).  
 (11) J. P. English, R. C. Clapp, Q. P. Cole, and J. Krapcho, *J. Amer. Chem. Soc.*, **67**, 2263 (1945).  
 (12) E. A. Steck and F. C. Nachod., *J. Amer. Chem. Soc.*, **79**, 4408 (1957).  
 (13) R. E. Lutz, P. S. Bailey, C.-K. Dien, and J. W. Rinker, *J. Amer. Chem. Soc.*, **75**, 5039 (1953).  
 (14) H. M. Holava, Jr., and R. A. Partyka, *J. Med. Chem.*, **14**, 262 (1971).

## 6-(Substituted phenyl)-5-substituted-4,5-dihydro-3(2H)-pyridazinones. Antihypertensive Agents


Francis J. McEvoy and George R. Allen, Jr.\*

Lederle Laboratories, A Division of American Cyanamid Company, Pearl River, New York 10965. Received July 19, 1973

The preparation and antihypertensive properties of a series of 6-(substituted phenyl)-5-substituted-4,5-dihydro-3(2H)-pyridazinones are described. The structure-activity relationship in this series is discussed further. The consistent antihypertensive activity of the 6-(alkylaminophenyl) compounds and their acyl derivatives is noteworthy.

Previous reports from these laboratories have described the preparation and antihypertensive effects of a series of 6-(substituted phenyl)-4,5-dihydro-3(2H)-pyridazinones (I).<sup>1</sup> These studies indicated that the compounds having amino, acylamino, cyano, and halogen substituents on the phenyl ring were among those having the more interesting antihypertensive activity. Moreover, this action persisted for a longer duration in those compounds also possessing a 5-methyl substituent. In the present paper we describe the preparation and biological properties of additional members of this series. Specifically, we have prepared those compounds of structure I in which the phenyl substituent is alkylamino, *N*-alkylacylamino, and dimethylamino. Moreover, the effect of other 5 substituents on activity was investigated. A cursory examination of the effect on activity caused by alteration of substituents at the 2 position was made,<sup>†</sup> and certain 6-(*o*-substituted phenyl) derivatives also were prepared.


**Chemistry.** Most of the compounds of type I (see Table I) having 6-(alkylaminophenyl) (1-12), 6-(dimethylaminophenyl) (13-18), 6-(ortho-substituted phenyl) (30, 32, 33), and 5-alkyl and aryl (41-51) substituents were prepared by treatment of the appropriate  $\gamma$ -keto acid or  $\gamma$ -keto ester with hydrazine (eq 1).<sup>2</sup> The 6-(dimethylaminophenyl) compounds also could be prepared by Eschweiler-Clarke alkylation of the corresponding 6-(aminophenyl)-4,5-dihydropyridazinone, but this procedure is limited to those compounds with a 2 substituent, and the yield is poor (*cf.* 16).



The yields for those dihydropyridazinones prepared by the procedure of eq 1 (methods A and A<sub>1</sub>) generally were excellent, but the 6-(ortho-substituted phenyl) derivatives 30, 32, and 33 were exceptions. Moderate yields were observed in the preparation of those dihydropyridazinones


<sup>†</sup>A more comprehensive study of this parameter was made by Drs. Goldman, Lin, and Stodja in these laboratories.

lacking a 5 substituent (30 and 33). However, the yield declined precipitously in the instance of the 6-(*o*-nitrophenyl)-5-methyl derivative 32, and the isopropylidene hydrazide 58 was a more significant product. Presumably 58 arises by interaction of 59 with acetone utilized in the experimental procedure, and isolation of 58 suggests that formation of the dihydropyridazinone nucleus proceeds in this instance *via* intramolecular condensation of the acyl hydrazide onto the carbonyl function. Moreover, the presence of only end absorption in the electronic spectrum of 58 indicates preference for an "out-of-plane" conformation with respect to the carbonyl function and the aryl system. This preference is the apparent result of limitations on the degrees of freedom imposed by the steric requirements of the nitro and methyl substituents, and these constrictions make the tetrahedral intermediate in the conversion of 59 into 32 less attainable.



Modification of appropriate dihydropyridazinones afforded other members of the series. Thus, catalytic reduction of certain 6-(nitrophenyl) compounds gave excellent yields of the corresponding 6-(aminophenyl) derivatives 31 and 35. The preparation of the 6-(*o*-aminophenyl)dihydropyridazinone (31) had been achieved earlier by treatment of  $\beta$ -(*o*-aminobenzoyl)propionic acid with hydrazine.<sup>3</sup> Acetylation of the requisite compounds gave the 6-(acylamino phenyl) derivatives 19-29, 48, and 49. The Sandmeyer procedure was used to prepare the *m*-hydroxy (39) and *m*-bromo (40) derivatives, and displacement<sup>4</sup> of bromide in the 6-bromophenyl derivatives 40 and 41 constituted an efficient alternative synthesis of the interesting *m*- (43) and *p*-cyanophenyl (44) compounds. Acid hydrolysis of 43 and 44 gave the carboxamides 55 and 56, respectively, which were converted into their carboxylic acids by treatment with nitrosonium hexafluorophosphate. The carboxamide and carboxylic acid derivatives were of particular interest, inasmuch as they are possible metabolites of the more interesting carbonitriles. In addition to these transformations, 6-(*p*-aminophenyl)-4,5-dihydro-5-methyl-3(2H)-pyridazinone<sup>1</sup> was converted into the sulfamoyl

Table I. 6-Phenyl-4,5-dihydro-3(2H)-pyridazinones



Compd	R	R'	R''	Meth- od	Yield, %	Recrystn solvent	Mp, °C	Formula	Analyses	Antihypertensive act., MABP <sup>a</sup>	
										4 hr	24 hr
A. 6-(Alkylaminophenyl) Derivatives											
1	<i>p</i> -CH <sub>3</sub> NH	H	H	A	90	EtOH	224-227	C <sub>11</sub> H <sub>13</sub> N <sub>3</sub> O	C, H, N	++	+
2	<i>p</i> -C <sub>2</sub> H <sub>5</sub> NH	H	H	A	44	EtOH	210-213	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O	C, H, N	++	+
3	<i>p</i> -CH <sub>3</sub> NH	CH <sub>3</sub>	H	A	64	MeOH	213-215	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O	C, H, N	+++	-
4	<i>m</i> -CH <sub>3</sub> NH	CH <sub>3</sub>	H	A	47	EtOH	218-221 dec	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O · HCl	C, H, Cl, N	++++	-
5	<i>p</i> -CH <sub>3</sub> NH	H	CH <sub>3</sub>	A	89	EtOH	199-200	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O	C, H, N	-	-
6	<i>p-n</i> -C <sub>3</sub> H <sub>7</sub> NH	H	H	A	83	EtOH	182-183	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	+++	-
7	<i>p</i> -CH <sub>3</sub> NH	CH <sub>3</sub>	CH <sub>3</sub>	A	63	EtOH	212-214	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	++	-
8	<i>p</i> -C <sub>2</sub> H <sub>5</sub> NH	CH <sub>3</sub>	H	A	82	EtOH	192-194	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	+++	+
9	<i>p</i> -CH <sub>3</sub> NH	C <sub>2</sub> H <sub>5</sub>	H	A	52 <sup>b</sup>	CH <sub>2</sub> Cl <sub>2</sub> -petr ether	142-144	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	-	-
10	<i>m</i> -CH <sub>3</sub> NH	CH <sub>3</sub>	CH <sub>3</sub>	A	83	EtOH	180-182 dec	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O · HCl	C, H, Cl, N	-	-
11	<i>p</i> -C <sub>2</sub> H <sub>5</sub> NH	CH <sub>3</sub>	CH <sub>3</sub>	A	75	EtOH	165-167	C <sub>14</sub> H <sub>19</sub> N <sub>3</sub> O	C, H, N	++	+
12	<i>p</i> -CH <sub>3</sub> NH	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	A	51	Acetone-hexane	142-144	C <sub>14</sub> H <sub>19</sub> N <sub>3</sub> O	C, H, N	-	-
B. 6-(Dimethylaminophenyl) Derivatives											
13	<i>p</i> -(CH <sub>3</sub> ) <sub>2</sub> N	H	H	A	86	CH <sub>2</sub> Cl <sub>2</sub> -petr ether	255-258	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O	C, H, N	-	-
14	<i>m</i> -(CH <sub>3</sub> ) <sub>2</sub> N	CH <sub>3</sub>	H	A	83	Acetone-hexane	146-148	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	+++	+
15	<i>p</i> -(CH <sub>3</sub> ) <sub>2</sub> N	CH <sub>3</sub>	H	A	81	MeOH	208-210	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	++	+
16	<i>p</i> -(CH <sub>3</sub> ) <sub>2</sub> N	H	CH <sub>3</sub>	H	9	Acetone-hexane	210-212	C <sub>13</sub> H <sub>17</sub> N <sub>3</sub> O	C, H, N	-	-
17	<i>p</i> -(CH <sub>3</sub> ) <sub>2</sub> N	CH <sub>3</sub>	CH <sub>3</sub>	A	90	CH <sub>2</sub> Cl <sub>2</sub> -petr ether	173-175	C <sub>14</sub> H <sub>19</sub> N <sub>3</sub> O	C, H, N	+	+
18	<i>m</i> -(CH <sub>3</sub> ) <sub>2</sub> N	CH <sub>3</sub>	CH <sub>3</sub>	A	68	MeOH-H <sub>2</sub> O	83-85	C <sub>14</sub> H <sub>19</sub> N <sub>3</sub> O	C, H, N	-	-
C. 6-(Acylaminophenyl) Derivatives											
19	<i>p</i> -N(CH <sub>3</sub> )CHO	H	H	G <sub>1</sub>	53	EtOH	191-193	C <sub>12</sub> H <sub>13</sub> N <sub>3</sub> O <sub>2</sub>	C, H, N	++++	-
20	<i>p</i> -CF <sub>3</sub> CONH	H	CH <sub>3</sub>	M	95	Acetone	275-276	C <sub>13</sub> H <sub>12</sub> F <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	C, H, F, N	-	-
21	<i>p</i> -CF <sub>3</sub> CONH	CH <sub>3</sub>	H	M	100	Acetone-hexane	254-256	C <sub>13</sub> H <sub>12</sub> F <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	C, H, F, N	+++	++
22	<i>p</i> -N(CH <sub>3</sub> )COCH <sub>3</sub>	H	H	G	82	Acetone-hexane	184-186	C <sub>13</sub> H <sub>15</sub> N <sub>3</sub> O <sub>2</sub> · 0.5H <sub>2</sub> O	C, H, N	++++	-
23	<i>p</i> -N(CH <sub>3</sub> )COCF <sub>3</sub>	CH <sub>3</sub>	H	M	94	Acetone-petr ether	160-161	C <sub>14</sub> H <sub>14</sub> F <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	C, H, F, N	+++	+
24	<i>p</i> -CF <sub>3</sub> CONH	CH <sub>3</sub>	CH <sub>3</sub>	M	80	Acetone-hexane	174-176	C <sub>14</sub> H <sub>14</sub> F <sub>3</sub> N <sub>3</sub> O <sub>2</sub>	C, H, F, N	++	++
25	<i>p</i> -N(CH <sub>3</sub> )COCH <sub>3</sub>	CH <sub>3</sub>	H	G	84	CH <sub>2</sub> Cl <sub>2</sub> -petr ether	218-220	C <sub>14</sub> H <sub>17</sub> N <sub>3</sub> O <sub>2</sub>	H, N; C <sup>c</sup>	++++	++
26	<i>p</i> -N(CH <sub>3</sub> )COC <sub>2</sub> H <sub>5</sub>	H	H	G	89	EtOH-H <sub>2</sub> O	108-110	C <sub>14</sub> H <sub>17</sub> N <sub>3</sub> O <sub>2</sub> · H <sub>2</sub> O	C, H, N	++++	-
27	<i>p</i> -N(C <sub>2</sub> H <sub>5</sub> )COCH <sub>3</sub>	H	H	G	83	Acetone-hexane	167-168	C <sub>14</sub> H <sub>17</sub> N <sub>3</sub> O <sub>2</sub>	C, H, N	++++	+
28	<i>p</i> -N(CH <sub>3</sub> )COCH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	G	83	Et <sub>2</sub> O-petr ether	110-112	C <sub>15</sub> H <sub>19</sub> N <sub>3</sub> O <sub>2</sub>	H, N; C <sup>d</sup>	++++	-
29	<i>p</i> -N(C <sub>2</sub> H <sub>5</sub> )COCH <sub>3</sub>	CH <sub>3</sub>	H	G	64	Acetone-hexane	148-150	C <sub>15</sub> H <sub>19</sub> N <sub>3</sub> O <sub>2</sub>	C, H, N	++++	+
D. (o-Substituted phenyl) Derivatives											
30	<i>o</i> -NO <sub>2</sub>	H	H	A <sub>1</sub>	47	Acetone-hexane	164-165	C <sub>10</sub> H <sub>9</sub> N <sub>3</sub> O <sub>3</sub>	C, H, N	+	-
31	<i>o</i> -NH <sub>2</sub>	H	H	I	82	EtOH	171-173	C <sub>10</sub> H <sub>11</sub> N <sub>3</sub> O	C, H, N	-	-
32	<i>o</i> -NO <sub>2</sub>	CH <sub>3</sub>	H	A <sub>1</sub>	5	Acetone-hexane	159-161	C <sub>11</sub> H <sub>11</sub> N <sub>3</sub> O <sub>3</sub>	C, H, N	++	-
33	<i>o</i> -NHCOCH <sub>3</sub>	H	H	A <sub>1</sub>	40	MeOH	257-258	C <sub>12</sub> H <sub>13</sub> N <sub>3</sub> O <sub>2</sub>	C, H, N	-	-
E. 2-Alkyl Derivatives											
34	<i>m</i> -NO <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	A	69	MeOH-H <sub>2</sub> O	196-197	C <sub>12</sub> H <sub>13</sub> N <sub>3</sub> O <sub>3</sub>	C, H, N	++++	+
35	<i>m</i> -NH <sub>2</sub>	CH <sub>3</sub>	CH <sub>3</sub>	I	93	H <sub>2</sub> O	123-124	C <sub>12</sub> H <sub>15</sub> N <sub>3</sub> O	C, H, N	-	-
36	<i>m</i> -CN	CH <sub>3</sub>	CH <sub>3</sub>	B	74	Et <sub>2</sub> O-petr ether	80-81	C <sub>13</sub> H <sub>13</sub> N <sub>3</sub> O	C, H, N	++++	+





within  $\pm 0.4\%$  of the theoretical values. The petroleum ether used was that fraction with bp 30–60°.

**Preparation of the 6-(Substituted phenyl)-4,5-dihydro-3(2H)-pyridazinones (I).** **Method A.** A solution of 34.85 g (0.128 mol) of  $\beta$ -(*p*-bromobenzoyl)butyric acid<sup>9</sup> and 10.0 g (0.21 mol, 10 ml) of hydrazine hydrate in 300 ml of EtOH was stirred at reflux temperature for 3 hr. A solid began separating after 5 min. The mixture was chilled and filtered to give 31.57 g (92%) of 6-(*p*-bromophenyl)-4,5-dihydro-5-methyl-3(2H)-pyridazinone (41) as white crystals, mp 197–199°. The characterization of this substance and others prepared in a similar manner is given in Table I.

**Method A<sub>1</sub>.** A solution of 4.68 g (20 mmol) of methyl 3-(*o*-nitrobenzoyl)propionate, 2.1 ml (40.5 mmol) of hydrazine hydrate, and 1.65 ml of HOAc in 25 ml of EtOH was heated at reflux temperature for 18 hr. Removal of the solvent gave a residue that was distributed between CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The material in the organic layer crystallized from CH<sub>2</sub>Cl<sub>2</sub>-petroleum ether to give 2.03 g (46%) of 4,5-dihydro-6-(*o*-nitrophenyl)-3(2H)-pyridazinone (30) as yellow crystals, mp 162–165°.

**Method B.** A stirred mixture of 1.00 g (4.65 mmol) of 6-(*p*-cyanophenyl)-4,5-dihydro-5-methyl-3(2H)-pyridazinone (44) and 1.29 g of powdered KOH in 30 ml of acetone was treated with 3.0 ml of MeI. The mixture was heated at reflux temperature for 40 min, and the solvent was removed. The residue was triturated with H<sub>2</sub>O and filtered to give 6-(*p*-cyanophenyl)-4,5-dihydro-2,5-dimethyl-3(2H)-pyridazinone (37), mp 172–174°. The characterization of this substance is given in Table I.

**Method B<sub>1</sub>.** Application of method B to 670 mg (2.14 mmol) of 4,5-dihydro-2,5-dimethyl-6-(*p*-2,2,2-trifluoroacetamidophenyl)-3(2H)-pyridazinone (24), 0.60 ml (8.5 mmol) of CH<sub>3</sub>I, and 475 mg (8.5 mmol) of powdered KOH in 12 ml of acetone afforded 490 mg of 4,5-dihydro-2,5-dimethyl-6-(*p*-methylaminophenyl)-3(2H)-pyridazinone (7). See Table I for its characterization.

**Method C.** A solution of 5.22 g (25.8 mmol) of 6-(*m*-anilino)-4,5-dihydro-5-methyl-3(2H)-pyridazinone<sup>1</sup> in 18 ml of H<sub>2</sub>O and 11 ml of 48% HBr was stirred at 0°. A solid separated, and 1.79 g (26 mmol) of NaNO<sub>2</sub> was added in portions to the slurry, the temperature being maintained at 0–3°. Solution occurs, and then a solid separates. The slurry was added dropwise to a cold solution of 4.05 g (14 mmol) of Cu<sub>2</sub>Br<sub>2</sub> in 13 ml of 48% HBr. The mixture was stirred at 0° for 30 min, allowed to warm to 20°, and then stirred at 10° for 1 hr. The mixture was cooled and diluted with 90 ml of H<sub>2</sub>O. The precipitate was collected, dried, and extracted with two 100-ml portions of CH<sub>2</sub>Cl<sub>2</sub>. Evaporation of the solvent gave a glass which was dissolved in 5% acetone in CH<sub>2</sub>Cl<sub>2</sub> and filtered through a synthetic magnesia-silica adsorbent. The solid was washed liberally with the same solvent mixture. Evaporation of the solvent from the filtrate gave a solid which was recrystallized from acetone-hexane to give 1.75 g of 6-(*m*-bromophenyl)-4,5-dihydro-5-methyl-3(2H)-pyridazinone (40) as white crystals, mp 160–163°. See Table I for its characterization.

**Method D.** A mixture of 20.43 g (76.5 mmol) of 6-(*p*-bromophenyl)-4,5-dihydro-5-methyl-3(2H)-pyridazinone (41) and 9.10 g (51 mmol) of Cu<sub>2</sub>(CN)<sub>2</sub> in 70 ml of DMF was stirred at reflux temperature for 5.5 hr. The hot mixture was poured into a stirred solution of 46 ml of ethylenediamine in 230 ml of H<sub>2</sub>O; stirring was continued for 20 min, whereafter the mixture was filtered to give 14.20 g (87%) of 6-(*p*-cyanophenyl)-4,5-dihydro-5-methyl-3(2H)-pyridazinone (44), mp 194–197°. See Table I for other compounds prepared similarly.

**Method E.** A solution of 500 mg (2.35 mmol) of *p*-(1,4,5,6-tetrahydro-4-methyl-6-oxo-3-pyridazinyl)benzotrile (41) in 5 ml of concentrated H<sub>2</sub>SO<sub>4</sub> was allowed to stand at room temperature for 18 hr and then was diluted with 45 ml of iced H<sub>2</sub>O to afford 460 mg (85%) of *p*-(1,4,5,6-tetrahydro-4-methyl-6-oxo-3-pyridazinyl)benzamide (56) as a white solid, mp 215–216°. See Table I for the characterization of this substance.

**Method F.** To a solution of 360 mg (2.06 mmol) of nitrosonium hexafluorophosphate in 10 ml of acetonitrile cooled to ice-bath temperature was added with stirring 395 mg (1.71 mmol) of *p*-(1,4,5,6-tetrahydro-4-methyl-6-oxo-3-pyridazinyl)benzamide (56). The solution was stirred at 0° for 10 min, at 25° for 20 min, and at 50° for 30 min. Water (0.3 ml) was added to the reaction, and the solid was collected to give 250 mg (63%) of *p*-(1,4,5,6-tetrahydro-4-methyl-6-oxo-3-pyridazinyl)benzoic acid (54) as white crystals, mp 266–270°.

**Method G.** A mixture of 500 mg (2.2 mmol) of 6-(*p*-ethylaminophenyl)-4,5-dihydro-3(2H)-pyridazinone (2) and 1 ml of acetic anhydride was stirred at room temperature for 1 hr. Water was added to destroy the excess anhydride, after which time the solid

*N*-ethyl-4'-(1,4,5,6-tetrahydro-6-oxo-3-pyridazinyl)acetanilide (27) (470 mg, 83%) was collected to give white crystals, mp 167–168°. See Table I for its characterization.

**Method G<sub>1</sub>.** A mixture of 400 mg (1.97 mmol) of 4,5-dihydro-6-(*p*-methylaminophenyl)-3(2H)-pyridazinone (1) and 1 ml of 97% HCO<sub>2</sub>H in 10 ml of toluene was heated at reflux temperature for 17 hr; separated H<sub>2</sub>O was collected in a modified Dean-Stark H<sub>2</sub>O trap. The solvent was removed, and the product, *N*-methyl-4'-(1,4,5,6-tetrahydro-6-oxo-3-pyridazinyl)formanilide (19), was purified as indicated in Table I.

**Method H.** A solution of 1.71 g (8.4 mmol) of 6-(*p*-anilino)-4,5-dihydro-2-methyl-3(2H)-pyridazinone, 0.55 g (18.5 mmol, 1.4 ml) of 37% formalin, and 1.83 g (42 mmol, 1.5 ml) of 97% HCO<sub>2</sub>H was heated on a steam bath for 17 hr. The volatile material was removed, and the residue was distributed between CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The material in the organic layer was chromatographed on a synthetic magnesia-silica adsorbent. The fractions eluted by CH<sub>2</sub>Cl<sub>2</sub> contained the product, the characterization of which is given in Table I.

**Method I.** A mixture of 2.56 g (10.4 mmol) of 4,5-dihydro-2,5-dimethyl-6-(*m*-nitrophenyl)-3(2H)-pyridazinone (34) and 250 mg of 10% Pd/C in 50 ml of EtOH was shaken under hydrogen until the pressure became constant (13 min). The solution was filtered and evaporated to furnish a gum which crystallized from Et<sub>2</sub>O-petroleum ether to give 2.10 g (93%) of 6-(*m*-anilino)-4,5-dihydro-2,5-dimethyl-3(2H)-pyridazinone (35) as white crystals, mp 120–122°.

**Preparation of Esters of 3-(Substituted benzoyl)alkanoic Acids.** **Method J.** A solution of 3.00 g (13.5 mmol) of 3-(*p*-aminobenzoyl)butyric acid<sup>9</sup> and 0.12 ml of H<sub>2</sub>SO<sub>4</sub> in 60 ml of MeOH was heated at reflux temperature for 18 hr. The solution was cooled and added to 1.0 g of anhydrous NaOAc and diluted with 60 ml of H<sub>2</sub>O, and the MeOH was removed until crystals separated. The mixture was cooled in an ice bath, and the solid was collected, affording 2.20 g (66%) of methyl 3-(*p*-aminobenzoyl)butyrate (67) as crystals, mp 116–118°. The characterization of this substance and the preparation of other esters by this procedure are given in Table II.

**Method K.** To a mixture of 1.49 g (7.20 mmol) of methyl 3-(*p*-aminobenzoyl)propionate (66) and 6.2 ml of 37% formalin in 30 ml of MeCN was added 1.37 g (23.2 mmol) of sodium cyanoborohydride. To the stirred mixture was added 0.77 ml of HOAc over a period of 7 min. Stirring was continued for 2 hr, and an additional 0.77 ml of HOAc added. After stirring for an additional 30 min, the solution was diluted with 100 ml of Et<sub>2</sub>O and washed with 1 N NaOH solution. The organic solution was dried and evaporated leaving 1.69 g of amber gum. This material was heated at reflux temperature for 90 min with 20 ml of 6 N HCl solution, and the hot solution was filtered through diatomaceous earth. Evaporation of the filtrate gave a residue that was triturated with 25 ml of acetone to give 1.62 g (100%) of crude 3-(*p*-dimethylaminobenzoyl)propionic acid (81) as a white solid, mp 172–175°. See Table II for the characterization of this substance and those prepared in similar fashion.

**Method L.** A solution of 4.08 g (20 mmol) of 3-(*o*-nitrobenzoyl)propionitrile<sup>9</sup> in 100 ml of methanol saturated with hydrogen chloride at 0° was heated at reflux temperature for 3 hr. Most of the solvent was removed, and the residue was distributed between methylene chloride and water. The dried organic solution was evaporated to give 4.68 g (99%) of yellow liquid; ir max 5.75–5.81, 6.60, 7.45  $\mu$ ; tlc in EtOAc-heptane (1:1) *R*<sub>f</sub> 0.65. The esters prepared in this manner were used for the preparation of the dihydropyridazinones without further purification.

**Method M.** A mixture of 11.80 g (53.4 mmol) of methyl 3-(*p*-aminobenzoyl)butyrate (66) and 25 ml of trifluoroacetic anhydride was stirred for 1 hr, diluted with 370 ml of ice-water, and stirred for an additional 1 hr. The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>, and the extracts were washed successively with 1 N HCl, saline, saturated NaHCO<sub>3</sub> solution, and saline. The dried solution was evaporated to give 15.97 g (100%) of methyl 3-(*p*-2,2,2-trifluoroacetamidobenzoyl)butyrate (71) as a gum that crystallized on trituration with Et<sub>2</sub>O-petroleum ether to give a white solid, mp 67–70°. See Table II for its characterization.

**Preparation of the 3-(Alkylaminobenzoyl)alkanoic Acids.** **Method N.** The following preparation of 3-(*p*-methylaminobenzoyl)butyric acid (77) illustrates the general procedure. To a stirred solution of 15.0 g (47.4 mmol) of methyl 3-(*p*-2,2,2-trifluoroacetamidobenzoyl)butyrate (71) and 30.8 g (13.5 ml, 0.21 mol) of iodomethane in 300 ml of acetone was added 14.2 g (0.25 mol) of powdered KOH. The mixture was stirred at reflux temperature for 5 min and then the solvent was removed. Water (200 ml) was

added to the residue, and the mixture was heated at reflux temperature for 10 min. The mixture was cooled and washed with  $\text{CH}_2\text{Cl}_2$ . The alkaline solution was stirred in an ice bath and rendered acidic (pH  $\sim$ 4) by the dropwise addition of 1 N HCl. The product was collected as tan crystals, mp 170–172°. The characterization of this acid and others prepared in a similar manner is described in Table II.

**3-(*p*-Bromobenzoyl)-2-methylcrotonic Acid Lactone (60).** A solution of 13.75 g (9.2 ml, 87.5 mmol) of bromobenzene in 10 ml of  $\text{CS}_2$  was added dropwise over a period of 60 min to a stirred mixture of 49.7 g (0.186 mol) of  $\text{AlBr}_3$  and 9.80 g (78 mmol) of 2,3-dimethylmaleic anhydride. The mixture was heated at reflux for 4 hr and then poured onto iced 37% HCl. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were washed with saline solution, dried, and evaporated. The residue was subjected to steam distillation, collecting 350 ml of distillate. The pot residue was extracted with  $\text{CH}_2\text{Cl}_2$  and the extracts were washed, dried, and evaporated. The residual gum was triturated with heptane to give 11.10 g (50%) of white solid, mp 92–96°. One crystallization from  $\text{CH}_2\text{Cl}_2$ -heptane gave 9.45 g of white crystals: mp 100–102°; uv max 228  $\text{m}\mu$  ( $\epsilon$  14,300); ir max 2.95, 5.73, 5.90, 6.25  $\mu$ ;  $\delta^{\text{TMS}}$  ( $\text{CDCl}_3$ ) 1.76 (s, 6,  $\text{CH}_3$ ), 3.42 (s, 1, OH), 7.27, 7.48 (d, 2 each, aryl  $\text{A}_2\text{B}_2$ ), exchange with  $\text{CD}_3\text{OD}$  erased the 3.42 signal.

***cis*- and *trans*-6-(*p*-Bromophenyl)-4,5-dihydro-4,5-dimethyl-3(2*H*)-pyridazinone.** A hot solution of 5.50 g (19.5 mmol) of 3-(*p*-bromobenzoyl)-2-methylcrotonic acid lactone (60) in 75 ml of HOAc and 30 ml of  $\text{H}_2\text{O}$  was treated with 2.73 g (41.8 mg-atoms) of Zn dust. The mixture was stirred at reflux temperature for 30 min. An additional 2.73 g of Zn dust was added and the heating continued for 3.5 hr. The mixture was cooled, and the supernatant was decanted from the Zn and concentrated. The concentrate was diluted with  $\text{H}_2\text{O}$ , and the mixture was extracted with  $\text{CH}_2\text{Cl}_2$ . The extracts were washed with saline, dried, and evaporated. Toluene was added to the residue and removed under reduced pressure to give 5.18 g of crude 3-(*p*-bromobenzoyl)-2-methylbutyric acids which were converted into the pyridazinones by method A.

The crude product (5.20 g) was subjected to partition chromatography on diatomaceous silica using a heptane-MeCN- $\text{H}_2\text{O}$  (200:55:45) system and monitoring the effluent for material with uv absorption at 292  $\text{m}\mu$ . The material with peak hold-back volume 4.4 ( $V_m/V_s = 3.46$ ) was crystallized from  $\text{Et}_2\text{O}$ -petroleum ether to give 840 mg (15%) of *cis* isomer 63: mp 174–176°;  $\delta^{\text{TMS}}$  ( $\text{CDCl}_3$ ) 1.10 (d, 3,  $J = 8.0$  Hz, 4- $\text{CH}_3$ ), 1.30 (d, 3,  $J = 8.0$  Hz, 5- $\text{CH}_3$ ), 2.78 (d of q, 1,  $J(\text{H}_a-\text{H}_b) \sim 7$  Hz,  $J(\text{H}_a-\text{CH}_3) = 8.0$  Hz, 4-H), 3.16 (d of q, 1,  $J(\text{H}_a-\text{H}_b) \sim 7$  Hz,  $J(\text{H}_b-\text{CH}_3) = 8.0$  Hz, 5-H), 7.57, 7.62 (each d, 1 each,  $J = 10$  Hz, aryl  $\text{A}_2\text{B}_2$ ), 9.15 (s, 1, NH). *Anal.* ( $\text{C}_{12}\text{H}_{13}\text{BrN}_2\text{O}$ ) C, H, Br, N.

The material eluted at peak hold-back volume 6.3 crystallized from  $\text{Et}_2\text{O}$ -petroleum ether to give 280 mg (5%) of *trans* isomer 62: mp 174–175°;  $\delta^{\text{TMS}}$  ( $\text{CDCl}_3$ ) 1.19, 1.22 (overlapping d, 3 each,  $J = 8$  Hz,  $\text{CH}_3$ ), 2.54 (d of q, 1,  $J(\text{H}_a-\text{CH}_3) = 8$  Hz,  $J(\text{H}_a-\text{H}_b) \sim 1.5$  Hz, 4-H), 3.02 (d of q, 1, 5-H), 7.56, 7.61 (aryl- $\text{A}_2\text{B}_2$ , 4), 9.11 (s, 1, NH). *Anal.* ( $\text{C}_{12}\text{H}_{13}\text{BrN}_2\text{O}$ ) C, H, N; Br: calcd, 28.43; found, 28.94.

***cis*-4,5-Dihydro-4,5-dimethyl-6-phenyl-3(2*H*)-pyridazinone (64)** was eluted at peak hold-back volume 10.0. Recrystallization from acetone-hexane gave 421 mg (11%) of white crystals; mp 124–125°;  $\delta^{\text{TMS}}$  ( $\text{CDCl}_3$ ) 1.12, 1.30 (each d, 3,  $J \sim 7$  Hz,  $\text{CH}_3$ ), 2.78 (d of q, 1,  $J(\text{H}_a-\text{CH}_3) = J(\text{H}_a-\text{H}_b) \sim 7$  Hz, 4-H), 3.22 (d of q, 1, 5-H), 7.42 (m, 3, *m*- and *p*-H of Ph), 7.77 (m, 2, *o*-H of Ph), 9.09 (s, 1, NH). *Anal.* ( $\text{C}_{12}\text{H}_{14}\text{N}_2\text{O}$ ) C, H, N.

**6-(*p*-Cyanophenyl)-4,5-dimethyl-3(2*H*)-pyridazinone (65).** A mixture of 1.28 g (4.55 mmol) of *cis*-6-(*p*-bromophenyl)-4,5-dihydro-4,5-dimethyl-3(2*H*)-pyridazinone (63) and 1.72 g (9.6 mmol) of  $\text{Cu}_2(\text{CN})_2$  in 5 ml of DMF was stirred and heated at reflux temperature for 5 hr. The hot solution was poured into a solution of 4.4 g of  $\text{FeCl}_3$  and 3.4 ml of 37% HCl in 12 ml of water and heated on the steam bath for 30 min. The solution was cooled, diluted with  $\text{H}_2\text{O}$ , and filtered to give 1.30 g of brown solid. This solid was extracted with  $\text{CH}_2\text{Cl}_2$ , and the extracts were evaporated. The residual solid was recrystallized from acetone to give 270 mg (26%) of white crystals: mp 258–262°; uv max 242  $\text{m}\mu$  ( $\epsilon$  23,900). *Anal.* ( $\text{C}_{13}\text{H}_{11}\text{N}_3\text{O}$ ) C, H, N.

**4'-(6-Amino-4,5-dihydro-4-methyl-3-pyridazinyl)acetanilide (82a).** A mixture of 1.28 g (5.6 mmol) of 4'-(3-cyano-2-methylpropionyl)acetanilide,<sup>9</sup> 0.58 ml of hydrazine hydrate, and 0.47 ml of HOAc in 7 ml of EtOH was stirred and heated at reflux temperature for 1 hr. The mixture was cooled, and the yellow crystals were collected and washed successively with methanol, methylene chloride, and acetone to give 635 mg (46%) of yellow crystals, mp 254–256°. *Anal.* ( $\text{C}_{13}\text{H}_{16}\text{N}_4\text{O}$ ) C, H, N.

**6-Amino-4,5-dihydro-4-methyl-3-(*m*-nitrophenyl)pyridazine (82b).** In the manner described above a mixture of 920 mg (4.2 mmol) of 3-*m*-nitrobenzoylbutyronitrile,<sup>9</sup> 0.44 ml of hydrazine hydrate, and 0.35 ml of HOAc in 10 ml of EtOH gave 500 mg (51%) of crystals, mp 200–210°. A sample recrystallized from acetone-hexane had mp 200–208°. *Anal.* ( $\text{C}_{11}\text{H}_{12}\text{N}_4\text{O}_2$ ) C, H, N.

**3-(*o*-Nitrobenzoyl)butyric Acid Isopropylidenehydrazide (58).** Using method A, 7.32 g (29.5 mmol) of methyl 3-(*o*-nitrobenzoyl)butyrate and 7 ml of hydrazine hydrate in 75 ml of MeOH gave 6.87 g of a gum after solvent removal. An acetone-ether solution of this material deposited 1.68 g (20%) of the hydrazide as yellow crystals, mp 140–143°, on standing at room temperature. Recrystallization of a sample from  $\text{CH}_2\text{Cl}_2$ -petroleum ether gave yellow crystals, mp 143–144°. *Anal.* ( $\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}_4$ ) C, H, N. The filtrate was evaporated to give 4.57 g of a residue from which 32 was isolated by partition chromatography.

**Reaction of Chlorosulfonyl Azide with 6-(*p*-Aminophenyl)-4,5-dihydro-2,5-dimethyl-3(2*H*)-pyridazinone.** A stirred, ice-cooled solution of 2.17 g (10 mmol) of 6-(*p*-aminophenyl)-4,5-dihydro-2,5-dimethyl-3(2*H*)-pyridazinone (crude material prepared from 3-(*p*-aminobenzoyl)butyric acid<sup>9</sup> and methylhydrazine using procedure A) and 1.4 ml (10.2 mmol) of triethylamine in 20 ml of  $\text{CH}_3\text{CN}$  was treated with a solution of chlorosulfonyl azide<sup>11</sup> in acetonitrile (prepared as described by Griffiths<sup>12</sup> from 5.0 mmol of sulfonyl chloride and 5.0 mmol of sodium azide). The reaction was concentrated to a volume of 5 ml, and the products were isolated with EtOAc in the usual manner. After removal of the solvent, the residue was crystallized from acetone-hexane to give 57 as tan crystals.

Evaporation of the filtrate gave a residue that was chromatographed on a synthetic magnesia-silica adsorbent. The material eluted by 8% acetone in methylene chloride crystallized from  $\text{Et}_2\text{O}$ -petroleum ether to give 52. Characterization of 52 and 57 is given in Table I.

**Acknowledgment.** We are indebted to Messrs. L. Brancone and W. Fulmor and their staffs for microanalyses and spectral data. Partition chromatography was performed by Mr. J. Baker with the cooperation of Dr. A. Streuli. The determination of the antihypertensive activity of candidate agents was performed by Miss L. Lipchuck and Mr. F. Beyer with the cooperation of Dr. R. Z. Gussin.

## References

- (1) W. V. Curran and A. Ross, *J. Med. Chem.*, **17**, 273 (1974).
- (2) E. A. Steck, R. P. Brundage, and L. T. Fletcher, *J. Amer. Chem. Soc.*, **75**, 1117 (1953), and references cited therein.
- (3) M. Morishita and F. Sakiyama, *Bull. Chem. Soc. Jap.*, **43**, 524 (1970).
- (4) (a) L. Friedman and H. Shechter, *J. Org. Chem.*, **26**, 2522 (1961); (b) M. S. Newman and H. Boden, *ibid.*, **26**, 2525 (1961).
- (5) W. L. Matier, W. T. Comer, and D. Deitchman, *J. Med. Chem.*, **15**, 538 (1972), and references cited therein.
- (6) R. E. Lutz and M. Couper, *J. Org. Chem.*, **6**, 77 (1941).
- (7) R. A. W. Johnstone, D. W. Payling, and C. Thomas, *J. Chem. Soc. C*, 2223 (1969).
- (8) R. F. Borch and A. I. Hassid, *J. Org. Chem.*, **37**, 1673 (1972).
- (9) F. J. McEvoy and G. R. Allen, Jr., *ibid.*, **38**, 4044 (1973).
- (10) J. R. Cummings, A. N. Welter, J. L. Grace, Jr., and L. M. Lipchuck, *J. Pharmacol. Exp. Ther.*, **161**, 88 (1968).
- (11) R. J. Shozda and J. A. Vernon, *J. Org. Chem.*, **32**, 2876 (1967).
- (12) J. Griffiths, *J. Chem. Soc. C*, 3191 (1971).