

- (15) N. W. Dunham and T. S. Miya, *J. Amer. Pharm. Ass.*, **46**, 208 (1957).
 (16) P. A. Janssen, C. Van De Westeringh, A. H. M. Jageneau, P. J. A. Demoen, B. K. F. Hermans, G. H. P. Van Daele, K. H. L. Schellekens, C. A. M. Van Der Eycken, and C. J. E. Niemegeers, *J. Med. Pharm. Chem.*, **1**, 281 (1959).
 (17) C. S. Scherrington and S. C. M. Sowton, *J. Physiol. (London)*, **51**, 404 (1917).
 (18) G. K. Suchowsky and L. Pegrassi, *Arzneim.-Forsch.*, **19**, 643 (1969).
 (19) F. E. Roth, S. Irwin, E. Eckhardt, I. I. A. Tabachnick, and W. M. Govier, *Arch. Int. Pharmacodyn. Ther.*, **118**, 375 (1959).


A Procedure for Preparing ^3H -Labeled Tertiary Amines. Synthesis of [^3H]-6,14-endo-Etheno-6,7,8,14-tetrahydrooripavine Derivatives

John W. Lewis,* Michael J. Rance, and G. Raymond Young

Reckitt & Colman Pharmaceutical Division, Hull, England.
 Received July 30, 1973

During radiosynthesis it is of paramount importance that the label is introduced at as late a stage as possible. The concept of a synthesis based on the unlabeled required product as starting material is very attractive; techniques such as the Wilzbach gas exposure method¹ and base-catalyzed exchange of aromatic protons² illustrate this approach. Both of these methods, however, suffer from distinct disadvantages, the former in that the labeling is unspecific and the latter in that exchange labeled material is susceptible to loss of activity by the reverse exchange process.

In the course of the study of the metabolism of the series of oripavine analgesics **1**, it has been necessary to develop methods of labeling these compounds with tritium. Initially, etorphine (**1a**), a very potent analgesic used for immobilization of animals,³ was labeled at C-8 by a long inefficient route.⁴ We here report a method of introducing a tritium label at carbon atoms α and/or β to a tertiary nitrogen by dehydrogenation to the enamine (*e.g.*, **2**) into which tritium is readily exchanged. The labeled tertiary base is regenerated by subsequent reduction with or without further tritiation.[†] Use of such a route has the advantages of high specific activity and cheapness inherent in exchange reactions, together with the regioselectivity and stability required for biological studies. Recently, Portoghesi⁶ has described a somewhat analogous method of la-



^oThebaine derivative.

[†]We have previously reported brief details of the synthesis of [^{15}H]etorphine from the 15,16-didehydro compound **2a**.⁵

Table I. Specific Activities of Labeled Oripavine Derivatives


Labeling method	Compd	Product	Sp act., mCi/mmol	
(i) T ₂ O (ca. 20 Ci)	1a	15- ³ H	100, 220, 260	
	1b	15- ³ H	160	
	(ii) NaBH ₄	1c	15- ³ H	110
		1d	15- ³ H	26 ^a
	1e	15- ³ H	230	
(i) T ₂ O (ca. 20 Ci)	1a	15,16- ³ H	900	
	(ii) NaBT ₄ (250 mCi)	1e	15,16- ³ H	1300
T ₂ (ca. 10 Ci); Pd/C (10%)		1a	15,16- ³ H	1500, 3600 ^b

^aT₂O used for this reaction was recovered from a previous experiment. ^bAfter isotope dilution.

belonging secondary amines in the α position by utilizing the acidity conferred on the α protons by the introduction of an *N*-nitroso function.

The enamines **2**, prepared by the dehydrogenation of the parent tertiary bases **1** with mercury(II) acetate,^{5,7} are strong bases and the proton at C-15 equilibrates rapidly with water. Nmr studies with **2e** showed that in CDCl₃ the AB quartet (δ 4.26 and 5.93, $J = 8$ Hz) due to H-15 and H-16 collapsed to a singlet (δ 5.95) on addition of D₂O. Thus, treatment of 15,16-didehydroetorphine (**2a**) with ³H₂O yielded the enamine specifically labeled at C-15. Subsequent reduction of the iminium salt with NaBH₄ yielded [^{15}H]etorphine. If the reduction step is carried out in the presence of ³H₂O, the resulting product contains, in theory, two labeled atoms at C-15 (Scheme I). Other tritiated compounds in the series have been synthesized in this way. The use of NaB³H₄ yielded 15,16-³H derivatives which showed increased specific activity (Table I).

Scheme I



Etorphine was also labeled in the 15 and 16 positions by hydrogenation of the enamine double bond with ³H₂ gas, using 10% palladized charcoal as catalyst; the 6,14-etheno bridge is not reduced under these conditions.⁸ The labeled etorphine produced by this route had a higher specific activity than the products of the previous experiments (Table I).

The limiting value for the specific activity of the product in the hydrogenation route is 58 Ci/mmol and though such activities would be difficult to obtain in practice it is apparent that the latter approach is the method of choice in view of the high pharmacological activity of the series and the consequent need for high specific activity.[‡] It may not, however, be applicable to compounds containing

[‡]It would be possible, in fact, to increase further the activity of the hydrogenation product by preliminary equilibration of the enamine with tritiated water to yield, after reduction, a triply labeled product.

the cyclopropylmethyl group as these undergo reductive ring opening under hydrogenation conditions.⁸

The methods outlined above should prove to be of use in other series of compounds in which similar enamines can be generated and isolated, *e.g.*, meperidine and its analogs,⁹ dihydromorphines,¹⁰ morphinans, and benzomorphans,¹¹ provided such compounds contain no functional groups reactive toward both catalytic hydrogenation and borohydride reduction.

Experimental Section

Specific activities were determined in a Packard Tricarb 3003 scintillation spectrometer. Samples were counted in a cocktail containing naphthalene (100 g), 2,5-diphenyloxazole (5.0 g), and 1,4-bis[2-(5-phenyl)oxazolyl]benzene (0.05 g) in 1,4-dioxane (1000 ml). Chemical and radiochemical purities were estimated by tlc on Kieselgel 60 F₂₅₄ plates (5 × 20 cm) supplied by Merck Ag using the following solvent systems: (i) *n*-BuOH-AcOH-H₂O (20:5:8), (ii) CHCl₃-MeOH-H₂O (8:5:1), (iii) EtOAc-MeOH (75:25). Distribution of radioactivity on the plates was determined with a Tracerlab 4π scanner. Radiochemical starting materials were supplied by the Radiochemical Centre, Amersham, England. Experimental details are given for etorphine only. Other compounds were labeled in an analogous manner.

15,16-Didehydrooripavine Derivatives. 15,16-Didehydrooripavine derivatives were prepared from the parent tertiary bases by the method of Lewis, *et al.*⁵

6,14-endo-Etheno-7α-(1-hydroxy-1-methylbutyl)tetrahydro-[15-³H]oripavine ([15-³H]Etorphine). **2a** (100 mg, 0.24 mmol) was placed in a round-bottomed flask (25 ml) attached to a vacuum bridge. ³H₂O (*ca.* 20 Ci, 80 μl) was frozen in liquid nitrogen, dry dioxane (3 ml) added, and the mixture vacuum transferred to the reaction flask. The vacuum was released to allow the addition of NaBH₄ (150 mg, 3.9 mmol). The reaction mixture was stirred at ambient temperature for 2 hr and taken to dryness *in vacuo*, and K₂CO₃ (0.5 g) in H₂O (3 ml) and dioxane (3 ml) was added. The mixture was heated at 90° for 10 min and cooled, and the volatile materials were again removed *in vacuo*. H₂O (5 ml) was added and the mixture was extracted with CHCl₃. The CHCl₃ solution was washed repeatedly with H₂O, dried (Na₂SO₄), and taken to dryness to yield the crude product (60 mg). A portion of this material (14 mg) was chromatographed on a silica column (30 × 1 cm; Merck Ag Kieselgel 60, 0.063-0.200 mm) with Et₂O as eluent. Fractions which contained only the required product were united and treated with a little ethereal HCl to yield [15-³H]etorphine hydrochloride (6 mg, 16%); sp act. 260 mCi/mmol; radiochemical purity >98%.

6,14-endo-Etheno-7α-(1-hydroxy-1-methylbutyl)tetrahydro-[15,16-³H]oripavine ([15,16-³H]Etorphine). **Method a.** [15,16-³H]-Etorphine was synthesized by a route analogous to the method used for the 15-³H derivative, NaB³H₄ being used in the reduction step; sp act. 900 mCi/mmol; radiochemical purity of product >98%.

Method b. **2a** (25 mg, 0.06 mmol) and 10% palladized charcoal (10 mg) in ethyl acetate (3 ml) was placed in a flask connected by a vacuum bridge to a break-seal ampoule containing ³H₂ gas (*ca.* 10 Ci; 4 ml at STP). The solution was cooled in liquid nitrogen and the system was evacuated and closed. The reaction mixture was exposed to ³H₂ gas by breaking the ampoule seal. The flask was allowed to warm to room temperature and the mixture stirred for 2 days, after which time hydrogen was introduced *via* the bridge to restore atmospheric pressure and the reaction was continued for a further 18 hr. The system was thoroughly evacuated, the catalyst removed by filtration, and the resulting solution was taken to dryness *in vacuo*. The crude product was chromatographed on a silica plate (20 × 20 × 0.2 cm) with *n*-BuOH-H₂O-AcOH (20:8:5) as eluent. The band corresponding to **1a** was eluted with CH₃OH (20 ml) and the resulting solution taken to dryness *in vacuo*. The residue was dissolved in cold 0.1 N HCl (5 ml) and the aqueous solution rapidly basified (K₂CO₃) and extracted with CHCl₃ (3 × 5 ml). The united CHCl₃ extracts were washed several times with water, dried (Na₂SO₄), and taken to dryness and the residue was dissolved in dry Et₂O (5 ml). Carrier **1a** (30 mg) was added and the solution was treated with a few drops of anhydrous ethereal HCl to yield [15,16-³H]etorphine hydrochloride (45.8 mg, 48%); sp act. *ca.* 3.6 Ci/mmol; radiochemical purity >98%.

§H. P. Crocker, unpublished results.

References


- (1) K. E. Wilzbach, *J. Amer. Chem. Soc.*, **79**, 1013 (1957).
- (2) G. W. Kirby and L. Agimkoya, *J. Chem. Soc.*, 5785 (1965).
- (3) K. W. Bentley and D. G. Hardy, *J. Amer. Chem. Soc.*, **89**, 3281 (1967); J. M. King and B. H. Carter, *East Afr. Wild Life J.*, **3**, 19 (1965); A. M. Harthoorn and J. Bligh, *J. Res. Vet. Sci.*, **6**, 290 (1965).
- (4) A. C. Lane, A. McCoubrey, and R. Peaker, *J. Label. Compounds*, **2**, 284 (1966).
- (5) D. I. Haddlesey, J. W. Lewis, P. A. Mayor, and G. R. Young, *J. Chem. Soc., Perkin Trans. 1*, 872 (1972).
- (6) P. S. Portoghese and D. L. Larson, *J. Med. Chem.*, **16**, 420 (1973).
- (7) N. J. Leonard, A. S. Hay, R. W. Fulmor, and D. W. Cash, *J. Amer. Chem. Soc.*, **77**, 439 (1955).
- (8) K. W. Bentley, D. G. Hardy, and B. Meek, *J. Amer. Chem. Soc.*, **89**, 3273 (1967).
- (9) J. W. Lewis and P. A. Mayor, *J. Chem. Soc. C*, 1074 (1970).
- (10) J. W. Lewis, British Patent No. 1,223,445 (1968).
- (11) J. W. Lewis, British Patent No. 1,224,918 (1968).

Synthesis and Biological Activity of 2'-Amino-2'-deoxy-5-fluorouridine

R. A. Sharma, M. Bobek, and A. Bloch*

Department of Experimental Therapeutics and J. T. Grace, Jr., Cancer Drug Center, Roswell Park Memorial Institute, New York State Department of Health, Buffalo, New York 14203. Received October 16, 1973

Recently, Verheyden, Wagner, and Moffatt¹ reported a useful synthesis of 2'-amino-2'-deoxyuridine and of its 5-iodo derivative. Although no information on the biological activity of these compounds has been published, it appeared worthwhile to us to prepare the 5-fluoro derivative of 2'-amino-2'-deoxyuridine, in view of the potent biological activity of other fluorinated pyrimidines.²



Chemical Results. Preparation of 2'-amino-2'-deoxy-5-fluorouridine (**4**) was first attempted by direct fluorination of 2'-amino-2'-deoxyuridine (**1**) with trifluoromethyl hypofluorite (CF₃OF), according to the methods described