

Derivatives of the Muscarinic Agent


N-Methyl-*N*-(1-methyl-4-pyrrolidino-2-butynyl)acetamide

Björn M. Nilsson,[†] Björn Ringdahl,[‡] and Uli Hacksell*[†]


Department of Organic Pharmaceutical Chemistry, Uppsala Biomedical Center, University of Uppsala, S-751 23 Uppsala, Sweden, and Department of Pharmacology, School of Medicine, University of California, Los Angeles, California 90024. Received June 19, 1987

A series of tertiary and quaternary analogues (acyclic imides, sulfonimides, *N*-acetyl sulfonamides, and trifluoroacetamides) of the selective partial muscarinic agonist *N*-methyl-*N*-(1-methyl-4-pyrrolidino-2-butynyl)acetamide (BM 5, **35**) was synthesized. The compounds were found to be muscarinic agonists, partial agonists, or antagonists in the isolated guinea pig ileum. Replacement of the acetyl group or the *N*-methyl group of **35** and its analogues by a methanesulfonyl group abolished efficacy and decreased affinity at ileal muscarinic receptors. Trifluoroacetamide analogues of **35** also had lower affinity and efficacy than **35**. Substitution of an acetyl group for the *N*-methyl group in compounds related to **35** decreased efficacy, but had no appreciable effect on affinity. Most of the tertiary amines showed central antimuscarinic activity as they antagonized oxotremorine-induced tremors in mice.

The oxotremorine analogue *N*-methyl-*N*-(1-methyl-4-pyrrolidino-2-butynyl)acetamide (BM 5, **35**)¹ has been reported to be a presynaptic antagonist and postsynaptic agonist at muscarinic receptors *in vitro*² and *in vivo*.³ Racemic **35**,¹ as well as its pure enantiomers,⁴ was found to block oxotremorine-induced tremors. The latter ob-



OXOTREMORINE



- 34**: R = H, Am = NC₄H₈
35: R = CH₃, Am = NC₄H₈
36: R = CH₃, Am = N(CH₃)₂
37: R = CH₃, Am = N⁺(CH₃)₃

servation suggests that **35** acts mainly as an antagonist in brain regions that are responsible for the tremor response. In other regions of the brain, e.g., those involved in the production of analgesia and hypothermia, **35** acts as a muscarinic agonist.⁵ Both enantiomers of **35** have been classified as partial muscarinic agonists in the guinea pig ileum, the *R* enantiomer being the more potent.⁴ The partial agonist properties of **35** have been invoked to explain its selective actions *in vivo*⁵ and *in vitro*.⁶

Compounds with pharmacological profiles similar to that of **35** have potential for the therapy of diseases associated with deficiencies in cholinergic transmission, e.g., Huntington's chorea and Alzheimer-type dementia.^{2,7}

Previous structure-activity relationship studies of flexible oxotremorine analogues have demonstrated that the muscarinic activity is modified by changes of the nature or size of the amide substituents.⁸ To investigate further the effects of structural modifications in the amide moiety in compounds related to **35**, we have now synthesized compounds **16-33** (Table II).

The new compounds were investigated for tremorogenic and tremorolytic activity in mice and for muscarinic and antimuscarinic activity on the isolated guinea pig ileum (Table III). Some compounds were partial agonists in the ileum and others were relatively potent tremorolytic agents, but none showed both of these effects (cf. **35**). However, **23**, **26**, and **30** were fairly potent antagonists to carbachol in the guinea pig ileum.

Chemistry

Synthesis. Standard methods starting from propargylamine or 1-methyl-2-propynylamine were used to obtain target compounds **16-33** (see the Experimental Section). The final step in the preparation of the tertiary amines involved a Mannich condensation (method H) between an acetylenic intermediate, the appropriate secondary amine, and paraformaldehyde in the presence of a catalytic amount of cuprous chloride. The quaternary ammonium derivatives were prepared by methylation of the corresponding tertiary Mannich bases (method I).

Various reaction conditions were investigated for the preparation of the *N,N*-diacetylated intermediates *N*-(2-propynyl)diacetamide (**10**) and *N*-(1-methyl-2-propynyl)diacetamide (**11**). Attempts to prepare **10** by use of the method described by Dahlbom et al.⁹ for the preparation of *N*-propargylsuccinimide (diacetamide, propargyl bromide, NaOEt in EtOH) were unsuccessful, probably due to decomposition of the diacetamide. However, **10** could be prepared by alkylation of diacetamide with propargyl bromide in the presence of potassium carbonate in acetonitrile (method G). Compound **11** was prepared from **9** by using acetyl chloride and triethylamine in a large excess (method F). It is noteworthy that attempts to prepare **11** from 1-methyl-2-propynyl *p*-toluenesulfonate by method G were unsuccessful.

In general, the acyclic imides appear to be considerably less stable than their previously reported^{9,10} cyclic analogues; partial decomposition was observed when crude **10** was passed through an alumina column eluted with

- (1) Resul, B.; Dahlbom, R.; Ringdahl, B.; Jenden, D. J. *Eur. J. Med. Chem.-Chim. Ther.* **1982**, *17*, 317.
- (2) Norström, Ö.; Alberts, P.; Westlind, A.; Undén, A.; Bartfai, T. *Mol. Pharmacol.* **1983**, *24*, 1.
- (3) Casamenti, F.; Cosi, C.; Pepeu, G. *Eur. J. Pharmacol.* **1986**, *122*, 288.
- (4) Dahlbom, R.; Jenden, D. J.; Resul, B.; Ringdahl, B. *Br. J. Pharmacol.* **1982**, *76*, 299.
- (5) Ringdahl, B.; Roch, M.; Jenden, D. J. *J. Pharmacol. Exp. Ther.* **1987**, *242*, 464.
- (6) Ringdahl, B. *Mol. Pharmacol.* **1987**, *31*, 351.
- (7) Hershenson, F.; Moos, W. H. *J. Med. Chem.* **1986**, *29*, 1125.
- (8) For reviews, see: Dahlbom, R. In *Cholinergic Mechanisms: Phylogenetic Aspects, Central and Peripheral Synapses, and Clinical Significance*; Pepeu, G., Ladinsky, H., Eds.; Plenum: New York, 1981; p 621. Ringdahl, B.; Jenden, D. J. *Life Sci.* **1983**, *32*, 2401.
- (9) Dahlbom, R.; Karlén, B.; George, R.; Jenden, D. J. *J. Med. Chem.* **1966**, *9*, 843.
- (10) Karlén, B.; Lindeke, B.; Lindgren, S.; Svensson, K.-G.; Dahlbom, R.; Jenden, D. J.; Giering, J. E. *J. Med. Chem.* **1970**, *13*, 651.

[†]University of Uppsala.

[‡]University of California.

Table I. Physical Data of the Intermediates

compd	R ¹	R ²	R ³	prepn meth ^a	yield, %	bp (mmHg) or mp, °C	formula
1	CH ₃ SO ₂	H	H	A	71	55.5–58 ^b	C ₄ H ₇ NO ₂ S
2	CH ₃ SO ₂	H	CH ₃	A	90	46–48 ^b	C ₅ H ₉ NO ₂ S
3	CH ₃ SO ₂	CH ₃	H	C	60	76–78 (0.3)	C ₆ H ₉ NO ₂ S ^{1/4} H ₂ O
4	CH ₃ SO ₂	CH ₃	CH ₃	C	97	73–74 ^c	C ₆ H ₁₁ NO ₂ S
5	CH ₃ SO ₂	CH ₃ SO ₂	H	D	55	111.5–112.5 ^c	C ₆ H ₉ NO ₄ S ₂
6	CH ₃ SO ₂	CH ₃ SO ₂	CH ₃	D	29	96–99 ^c	C ₆ H ₁₁ NO ₄ S ₂
7	CH ₃ SO ₂	CH ₃ CO	H	E	91	45–46 ^d	C ₆ H ₉ NO ₃ S
8	CH ₃ SO ₂	CH ₃ CO	CH ₃	E	80	73–75 (0.3)	C ₇ H ₁₁ NO ₃ S
9	CH ₃ CO	H	CH ₃	e	81	91–92 ^c	
10	CH ₃ CO	CH ₃ CO	H	G	94	136–137 (48)	C ₇ H ₉ NO ₂ ^f
11	CH ₃ CO	CH ₃ CO	CH ₃	F	65	94–95 (10)	C ₈ H ₁₁ NO ₂ ^g
12	CF ₃ CO	H	H	B	65	h	
13	CF ₃ CO	H	CH ₃	B	74	70 (18)	C ₆ H ₆ F ₃ NO ⁱ
14	CF ₃ CO	CH ₃	H	C	56	57 (15)	C ₆ H ₆ F ₃ NO ^j
15	CF ₃ CO	CH ₃	CH ₃	C	72	50–51 (10)	C ₇ H ₈ F ₃ NO ^k

^a See the Experimental Section. ^b From ether–petroleum ether. ^c From ether. ^d From ether–cyclohexane. ^e Previously described (ref 1), mp 92.5–93 °C. ^f Anal. Calcd for C₇H₉NO₂: C, 60.4; N, 10.1. Found: C, 59.2; N, 9.5. MS (12 eV) *m/z* 97 (M⁺ – CH₂CO). ^g Anal. Calcd for C₈H₁₁NO₂: C, 62.7. Found: C, 62.2. MS (12 eV) *m/z* 111 (M⁺ – CH₂CO). ^h Previously reported (ref 30), bp 95 °C (11 mmHg). ⁱ Anal. Calcd for C₆H₆F₃NO: C, 43.6. Found: C, 42.0. MS (70 eV) *m/z* 150 (M⁺ – CH₃). ^j Anal. Calcd for C₆H₆F₃NO: N, 8.5. Found: N, 9.4. MS (70 eV) *m/z* 165 (M⁺). ^k Anal. Calcd for C₇H₈F₃NO: C, 46.9; N, 7.8. Found: C, 45.3; N, 7.3. MS (70 eV) *m/z* 165 (M⁺ – CH₃).

Table II. Physical Data for the Compounds Tested

compd	R ¹	R ²	R ³	R ^{4a}	yield, %	recrystn solvents ^b	mp, °C	formula ^c
16	CH ₃ SO ₂	CH ₃	H	NC ₄ H ₈	65	A	76–79	C ₁₀ H ₁₈ N ₂ O ₂ S·C ₂ H ₂ O ₄
17	CH ₃ SO ₂	CH ₃	CH ₃	NC ₄ H ₈	64	A	112–114.5	C ₁₁ H ₂₀ N ₂ O ₂ S·C ₂ H ₂ O ₄
18	CH ₃ SO ₂	CH ₃ SO ₂	H	NC ₄ H ₈	80	B	145.5–146.5	C ₁₀ H ₁₈ N ₂ O ₄ S ₂ ·C ₂ H ₂ O ₄
19	CH ₃ SO ₂	CH ₃ SO ₂	CH ₃	NC ₄ H ₈	84	B	162–163	C ₁₁ H ₂₀ N ₂ O ₄ S ₂ ·C ₂ H ₂ O ₄
20	CH ₃ SO ₂	CH ₃ CO	H	NC ₄ H ₈	90	C	124–126	C ₁₁ H ₁₈ N ₂ O ₃ S·C ₂ H ₂ O ₄
21	CH ₃ SO ₂	CH ₃ CO	CH ₃	NC ₄ H ₈	70	A	120–122	C ₁₂ H ₂₀ N ₂ O ₃ S·C ₂ H ₂ O ₄
22	CF ₃ CO	CH ₃	H	NC ₄ H ₈	75	A	83–84	C ₁₁ H ₁₆ F ₃ N ₂ O·C ₂ H ₂ O ₄
23	CF ₃ CO	CH ₃	CH ₃	NC ₄ H ₈	40	A	60–62.5	C ₁₂ H ₁₇ F ₃ N ₂ O·C ₂ H ₂ O ₄ ^{1/4} H ₂ O
24	CF ₃ CO	CH ₃	CH ₃	N(CH ₃) ₂	28	A	117–119	C ₁₀ H ₁₆ F ₃ N ₂ O·C ₂ H ₂ O ₄
25	CF ₃ CO	CH ₃	CH ₃	N(CH ₃) ₃ ⁺	54	C	163–165	C ₁₁ H ₁₆ F ₃ N ₂ O·I
26	CH ₃ CO	CH ₃ CO	H	NC ₄ H ₈	85	C	70–72	C ₁₂ H ₁₈ N ₂ O ₂ ·C ₂ H ₂ O ₄
27	CH ₃ CO	CH ₃ CO	H	N(C ₂ H ₅) ₂	65	B	120–121.5	C ₁₂ H ₂₆ N ₂ O ₂ ·C ₂ H ₂ O ₄
28	CH ₃ CO	CH ₃ CO	H	N(CH ₃) ₂	46	A	108–110	C ₁₀ H ₁₆ N ₂ O ₂ ·C ₂ H ₂ O ₄
29	CH ₃ CO	CH ₃ CO	H	N(CH ₃) ₃ ⁺	38	D	109–111	C ₁₁ H ₁₆ N ₂ O ₂ ·I
30	CH ₃ CO	CH ₃ CO	CH ₃	NC ₄ H ₈	43	E	78.5–79.5	C ₁₃ H ₂₀ N ₂ O ₂ ·C ₂ H ₂ O ₄ ^{1/4} H ₂ O
31	CH ₃ CO	CH ₃ CO	CH ₃	N(C ₂ H ₅) ₂	55	A	81.5–82.5	C ₁₃ H ₂₂ N ₂ O ₂ ·C ₂ H ₂ O ₄ ^{1/4} H ₂ O
32	CH ₃ CO	CH ₃ CO	CH ₃	N(CH ₃) ₂	66	A	87.5–89.5	C ₁₁ H ₁₈ N ₂ O ₂ ·C ₂ H ₂ O ₄
33	CH ₃ CO	CH ₃ CO	CH ₃	N(CH ₃) ₃ ⁺	87	D	109–110.5	C ₁₂ H ₂₁ N ₂ O ₂ ·I

^a NC₄H₈ = pyrrolidino. ^b Recrystallization solvents: A, EtOH–ether; B, EtOH; C, MeOH–ether; D, acetone–ether; E, acetone. ^c All compounds were analyzed for C, H, and N. The analytical results obtained for those elements were within ±0.4% of the theoretical value.

CH₂Cl₂, and aminolysis occurred when excess of the secondary amine was used in the Mannich reactions of compounds 10 and 11. For example, a 6:1 mixture of imide 27 and the corresponding secondary amide was formed when compound 10 was subjected to a Mannich reaction in the presence of 1.25 equiv of diethylamine.

Physical data of the synthetic intermediates 1–15 and of the target compounds 16–33 are presented in Tables I and II, respectively.

Spectroscopy. NMR spectroscopic data of the compounds are given in the supplementary material. Some compounds gave informative spectra that allowed determination of certain conformational characteristics.

It was evident from ¹H, ¹³C, and ¹⁹F NMR spectroscopy that the trifluoroacetamides exist as an equilibrium mixture of *E* and *Z* conformations in solution. Resonances could be assigned to *E* or *Z* conformations on the basis of

the magnitude of the fluorine–proton couplings observed in the ¹H and ¹⁹F NMR spectra and on the magnitude of the carbon–fluorine couplings in the ¹³C NMR spectra. The relative proportions of the two conformations were determined by integration over appropriate resonances in the ¹H and ¹⁹F NMR spectra (information obtained from ¹⁹F NMR spectroscopy is collected in Table IV). For example, integration of the *N*-Me resonances due to the *Z* (δ 3.17; ⁵J_{H,F} = 1.7 Hz) and *E* (δ 3.05; ⁵J_{H,F} < 1 Hz) conformations of the oxalate of 23 in CD₃OD indicated that the *Z* conformation predominated (4:1) at 20 °C. Corroborating results were obtained by integration of the quartet (δ –69.7; *Z* conformation) and the broad singlet (δ –68.5; *E* conformation), which were observed as the sole resonances in the ¹⁹F NMR spectrum of 23. The *N*-Me resonance due to the *Z* conformation appeared as a well-resolved quartet (δ 30.7; J_{C,F} ≈ 4 Hz) in the proton de-

Table III. Muscarinic and Antimuscarinic Effects of Some Oxotremorine Analogues^a

compd	guinea pig ileum				relative efficacy ^e	tremorolytic dose in mice, ^f μmol/kg
	EC ₅₀ , ^b μM	E _{max} ^c	EPMR ^d	K _D , μM		
16				68.0 ± 8.6 (4)		174 ± 51
17				13.0 ± 3.5 (4)		82.0 ± 17.6
18				15.9 ± 4.3 (5)		>200
19				5.4 ± 0.06 (4)		157 ± 44
20				17.9 ± 2.0 (5)		>200
21				3.1 ± 0.3 (5)		99.7 ± 23.0
22	16.9 ± 5.2 (5)	0.16 ± 0.03 (5)	145 ± 38 (5)	15.6 ± 3.4 (4)	0.0030 ± 0.0004 (4)	150 ± 41
23		0		0.57 ± 0.03 (7)		21.4 ± 2.5
24		<0.1		95.4 ± 12.6 (5)		>200
25	27.1 ± 3.5 (7)	0.89 ± 0.04 (7)	214 ± 24 (7)	91.9 ± 22.1 (5)	0.026 ± 0.004 (5)	
26		0		0.44 ± 0.06 (5)		2.9 ± 0.5
27		0		7.2 ± 0.6 (5)		107 ± 19
28	38.8 ± 10.1 (6)	0.84 ± 0.07 (6)	320 ± 12 (6)	65.6 ± 11.9 (5)	0.013 ± 0.003 (5)	>200
29	2.0 ± 0.3 (8)	1.00 ± 0.02 (8)	19.3 ± 1.8 (8)	8.2 ± 2.1 (7)	0.034 ± 0.008 (7)	
30		0		0.28 ± 0.04 (7)		3.2 ± 1.0
31		0		1.6 ± 0.2 (6)		39.0 ± 5.3
32	5.5 ± 0.7 (6)	0.99 ± 0.03 (6)	57.9 ± 2.9 (6)	21.6 ± 6.5 (5)	0.030 ± 0.006 (5)	109 ± 11.3
33	1.2 ± 0.1 (8)	1.01 ± 0.02 (8)	10.7 ± 0.7 (8)	8.6 ± 1.9 (7)	0.055 ± 0.01 (7)	
34	0.069 ± 0.002 (7)	1.01 ± 0.01 (7)		2.2 ± 0.3 (6)	0.15 ± 0.02 (6)	g
35	0.19 ± 0.03 (4)	0.83 ± 0.003 (4)		0.24 ± 0.07 (4)	0.013 ± 0.0002 (4)	0.6
38	1.0 ± 0.03 (8)	0.98 ± 0.01 (8)		5.0 ± 0.5 (5)	0.027 ± 0.002 (5)	23
40	2.5 ± 0.1 (9)	1.01 ± 0.01 (9)		208 ± 30 (6)	0.35 ± 0.07 (6)	g
42				0.15 ± 0.01 (4) ^h		1.2 ⁱ
carbachol	0.11 ± 0.01 (6)	1.00	1.00	15.4 ± 3.1 (6)	1.00	

^a Values represent means plus or minus standard errors. ^b The concentration of an agonist or a partial agonist that elicits 50% of its own maximum contractile response. ^c The maximum contractile response relative to that elicited by carbachol. ^d Equipotent molar ratio relative to that of carbachol, which equals 1.00. ^e Efficacy relative to that of carbachol; calculated from EC₅₀ and K_D values as described in ref 6 and 15. ^f Dose required to double the dose of oxotremorine inducing a predetermined (grade 2) tremor intensity in mice. ^g Agonist. ^h Unpublished. ⁱ From ref 10.

Table IV. ¹⁹F NMR Spectral Data of Some Trifluoroacetamides^a

compd	solvent	chemical shifts, δ		
		Z conformer ^b	E conformer ^b	% Z
12	CDCl ₃		-76.7	
13	CDCl ₃		-76.3	
14	CDCl ₃	-70.4	-69.6	72
15	CDCl ₃	-70.2	-69.0	77
22	CD ₃ OD	-69.5	-68.6	74
23	CD ₃ OD	-69.7	-68.5	79
24	CD ₃ OD	-69.8	-68.6	81
25	CD ₃ OD	-69.7	-68.5	83

^a Spectra were referenced to internal CFC1₃. ^b Assignment was based on the magnitude of the ⁵J_{F,H} coupling constants.

coupled ¹³C NMR spectrum of the oxalate of 23. In the *E* conformation of 23, this carbon was observed as a broad unresolved resonance (δ 30.0). The other trifluoroacetamides (22, 24, and 25) were found to have similar conformational preferences (*Z/E* around 4:1; compare Figure 1). In CDCl₃, the acetamide analogue (base) of 22 exists to about 60% in the *Z* conformation at 25 °C,¹¹ and the base form of acetamide 35 assumes a 3:1 ratio of the *Z* and *E* conformations at 37 °C in the same solvent.¹ Integration of the *N*-Me resonances in the spectrum of the oxalate of 35 in CD₃OD at 25 °C shows that the *Z* conformation is considerably more favored under these latter conditions (*Z/E* = 4:1).

Acyclic imides can assume *E,E*, *E,Z*, or *Z,Z* conformations.¹² In ¹H NMR spectra of CD₃OD solutions of the oxalates of 26–28 and 30–32, the two acetyl groups appeared as a single resonance at 20 °C. These observations indicate either that 26–28 and 30–32 assume mainly one




Figure 1. Partial ¹⁹F NMR spectrum of 24 in CD₃OD (at 20 °C), which demonstrates the predominance of the *Z* conformation.

symmetric imide conformation or that the barrier to conformational interconversion is low. In contrast, ¹H NMR spectra of the methiodides 29 and 33 in CD₃OD at 20 °C exhibited two resonances (29, δ 1.97 and 2.02; 33, δ 1.96 and 2.02) due to the acetyl groups that were of equal intensity and were assigned to the unsymmetrical *E/Z* conformation.

The IR spectra of 29 and 33 (KBr pellets) showed two carbonyl absorption bands, one of stronger intensity at ν 1715 cm⁻¹ and one of medium intensity at ν 1695–1700

- (11) Resul, B.; Ringdahl, B.; Hacksell, U.; Svensson, U.; Dahlbom, R. *Acta Pharm. Suec.* 1979, 16, 225.
 (12) (a) Noe, E.; Raban, M. *J. Am. Chem. Soc.* 1973, 95, 6118; 1975, 97, 5811. (b) Tetu, G.; Duplan, J.-C.; Pellissier, N.; Laurent, E. *J. Chem. Res. Synopses* 1978, 98.

cm^{-1} . Only a single strong carbonyl absorption band at ν 1710 cm^{-1} was observed for the bases of the corresponding tertiary amines (liquid film).

Pharmacology

Compounds 16–33 were tested for muscarinic and antimuscarinic activity in the isolated guinea pig ileum and in intact mice. The results are presented in Table III. For comparison, relevant data for 34, 35, 38, 40, and 42 are also included in the table. In Table III, E_{max} values are equivalent to intrinsic activities in the terminology of Ariens.¹³ K_D is the equilibrium dissociation constant of the drug–receptor complex and is a measure of affinity.¹⁴ K_D values were determined on the ileum by methods¹⁵ that also provided reliable estimates of agonist efficacy.¹⁶

None of the *N*-methanesulfonyl-substituted analogues (16–21) showed spasmogenic activity on the guinea pig ileum. Instead, they were antagonists to carbachol on this tissue. Spasmolytic activity as well as tremorolytic activity among 16–21 increased with methyl substitution in position 1 of the butynyl chain (Table III).

The trifluoroacetamides 22 and 25 were about 100-fold less potent than carbachol on the ileum. They elicited contractile responses that were significantly ($P < 0.05$) lower than those produced by carbachol. Compound 24 also appeared to be a partial agonist, but its efficacy was too low to be measured. In contrast, 23 was a pure antagonist having relatively high affinity for muscarinic receptors.

Among the acyclic imides (26–33), the dimethylamino (28 and 32) and trimethylammonium (29 and 33) derivatives were full or nearly full agonists on the ileum, but they were more than 10-fold less potent than carbachol. This was due primarily to lower efficacy, since the affinities of 28, 29, 32, and 33 were similar to that of carbachol. The pyrrolidine derivatives 26 and 30 were potent antagonists to carbachol on the ileum and to the central effects of oxotremorine in vivo. Trifluoroacetamides and acyclic imides are known to be sensitive to hydrolysis.^{17,18} The possibility that the compounds studied were broken down in the tissue bath was not investigated in detail. However, the high potency of 23, 26, and 30 in vitro suggests that hydrolysis was not extensive.



Among the tertiary amines, only 32 showed visible muscarinic effects in vivo; doses of 32 that produced tremor blockade also gave profuse salivation. The trimethylammonium salts were not tested in vivo.

Discussion

Replacement of the acetyl or *N*-methyl group (or both) of 35 by a methanesulfonyl group decreased affinity and abolished efficacy at muscarinic receptors in the guinea pig ileum as the resulting compounds (17, 19, and 21) were relatively weak antagonists (Table III). Similarly, substitution of a trifluoroacetyl group for the acetyl group of 35 (yielding 23) abolished efficacy, but caused only a 2-fold reduction in affinity. The trifluoroacetamides 22 and 25 were considerably less potent in causing contractions of the ileum than the previously reported acetamide analogues 34¹⁹ and 37.⁴ Compounds 22–24 were relatively poor


tremorolytic agents in vivo. This may be due partly to the hydrolytic instability of these compounds.¹⁷ The trifluoroacetamides and the acetamides appear to have similar conformational preferences (see above). Thus, the lower in vitro potency of the trifluoroacetamides as compared to the analogous acetamides does not appear to be related to conformational differences. Instead, it may be related to the decreased oxygen-electron density (increased C=O stretching frequency; 23, ν 1695 cm^{-1} , and 35, ν 1640 cm^{-1}) in the fluorinated amides (compare ref 20).

Substitution of an acetyl group for the *N*-methyl group of 34–37 decreased (32 and 33) or abolished (26 and 30) efficacy without having much effect on affinity. The acyclic imides 28 and 29 were about 15-fold less potent on the ileum than the corresponding succinimides 40 and 41.²¹



This was due entirely to lower efficacy of 28 and 29 as they had somewhat higher affinity than their succinimide analogues. Similarly, the antagonists 26 and 27 had 5- to 10-fold higher affinity than the succinimides 38 and 39.²¹

It is noteworthy that 38 was an agonist, albeit of low efficacy,²¹ whereas 26 was an antagonist, in agreement with the lower efficacy of the acyclic imides. It is tempting to suggest that the 10-fold higher affinity of 26 as compared to 38 is due to the presence of the *E/Z* conformer 26a which resembles closely the potent antagonist 43.²² The



10-fold higher affinity of 43 as compared to oxotremorine was explained by additional binding of the methyl group to the receptor.^{23,24} It seems likely that the *Z*-methyl group of 26a also can participate in such binding.

Methyl substitution in the butynyl chain of 26–29 had little effect on affinity and efficacy. In contrast, analogous methyl substitution in the succinimides 38–41 in general leads to substantial increase in affinity and sharp decrease in efficacy. For example, addition of a methyl group in position 1 of the butynyl chain of 38, giving 42, increased affinity for muscarinic receptors 30-fold and abolished efficacy (Table III).^{10,25} The tremorolytic potencies of the acyclic imides were similar to or lower than those of the

- (13) Ariens, E. J., Ed. *Molecular Pharmacology. The Mode of Action of Biologically Active Compounds* Academic: New York, 1964; Vol. 1.
 (14) Furchgott, R. F. *Adv. Drug Res.* 1966, 3, 21.
 (15) Ringdahl, B. *J. Pharmacol. Exp. Ther.* 1984, 229, 199.
 (16) Stephenson, R. P. *Br. J. Pharmacol. Chemother.* 1956, 11, 379.
 (17) Greene, T. W. *Protective Groups in Organic Synthesis*; Wiley: New York, 1981; p 254.
 (18) Corey, E. J.; Sheehan, J. C. *J. Am. Chem. Soc.* 1952, 74, 4555.

- (19) Bebbington, A.; Brimblecombe, R. W.; Shakeshaft, D. *Br. J. Pharmacol. Chemother.* 1966, 26, 56. Ringdahl, B.; Resul, B.; Jenden, D. J.; Dahlbom, R. *Eur. J. Pharmacol.* 1982, 85, 79.
 (20) Bebbington, A.; Brimblecombe, R. W. *Adv. Drug Res.* 1965, 2, 143.
 (21) Ringdahl, B. *Eur. J. Pharmacol.* 1987, 140, 13.
 (22) Ringdahl, B.; Muhi-Eldeen, Z.; Ljunggren, C.; Karlén, B.; Resul, B.; Dahlbom, R.; Jenden, D. *Acta Pharm. Suec.* 1979, 16, 89.
 (23) Ringdahl, B.; Jenden, D. *J. Mol. Pharmacol.* 1983, 23, 17.
 (24) Amstutz, R.; Ringdahl, B.; Karlén, B.; Roch, M.; Jenden, D. *J. Med. Chem.* 1985, 28, 1760.
 (25) Ringdahl, B. *Proceedings of the 8th International Symposium on Medicinal Chemistry*; Dahlbom, R., Nilsson, J. L. G., Eds.; Swedish Pharmaceutical: Stockholm, 1985; Vol. 2, p 391.

corresponding succinimides with the exception of **26**, which was almost 10-fold more potent than **38**. It should be noted that acyclic imides are less stable toward solvolysis than cyclic imides¹⁸ and that this may influence considerably the in vivo data.

In summary, the structural modifications of **35**, which are described herein, generally gave compounds with lower efficacies than that of **35**. Compounds showing agonist or partial agonist properties had lower affinity than **35**. However, some of the compounds were fairly potent antagonists.

Experimental Section

Chemistry. General Comments. Melting points (uncorrected) were determined in open glass capillaries on a Thomas-Hoover apparatus. IR spectra were recorded on a Perkin-Elmer 298 infrared spectrophotometer. ¹H, ¹⁹F, and ¹³C NMR spectra were recorded on a JEOL FX 90Q spectrometer at 90, 84.3, and 22.5 MHz, respectively. ¹H NMR and ¹³C NMR spectra were referenced to internal tetramethylsilane. ¹⁹F NMR spectra were referenced to internal CFCl₃. Mass spectra were obtained at 70 eV on a LKB 9000 spectrometer using a direct insertion probe. All spectra were in accordance with the assigned structures. The elemental analyses, which were performed by Mikro Kemi AB, Uppsala, Sweden, were within ±0.4% of the calculated values unless otherwise noted.

N-(2-Propynyl)methanesulfonamide (1). Method A. A solution of methanesulfonyl chloride (4.0 g, 35 mmol) in dry ether was carefully added to a cooled (0 °C) solution of propargylamine (3.8 g, 69 mmol) in dry ether (60 mL) under N₂. The resulting mixture was stirred for 30 min, and the precipitate that formed was filtered off and washed with ether. Concentration of the combined ether layers afforded the crude sulfonamide, which was purified by recrystallization: ¹H NMR (CDCl₃) δ 5.30 (br s, NH), 3.96 (dd, N-CH₂), 3.11 (s, CH₃SO₂), 2.43 (t, J = 2.5 Hz, CCH); ¹³C NMR (CDCl₃) δ 79.06 (C2), 73.32 (C3), 41.35 (CH₃SO₂), 32.55 (C1).

When triethylamine or pyridine was used as acid scavengers, the yield of **1** was lower, and some dimesylation occurred; for example, when **1** equiv of pyridine was used, the crude reaction product consisted of **1** and dimesylated product (**5**) in a 5:1 ratio.

N-(1-Methyl-2-propynyl)trifluoroacetamide (13). Method B. A solution of trifluoroacetic anhydride (9.0 g, 43 mmol) in dry ether (30 mL) was added to a stirred solution of 1-methyl-2-propynylamine²⁶ (2.7 g, 39 mmol) and pyridine (3.1 g, 39 mmol) in dry ether (40 mL) under N₂ at 0 °C. After 30 min, the pyridinium trifluoroacetate was filtered off, and the ether was evaporated. The oily residue was purified by distillation: ¹H NMR (CDCl₃) δ 6.70 (br s, NH), 4.82 (m, N-CH), 1.51 (d, J = 6.9 Hz, CCH₃), 2.37 (d, J = 2.4 Hz, CCH); ¹³C NMR (CDCl₃) δ 158.06 (q, J_{C,F} = 38 Hz, CF₃CO), 116.06 (q, J_{C,F} = 287 Hz, CF₃CO), 38.11 (C1), 21.37 (C1 Me), 82.00 (C2), 71.90 (C3).

N-Methyl-N-(2-propynyl)methanesulfonamide (3). Method C. A solution of iodomethane (0.65 g, 4.6 mmol) in dry acetone was added to a stirred mixture of **1** (0.52 g, 3.9 mmol) and a large excess of anhydrous powdered K₂CO₃ (8.0 g, 58 mmol) in dry acetone (75 mL). The reaction mixture was heated under N₂ for 5 h. Ether (25 mL) was added, and the mixture was filtered. Concentration of the ether layer afforded **3** as an oil, which was purified by distillation: ¹H NMR (CDCl₃) δ 4.08 (d, J = 2.5 Hz, N-CH₂), 2.95, 2.92 (s's, CH₃SO₂, N-CH₃), 2.40 (t, J = 2.5 Hz, CCH); ¹³C NMR (CDCl₃) δ 76.93 (C2), 75.17 (C3), 39.41, 36.16 (N-CH₃, CH₃SO₂), 34.25 (C1).

N-(1-Methyl-2-propynyl)dimethanesulfonamide (6). Method D. Methanesulfonyl chloride (3.7 g, 32 mmol) was added dropwise to a mixture of N-(1-methyl-2-propynyl)sulfonamide (**2**; 1.0 g, 6.8 mmol), sodium hydroxide (0.31 g, 7.8 mmol), and water (7 mL). The mixture was stirred at room temperature for 12 h. NaOH (5 M, 10 mL) was added, and the solution was extracted twice with ether. The combined ether layers were dried (K₂CO₃) and concentrated to give crude **6**, which was purified by recrystallization: ¹H NMR (CDCl₃) δ 5.30 (dq, N-CH), 3.38 (s,

2 × CH₃SO₂), 2.57 (d, J = 2.4 Hz, CCH), 1.77 (d, J = 7.0 Hz, CCH₃); ¹³C NMR (CDCl₃) δ 80.73 (C2), 73.84 (C3), 47.62 (C1), 44.75 (2 × CH₃SO₂), 22.30 (C1 Me).

The reaction affording **5** was complete in 30 min.

N-Acetyl-N-(2-propynyl)methanesulfonamide (7). Method E. A solution of acetyl chloride (3.3 g, 42 mmol) in dry acetone (5 mL) was added to a mixture of **1** (2.3 g, 17 mmol) and anhydrous, powdered K₂CO₃ (30.0 g, 217 mmol) in dry acetone (100 mL). The reaction mixture was kept under N₂ at room temperature for 24 h. Filtration followed by evaporation of volatiles yielded crude **7** as an oil, which solidified upon cooling. Pure **7** was obtained after recrystallization: ¹H NMR (CDCl₃) δ 4.57 (d, J = 2.4 Hz, N-CH₂), 3.33 (s, CH₃SO₂), 2.46 (s, CH₃CO), 2.42 (t, J = 2.4 Hz, CCH); ¹³C NMR (CDCl₃) δ 170.23 (CH₃CO), 77.80 (C2), 73.23 (C3), 42.50 (CH₃SO₂), 35.30 (C1), 24.43 (CH₃CO).

N-(1-Methyl-2-propynyl)diacetamide (11). Method F. A solution of acetyl chloride (7.7 g, 98 mmol) in dry ether (10 mL) was slowly added to a precooled (0 °C) mixture of **9** (1.0 g, 9 mmol) and triethylamine (4.5 g, 45 mmol) in dry ether (100 mL). The reaction mixture was allowed to reach room temperature and was then stirred under N₂ for 24 h. The resulting mixture was filtered, and the volatiles were evaporated. Ether (100 mL) was added to the residue, and the ether solution was extracted with five 15-mL portions of saturated aqueous NaHCO₃. Concentration of the dried (K₂CO₃) organic layer, followed by chromatography on silica gel using CH₂Cl₂ as eluant, afforded **11** as a colorless oil: ¹H NMR (CDCl₃) δ 5.61 (dq, N-CH), 2.48 (s, 2 × CH₃CO), 2.39 (d, J = 2.6 Hz, CCH), 1.57 (d, J = 7.0 Hz, CCH₃); ¹³C NMR (CDCl₃) δ 172.89 (2 × CH₃CO), 82.40 (C2), 72.33 (C3), 40.80 (C1), 26.78 (2 × CH₃CO), 20.75 (C1 Me).

N-(2-Propynyl)diacetamide (10). Method G. Propargyl bromide (3.3 g, 28 mmol) was added to a stirred mixture of diacetamide²⁷ (2.0 g, 20 mmol) and dry powdered K₂CO₃ (20 g, 145 mmol) in acetonitrile (40 mL). The mixture was stirred under N₂ for 3 days at room temperature. The mixture was filtered, and the volatiles were evaporated. The oily residue was purified by flash chromatography on silica gel using CH₂Cl₂ as eluant: ¹H NMR (CDCl₃) δ 4.49 (d, J = 2.4 Hz, N-CH₂), 2.50 (s, 2 × CH₃CO), 2.27 (t, J = 2.4 Hz, CCH); ¹³C NMR (CDCl₃) δ 171.28 (2 × CH₃CO), 77.61 (C2), 70.91 (C3), 32.80 (C1), 25.23 (2 × CH₃CO).

N-Acetyl-N-[4-(diethylamino)-1-methyl-2-butynyl]acetamide (31). Method H. A solution of **11** (0.30 g, 1.96 mmol) in dioxane (5 mL) was added to a stirred mixture of paraformaldehyde (0.07 g, 2.33 mmol), diethylamine (0.14 g, 1.91 mmol), glacial acetic acid (0.12 g, 2 mmol), and a few grains of cuprous chloride in dioxane (10 mL). The mixture was stirred under N₂ at room temperature for 20 h. The volatiles were evaporated, and the oily residue was dissolved in 1 M HCl (40 mL). The water solution was extracted with ether (3 × 30 mL). Saturated aqueous NaHCO₃ was added to render the aqueous layer alkaline (pH ~8). Extraction with CH₂Cl₂ (6 × 45 mL) and drying (K₂CO₃) and concentration of the organic layer yielded crude **31** as an oily residue. The base was converted into the oxalate and recrystallized: ¹H NMR (CD₃OD) δ 5.31 (m, N-CH), 4.11 (d, J = 1.7 Hz, CH₂N), 3.30 (q, J = 7.3 Hz, N(CH₂C)₂), 2.41 (s, 2 × CH₃CO), 1.59 (d, J = 7.0 Hz, CCH₃), 1.32 (t, J = 7.3 Hz, N(CCH₃)₂); ¹³C NMR (CD₃OD) δ 174.40 (2 × CH₃CO), 90.12 (C2), 72.36 (C3), 48.92 (C4), 44.35 (C1), 41.69 (N(CH₂C)₂), 26.81 (2 × CH₃CO), 20.41 (C1 Me), 9.61 (N(CH₂C)₂).

Further Comments. Excess secondary amine (1.2 equiv) was added in the preparation of **16**–**24**. The reaction mixtures giving **16** and **17** were heated at 80 °C for 2.5–5 h. Aqueous NaOH was used to alkalize the acidic water layer in the workup of **16**–**19**. Crude **27** was contaminated with around 15% of the corresponding secondary amide (¹H NMR); IR spectroscopy of crude **27** revealed absorptions at ν 3300 cm⁻¹ (N—H stretching), 1650–1670 cm⁻¹ (C=O stretching), and 1540–1550 cm⁻¹ (N—H bending) due to the impurity. An IR spectrum of pure **27** (obtained after repetitive recrystallization from ethanol) showed a single carbonyl absorption at ν 1710 cm⁻¹.

(4-Diacetamidopent-2-ynyl)trimethylammonium Iodide (33). Method I. An excess of iodomethane (0.5 mL, 8 mmol) was added to the free base of **32** (generated from 0.26 g, 0.86 mmol,

of the oxalate) in dry acetone (8 mL). After 1.5 h at room temperature, ether was added, and the precipitate that formed was recrystallized: ^1H NMR (CD_3OD) δ 4.73 (m, N-CH), 4.37 (d, $J = 1.7$ Hz, CH_2N^+), 3.24 (s, $\text{N}^+(\text{CH}_3)_3$), 2.02 and 1.96 (2 s, CH_3CO), 1.44 (d, $J = 7.1$ Hz, CCH_3); ^{13}C NMR (CD_3OD) δ 173.32 and 172.18 (CH_3CO), 94.10 (C2), 70.60 (C3), 57.54 (C4), 53.58 ($\text{N}^+(\text{CH}_3)_3$), 38.02 (C1), 21.28 (C1 Me), 22.70 and 20.69 (CH_3CO).

Pharmacology. Guinea Pig Ileum. A standard guinea pig ileum preparation was set up in Tyrode solution (pH 7.4) at 37 °C as described previously.¹⁵ The Tyrode solution contained hexamethonium (0.03 mM). Contractions were recorded isotonically at 1 g of tension. Concentration–response curves were constructed by the cumulative dose–response technique by increasing stepwise the concentration of agonist by a factor of 2.15. Desensitization during cumulative addition of drugs was unlikely to occur since preliminary experiments showed that there was no difference in agonist potency measured by cumulative and intermittent administration.

Spasmogenic activity (EC_{50} values) was estimated by interpolation at the 50% response level of each compound. Maximal contractile responses (E_{max}) and equipotent molar ratios (EMPR) were expressed with reference to carbachol. Statistical significance ($P < 0.05$) of differences between mean E_{max} and EMPR values was determined by the Student's *t* test.

Dissociation constants (K_D) and relative efficacies of **22**, **25**, **28**, **29**, **32**, and **33** at muscarinic receptors in the ileum were determined by comparison of their concentration–response curves with that of carbachol as described previously for compounds acting as partial agonists.^{6,15} K_D values of antagonists were estimated by using carbachol as the agonist.^{15,21} The antagonist was allowed to equilibrate with the tissue for 15 min before the addition of carbachol. At least four different concentrations of each antagonist were used.

Tremorolytic Activity in Mice. Male Swiss–Webster mice (24–32 g body weight) were used. The test compounds were administered ip at a dose of 200 $\mu\text{mol}/\text{kg}$ to groups of six or more mice, while six control animals remained untreated. Twenty minutes after drug administration, the ED_{50} value of oxotremorine, injected iv, was estimated by the up-and-down method²⁸ with

intermittent spontaneous (grade 2) tremor²⁹ as the end point. Compounds that showed no significant antagonism of oxotremorine under these conditions were not tested further. Test of significance ($P < 0.05$) of differences between ED_{50} values of oxotremorine in the presence and absence of antagonist was performed, assuming that the ratio of the difference to its standard error was normally distributed. Compounds that significantly blocked oxotremorine at a dose of 200 $\mu\text{mol}/\text{kg}$ were tested at two additional doses. For the latter compounds, the ED_{50} value of oxotremorine was plotted against the dose of antagonist (including zero) used for premedication. That dose of antagonist that doubled the ED_{50} value of oxotremorine was estimated by linear regression analysis.^{5,24} Under these conditions, atropine and **35** had tremorolytic doses of 0.9 and 0.6 $\mu\text{mol}/\text{kg}$, respectively.

Acknowledgment. Support for this study was provided by grants from the Swedish Natural Science Research Council, Astra Alab Ab, and U.S. Public Health Service (Grants MH-17691 and GM-37816).

Registry No. 1, 93501-84-9; 2, 111903-20-9; 3, 111903-21-0; 4, 111903-22-1; 5, 111903-23-2; 6, 111903-24-3; 7, 111903-25-4; 8, 111903-26-5; 9, 42105-25-9; 10, 111903-27-6; 11, 111903-28-7; 12, 14719-21-2; 13, 111903-29-8; 14, 111903-30-1; 15, 111903-31-2; 16, 111903-33-4; 17, 111903-35-6; 18, 111903-37-8; 19, 111903-39-0; 20, 111903-41-4; 21, 111903-43-6; 22, 111903-45-8; 23, 111903-47-0; 24, 111903-49-2; 25, 111903-50-5; 26, 111903-52-7; 27, 111903-54-9; 28, 111903-56-1; 29, 111903-58-3; 30, 111903-60-7; 31, 111903-62-9; 32, 111903-64-1; 33, 111903-65-2; butynylamine, 41282-40-0; dimethylamine, 124-40-3; trimethylamine, 75-50-3; diethylamine, 109-89-7; methanesulfonyl chloride, 124-63-0; propargylamine, 2450-71-7; trifluoroacetic anhydride, 407-25-0; 1-methyl-2-propynylamine, 30389-17-4; iodomethane, 74-88-4; acetyl chloride, 75-36-5; propargyl bromide, 106-96-7; diacetamide, 625-77-4; paraformaldehyde, 30525-89-4.

Supplementary Material Available: Four tables with ^1H and ^{13}C NMR spectral data of compounds 1–35 (6 pages). Ordering information is given on any current masthead page.

(28) Dixon, W. J. *J. Am. Stat. Assoc.* 1965, 60, 967.

(29) Cho, A. K.; Jenden, D. J. *Int. J. Neuropharmacol.* 1964, 3, 27.
(30) Pailer, M.; Hübsch, W. J. *Monatsh. Chem.* 1966, 97, 1541.