

Novel Non-Nucleoside Inhibitors of HIV-1 Reverse Transcriptase. 1. Tricyclic Pyridobenzo- and Dipyridodiazepinones


Karl D. Hargrave,*† John R. Proudfoot,† Karl G. Grozinger,† Ernest Cullen,† Suresh R. Kapadia,† Usha R. Patel,† Victor U. Fuchs,† Scott C. Mauldin,† Jana Vitous,† Mark L. Behnke,† Janice M. Klunder,† Kollol Pal,† Jerry W. Skiles,† Daniel W. McNeil,† Janice M. Rose,† Grace C. Chow,† Mark T. Skoog,† Joe C. Wu,† Günther Schmidt,†‡ Wolfhard W. Engel,† Wolfgang G. Eberlein,† Tracy D. Saboe,† Scot J. Campbell,† Alan S. Rosenthal,† and Julian Adams†

Research and Development, Boehringer Ingelheim Pharmaceuticals, Inc., 90 East Ridge, Ridgefield, Connecticut 06877, and Department of Chemical Research, Dr. Karl Thomae GmbH, D-7950 Biberach 1, Federal Republic of Germany.
Received October 1, 1990

Novel pyrido[2,3-*b*][1,4]benzodiazepinones (I), pyrido[2,3-*b*][1,5]benzodiazepinones (II), and dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepinones (III) were found to inhibit human immunodeficiency virus type 1 (HIV-1) reverse transcriptase in vitro at concentrations as low as 35 nM. In all three series, small substituents (e.g., methyl, ethyl, acetyl) are preferred at the lactam nitrogen, whereas slightly larger alkyl moieties (e.g., ethyl, cyclopropyl) are favored at the other (N-11) diazepinone nitrogen. In general, lipophilic substituents are preferred on the A ring, whereas substitution on the C ring generally reduces potency relative to the corresponding compounds with no substituents on the aromatic rings. Maximum potency is achieved with methyl substitution at the position ortho to the lactam nitrogen atom; however, in this case an unsubstituted lactam nitrogen is preferred. Additional substituents on the A ring can be readily tolerated. The dipyridodiazepinone derivative 11-cyclopropyl-5,11-dihydro-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (96, nevirapine) is a potent (IC₅₀ = 84 nM) and selective non-nucleoside inhibitor of HIV-1 reverse transcriptase, and has been chosen for clinical evaluation.

It is widely accepted that the primary etiological agent for the acquired immunodeficiency syndrome (AIDS) is the human immunodeficiency virus type 1 (HIV-1). An essential component of the life cycle of this retrovirus is the transcription of single-stranded viral genomic RNA into double-stranded DNA by the enzyme reverse transcriptase (RT). Inhibition of this viral enzyme should therefore provide an effective means of blocking the spread of HIV-1 infection. In fact, the clinical potential of compounds that interfere with the function of the RT enzyme has already been demonstrated by zidovudine (AZT), the

side effects such as bone marrow suppression, anemia, neutropenia,⁵ mitochondrial myopathy,⁶ peripheral neuropathy, and pancreatitis.⁷ Less severe side effects such



only drug thus far approved for use in the treatment of AIDS and AIDS-related complex (ARC).¹ It has been shown that AZT is phosphorylated by kinases within the cell (a process known as anabolic phosphorylation) to form AZT-5'-triphosphate.² It appears that this substance effectively competes with the natural substrate thymidine triphosphate and acts as a chain terminator in the synthesis of DNA since it lacks the 3'-hydroxyl group necessary for the formation of phosphodiester linkages.^{2b,3} Structurally similar to the natural nucleotide substrates, AZT and other nucleoside analogues that are in clinical trials have been claimed to be selective for RT, but also inhibit other enzymes involved in DNA synthesis, such as mammalian DNA polymerases α , β , γ , and δ .^{2b,4} These dideoxynucleoside analogues often exhibit dose limiting

- (1) (a) Yarchoan, R.; Klecker, R. W.; Weinhold, K. J.; Markham, P. D.; Lyerly, H. K.; Durack, D. T.; Gelmann, E.; Lehrman, S. N.; Blum, R. M.; Barry, D. W.; Shearer, G. M.; Fischl, M. A.; Mitsuya, H.; Gallo, R. C.; Collins, J. M.; Bolognesi, D. P.; Myers, C. E.; Broder, S. *Lancet* 1986, *i*, 575. (b) Yarchoan, R.; Berg, G.; Brouwers, P.; Fischl, M. A.; Spitzer, A. R.; Wichman, A.; Grafman, J.; Thomas, R. V.; Safai, B.; Brunetti, A.; Perno, C. F.; Schmidt, P. J.; Larson, S. M.; Myers, C. E.; Broder, S. *Lancet* 1987, *i*, 132. (c) Fischl, M. A.; Richman, D. D.; Grieco, M. H.; Gottlieb, M. S.; Volberding, P. A.; Laskin, O. L.; Leedom, J. M.; Groopman, J. E.; Mildvan, D.; Schooley, R. T.; Jackson, G. G.; Durack, D. T.; King, D. N. *Engl. J. Med.* 1987, *317*, 185.
- (2) (a) Furman, P. A.; Fyfe, J. A.; St. Clair, M. H.; Weinhold, K. J.; Rideout, J. L.; Freeman, G. A.; Lehrman, S. N.; Bolognesi, D. P.; Broder, S.; Mitsuya, H.; Barry, D. W. *Proc. Natl. Acad. Sci. U.S.A.* 1986, *83*, 8333. (b) Balzarini, J.; Herdewijn, P.; De Clercq, E. *J. Biol. Chem.* 1989, *264*, 6127.
- (3) (a) St. Clair, M. H.; Richards, C. A.; Spector, T.; Weinhold, K. J.; Miller, W. H.; Langlois, A. J.; Furman, P. A. *Antimicrob. Agents Chemother.* 1987, *31*, 1972. (b) Cheng, Y.-C.; Dutschman, G. E.; Bastow, K. F.; Sarngadharan, M. G.; Ting, R. Y. *C. J. Biol. Chem.* 1987, *262*, 2187. (c) Mitsuya, H.; Broder, S. *Proc. Natl. Acad. Sci. U.S.A.* 1986, *83*, 1911. (d) Reardon, J. E.; Miller, W. H. *J. Biol. Chem.* 1990, *265*, 20302.
- (4) (a) Mitsuya, H.; Dahlberg, J. E.; Spigelman, Z.; Matsushita, S.; Jarrett, R. F.; Matsukura, M.; Currens, M. J.; Aaronson, S. A.; Reitz, M. A.; McCaffrey, R. S.; Broder, S. In *Human Retroviruses, Cancer, and AIDS: Approaches to Prevention and Therapy*; Bolognesi, D., Ed.; Alan R. Liss: New York, 1988; pp 407-421. (b) Starnes, M. C.; Cheng, Y.-C. *J. Biol. Chem.* 1987, *262*, 988. (c) Waqar, M. A.; Evans, M. J.; Manly, K. F.; Hughes, R. G.; Huberman, J. A. *J. Cell Physiol.* 1984, *121*, 402. (d) Huang, P.; Farquhar, D.; Plunkett, W. *J. Biol. Chem.* 1990, *265*, 11914.
- (5) (a) Richman, D. D.; Fischl, M. A.; Grieco, M. H.; Gottlieb, M. S.; Volberding, P. A.; Laskin, O. L.; Leedom, J. M.; Groopman, J. E.; Mildvan, D.; Hirsch, M. S.; Jackson, G. G.; Durack, D. T.; Nusinoff-Lehrman, S. N. *Engl. J. Med.* 1987, *317*, 192. (b) Fischl, M. A.; Richman, D. D.; Causey, D. M.; Grieco, M. H.; Bryson, Y.; Mildvan, D.; Laskin, O. L.; Groopman, J. E.; Volberding, P. A.; Schooley, R. T.; Jackson, G. G.; Durack, D. T.; Andrews, J. C.; Nusinoff-Lehrman, S.; Barry, D. W. *J. Am. Med. Assoc.* 1989, *262*, 2405.
- (6) Dalakas, M. C.; Ila, I.; Pezeshkpour, G. H.; Laukaitis, J. P.; Cohen, B.; Griffin, J. L. N. *Engl. J. Med.* 1990, *322*, 1098.


*Boehringer Ingelheim Pharmaceuticals, Inc.

†Dr. Karl Thomae GmbH.


‡Deceased.

as nausea, headaches, and cutaneous eruptions have also been observed.^{5a,7a} Moreover, AZT-resistant strains of HIV-1 have emerged.⁸

Wishing to avoid the toxicological disadvantages inherent in many nucleoside inhibitors, we initiated a search for non-nucleoside inhibitors of HIV-1 RT. Lacking suitable structural leads on which to base a program of rational drug design, we screened a structurally diverse group of compounds on a random basis and found a pyrido[2,3-*b*][1,4]benzodiazepinone (I) to be weakly active. Since this compound is structurally related to the M₁-selective antimuscarinic compound pirenzepine,⁹ a large number of pyrido[2,3-*b*][1,4]benzodiazepinones (I) and pyrido[2,3-*b*][1,5]benzodiazepinones (II) and a few dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepinone (III) analogues were available for testing. Screening of these three series led to the selection of 6,11-dihydro-11-ethyl-6-methyl-5*H*-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (11) as a lead compound (IC₅₀ = 350 nM). The preliminary structure-activity relationship (SAR) derived from all of these compounds thus formed a basis from which to explore the therapeutic potential of the series, the synthesis and SAR of which are herein described. On the basis of this information, we focused our synthetic efforts on the pyrido[2,3-*b*][1,5]benzodiazepinones (II) and dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepinones (III).



The criteria set for the selection of a preclinical candidate was that the compound have an IC₅₀ of less than 100 nM in both the enzyme and cellular (vide infra) assays, that it be nontoxic and specific for RT, that it exhibit suitable solubility and bioavailability, and that it be relatively stable to metabolic degradation. Furthermore, it was deemed necessary that the selected preclinical candidate distribute between the brain and plasma since the virus infects the CNS, resulting in AIDS dementia complex, an often fatal consequence of viral infection.¹⁰ Preliminary measurements of solubility in aqueous solu-

Scheme I^a

tions at various pH's and, more importantly, bioavailability and metabolism of selected compounds in rats provided important information which influenced the selection of synthetic targets. On the basis of our SAR studies and the above criteria, 11-cyclopropyl-5,11-dihydro-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (96, nevirapine) was selected for clinical evaluation.

Chemistry

The pyrido[2,3-*b*][1,5]benzodiazepinones (II)¹¹ were prepared as outlined in Scheme I (method A). Condensation of *o*-phenylenediamines IV with 2-halonicotinic acids V in an inert solvent at $\sim 150^\circ\text{C}$ gave the diazepinones VI in one step and not, as might have been expected, the corresponding 2-(2-chloro-3-pyridinyl)-1*H*-benzimidazoles. Mixtures of isomers (ring A) often resulted when the reaction was carried out with substituted *o*-


- (7) (a) Yarchoan, R.; Perno, C. F.; Thomas, R. V.; Klecker, R. W.; Allain, J.-P.; Wills, R. J.; McAtee, N.; Fischl, M. A.; Dubinsky, R.; McNeely, M. C.; Mitsuya, H.; Pluda, J. M.; Lawley, T. J.; Leuther, M.; Safai, B.; Collins, J. M.; Myers, C. E.; Broder, S. *Lancet* 1988, *i*, 76. (b) Broder, S.; Yarchoan, R. *Am. J. Med.* 1990, 88 (suppl. 5B), 31S. (c) Lambert, J. S.; Seidlin, M.; Reichman, R. C.; Plank, C. S.; Laverty, M.; Morse, G. D.; Knupp, C.; McLaren, C.; Pettinelli, C.; Valentine, F. T.; Dolin, R. *N. Engl. J. Med.* 1990, 322, 1335. (d) Cooley, T. P.; Kunches, L. M.; Saunders, C. A.; Ritter, J. K.; Perkins, C. J.; McLaren, C.; McCaffrey, R. P.; Liebman, H. A. *N. Engl. J. Med.* 1990, 322, 1340.
- (8) Larder, B. A.; Darby, G.; Richman, D. D. *Science* 1989, 243, 1731.
- (9) Carmine, A. A.; Brogden, R. N. *Drugs* 1985, 30, 85.
- (10) (a) Price, R. W.; Brew, B.; Sidtis, J.; Rosenblum, M.; Scheck, A. C.; Cleary, P. *Science* 1988, 239, 586. (b) Navia, B. A.; Jordan, B. D.; Price, R. W. *Ann. Neurol.* 1986, 19, 517.

- (11) (a) Schmidt, G. U.S. Patent 3316251, 1967. (b) Schmidt, G. U.S. Patent 3326900, 1967. (c) Hargrave, K. D.; Schmidt, G.; Engel, W.; Trummelitz, G.; Eberlein, W. Eur. Patent Appl. E.P. 393604, 1990.

phenylenediamines. In many cases, the major isomer could be isolated by fractional recrystallization, either at this stage or following alkylation at N-6 and/or N-11. Occasionally, however, separation of the isomers was not practically achieved at any stage. Monoalkylation of the tricyclic ring system occurred at the lactam nitrogen (N-6) with 1 equiv each of a suitable base and alkylating agent, and subsequent alkylation provided the desired N,N'-disubstituted diazepinones (II, $R^1 \neq R^2$). Monoalkylation at N-11 required protection (e.g., benzyl, benzoyl) of the lactam nitrogen, though it could be achieved in low yields by the use of two equivalents of a suitable base and 1 equiv of alkylating agent. In those cases wherein $R^1 = R^2$, dialkylation was achieved in one step using 2 equiv each of base and alkylating agent. The choice of base used in the alkylation of N-6 was often critical in order to minimize the formation of side products such as X and XI. In some cases, 30% aqueous sodium hydroxide in DMSO or potassium *tert*-butoxide in 1,4-dioxane proved more convenient than sodium hydride in DMF, which was generally preferred for alkylation at N-11. Occasionally, the formation of some N-1 alkylated product (XIa or XIb) was observed following the alkylation of VI or VII, respectively. Conversion of II to the corresponding thiolactams (IX) was effected by treatment with Lawesson's reagent¹² in toluene at reflux.

In addition to the synthetic sequence depicted in Scheme I, further functional group manipulation was required for the preparation of several of the compounds listed in Table I. Thus, carboxylic acid 47 was obtained by hydrolysis of the corresponding ester 46, and carboxamides 48 and 49 were prepared by the mixed anhydride method from 47 and the appropriate amines. Treatment of the 8-methoxy derivative 36 with boron tribromide in methylene chloride gave the corresponding phenol 37. Reduction of the nitro compound 42 with stannous chloride in concentrated HCl afforded the amine 43, which was treated with acetic anhydride to provide the acetamide 44 or diazotized and reacted with sodium azide to give the azido compound 45. Treatment of 11 with chlorine in trimethyl phosphate¹³ unexpectedly provided the 3-chloro derivative 38 (see Experimental Section). Although we have no explanation for the regioselectivity of this reaction, it should be noted that electrophilic aromatic substitution would be expected to occur preferentially on the phenyl (A) ring. Indeed, Tscherniac-Einhorn reaction¹⁴ of 11 provided the expected regioisomeric phthalamidomethyl intermediate, which was deprotected to give the desired 8-aminomethyl derivative 50. The hydroxymethyl analogue 52 was subsequently prepared by diazotization of 50 followed by treatment with water.


The dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepinones (III) were synthesized as shown in Scheme II.¹⁵ In method B, condensation of 3-amino-2-chloropyridines (XII; X = Cl) with 2-chloronicotinoyl chlorides XIII provided the 2,2'-dihalo amides XIV. These were converted to the 2'-alkylamino compounds XV by treatment with the appropriate amine, usually at >100 °C in a sealed tube. In general, activation by the adjacent carbonyl group was sufficient to ensure

Scheme II^a

^a (a) Inert solvent; (b) NaH, DMSO, R¹X; (c) R²NH₂, 1,4-dioxane or xylene or neat; (d) NaH, pyridine or bis(2-methoxyethyl)ether or xylene; (e) Lawesson's reagent, toluene; (f) trifluoroacetic acid.

that only the 2'-chlorine atom was displaced, and excellent yields of the desired amines XV were obtained. If, however, electron-withdrawing groups (e.g., Cl or CF₃) were present on the other pyridine ring, then competing substitution of the 2-chloro substituent was observed and the yield of the desired XV was decreased. Ring closure to the dipyridodiazepinones IIIa was effected by heating the dianion of XVa, generated with 2 equiv of sodium hydride, under reflux in either pyridine or bis(2-methoxyethyl) ether. Dimethylformamide could also be used as a solvent for this reaction, but led to lower yields of less pure product. Alkylation of the lactam nitrogen, if desired, was achieved by procedures described above. The dipyrido-diazepinones IIIb could also be obtained by a variation of this sequence involving alkylation of the amides XIVa to give XIVb, which were often observed by NMR as a mixture of conformers (see Experimental Section). Subsequent conversion of XIVb to XVb with the appropriate alkylamine as described above, and cyclization of the monoanion of XVb in xylene at reflux provided IIIb. In some cases, the desired diazepinones were synthesized by reaction of the anion of XIXb with the appropriate elec-


- (12) (a) Pedersen, B. S.; Scheibye, S.; Nilsson, N. H.; Lawesson, S.-O. *Bull. Soc. Chim. Belg.* 1978, 87, 223. (b) Pedersen, B. S.; Lawesson, S.-O. *Tetrahedron* 1979, 35, 2433.
 (13) Pearson, D. E.; Frazer, M. G.; Frazer, V. S.; Washburn, L. C. *Synthesis* 1976, 621.
 (14) For a review, see: Zaugg, H. E.; Martin, W. B. *Org. Reactions* 1965, 14, 52.
 (15) Hargrave, K. D.; Schmidt, G.; Engel, W.; Trummlitz, G.; Eberlein, W. Eur. Patent Appl. E.P. 393529, 1990.

Table I. 6,11-Dihydro-5H-pyrido[2,3-b][1,5]benzodiazepin-5-ones^a

no.	R ¹	R ²	R	recryst solvent	mp, °C	formula	% inh @ 10 µg mL ^b	IC ₅₀ ^b µM
1	H	H	H	5% DMF/dioxane	283-287	C ₁₂ H ₉ N ₃ O	22	>>1
2	H	CH ₃	H	1-propanol	235-237	C ₁₃ H ₁₁ N ₃ O	75	>1
3	H	CH ₂ CH ₃	H	EtOAc	210-220	C ₁₄ H ₁₃ N ₃ O	99	0.38
4	CH ₃	CH ₂ C ₆ H ₅	H	CH ₂ Cl ₂ /pet. ether	168-170	C ₂₀ H ₁₇ N ₃ O	93	9
5	CH ₃	H	H	EtOH	171	C ₁₃ H ₁₁ N ₃ O	43	>>1
6	CH ₂ CH ₃	H	H	MeOH	149-151	C ₁₄ H ₁₃ N ₃ O	87	>1
7	CH ₂ COO- <i>t</i> -Bu	H	H	EtOAc/hexane	200-202	C ₁₈ H ₁₅ N ₃ O ₃	39	>>1
8	CH ₂ CH=CH ₂	H	H	EtOH	105-107	C ₁₅ H ₁₃ N ₃ O	96	1.4
9	CH ₂ CH ₂ OH	H	H	EtOH	190-192 dec	C ₁₄ H ₁₃ N ₃ O ₂ ·CHI	46	>>1
10	CH ₂ CH ₂ OCH ₃	H	H	70% aq EtOH	125-126	C ₁₅ H ₁₅ N ₃ O ₂	36	>>1
11	CH ₃	CH ₂ CH ₃	H	cyclohexane	106-111	C ₁₆ H ₁₆ N ₃ O	100	0.35
12	CH ₃	CH ₃	H	cyclohexane	126-128	C ₁₄ H ₁₃ N ₃ O	85	>1
13	CH ₂ CH ₃	CH ₃	H	cyclohexane	118-119	C ₁₆ H ₁₆ N ₃ O	84	>1
14	CH ₂ CH ₃	CH ₂ CH ₂	H	heptane	108-110	C ₁₆ H ₁₇ N ₃ O	99	0.38
15	CH ₃	CH ₂ CH ₂ CH ₃	H	pet. ether	96-98	C ₁₆ H ₁₇ N ₃ O	99	0.23
16	CH ₃	CH(CH ₃) ₂	H	cyclohexane	144-147	C ₁₆ H ₁₇ N ₃ O	98	0.20
17	CH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	pet. ether	56-58	C ₁₇ H ₁₉ N ₃ O	81	1
18	CH ₃	CH ₂ SCH ₃	H	ether/pet. ether	127-129	C ₁₅ H ₁₅ N ₃ OS	84	6.6
19	CH ₃	CH ₂ COO- <i>t</i> -Bu	H	CH ₂ Cl ₂ /EtOAc	64-65	C ₁₉ H ₂₁ N ₃ O ₃	87	>1
20	CH ₃	CH ₂ COOC ₂ H ₅	H	EtOAc/hexane	99-100	C ₁₇ H ₁₇ N ₃ O ₃	56	16
21	CH ₃	CH ₂ CH ₂ OH	H	xylene	133-134	C ₁₆ H ₁₆ N ₃ O ₂	58	>1
22	CH ₃	COCH ₃	H	cyclohexane	140-142	C ₁₅ H ₁₃ N ₃ O ₂	72	>1
23	CH ₃	CH ₂ CH ₂ F	H	EtOAc/pet. ether	118-120	C ₁₆ H ₁₄ FN ₃ O	97	1
24	CH ₂ F	COCH ₃	H	CH ₂ Cl ₂ /pet. ether	113-116	C ₁₅ H ₁₄ FN ₃ O	85	>1
25	CH ₃	COOC ₂ H ₅	H	CH ₂ Cl ₂ /ether	123-125	C ₁₆ H ₁₆ N ₃ O ₃	13	>>1
26	CH ₂ CON(CH ₃) ₂	H	H	CH ₃ CN	199-201	C ₁₆ H ₁₆ N ₄ O ₂	24	>>1
27	CH ₃	CH ₂ CO ₂ H	H	c	189-192	C ₁₆ H ₁₃ N ₃ O ₃	60	>1
28	CH ₃	C≡CH	H	c	142-143	C ₁₅ H ₁₁ N ₃ O	97	0.65
29	CH ₂ CH ₃	CH ₂ CH ₃	7-CH ₃	pet. ether	120-122	C ₁₇ H ₁₉ N ₃ O	90	8
30	H	H	7-CH ₃	1,4-dioxane/H ₂ O	233-234	C ₁₅ H ₁₁ N ₃ O·0.1H ₂ O	77	>1
31	CH ₃	CH ₂ CH ₃	7-CH ₃	CH ₃ CN	132-134	C ₁₆ H ₁₇ N ₃ O	100	0.24
32	CH ₂ C ₆ H ₅	CH ₂ CH ₃	7-CH ₃	EtOH	145-147	C ₂₂ H ₂₁ N ₃ O	40	>>1
33	H	CH ₂ CH ₃	7-CH ₃	EtOAc	225-227	C ₁₆ H ₁₆ N ₃ O	100	0.038
34	CH ₃	CH ₂ CH ₃	8-CH ₃	c	oil	C ₁₆ H ₁₇ N ₃ O·0.2H ₂ O	100	0.19
35	CH ₃	CH ₂ CH ₂ N(CH ₃) ₂	8,9-(CH ₃) ₂	EtOAc	185-188	C ₁₉ H ₂₄ N ₄ O	0	>>1
36	CH ₃	CH ₂ CH ₃	8-OCH ₃	c	oil	C ₁₆ H ₁₇ N ₃ O ₂	95	2.1
37	CH ₃	CH ₂ CH ₃	8-OH	c	207-209	C ₁₆ H ₁₆ N ₃ O ₂	47	>>1
38	CH ₃	CH ₂ CH ₃	3-Cl	pet. ether	131-132	C ₁₅ H ₁₄ ClN ₃ O	91	7.6
39	CH ₃	CH ₂ CH ₃	9-CF ₃	c	104-106	C ₁₆ H ₁₄ F ₃ N ₃ O	97	0.23
40	CH ₃	CH ₂ CH ₃	2-F	EtOAc/hexane	100-101	C ₁₅ H ₁₄ FN ₃ O	84	2.9
41	CH ₃	CH ₂ CH ₃	8-NO ₂	EtOAc/hexane	173-176	C ₁₆ H ₁₄ N ₄ O ₃	56	>1
42	CH ₃	CH ₂ CH ₃	9-NO ₂	EtOAc/hexane	179.5-180.5	C ₁₆ H ₁₄ N ₄ O ₃	99	0.4
43	CH ₃	CH ₂ CH ₃	9-NH ₂	EtOAc	193-195	C ₁₆ H ₁₆ N ₄ O	89	4.5
44	CH ₃	CH ₂ CH ₃	9-NHAc	toluene/EtOAc/CH ₂ Cl ₂	105-107	C ₁₇ H ₁₈ N ₄ O ₂	49	>>1
45	CH ₃	CH ₂ CH ₃	9-N ₃	ether	115-118	C ₁₆ H ₁₄ N ₆ O	100	0.14
46	CH ₃	CH ₂ CH ₃	8-COOCH ₃	c	132-136	C ₁₇ H ₁₇ N ₃ O ₃	94	1.1
47	CH ₃	CH ₂ CH ₃	8-COOH	EtOH	290-291	C ₁₆ H ₁₆ N ₃ O ₃	18	>>1
48	CH ₃	CH ₂ CH ₃	8-CONH ₂	c	104-105	C ₁₆ H ₁₆ N ₄ O ₂ ·H ₂ O ^d	66	27
49	CH ₃	CH ₂ CH ₃	8-CONEt ₂	c	oil	C ₁₆ H ₂₄ N ₄ O ₂ ·H ₂ O ^e	11	>>1
50	CH ₃	CH ₂ CH ₃	8-CH ₂ NH ₂	c	oil	C ₁₆ H ₁₈ N ₄ O·0.5H ₂ O	84	>1
51	CH ₃	CH ₂ CH ₃	8-CH ₂ NHCO- <i>t</i> -Bu	ether/hexane	118-120	C ₂₁ H ₂₆ N ₄ O ₂	15	>>1
52	CH ₃	CH ₂ CH ₃	8-CH ₂ OH	c	oil	C ₁₆ H ₁₇ N ₃ O ₂	78	>1
53	CH ₃	CH ₂ CH ₃	8,9-(CH ₃) ₂	pet. ether	140-144	C ₁₇ H ₁₉ N ₃ O	100	0.04
54	CH ₃	COCH ₃	8,9-(CH ₃) ₂	cyclohexane	168-170	C ₁₇ H ₁₇ N ₃ O ₂	96	0.42
55	CH ₃	CH ₂ CH ₃	7,8-(CH ₃) ₂	c	140-144	C ₁₇ H ₁₉ N ₃ O	99	0.036
56	CH ₃	CH ₂ CH ₃	8,9-Cl ₂	c	146-147	C ₁₆ H ₁₃ Cl ₂ N ₃ O	100	0.062
57	CH ₃	CH ₂ CH ₃	7-Cl, 9-CF ₃	c	109-111	C ₁₆ H ₁₃ ClF ₃ N ₃ O	96	0.29
58	CH ₃	CH ₂ CH ₃	7-Br, 9-CF ₃	c	oil	C ₁₆ H ₁₃ BrF ₃ N ₃ O·0.2H ₂ O	71	>1
59	CH ₃	CH ₃	2,4,8-(CH ₃) ₃	pet. ether (100-140 °C)	174-176	C ₁₇ H ₁₉ N ₃ O	49	>>1
60	CH ₂ CH=CH ₂	COCH ₃	H	cyclohexane	152-154	C ₁₇ H ₁₅ N ₃ O ₂	83	>1
61	CH ₂ CH=CH ₂	COCH ₂ CH ₃	H	cyclohexane	118-120	C ₁₈ H ₁₇ N ₃ O ₂	72	>1
62	CH ₃	CH ₂ CH ₃	3-CF ₂ CF ₃	c	oil	C ₁₇ H ₁₄ F ₅ N ₃ O	5	>>1
63	CH ₃	CH ₂ CH ₃	3-Br, 9-NO ₂	c	156-158	C ₁₆ H ₁₃ BrN ₄ O ₃	95	0.89
64	CH ₃	CH ₂ CH ₃	3-Br, 8-NO ₂	c	171-173	C ₁₆ H ₁₃ BrN ₄ O ₃	64	>1
65	CH ₃	CH ₂ CH ₃	8-Cl	cyclohexane/pet. ether	158-159	C ₁₆ H ₁₄ ClN ₃ O	99	0.17
66	CH ₂ CH=CH ₂	H	8-Cl	EtOH	167-169	C ₁₆ H ₁₂ ClN ₃ O	79	>1
67	CH ₃	CH ₂ CH ₃	9-COC ₂ H ₅	c	67-72	C ₂₂ H ₁₉ N ₃ O ₂	73	>1
68	CH ₃	CH ₂ CH ₃	9-Cl	pet. ether	169-171	C ₁₆ H ₁₄ ClN ₃ O	100	0.17

^aAll compounds were prepared by method A. ^bInhibition of HIV-1 RT. See Experimental Section. ^cSample purified by column chromatography and not requiring recrystallization. ^dN: calcd, 17.82; found, 17.32. ^eN: calcd, 15.12; found, 14.38. ^fH: calcd, 4.90; found, 5.41.

Table II. 5,11-Dihydro-6H-dipyrido[3,2-b:2',3'-e][1,4]diazepin-6-one



no.	R ¹	R ²	R	method	recryst solvent	mp, °C	formula	% inh @ 10 µg/mL ^a	IC ₅₀ ^a µM
69	CH ₃	C ₂ H ₅	H	B	EtOAc/hexane	130-132	C ₁₄ H ₁₄ N ₄ O	100	0.125
70	H	C ₂ H ₅	H	B	EtOAc/EtOH	211-212	C ₁₃ H ₁₂ N ₄ O	99	0.44
71	H	c-C ₃ H ₅	H	B	EtOAc	240-250	C ₁₄ H ₁₂ N ₄ O	99	0.45
72	H	c-C ₃ H ₇	H	B	EtOAc	241-243	C ₁₅ H ₁₄ N ₄ O	99	0.43
73	H	c-C ₃ H ₉	H	B	EtOAc/ether	225-228	C ₁₆ H ₁₆ N ₄ O	94	>1
74	CH ₃	n-C ₃ H ₇	H	B	b	oil	C ₁₅ H ₁₆ N ₄ O	100	0.45
75	H	i-C ₃ H ₇	H	B	EtOAc/hexane	204-206	C ₁₄ H ₁₄ N ₄ O	99	0.59
76	CH ₃	t-C ₃ H ₉	H	B	hexane	192-194	C ₁₆ H ₁₈ N ₄ O	87	11
77	CH ₃	c-C ₃ H ₁₁	H	B	ether/pet. ether	145-146	C ₁₈ H ₂₀ N ₄ O ^{1/4} H ₂ O	73	>1
78	CH ₃	CH ₂ CH=CH ₂	H	C	EtOAc/hexane	93-95	C ₁₅ H ₁₄ N ₄ O	99	0.85
79	CH ₃	CH ₂ C=CH	H	B	EtOAc	169-170	C ₁₅ H ₁₂ N ₄ O ^c	94	>1
80	H	(R)-C ₄ H ₉	H	B	EtOAc/hexane	172-174	C ₁₅ H ₁₆ N ₄ O	85	>1
81	H	(S)-C ₄ H ₉	H	B	EtOAc/hexane	173-175	C ₁₅ H ₁₆ N ₄ O	88	>1
82	H	C ₆ H ₅	H	B	EtOAc/hexane	220-222	C ₁₇ H ₁₂ N ₄ O	71	>1
83	COCH ₃	C ₂ H ₅	H	B	EtOAc/hexane	123-124.5	C ₁₅ H ₁₄ N ₄ O ₂	99	0.36
84	CH ₃	COCH ₃	H	C	EtOAc/hexane	138-143	C ₁₄ H ₁₂ N ₄ O ₂	73	15.3
85	CH ₃	CH ₂ CH ₂ F	H	C	pet. ether/ether	117-118	C ₁₄ H ₁₃ FN ₄ O	96	2.9
86	CH ₃	SO ₂ CH ₃	H	C	EtOAc/hexane	239-241	C ₁₃ H ₁₂ N ₄ O ₃ S	77	>1
87	CH ₃	CH ₂ SCH ₃	H	C	ether/pet. ether	109-110	C ₁₄ H ₁₄ N ₄ OS	99	0.85
88	CH ₃	C ₂ H ₅	2-CH ₃	B	hexane	124-126	C ₁₅ H ₁₆ N ₄ O	100	0.17
89	CH ₃	C ₂ H ₅	2-Cl	B	heptane	125-126	C ₁₄ H ₁₃ ClN ₄ O	100	0.15
90	CH ₃	C ₂ H ₅	3-CH ₃	B	hexane	94-96	C ₁₅ N ₁₆ N ₄ O	100	0.76
91	CH ₃	C ₂ H ₅	3-Cl	B	heptane	124-125	C ₁₄ H ₁₃ ClN ₄ O	99	>1
92	CH ₃	C ₂ H ₅	3-NO ₂	B	EtOAc/hexane	154-156	C ₁₄ H ₁₃ N ₆ O ₃	64	>1
93	CH ₃	C ₂ H ₅	3-NH ₂	B	EtOAc/MeOH	235-240	C ₁₄ H ₁₆ N ₅ O	92	>1
94	H	C ₂ H ₅	4-CH ₃	B	CH ₂ ClCH ₂ Cl	212-214	C ₁₄ H ₁₄ N ₄ O	100	0.035
95	CH ₃	C ₂ H ₅	4-CH ₃	B	EtOAc/hexane	157-159	C ₁₅ H ₁₆ N ₄ O	98	1.9
96	H	c-C ₃ H ₅	4-CH ₃	B	EtOAc	247-249	C ₁₅ H ₁₄ N ₄ O	100	0.084
97	CH ₃	c-C ₃ H ₅	4-CH ₃	B	EtOAc/ether	244-245	C ₁₆ H ₁₆ N ₄ O	96	>1
98	H	C ₂ H ₅	4-Cl	B	CH ₃ CN	184-186	C ₁₃ H ₁₁ ClN ₄ O	100	0.095
99	H	C ₂ H ₅	4-C ₂ H ₅	B	EtOAc	218-219	C ₁₅ H ₁₆ N ₄ O	100	0.11
100	H	C ₂ H ₅	4-OCH ₃	B	EtOAc/hexane	156-157	C ₁₄ H ₁₄ N ₄ O ₂	96	>1
101	H	C ₂ H ₅	4-OH	B	HOAc	295-296	C ₁₃ H ₁₂ N ₄ O ₂ ·2HBr·H ₂ O	94	>1
102	H	c-C ₃ H ₅	4-CH ₂ OH	B ^d	EtOAc/hexane	243-245	C ₁₅ H ₁₄ N ₄ O ₂	94	3.0
103	H	c-C ₃ H ₅	4-CN	B ^d	EtOAc/hexane	243-245	C ₁₅ H ₁₁ N ₅ O ^e	96	1.25
104	H	C ₂ H ₅	2,3-(CH ₃) ₂	B	EtOAc/hexane	212-214	C ₁₅ H ₁₆ N ₄ O	100	0.41
105	CH ₃	C ₂ H ₅	2,3-(CH ₃) ₂	B	hexane	143-145	C ₁₆ H ₁₈ N ₄ O	100	0.24
106	CH ₃	C ₂ H ₅	2-NO ₂ , 3-Cl	B	EtOAc/hexane	153-157	C ₁₄ H ₁₂ ClN ₅ O ₃	98	>1
107	CH ₃	C ₂ H ₅	2-NH ₂ , 3-Cl	B	EtOAc/hexane	160-162	C ₁₄ H ₁₄ ClN ₅ O	98	0.85
108	H	C ₂ H ₅	7-CH ₃	B	EtOAc/hexane	193-194	C ₁₄ H ₁₄ N ₄ O·H ₂ O ^f	80	>1
109	H	C ₂ H ₅	8-CH ₃	B	EtOAc/hexane	182-183	C ₁₄ H ₁₄ N ₄ O·0.1H ₂ O	89	>1
110	CH ₃	C ₂ H ₅	8-Cl	B	hexane	105-106	C ₁₄ H ₁₃ ClN ₄ O	87	3.5
111	H	c-C ₃ H ₇	4-CH ₃ , 7-OH	B ^d	EtOAc/hexane	225-227	C ₁₅ H ₁₄ N ₄ O ₂ ^g /4H ₂ O ^h	96	>1
112	CH ₃	C ₂ H ₅	9-CH ₃	B	hexane	79-93	C ₁₅ H ₁₆ N ₄ O	65	>1
113	CH ₂ CH ₂ N(CH ₃) ₂	c-C ₃ H ₇	H	B	CH ₂ Cl ₂ /MeOH	88-89	C ₁₈ H ₂₁ N ₅ O	34	>>1

^a Inhibition of HIV-1 RT. See Experimental Section. ^b Sample purified by column chromatography and not requiring recrystallization. ^c N: calcd, 21.20; found, 20.61. ^d Derived from compound 90. ^e No elemental analysis available; characterized by NMR and MS. ^f C: calcd, 67.15; found, 66.54. ^g N: calcd, 21.65; found, 20.74. ^h N: calcd, 18.94; found, 18.31.

trophile (method C). Sulfurization to provide the thiolactam (XVI) from III was achieved with Lawesson's reagent,¹² as described earlier.


The chemistry of the dihydropyridodiazepinones III is currently being explored further, and we have found, for example, that nitration of the pyridine rings can be achieved using nitronium tetrafluoroborate in acetonitrile. Thus, 69 gives the 3-nitro and 3,8-dinitro derivatives upon treatment with 1 equiv of the nitrating agent. Interestingly, the 3-chloro analogue of 69 (91) gives the 2-nitro and 2,8-dinitro derivatives under the same conditions, with the initial nitration occurring on ring A.

Results and Discussion

The in vitro HIV-1 RT enzyme assay was the primary screen used to evaluate these compounds for their potential

as anti-AIDS drugs; the test results are presented in Tables I-III. Selected compounds were further tested for their ability to block the proliferation of HIV-1 infection in vitro. This was demonstrated by their inhibition of the formation of syncytia and p24 antigen in cultures of CD4⁺ human T-cells (c8166) infected with HIV-1. A direct correlation between the enzymatic and cellular potencies of representatives of II and III is observed. These and other data support the hypothesis that the compounds reported here inhibit the spread of HIV-1 infection in vitro as a result of their ability to inhibit RT.¹⁶ Utilizing cells from the same c8166 cell line, these compounds were also tested in an MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-

(16) Hargrave, K. D.; Sullivan, J. L.; Eckner, R. J.; Merluzzi, V. J.; Koup, R. A. Unpublished results.

Table III. 5,11-Dihydro-6*H*-pyrido[2,3-*b*][1,4]benzodiazepin-6-ones and 5,10-Dihydro-11*H*-dibenzo[*b,e*][1,4]diazepin-11-ones

no.	R ¹	R ²	R	X	recryst solvent	mp, °C	formula	% inh @ 10 µg/mL ^a	IC ₅₀ , ^a µM
114	H	H	H	N	DMF	282.6–282.9	C ₁₂ H ₉ N ₃ O	23	>>1
115	CH ₃	CH ₃	H	N	EtOH	113.5–114.5	C ₁₄ H ₁₃ N ₃ O	74	>1
116	CH ₃	CH ₂ CH ₃	H	N	EtOAc	127–128	C ₁₅ H ₁₆ N ₃ O	99	0.50
117	CH ₂ CH ₃	CH ₂ CH ₃	H	N	cyclohexane	95–97	C ₁₆ H ₁₇ N ₃ O	96	0.69
118	CH ₃	CH ₂ CH ₃	8-CH ₃	N	EtOH	137–138	C ₁₆ H ₁₇ N ₃ O	99	0.58
119	CH ₂ CH ₃	CH ₂ CH ₃	10-CH ₃	N	pet. ether (100–140 °C)	143–144	C ₁₇ H ₁₉ N ₃ O	99	0.65
120	CH ₂ CH ₃	CH ₃	2-Cl	N	cyclohexane	162–164	C ₁₅ H ₁₄ ClN ₃ O	76	3.3
121	H	H	2-CH ₃	N	80% aq. HOAc	260–262	C ₁₃ H ₁₁ N ₃ O	46	>1
122	H	H	2,4-(CH ₃) ₂	N	dimethylacetamide	281–283	C ₁₄ H ₁₃ N ₃ O	51	>1
123	CH ₃	CH ₃	H	CH	EtOH	137.5–139	C ₁₅ H ₁₄ N ₂ O	52	>1
124	CH ₃	CH ₂ CH ₃	H	CH	heptane	189.5–190.5	C ₁₆ H ₁₆ N ₂ O	80	>1

^a Inhibition of HIV-1 RT. See Experimental Section.

tetrazolium bromide] cytotoxicity assay¹⁷ and were found to be nontoxic at the doses required for activity (therapeutic index >1000). Specificity was established by the inactivity of selected compounds at relatively high doses in an HIV-2 cell culture assay, and by their lack of inhibition of HIV-2 RT and various non-HIV RT enzymes, including those of the simian immunodeficiency virus, feline leukemia virus, Moloney murine leukemia virus, and avian myeloblastosis virus. In addition, nevirapine does not inhibit other mammalian DNA polymerases (*vide infra*).

Evaluation of representative examples from the two monopyridodiazepinone series (I and II) and the dipyrindodiazepinone series (III) suggests that, in general, the dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepinones (III) inhibit RT with equivalent or slightly enhanced potency relative to the corresponding pyrido[2,3-*b*][1,5]benzodiazepinones (II), which are, in turn, slightly more potent than the corresponding pyrido[2,3-*b*][1,4]benzodiazepinones (I). Moreover, the corresponding dibenzo[*b,e*][1,4]diazepinones are less active relative to the mono- or dipyrindodiazepinones. The pyrido[2,3-*b*][1,4]benzodiazepinones¹⁸ (I) and dibenzo[*b,e*][1,4]diazepinones¹⁹ were synthesized as previously described. The test results (Table III) suggest that compounds with a methyl group in the 2-, 4-, 8-, or 10-positions or a chloro group in the 2-position generally exhibit potencies similar to or greater than those of the corresponding unsubstituted analogues. In addition to the finding that at least one of the aromatic rings in this series must be a pyridine ring for potent activity, the position of the pyridine nitrogen is also critical. For example, isomers of III in which the A-ring nitrogen is at position 3 are very weakly active. As a result, our synthetic efforts focused primarily on the pyrido[2,3-*b*][1,5]benzodiazepinones (II) and dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepinones (III).

6,11-Dihydro-5*H*-pyrido[2,3-*b*][1,5]benzodiazepin-5-ones (Table I). In the following evaluation of the pyrido[2,3-*b*][1,5]benzodiazepinone (II) series, compound 11, which is unsubstituted on the aromatic rings, is used as the reference standard. Derivatives with substituents present on the C ring all exhibit less activity than the corresponding unsubstituted analogues. For example, the 2-fluoro (40) and 3-chloro (38) analogues are less potent than the corresponding unsubstituted compound (11). A methyl group at position 7 enhances potency if the lactam nitrogen (N-6) is unsubstituted (compound 33). In the 8-position, potency of 6-methylated compounds is enhanced by methyl (34) and chloro (65) substitution. In contrast, 8-substituted compounds with electronegative groups such as nitro, methoxycarbonyl, carboxy, aminocarbonyl, and (*N,N*-diethylamino)carbonyl are significantly less potent (compare compounds 41, 46, 47, 48, and 49 to 11). Also, nonlipophilic substituents in the 8-position, such as methoxy, hydroxy, hydroxymethyl, and aminomethyl, all lower activity (compare compounds 36, 37, 52, and 50 to 11). At the 9-position, chloro, trifluoromethyl, and azido²⁰ groups enhance potency, whereas amino and acetamido groups reduce activity, and nitro substitution is relatively neutral (compare compounds 68, 39, 45, 43, 44, 42, and 11). The most potent compounds, with IC₅₀'s in the 35–65 nM range, are derivatives disubstituted with the lipophilic methyl (53 and 55) or chloro (56) groups at positions 7 and 8 or 8 and 9, or the analogue monomethylated at position 7 (33).

5,11-Dihydro-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-ones (Table II). Synthetic efforts in the dipyrindodiazepinone (III) series focused on substitutions at the six available positions of the aromatic rings. Substituents at the 2-position generally appear to have no dramatic effect on activity, with methyl, chloro, nitro, and amino derivatives exhibiting potencies similar to those of the corresponding 2-unsubstituted analogues. On the other hand, the same groups placed at the 3-position have a detrimental effect on potency relative to the corresponding unsubstituted compound 69. In the 4-position, methyl substitution produces the best activity, provided the lactam nitrogen (N-5) is unsubstituted. If the lactam nitrogen is substituted, then a 4-methyl group is detrimental. This

- (17) (a) Mosmann, T. J. *Immunol. Methods* 1983, 65, 55. (b) Pauwels, R.; Balzarini, J.; Baba, M.; Snoeck, R.; Schols, D.; Herdewijn, P.; Desmyter, J.; De Clercq, E. *J. Virol. Methods* 1988, 20, 309.
- (18) (a) Schmidt, G. U.S. Patent 3 406 168, 1968. (b) Schmidt, G.; Engelhardt, G.; Machleidt, H.; Koenigsdorfer, K. So. African Patent 68 05 942, 1969. (c) Engel, W. W.; Eberlein, W. G.; Mihm, G.; Hammer, R.; Trummelitz, G. *J. Med. Chem.* 1989, 32, 1718. (d) Hargrave, K. D.; Schmidt, G.; Engel, W.; Austel, V. *Eur. Patent Appl. E.P.* 393530, 1990.
- (19) Clemo, G. R.; Perkin, W. H., Jr.; Robinson, R. *J. Chem. Soc.* 1924, 125, 1751.

- (20) Wu, J. C.; Warren, T. C.; Adams, J.; Proudfoot, J.; Skiles, J.; Raghavan, P.; Perry, C.; Potocki, I.; Farina, P. R.; Grob, P. M. *Biochemistry* 1991, 30, 2022.

combination of a 4-methyl group and an unsubstituted lactam is present in all of the most active compounds, although 4-chloro and -ethyl compounds are also highly active. In contrast, potency is reduced by the presence in the 4-position of electron-donating, electron-withdrawing, and/or hydrophilic groups such as methoxy, hydroxy, cyano, or hydroxymethyl. Methyl substitution at positions 7, 8, or 9 has a detrimental effect on potency; and similar results were obtained with 7-hydroxy or with 8-chloro derivatives.

Summarizing the SAR of the dipyrindodiazepinones based on available data, maximum enzyme inhibition is obtained with compounds containing a 4-methyl substituent in the absence of substitution at the lactam nitrogen (N-5); however, 5-methyl substitution is preferred if the 4-position is not substituted. Substitution at positions 2, 3, 7, 8, and 9 appears to have little effect or to reduce activity of the analogues evaluated thus far. At N-11, small alkyl groups such as ethyl and cyclopropyl are preferred.

Although there are some differences between series in the SAR involving substitution of the aromatic rings, a similar SAR is operative for the two diazepinone nitrogen substituents. Specifically, small substituents such as methyl, ethyl, or acetyl at the lactam nitrogen atom are generally preferred. Larger substituents such as allyl, methoxyethyl, or (*N,N*-dimethylamino)ethyl, which is present on some tricyclic antidepressants, reduce activity. In contrast, an unsubstituted lactam nitrogen ($R^1 = H$) is much preferred when the position ortho to the lactam nitrogen in II or III is occupied by a methyl or chloro substituent. In general, thiolactams are more potent than the corresponding lactams. However, due to poor solubility and rapid metabolism, the thiolactams were not considered for further development. At the N-11 position, relatively small lipophilic groups are required for optimum activity ($R^2 = Et, c\text{-}Pr$); analogues with smaller ($R^2 = H, Me$) or larger ($R^2 = Bu, Ph$) groups are less potent. Furthermore, basic aminoalkyl substituents at N-11, which are necessary for antimuscarinic activity,²¹ result in very weak or inactive compounds in the RT assay. Also, hydrophilic or electron-withdrawing substituents reduce potency relative to that of the *N*-ethyl derivative.

Studies with representatives of the three series suggest that, as a class, the dipyrindodiazepinones (III) are not only more potent, but are also more water-soluble, less cytotoxic, and more resistant to metabolic *N*-dealkylation than either of the monopyrindodiazepinone series (I or II). The most active compounds of series I and II are poorly soluble in aqueous solution. When the more soluble unsubstituted compound 11 was evaluated in rats following oral administration, it was found to be dealkylated quite rapidly at both diazepinone nitrogen atoms. Metabolism studies with 69, the dipyrido analogue of 11, suggested that *N*-dealkylation would also be a potential problem in series III, although the degradation was not as rapid. However, the finding that the combination of a 4-methyl substituent and an unsubstituted lactam results in maximal activity eliminated the dealkylation problem at N-5. In fact, metabolism studies in the rat and monkey with 94 and 96 indicated that the presence of a 4-methyl substituent also significantly reduces dealkylation at N-11.²²

Nevirapine has been found to be a potent ($IC_{50} = 84 \text{ nM}$) and specific noncompetitive ($K_i = 200 \text{ nM}$) inhibitor of HIV-1 RT which acts at an allosteric site of the enzyme.^{20,23}

It blocks HIV-1 replication in vitro ($IC_{50} = 40 \text{ nM}$) in CD4⁺ human T-cells (c8166), but is noncytotoxic except at significantly higher doses ($CC_{50} = 321\,000 \text{ nM}$).^{23a} In comparison, AZT is a more potent inhibitor of HIV-1 replication in vitro ($IC_{50} = 6 \text{ nM}$), but is more cytotoxic ($CC_{50} = 66\,000 \text{ nM}$) than nevirapine in the same assays.¹⁶ Furthermore, nevirapine has been found to be effective against all clinical isolates of HIV-1 which have been tested, including those which were AZT-sensitive or AZT-resistant. Nevirapine is specific for HIV-1; it is ineffective against HIV-2 and is inactive against simian and feline RT. Moreover, it does not inhibit human DNA polymerases α , β , γ , and δ , and calf thymus DNA polymerase α .²³ Similar results in many of these tests were obtained for 94; however, the 11-cyclopropyl compound (96) exhibited greater bioavailability in the rat and monkey compared to the corresponding 11-ethyl analogue (94).²² Thus, on the basis of these and other results, and on the basis of its favorable levels of specificity, bioavailability, and metabolic stability, nevirapine was selected for clinical evaluation. Details of pharmacology,²⁴ toxicology,²⁵ kinetics,²⁶ metabolism,²² and pharmacokinetics²⁷ studies, and the proposed mechanism of action of this compound will be reported elsewhere.

Tetrahydroimidazobenzodiazepinone (TIBO) compounds were recently reported²⁸ to block in vitro replication of HIV-1. These substances, which were discovered by random screening, also act by inhibiting the HIV-1 RT enzyme, and are ineffective against HIV-2 replication. Although structurally distinct from the compounds reported here, a representative of these TIBO analogues acts on the enzyme at the same allosteric site as the tricyclic diazepinones.²⁰

Experimental Section

General. Compounds were purified by gravity chromatography over silica gel. Melting points were determined on a Büchi 510 melting point apparatus and are uncorrected. ¹H NMR spectra were recorded on a Bruker WM-250 (250-MHz) spectrometer and ¹³C NMR spectra were recorded on a Bruker AC-270 (69.2 MHz) spectrometer with Me₄Si as the internal standard. NOE enhancements were measured by saturating the resonance of interest for one second prior to the observation pulse. All NOE enhancements reported were greater than 5%. Mass spectra were recorded on a Finnegan 4023 GC/MS/DS spectrometer. Elemental analyses were determined by Midwest Laboratories, Indianapolis, IN, and are within 0.4% of theoretical values unless otherwise noted.

Method A. 6,11-Dihydro-7-methyl-5*H*-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (30). A well-stirred mixture of 3-methyl-*o*-phenylenediamine (15 g, 0.123 mol) and 2-chloronicotinic

- (21) Eberlein, W. G.; Trummelitz, G.; Engel, W. W.; Schmidt, G.; Pelzer, H.; Mayer, N. *J. Med. Chem.* 1987, 30, 1378.
 (22) Riska, P.; Erickson, D.; Dinallo, R.; Hansen, G.; Hattox, S. Unpublished results.

- (23) (a) Merluzzi, V. J.; Hargrave, K. D.; Labadia, M.; Grozinger, K.; Skoog, M.; Wu, J.; Shih, C.-K.; Eckner, K.; Hattox, S.; Adams, J.; Rosenthal, A. S.; Faanes, R.; Eckner, R. J.; Koup, R. A.; Sullivan, J. L. *Science* 1990, 250, 1411. (b) Richman, D.; Rosenthal, A. S.; Skoog, M.; Eckner, R. J.; Chou, T.-C.; Sabo, J. P.; Merluzzi, V. J. *Antimicrob. Agents Chemother.* 1991, 35, 305. (c) Koup, R. A.; Merluzzi, V. J.; Hargrave, K. D.; Adams, J.; Grozinger, K.; Eckner, R. J.; Sullivan, J. L. *J. Inf. Dis.*, in press.
 (24) Frei, P.; Winquist, R.; Shao, J.; LeClercq, T.; Letts, G.; Panzenbeck, M.; Harrison, P. Unpublished results.
 (25) Andrews, L.; Tigner, J.; Pack, F. Unpublished results.
 (26) Kopp, E. B.; Miglietta, J. J.; Shrutkowski, A. G.; Shih, C.-K.; Grob, P. M.; Skoog, M. T. *Nuc. Acids Res.*, in press.
 (27) Riska, P.; Silverstein, H.; Johnstone, N.; St. George, R.; Cohn, D.; Erickson, D.; Norris, S.; Keirns, J.; Hattox, S. Unpublished results.
 (28) Pauwels, R.; Andries, K.; Desmyter, J.; Schols, D.; Kukla, M. J.; Breslin, H. J.; Raeymaeckers, A.; Van Gelder, J.; Woestenborghs, R.; Heykants, J.; Schellekens, K.; Janssen, M. A. C.; De Clercq, E.; Janssen, P. A. J. *Nature* 1990, 343, 470.

acid (23.2 g, 0.147 mol) in 2-butoxyethanol (200 mL) was heated at 140–150 °C for 6 h. The dark solution was then poured over crushed ice, and the resulting dark brown solid was collected, washed with water, and dried. Recrystallization from 50% 1,4-dioxane/water with charcoal treatment afforded **30** (13.7 g, 50%) as a light yellow crystalline solid: mp 231–232 °C; ¹H NMR (*d*₆-DMSO) δ 9.35 (s, 1 H, NH), 8.6 (s, 1 H, NH), 8.26 (dd, *J* = 1.8, 4.8 Hz, 1 H), 8.0 (dd, *J* = 1.8, 7.7 Hz, 1 H), 7.05 (dd, *J* = 1.8, 7.7 Hz, 1 H), 6.96 (dd, *J* = 4.8, 7.7 Hz, 1 H), 6.92 (apparent t, *J* = 7.7 Hz, 1 H), 6.84 (dd, *J* = 1.8, 7.7 Hz, 1 H), 2.3 (s, 3 H). The position of the methyl group was assigned on the basis of the observed NOE between δ 9.3 (–NHCO–) and δ 2.3 (CH₃). Anal. (C₁₃H₁₁N₃O·0.1H₂O) C, H, N.

8,9-Dichloro-6,11-dihydro-6-methyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (VII; R¹ = Me, R³ = 8,9-Cl₂, R⁴ = H). A 30% aqueous NaOH solution (2.9 mL, 22 mmol) was added to a suspension of 6,11-dihydro-8,9-dichloro-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (5.6 g, 20 mmol), obtained in 83% yield from 3,4-dichloro-*o*-phenylenediamine and 2-chloronicotinic acid according to the above procedure, in DMSO (40 mL). After 2 h, methyl iodide (5.0 mL, 80 mmol) was added and stirring was continued for 3 days at ambient temperature. The reaction mixture was then diluted with water, and the resulting solid was collected, washed with water, and air-dried. Trituration with EtOH afforded the title compound (5.3 g, 91%) as a light yellow solid: mp 294–297 °C; ¹H NMR (*d*₆-DMSO) δ 8.97 (s, 1 H, NH), 8.32 (dd, *J* = 4.8, 1.8 Hz, 1 H), 8.08 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.61 (s, 1 H), 7.49 (s, 1 H), 7.07 (dd, *J* = 7.7, 4.8 Hz, 1 H), 3.38 (s, 3 H, CH₃). Anal. (C₁₃H₉Cl₂N₃O·0.2H₂O) C, H, N.

6,11-Dihydro-6-ethyl-7-methyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (VII; R¹ = Et, R³ = 7-Me, R⁴ = H) and 5-Ethoxy-7-methyl-11H-pyrido[2,3-*b*]-[1,5]benzodiazepine (X; R¹ = Et, R³ = 7-Me, R⁴ = H). Sodium hydride (0.13 g, 5.5 mmol) was added at room temperature to a suspension of **30** (1.12 g, 5.0 mmol) in DMF (20 mL) under argon. The reaction mixture was heated at 60 °C for 1 h and then was cooled in an ice bath before the addition of ethyl iodide (0.86 g, 5.5 mmol). Stirring was continued at room temperature overnight, followed by warming to 60 °C for 1 h. The solution was concentrated in vacuo to near dryness, the residue was poured over ice-water, and the resulting solid was collected, washed with water, and dried. Purification by chromatography (elution with 1% MeOH/CH₂Cl₂) afforded, in order of elution, X (R¹ = Et, R³ = 7-Me, R⁴ = H) (0.23 g, 18%) and VII (R¹ = Et, R³ = 7-Me, R⁴ = H) (0.61 g, 49%). The *O*-alkylated product, X, crystallized as yellow needles from petroleum ether (30–60 °C): mp 135–136 °C; ¹H NMR (*d*₆-DMSO) δ 8.18 (dd, *J* = 4.8, 1.8 Hz, 1 H), 7.94 (s, 1 H, NH), 7.72 (dd, *J* = 7.7, 1.8 Hz, 1 H), 6.91 (dd, *J* = 7.7, 4.8 Hz, 1 H), 6.82 (m, 3 H), 4.34 (q, *J* = 7.0 Hz, 2 H), 2.21 (s, 3 H), 1.36 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (*d*₆-DMSO) δ 163.06, 155.97 (N=CO), 151.13, 137.38, 138.32, 134.70, 134.55, 125.13, 124.57, 61.83 (OCH₂), 18.33, 14.02; MS (EI) *m/z* 253 (M⁺), 238 ([M – CH₃]⁺), 225 ([M – C₂H₄]⁺), 208 ([M – OC₂H₅]⁺), 196 ([M – COC₂H₅]⁺). Anal. (C₁₅H₁₅N₃O·0.1H₂O) C, H, N.

The *N*-alkylated product VII (R¹ = Et, R³ = 7-Me, R⁴ = H) crystallized from 1,4-dioxane/water: mp 186–188 °C; ¹H NMR (*d*₆-DMSO) δ 8.59 (s, 1 H, NH), 8.25 (dd, *J* = 4.8, 1.8 Hz, 1 H), 8.01 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.08 (m, 4 H), 4.38 (dq, *J* = –14.0, 7.0 Hz, 1 H), 3.09 (dq, *J* = –14.0, 7.0 Hz, 1 H), 2.25 (s, 3 H), 0.97 (t, *J* = 7.0 Hz, 3 H); ¹³C NMR (*d*₆-DMSO) δ 167.50, 161.09 (C=O), 150.44, 145.26, 140.86, 134.44, 132.00, 126.45, 118.62, 117.79, 44.53 (NCH₂), 18.55, 13.11; MS (CI) 254 (MH⁺). Anal. (C₁₅H₁₅N₃O) C, H, N.

6,11-Dihydro-11-ethyl-6,7-dimethyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (31). Sodium hydride (80% oil dispersion, 0.165 g, 5.5 mmol) was added in several portions to a stirred suspension of 5,6-dihydro-6,7-dimethyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (VII; R¹ = Me, R³ = 7-Me, R⁴ = H) (1.2 g, 5.0 mmol) in DMF (10 mL) under argon. The resulting mixture was stirred at room temperature for 3 h and at 60 °C for 1 h, and then was cooled to room temperature prior to the addition of ethyl iodide (0.86 g, 5.5 mmol). After 2 days, the reaction mixture was concentrated in vacuo and the residue was dissolved in EtOAc, washed successively with saturated aqueous NaHCO₃ and saturated aqueous NaCl, dried (MgSO₄), and concentrated in vacuo. Purification by chromatography (elution with 10–30%

EtOAc/hexane) and recrystallization from CH₂CN afforded **31** (1.0 g, 78%): mp 132–134 °C; ¹H NMR (*d*₆-DMSO) δ 8.34 (dd, *J* = 4.8, 1.8 Hz, 1 H), 7.95 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.10 (m, 4 H), 4.10 (dq, *J* = –14.0, 7.0 Hz, 1 H), 3.64 (dq, *J* = –14.0, 7.0 Hz, 1 H), 3.28 (s, 3 H), 2.28 (s, 3 H), 1.16 (t, *J* = 7.0 Hz, 3 H). Anal. (C₁₆H₁₇N₃O) C, H, N.

6-Benzyl-6,11-dihydro-7-methyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (VII; R¹ = CH₂C₆H₅, R³ = 7-Me, R⁴ = H). Potassium (0.215 g, 5.5 mmol) was dissolved in *t*-BuOH (8 mL) and the resulting solution was added in portions to a stirred solution of **30** (1.18 g, 5.0 mmol) in dry dioxane (10 mL) under argon. The mixture was heated at 60 °C for 3 h, and then cooled to room temperature. Benzyl bromide (0.94 g, 5.5 mmol) was added and the reaction mixture was stirred overnight at room temperature. Dilution with ice-water produced an oil, which solidified upon standing. The solid was collected, air-dried, and recrystallized from EtOH/water to give VII (R¹ = CH₂C₆H₅, R³ = 7-Me, R⁴ = H) (1.11 g, 69%): mp 174–175 °C; ¹H NMR (*d*₆-DMSO) δ 8.72 (s, 1 H, NH), 8.26 (dd, *J* = 4.8, 1.8 Hz, 1 H), 8.12 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.18 (m, 9 H), 5.52 (d, *J* = –15.1 Hz, 1 H), 4.34 (d, *J* = –15.1 Hz, 1 H), 2.37 (s, 3 H). Anal. (C₂₀H₁₇N₃O) C, H, N.

6-Benzyl-6,11-dihydro-11-ethyl-7-methyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (32). This product was prepared in 80% yield as described above for **31**. It was crystallized from EtOH: mp 145.5–147 °C; ¹H NMR (*d*₆-DMSO) δ 8.34 (dd, *J* = 4.8, 1.8 Hz, 1 H), 7.97 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.20 (m, 9 H), 5.61 (d, *J* = –15.1 Hz, 1 H), 4.39 (d, *J* = –15.1 Hz, 1 H), 4.14 (dq, *J* = –14.0, 7.0 Hz, 1 H), 3.54 (dq, *J* = –14.0, 7.0 Hz, 1 H), 2.39 (s, 3 H), 1.08 (t, *J* = 7.0 Hz, 3 H). Anal. (C₂₂H₂₁N₃O) C, H, N.

6,11-Dihydro-11-ethyl-7-methyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (33). A stirred mixture of **32** (79 mg, 0.2 mmol) and anisole (0.05 mL, 0.02 mmol) in trifluoroacetic acid (5 mL) containing concentrated H₂SO₄ (0.25 mL) was heated at 110 °C for 8 h. The resulting dark solution was poured into ice-water (100 mL), and the milky suspension was extracted with EtOAc (3 × 50 mL). The combined extract was washed successively with saturated aqueous NaHCO₃ and saturated aqueous NaCl, dried (MgSO₄), and concentrated in vacuo. The residue was purified by chromatography (elution with 10–20% EtOAc/hexane) to give **33** (30 mg, 52%): mp 225–227 °C; ¹H NMR (*d*₆-DMSO) δ 9.81 (s, 1 H, NH), 8.41 (dd, *J* = 4.8, 1.8 Hz, 1 H), 7.97 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.12 (dd, *J* = 7.7, 4.8 Hz, 1 H), 7.04 (m, 3 H), 4.08 (br, 1 H), 3.69 (br, 1 H), 2.31 (s, 3 H), 1.13 (t, *J* = 7.0 Hz, 3 H). Anal. (C₁₅H₁₅N₃O) C, H, N.

6,11-Dihydro-11-ethyl-6,7-dimethyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepine-5-thione (IX; R¹ = Me, R² = Et, R³ = 7-Me, R⁴ = H). Lawesson's reagent¹² (0.25 g, 0.625 mmol) was added to a solution of **31** (0.32 g, 1.25 mmol) in toluene (10 mL), and the reaction mixture was stirred at reflux for 3 h. Concentration in vacuo followed by chromatography (elution with CH₂Cl₂) of the residue and recrystallization from CH₂CN afforded the title compound (0.3 g, 86%): mp 196–199 °C; ¹H NMR (*d*₆-DMSO) δ 8.28 (dd, *J* = 4.8, 1.8 Hz, 1 H), 8.16 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.15 (m, 4 H), 4.02 (dq, *J* = –14.0, 7.0 Hz, 1 H), 3.71 (s, 3 H), 3.61 (dq, *J* = –14.0, 7.0 Hz, 1 H), 2.28 (s, 3 H), 1.16 (t, *J* = 7.0 Hz, 3 H). Anal. (C₁₆H₁₇N₃S) C, H, N, S.

6,11-Dihydro-3-chloro-11-ethyl-6-methyl-5H-pyrido[2,3-*b*]-[1,5]benzodiazepin-5-one (38). Chlorine gas (0.15 g, 2.2 mmol) was absorbed in trimethyl phosphate (TMP) (1.5 mL) and added to an ice-cold, stirred solution of **11** (0.50 g, 1.97 mmol) in TMP (5.0 mL). The resulting mixture was allowed to warm to room temperature overnight. An additional equivalent of the TMP-Cl₂ solution was added at 24 h, and then a third equivalent at 48 h. On the 4th day, excess chlorine was removed in vacuo and the reaction mixture was then poured into ice-water. The resulting white solid was collected, washed, dried, and purified by chromatography (elution with 0–30% EtOAc/hexane). The product crystallized from petroleum ether (30–60 °C) in white needles (130 mg, 23%): mp 131–132 °C; ¹H NMR (*d*₆-DMSO) δ 8.45 (d, *J* = 2.6 Hz, 1 H), 8.00 (d, *J* = 2.6 Hz, 1 H), 7.41–7.44 (m, 1 H), 7.21–7.31 (m, 3 H), 4.15 (dq, *J* = –14, 7.0 Hz, 1 H), 3.60 (dq, *J* = 14, 7.0 Hz, 1 H), 3.45 (s, 3 H), 1.15 (t, *J* = 7.0 Hz, 3 H). Anal. (C₁₅-H₁₄ClN₃O) C, H, N.

Synthesis of the Photoaffinity Probe (45). **6,11-Dihydro-6-methyl-9-nitro-5H-pyrido[2,3-*b*]-[1,5]benzo-**

diazepin-5-one (VII; R¹ = CH₃, R³ = 9-NO₂, R⁴ = H) and 6,11-Dihydro-6-methyl-8-nitro-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (VII; R¹ = CH₃, R³ = 8-NO₂, R⁴ = H). A mixture of 2-chloronicotinic acid (15.7 g, 0.1 mol) and 4-nitro-*o*-phenylenediamine (15.3 g, 0.1 mol) was heated in sulfolane (100 mL) at 170 °C for 5 h. After cooling to room temperature, the solid material was removed by filtration, washed with hot EtOH, and air-dried to give a mixture of the 8- and 9-nitro isomers (21 g, 0.079 mol, 79%). This mixture was added, as a solution in DMSO (100 mL), to a solution of dimethyl sodium (200 mL, 0.42 M). After the reaction mixture had stirred for 1 h, methyl iodide (12 g, 0.08 mol) was added and stirring was continued for an additional 12 h. Water (200 mL) was added and the precipitate was removed by filtration. Fractional recrystallization from EtOH afforded VII (R¹ = CH₃, R³ = 9-NO₂, R⁴ = H) (6.8 g, 0.025 mol, 31%): mp 278–280 °C; ¹H NMR (*d*₆-DMSO) δ 9.20 (br s, 1 H, NH), 8.34 (dd, *J* = 2.0, 4.7 Hz, 1 H), 8.18 (d, *J* = 3.0 Hz, 1 H), 8.10 (dd, *J* = 2.0, 7.7 Hz, 1 H), 7.96 (dd, *J* = 3.0, 9.0 Hz, 1 H), 7.53 (d, *J* = 9.0 Hz, 1 H), 7.08 (dd, *J* = 4.7, 7.7 Hz, 1 H), 3.42 (s, 3 H). An NOE was observed between the resonances at δ 3.42 and 7.53, and between the resonances at δ 9.2 and 8.18; MS (CI) *m/z* 271 (MH⁺). Anal. (C₁₃H₁₀N₄O₃) C, H, N. The combined supernatants were concentrated in vacuo, and the residue was purified by chromatography (elution with 0.5% CH₂Cl₂/hexane). A second chromatographic purification (elution with 30–50% EtOAc/hexane) gave VII (R¹ = CH₃, R³ = 8-NO₂, R⁴ = H) (0.3 g, 1.4%): mp 245–247 °C; ¹H NMR (*d*₆-DMSO) δ 9.54 (br s, 1 H, NH), 8.35 (dd, *J* = 1.9, 4.7 Hz, 1 H), 8.11–8.14 (m, 2 H), 8.03 (dd, *J* = 2.5, 9 Hz, 1 H), 7.43 (d, *J* = 9 Hz, 1 H), 7.10 (dd, *J* = 4.7, 7.7 Hz, 1 H), 3.44 (s, 3 H). An NOE was observed between the resonances at δ 3.44 and 8.11–8.14, and between the resonances at δ 9.54 and 7.43; MS (CI) *m/z* 271 (MH⁺). Anal. (C₁₃H₁₀N₄O₃) C, H, N.

6,11-Dihydro-11-ethyl-6-methyl-9-nitro-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (42). A solution of VII (R¹ = CH₃, R³ = 9-NO₂, R⁴ = H) (1.5 g, 5.3 mmol) in DMSO (5 mL) was added to a solution of dimethyl sodium (5 mL, 1.1 M), and the resulting mixture was stirred for 20 min. Ethyl iodide (0.98 g, 6.3 mmol) was added and stirring was continued for an additional 20 min. Water (100 mL) was then added, and the mixture was extracted with CH₂Cl₂. The organic phase was washed with water, dried (Na₂SO₄), filtered, and concentrated. The residue was purified by chromatography (elution with CH₂Cl₂/EtOH) to give 42, which crystallized from diisopropyl ether (0.5 g, 31%): mp 175–178 °C; ¹H NMR (*d*₆-DMSO) δ 8.47 (dd, *J* = 1.9, 4.8 Hz, 1 H), 7.99–8.09 (m, 3 H), 7.66 (d, *J* = 8.9 Hz, 1 H), 7.20 (dd, *J* = 4.8, 7.7 Hz, 1 H), 4.30 (m, 1 H), 3.67 (m, 1 H), 3.50 (s, 3 H), 1.18 (t, *J* = 7.0 Hz, 3 H); MS (EI) *m/z* 298 (M⁺). Anal. (C₁₈H₁₄N₄O₃) C, H, N. The supernatant was concentrated and the residue was purified by chromatography (elution with 30% EtOAc/hexane) to give the 1-ethylated product (XIb; R = Et, R¹ = Me, R³ = 9-NO₂, R⁴ = H), which crystallized from EtOAc/hexane (0.09 g, 6%): mp 179–180.5 °C; ¹H NMR (*d*₆-DMSO) δ 7.92 (dd, *J* = 1.9, 7.2 Hz, 1 H), 7.88 (dd, *J* = 1.9, 6.5 Hz, 1 H), 7.80 (dd, *J* = 2.8, 9.0 Hz, 1 H), 7.59 (d, *J* = 2.8 Hz, 1 H), 7.24 (d, *J* = 9.0 Hz, 1 H), 6.16 (dd, *J* = 6.5, 7.2 Hz, 1 H), 4.15 (q, *J* = 7.0 Hz, 2 H), 3.25 (s, 3 H), 1.35 (t, *J* = 7.0 Hz, 3 H); MS (CI) *m/z* 299 (MH⁺). Anal. (C₁₈H₁₄N₄O₃) C, H, N.

9-Amino-6,11-dihydro-11-ethyl-6-methyl-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (43). To a solution of 42 (0.3 g, 1.0 mmol) in HOAc (5 mL) was added a solution of SnCl₂·2H₂O (1.7 g, 7.5 mmol) in concentrated HCl (2.2 mL). The mixture was stirred at room temperature for 6 h and then basified with saturated aqueous NaHCO₃. Extraction with EtOAc and concentration in vacuo afforded a solid residue, which recrystallized from EtOAc to give 43 (0.12 g, 45%): mp 193–195 °C; ¹H NMR (*d*₆-DMSO) δ 8.35 (dd, *J* = 2.0, 4.8 Hz, 1 H), 7.94 (dd, *J* = 2.0, 7.5 Hz, 1 H), 7.09 (dd, *J* = 4.8, 7.5 Hz, 1 H), 7.02 (d, *J* = 8.5 Hz, 1 H), 6.35–6.41 (m, 2 H), 5.17 (br s, 2 H, NH₂), 4.08 (m, 1 H), 3.48 (m, 1 H), 3.35 (s, 3 H), 1.15 (t, *J* = 7.0 Hz, 3 H); MS (CI) *m/z* 269 (MH⁺). Anal. (C₁₅H₁₆N₄O) C, H, N.

9-Azido-6,11-dihydro-11-ethyl-6-methyl-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (45). To a solution of 43 (0.21 g, 0.78 mmol) in water (3.5 mL) and concentrated HCl (0.5 mL) was added over 40 min (at such a rate that the internal temperature remained below 5 °C) a solution of NaNO₂ (0.24 g, 2.4 mmol) in

water (10 mL). The mixture was stirred at 5 °C for an additional 20 min. A solution of NaN₃ (0.41 g, 6.3 mmol) in water (3 mL) was then added all at once, and stirring was continued for 15 min. The reaction mixture was extracted with CH₂Cl₂, and the organic phase was dried (Na₂SO₄) and concentrated. The residue was recrystallized from Et₂O to give 45 (0.13 g, 56%): mp 115–118 °C; ¹H NMR (*d*₆-DMSO) δ 8.41 (dd, *J* = 2.0, 4.8 Hz, 1 H), 7.99 (dd, *J* = 2.0, 7.7 Hz, 1 H), 7.43 (d, *J* = 8.6 Hz, 1 H), 7.15 (dd, *J* = 4.8, 7.7 Hz, 1 H), 6.98 (dd, *J* = 2.4, 8.6 Hz, 1 H), 6.91 (d, *J* = 2.4 Hz, 1 H), 4.19 (m, 1 H), 3.60 (m, 1 H), 3.42 (s, 3 H), 1.15 (t, *J* = 7.0 Hz, 3 H); MS (CI) *m/z* 295 (MH⁺). Anal. (C₁₅H₁₄N₆O) C, H, N.

6,11-Dihydro-11-ethyl-6-methyl-8-(phthalimidomethyl)-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (II; R¹ = Me, R² = Et, R³ = 8-phthalimidomethyl, R⁴ = H). Into a stirred solution of *N*-(hydroxymethyl)phthalimide (26.57 g, 0.15 mol) in H₂SO₄ (200 mL) at 5 °C was added a solution of 11 (38 g, 0.15 mol) in H₂SO₄ (300 mL), and the resulting mixture was stirred at room temperature for 5 days. The reaction mixture was then poured onto ice, neutralized to pH 8 with NH₄OH, and extracted with CH₂Cl₂. The organic layer was concentrated in vacuo and the residue was purified by chromatography (elution with 2% MeOH/CH₂Cl₂). Recrystallization from CH₂Cl₂/Et₂O afforded the title compound (42 g, 68%): mp 190–192 °C; ¹H NMR (*d*₆-DMSO) 8.38 (dd, *J* = 4.8, 1.8 Hz, 1 H), 7.96 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.86 (m, 4 H), 7.39 (d, *J* = 1.5 Hz, 1 H), 7.22 (d, *J* = 8.1 Hz, 1 H), 7.11 (m, 2 H), 4.75 (s, 2 H), 4.16 (dd, *J* = -14.0, 7.0 Hz, 1 H), 3.57 (dq, *J* = -14.0, 7.0 Hz, 1 H), 3.43 (s, 3 H), 1.13 (t, *J* = 7.0 Hz, 3 H); MS (CI) *m/z* 413 (MH⁺). Anal. (C₂₄H₂₀N₄O₃) C, H, N.

8-(Aminomethyl)-6,11-dihydro-11-ethyl-6-methyl-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (50). A suspension of II (R¹ = Me, R² = Et, R³ = 8-phthalimidomethyl, R⁴ = H) (38 g, 0.09 mol) and hydrazine hydrate (23 g, 0.46 mol) in EtOH (900 mL) was stirred at reflux for 1 h. The mixture was then filtered and the filtrate was concentrated to 150 mL. Water (500 mL) was added and the solution was basified with 2 N NaOH. Following extraction into CH₂Cl₂ and concentration in vacuo, the crude product was purified by chromatography (elution with 10% MeOH/CH₂Cl₂) to give 50 (25 g, 96%) as a colorless, hygroscopic resin: ¹H NMR (*d*₆-DMSO) δ 8.38 (m, 1 H), 7.9 (m, 1 H), 7.2–7.3 (m, 4 H), 4.2 (m, 1 H), 3.5–3.6 (m, 1 H), 3.7 (s, 2 H), 3.4 (s, 3 H), 1.1 (m, 3 H); MS (CI) *m/z* 283 (MH⁺). Anal. (C₁₆H₁₈N₄O·0.5H₂O) C, H, N.

6,11-Dihydro-11-ethyl-8-(hydroxymethyl)-6-methyl-5H-pyrido[2,3-*b*][1,5]benzodiazepin-5-one (52). A solution of 50 (1.9 g, 7 mmol) in 50% HOAc (50 mL) was treated with NaNO₂ (5.7 g, 80 mmol), and the resulting mixture was stirred at room temperature for 24 h. The reaction mixture was then basified with 40% aqueous NaOH and extracted with Et₂O. The organic extract was treated with a solution of 5% KOH/MeOH (50 mL) for 2 h in order to effect hydrolysis. The reaction mixture was washed with water and the organic layer was concentrated in vacuo. Chromatographic purification (elution with 1.5% MeOH/CH₂Cl₂) of the residue afforded 52 (1.75 g, 92%) as a colorless oil: ¹H NMR (*d*₆-DMSO) δ 8.38 (dd, *J* = 4.8, 1.8 Hz, 1 H), 7.98 (dd, *J* = 7.7, 1.8 Hz, 1 H), 7.1–7.2 (m, 4 H), 5.20 (t, *J* = 5.5 Hz, 1 H), 4.46 (d, *J* = 5.5 Hz, 1 H), 4.17 (dq, *J* = -14.0, 7.0 Hz, 1 H), 3.60 (dq, *J* = -14.0, 7.0 Hz, 1 H), 3.45 (s, 3 H), 1.15 (t, *J* = 7.0 Hz, 3 H); MS (CI) *m/z* 284 (MH⁺). Anal. (C₁₆H₁₇N₃O₂) C, H, N.

Method B. 2-Chloro-*N*-(2-chloro-3-pyridinyl)-3-pyridinocarboxamide (XIVa; R³ = R⁴ = H). To a solution of 3-amino-2-chloropyridine (215 g, 1.672 mol) in a mixed solvent system of 1,4-dioxane (400 mL), cyclohexane (500 mL), and pyridine (130 mL) was added over 30 min a solution of 2-chloronicotinoyl chloride (292.2 g, 1.7 mol) in 1,4-dioxane (200 mL). The reaction mixture was stirred overnight at ambient temperature under N₂ atmosphere. The solid product was removed by filtration, dissolved in CH₂Cl₂, washed with water, dried (MgSO₄), and concentrated in vacuo. The resultant solid was washed with EtOAc, filtered, and dried in vacuo to give the desired product (386 g, 84%): mp 156–159 °C; ¹H NMR (*d*₆-DMSO) δ 10.61 (br s, 1 H), 8.56 (dd, *J* = 1.8, 4.8 Hz, 1 H), 8.32 (dd, *J* = 1.8, 4.8 Hz, 1 H), 8.23 (dd, *J* = 1.8, 7.7 Hz, 1 H), 8.1 (dd, *J* = 1.8, 7.7 Hz, 1 H), 7.59 (dd, *J* = 4.8, 7.7 Hz, 1 H), 7.53 (dd, *J* = 4.8,

7.7 Hz, 1 H); MS (CI) m/z 268 (MH⁺). Anal. (C₁₁H₇Cl₂N₃O) C, H, Cl, N.

2-Chloro-*N*-methyl-*N*-(2-chloro-3-pyridinyl)-3-pyridinecarboxamide (XIVb; R¹ = Me, R³ = R⁴ = H). Sodium hydride (60% dispersion in oil, 5.0 g, 0.125 mol) was added in several portions over 30 min to a solution of 2-chloro-*N*-(2-chloro-3-pyridinyl)-3-pyridinecarboxamide (27 g, 0.1 mol) in DMSO (100 mL) under nitrogen, and the mixture was heated at 60 °C for 1 h. The solution was then cooled to room temperature, methyl iodide (8.0 mL, 0.125 mol) was added, and the reaction mixture was stirred overnight at room temperature. Excess base was destroyed by the addition of ice, and the mixture was partitioned between EtOAc and water. The aqueous phase was extracted with EtOAc, and the combined organic extract was concentrated in vacuo. The residue was taken up in Et₂O, washed with water, dried (MgSO₄), and concentrated in vacuo to give an oil. Trituration with Et₂O/petroleum ether (30–60 °C) afforded the title compound (22.0 g, 77%): ¹H NMR (CDCl₃) (2:1 mixture of conformers) δ 8.58 (dd, J = 1.8, 4.8 Hz, 1/3 H), 8.48 (dd, J = 1.8, 4.7 Hz, 1/3 H), 8.28–8.32 (m, 2 × 2/3 H), 8.05–8.11 (m, 2 × 1/3 H), 7.89 (dd, 1.9, 7.6 Hz, 2/3 H), 7.59–7.65 (m, 2 × 1/3 H), 7.42 (dd, J = 4.7, 7.9 Hz, 2/3 H), 7.35 (dd, J = 4.8, 7.6 Hz, 2/3 H), 3.34 (s, 3 × 2/3 H), 3.12 (s, 3 × 1/3 H).

Synthesis of 11-Cyclopropyl-5,11-dihydro-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (96). 3-Amino-2-chloro-4-methylpyridine. 2-Chloro-4-methyl-3-nitropyridine (21.4 g, 0.124 mol) in EtOH (100 mL) was hydrogenated under pressure (50 psi) in the presence of 5% Rh/C (1.3 g) for 3 h. The catalyst was removed by filtration through Celite, and the filtrate was concentrated in vacuo. The residue was treated with hot EtOAc (100 mL) and the resultant solution was filtered through Celite. Concentration of the clear filtrate afforded the title compound (13.3 g, 75%): mp 65–66 °C; ¹H NMR (d₆-DMSO) δ 7.51 (d, J = 4.6 Hz, 1 H), 7.00 (d, J = 7.6 Hz, 1 H), 5.23 (br s, 2 H), 2.16 (s, 3 H); MS (CI) m/z 143 (MH⁺). Anal. (C₈H₇ClN₂) C, H, Cl, N.

2-Chloro-*N*-(2-chloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (XIVa; R³ = Me, R⁴ = H). To a solution of 3-amino-2-chloro-4-methylpyridine (260 g, 1.82 mol) in a mixture of cyclohexane (600 mL), 1,4-dioxane (100 mL), and pyridine (575 mL) was added in portions over 30 min a solution of 2-chloro-nicotinoyl chloride (321 g, 1.28 mol) in 1,4-dioxane (500 mL). The resulting mixture was stirred at ambient temperature for 48 h to produce a crystalline precipitate, which was removed by filtration and washed with water. The wet filter cake was added to a mixture of EtOH (1.75 L) and aqueous NaOH (0.1 N, 360 mL), heated under reflux for 2 h, and then stirred overnight at ambient temperature. The solvent was evaporated in vacuo and water (1 L) was added, with stirring, to the residue. The mixture was cooled to 10 °C and the crystalline product was collected by filtration, washed with cold water and dried under vacuum to give the title compound (337 g, 65.5%): mp 189–191 °C; ¹H NMR (d₆-DMSO) δ 10.60 (br s, 1 H), 8.58 (dd, J = 1.9, 5.0 Hz, 1 H), 8.27 (d, J = 5.0 Hz, 1 H), 8.09 (dd, J = 1.9, 7.6 Hz, 1 H), 7.61 (dd, J = 5.0, 7.6 Hz, 1 H), 7.43 (d, J = 5.0 Hz, 1 H), 2.38 (s, 3 H); MS (CI) m/z 282 (MH⁺). Anal. (C₁₂H₉Cl₂N₃O) C, H, Cl, N.

***N*-(2-Chloro-4-methyl-3-pyridinyl)-2-(cyclopropylamino)-3-pyridinecarboxamide (XVa; R² = *c*-Pr, R³ = Me, R⁴ = H).** A mixture of 2-chloro-*N*-(2-chloro-4-methyl-3-pyridinyl)-3-pyridinecarboxamide (167 g, 0.59 mol) and cyclopropylamine (135 g, 2.35 mol) in xylene (1 L) was heated with stirring at 110 °C in a 2-L autoclave for 18 h. After cooling to room temperature, the mixture was diluted with CH₂Cl₂ (2 L), washed with water, dried (Na₂SO₄), filtered, and concentrated in vacuo. The residual oil was triturated with CH₂Cl₂ and Et₂O to give the title compound (147 g, 83%): mp 127–129 °C; ¹H NMR (d₆-DMSO) δ 10.2 (br s, 1 H), 8.33 (dd, J = 1.7, 4.8 Hz, 1 H), 8.19–8.25 (m, 3 H), 7.41 (d, J = 5.2 Hz, 1 H), 6.75 (dd, J = 4.8, 7.7 Hz, 1 H), 2.84 (m, 1 H), 2.25 (s, 3 H), 0.72 (m, 2 H), 0.43 (m, 2 H); MS (CI) m/z 303 (MH⁺). Anal. (C₁₅H₁₅ClN₄O) C, H, Cl, N.

11-Cyclopropyl-5,11-dihydro-4-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (96). To a solution of *N*-(2-chloro-4-methyl-3-pyridinyl)-2-(cyclopropylamino)-3-pyridinecarboxamide (145 g, 0.48 mol) in bis(2-methoxyethyl) ether (870 mL) under nitrogen was added, in portions, NaH (50% dispersion

in oil, 46 g, 0.96 mol). The mixture was heated slowly to 160 °C (Caution! Potential exotherm with vigorous H₂ evolution at 150 °C) and then heated under gentle reflux for 1.5 h. The mixture was cooled to 65 °C, poured onto ice (2 kg) and water (1 L), and left stirring overnight. The precipitate was removed by filtration and dissolved in hot pyridine (1 L). The product was precipitated by the slow addition of water (10 L), collected by suction filtration, and stirred in MeOH (1 L) at reflux. Filtration, followed by recrystallization of the collected solid from pyridine/water, gave 96 (85.6 g, 67%): mp 247–249 °C; ¹H NMR (d₆-DMSO) δ 9.90 (br s, 1 H), 8.51 (dd, J = 1.9, 4.7 Hz, 1 H), 8.08 (d, J = 5.0 Hz, 1 H), 8.02 (dd, J = 1.9, 8.0 Hz, 1 H), 7.20 (dd, J = 4.7, 8.0 Hz, 1 H), 7.07 (d, J = 5.0 Hz, 1 H), 3.62 (m, 1 H), 2.34 (s, 3 H), 0.88 (m, 2 H), 0.35 (m, 2 H); MS (CI) m/z 267 (MH⁺). Anal. (C₁₅H₁₄N₄O) C, H, N.

Method C. *N*-(2-Chloro-3-pyridinyl)-2-[[4-methoxyphenyl)methyl]amino]-3-pyridinecarboxamide (XVII; R¹ = R³ = R⁴ = H). A mixture of 2-chloro-*N*-(2-chloro-3-pyridinyl)-3-pyridinecarboxamide (67.4 g, 0.251 mol) and 4-methoxybenzylamine (34.5 g, 0.251 mol) in xylene (400 mL) was heated under reflux for 2 h. The solvent was evaporated, and the residue was diluted with water and extracted with CH₂Cl₂. The organic phase was dried (Na₂SO₄), filtered, and concentrated in vacuo to give a brown oil. This crystallized upon addition of Et₂O to provide XVII (R¹ = R³ = R⁴ = H) (78.0 g, 91%) as a beige powder: mp 121–122 °C; ¹H NMR (d₆-DMSO) δ 10.2 (br s, 1 H, NH), 8.44 (br t, J = 5.5 Hz, 1 H, NH), 8.32 (dd, J = 1.8, 4.8 Hz, 1 H), 8.27 (dd, J = 1.8, 4.8 Hz, 1 H), 8.19 (dd, J = 1.8, 7.7 Hz, 1 H), 8.01 (dd, J = 1.8, 7.7 Hz, 1 H), 7.49 (dd, J = 4.8, 7.7 Hz, 1 H), 7.24–7.29 (m, 2 H), 6.85–6.90 (m, 2 H), 6.70 (dd, J = 4.8, 7.7 Hz, 1 H), 4.56 (d, J = 5.5 Hz, 1 H), 3.72 (s, 3 H); MS (CI) m/z 369 (MH⁺).

5,11-Dihydro-11-[[4-methoxyphenyl)methyl]-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (XVIIIa; R³ = R⁴ = H). Sodium hydride (50% dispersion in oil, 13.3 g, 270 mmol) was added to a solution of *N*-(2-chloro-3-pyridinyl)-2-[[4-methoxyphenyl)methyl]amino]-3-pyridinecarboxamide (34.03 g, 90 mmol) in DMF (500 mL). When hydrogen evolution had ceased, the reaction mixture was heated to 140 °C and left stirring overnight. The reaction mixture was quenched with ice, diluted with water, and extracted with Et₂O and CH₂Cl₂. The combined organic extract was concentrated in vacuo, redissolved in water, and extracted with Et₂O. The product crystallized from the Et₂O layer and was collected by filtration to give XVIIIa (R³ = R⁴ = H) (12.63 g, 42%): mp 209–210 °C; ¹H NMR (d₆-DMSO) δ 10.50 (s, 1 H, NH), 8.44 (dd, J = 1.8, 4.8 Hz, 1 H), 8.11 (dd, J = 1.8, 4.8 Hz, 1 H), 8.07 (dd, J = 1.8, 7.7 Hz, 1 H), 7.47 (dd, J = 1.8, 5.9 Hz, 1 H), 7.28 (d, J = 8.8 Hz, 2 H), 7.14 (m, 2 H), 6.79 (d, J = 8.8 Hz, 2 H), 5.23 (s, 2 H), 3.67 (s, 3 H). Anal. (C₁₉H₁₈N₄O₂) C, H, N.

5,11-Dihydro-11-[[4-methoxyphenyl)methyl]-5-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (XVIIIb; R¹ = Me, R³ = R⁴ = H). To a stirred slurry of XVIIIa (R³ = R⁴ = H) (2.92 g, 8.79 mmol) in DMF (20 mL) was added NaH (50% dispersion in oil, 0.464 g, 9.66 mmol) in one portion. After 2 h, methyl iodide (2.49 g, 17.6 mmol) was added and the reaction mixture was stirred for an additional 2 h. Diethyl ether (200 mL) was added, and the organic phase was washed with water, dried (MgSO₄), filtered, and concentrated in vacuo to give an oil. Purification by chromatography (elution with CH₂Cl₂) afforded XVIIIb (R¹ = Me, R³ = R⁴ = H) (0.85 g, 28%): mp 120–121.5 °C; ¹H NMR (d₆-DMSO) δ 8.36 (dd, J = 4.8, 1.9 Hz, 1 H), 8.16 (dd, J = 4.6, 1.6 Hz, 1 H), 8.11 (dd, J = 7.7, 1.9 Hz, 1 H), 7.48 (dd, J = 7.9, 1.6 Hz, 1 H), 7.34 (d, J = 8.7 Hz, 2 H), 7.05 (dd, J = 7.9, 4.7 Hz, 1 H), 7.00 (dd, J = 7.7, 4.8 Hz, 1 H), 6.76 (d, J = 8.7 Hz, 2 H), 3.73 (s, 3 H), 3.53 (s, 3 H); MS (CI) m/z 347 (MH⁺). Anal. (C₂₀H₁₈N₄O₂) C, H, N.

5,11-Dihydro-5-methyl-6*H*-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (XIXb, R¹ = Me, R³ = R⁴ = H). A solution of XVIIIb (R¹ = Me, R³ = R⁴ = H) (2.6 g, 7.5 mmol) in trifluoroacetic acid (15 mL) was stirred at room temperature for 1.5 h. After removal of the solvent, the residue was stirred with 0.5% NH₄OH for 2 h. The precipitated solid was removed by filtration, washed with water, air-dried, and recrystallized from EtOH to give XIXb (R¹ = Me, R³ = R⁴ = H) (1.51 g, 89%): mp 227–229 °C; ¹H NMR (d₆-DMSO) δ 8.83 (br s, 1 H, NH), 8.35 (dd, J = 1.8, 4.8 Hz, 1

H), 8.10 (dd, $J = 1.0, 4.8$ Hz, 1 H), 8.11 (dd, $J = 1.8, 7.7$ Hz, 1 H), 7.78 (dd, $J = 1.0, 7.7$ Hz, 1 H), 7.21 (dd, $J = 4.8, 7.7$ Hz, 1 H), 7.09 (dd, $J = 4.8, 7.7$ Hz, 1 H), 3.38 (s, 3 H); MS (CI) m/z 227 (MH⁺). Anal. (C₁₂H₁₀N₄O) C, H, N.

11-Allyl-5,11-dihydro-5-methyl-6H-dipyrido[3,2-*b*:2',3'-*e*][1,4]diazepin-6-one (78). To a suspension of XIXb (R¹ = Me, R³ = R⁴ = H) (0.5 g, 2.2 mmol) in DMF (20 mL) at 60 °C was added NaH (50% dispersion in oil, 0.2 g, 4.2 mmol), and the resultant mixture was stirred at 50 °C for 1 h. Allyl bromide (0.35 mL, 4.0 mmol) was added and stirring was continued at 50 °C for an additional 2 h and at room temperature overnight. The reaction was quenched with alcohol and the solvent was removed in vacuo. The residue was dissolved in CH₂Cl₂, washed with water, dried (MgSO₄), and concentrated in vacuo to give the crude product. Chromatography (elution with 50% EtOAc/hexane) and recrystallization from EtOAc/hexane afforded 78 (0.42 g, 72%): mp 93–95 °C; ¹H NMR (*d*₆-DMSO) δ 8.43 (dd, $J = 1.8, 4.8$ Hz, 1 H), 8.20 (dd, $J = 1.8, 4.8$ Hz, 1 H), 8.06 (dd, $J = 1.8, 7.7$ Hz, 1 H), 7.87 (dd, $J = 1.8, 8.1$ Hz, 1 H), 7.29 (dd, $J = 4.8, 8.1$ Hz, 1 H), 7.18 (dd, $J = 4.8, 7.7$ Hz, 1 H), 5.92 (m, 1 H), 5.19 (m, 1 H), 5.02 (m, 1 H), 4.76 (br m, 1 H), 3.44 (s, 3 H); MS (CI) m/z 267 (MH⁺). Anal. (C₁₅H₁₄N₄O) C, H, N.

HIV-1 Reverse Transcriptase Enzyme Assay. Recombinant HIV-1 reverse transcriptase was obtained from T. Steitz (Yale University). This enzyme was homogeneous (>95%) p66/p51 heterodimer. Reverse transcriptase was assayed by a modification of the previously described procedure.^{23a} The reaction mixture consisted of 50 mM Tris (pH 7.8), 50 mM glutamic acid, 1 mM DTT, 2 mM MgCl₂, 0.02% CHAPS, 0.8 μ g/mL poly(rC):oligo(dG) (Pharmacia), and 500 nM [³H]dGTP (NEN Du Pont), adjusted to a final volume of 60 μ L. Inhibitors were assayed at a concentration of 10 μ g/mL in the presence of RT (0.5 nM). The inhibitors, which were dissolved in DMSO and diluted many-fold with buffer, were added as solutions, and reaction was initiated by the addition of enzyme. After 1 h at room temperature, 50 μ L each of ice-cold 10% aqueous trichloroacetic acid and 2% aqueous sodium pyrophosphate was added and the mixture was cooled at 4 °C for 15 min. Acid insoluble products were harvested onto #30 glass fiber filters (Schleicher and Schuell) using a Skatron cell harvester. Dried filters were counted in an LKB 1205 Betaplate liquid scintillation counter. Inhibition was determined by comparison of the amount of product in reactions run with and without test compound. IC₅₀s were determined by comparison of the amounts of product produced in the presence and absence of test compound. Plots of per cent inhibition versus log ([compound]) yielded the concentration at which half maximal inhibition was observed.

Acknowledgment. We wish to express our sincere appreciation to T. Phil Pitner for assistance in NMR analyses and to Roger Dinallo, Keith McKellop, Walter Davidson, and Gordon Hansen for mass spectral determinations. Susan Hattox, Paul Riska, and Helga Silverstein performed metabolism studies, and Nita Pandit and Gerry Bell provided solubility data. Günter Trummelitz

(Dr. Karl Thomae GmbH) kindly provided compounds 65 and 68.

Registry No. 1, 10189-78-3; 2, 132686-74-9; 3, 132686-96-5; 4, 132686-97-6; 5, 16287-28-8; 6, 16287-29-9; 7, 133626-57-0; 8, 16287-49-3; 9-HCl, 133626-58-1; 10, 16287-50-6; 11, 132686-75-0; 12, 132686-83-0; 13, 132686-84-1; 14, 132686-94-3; 15, 132686-76-1; 16, 132686-78-3; 17, 132686-79-4; 18, 133626-59-2; 19, 132686-98-7; 20, 132686-99-8; 21, 132686-80-7; 22, 132686-91-0; 23, 132687-04-8; 24, 132687-05-9; 25, 133626-60-5; 26, 133626-61-6; 27, 133626-62-7; 28, 133626-63-8; 29, 133626-64-9; 30, 132687-03-7; 31, 133626-65-0; 32, 133626-66-1; 33, 133626-67-2; 34, 133626-68-3; 35, 133626-69-4; 36, 133626-70-7; 37, 133626-71-8; 38, 133626-72-9; 39, 133626-73-0; 40, 133626-74-1; 41, 133626-75-2; 42, 132687-06-0; 43, 132687-07-1; 44, 133626-76-3; 45, 132377-83-4; 46, 132687-01-5; 47, 133626-77-4; 48, 133626-78-5; 49, 133626-79-6; 50, 133626-80-9; 51, 133626-81-0; 52, 133626-82-1; 53, 132687-00-4; 54, 132686-92-1; 55, 133626-83-2; 56, 133626-84-3; 57, 133626-85-4; 58, 133626-86-5; 59, 132707-73-4; 60, 133626-87-6; 61, 133626-88-7; 62, 133626-89-8; 63, 133626-90-1; 64, 133626-91-2; 65, 133626-92-3; 66, 133626-93-4; 67, 133626-94-5; 68, 133626-95-6; 69, 132312-85-7; 70, 133626-96-7; 71, 133626-97-8; 72, 133626-98-9; 73, 133626-99-0; 74, 132312-81-3; 75, 133627-00-6; 76, 133627-01-7; 77, 133627-02-8; 78, 133627-03-9; 79, 133627-04-0; 80, 133627-05-1; 81, 133627-06-2; 82, 133627-07-3; 83, 133648-15-4; 84, 133648-16-5; 85, 133627-08-4; 86, 133627-09-5; 87, 133627-10-8; 88, 133627-11-9; 89, 133627-12-0; 90, 133627-13-1; 91, 133627-14-2; 92, 133627-15-3; 93, 133627-16-4; 94, 133627-17-5; 95, 133627-18-6; 96, 129618-40-2; 97, 133627-19-7; 98, 133627-20-0; 99, 133627-21-1; 100, 133627-22-2; 101-2HBr, 133627-23-3; 102, 133627-24-4; 103, 133627-25-5; 104, 133627-26-6; 105, 133627-27-7; 106, 133627-28-8; 107, 133627-29-9; 108, 133627-30-2; 109, 133627-31-3; 110, 133627-32-4; 111, 133627-33-5; 112, 133627-34-6; 113, 133627-35-7; 114, 885-70-1; 115, 24000-48-4; 116, 24000-52-0; 117, 24000-53-1; 118, 132932-27-5; 119, 132932-26-4; 120, 132932-29-7; 121, 114368-10-4; 122, 114368-13-7; 123, 133627-36-8; 124, 133627-37-9; II (R¹ = Me, R² = Et, R³ = 8-phthalimidomethyl, R⁴ = H), 133627-38-0; IV (R³ = Me), 2687-25-4; V (R⁴ = H), 2942-59-8; VII (R¹ = Me, R³ = 8,9-Cl₂, R⁴ = H), 133627-39-1; VII (R¹ = Et, R³ = 7-Me, R⁴ = H), 132687-02-6; VII (R¹ = Me, R³ = 7-Me, R⁴ = H), 133627-40-4; VII (R¹ = CH₂Ph, R³ = 7-Me, R⁴ = H), 133627-41-5; VII (R¹ = Me, R³ = 9-NO₂, R⁴ = H), 132687-24-2; VII (R¹ = Me, R³ = 8-NO₂, R⁴ = H), 133627-42-6; VIII (R², R⁴ = H, R³ = 8,9-Cl₂), 10321-11-6; IX (R¹ = Me, R² = Et, R³ = 7-Me, R⁴ = H), 133627-43-7; X (R¹ = Et, R³ = 7-Me, R⁴ = H), 133627-44-8; XIb (R = Et, R¹ = Me, R³ = NO₂, R⁴ = H), 133648-17-6; XII (R³ = 4-Me, X = Cl), 133627-45-9; XII (R³ = H, X = Cl), 6298-19-7; XIII (R⁴ = H), 49609-84-9; XIVa (R³, R⁴ = H), 132312-86-8; XIVa (R³ = 4-Me, R⁴ = H), 133627-46-0; XIVb (R¹ = Me, R³, R⁴ = H), 132312-87-9; XVa (R² = *c*-C₃H₅, R³ = 4-Me, R⁴ = H), 133627-47-1; XVII (R¹, R³, R⁴ = H), 132312-45-9; XVIIIa (R³, R⁴ = H), 132312-46-0; XVIIIb (R¹ = Me, R³, R⁴ = H), 132312-91-5; XIXb (R¹ = Me, R³, R⁴ = H), 132312-92-6; *c*-C₃H₅-NH₂, 765-30-0; MeO-*p*-C₆H₄-CH₂NH₂, 2393-23-9; *N*-(hydroxymethyl)phthalimide, 118-29-6; 2-chloro-4-methyl-3-nitropyridine, 23056-39-5; reverse transcriptase, 9068-38-6.