

* To whom correspondence should be addressed at the Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208-3113.

† Department of Chemistry, Northwestern University.

‡ The Institute for Neuroscience, Northwestern University.

§ Department of Pharmacology, Warner-Lambert.

¶ Department of Biochemistry, Molecular Biology, and Cell Biology.

Richard B. Silverman,^{*,†,‡,§,¶} Ryszard Andruszkiewicz[†]
Shrenik M. Nanavati,[†] Charles P. Taylor[‡]
Mark G. Vartanian[§]

Department of Chemistry
Department of Biochemistry, Molecular Biology, and Cell
Biology

The Institute for Neuroscience

Northwestern University

Evanston, Illinois 60208-3113

Department of Pharmacology

Parke-Davis Pharmaceutical Research Division


Warner-Lambert

Ann Arbor, Michigan 48105-2430

Received April 9, 1991


New Pepstatin Analogues: Synthesis and Pepsin Inhibition

Since the isolation of pepstatin, isovaleryl-L-valyl-L-valyl-(3S,4S)-statyl-L-alanyl-(3S,4S)-statine (1), by Umezawa,



- (1) Aoyagi, T.; Umezawa, H. *Proteinases and Biological Control*; Reich, E., Rifkin, D. B., Shaw, E., Ed.; Cold Spring Harbor Laboratory: Cold Spring Harbor, NY, 1975; pp 429-454.
- (2) Umezawa, H.; Aoyagi, T. *Proteinases in Mammalian Cells and Tissues*; Barrett, A. J., Ed.; North Holland: New York, 1977; pp 637-662.
- (3) Rich, D. H.; Bernatowicz, M. S.; Agarwal, N. S.; Kawai, M.; Salituro, F. G.; Schmidt, P. G. Inhibition of Aspartyl Proteinases by Pepstatin and 3-methylstatine Derivatives of Pepstatin. *Biochemistry* 1985, 24, 3165-3173.
- (4) Nisato, D.; Wagon, J.; Callet, G.; Mettefeu, D.; Assens, J. L.; Plouzane, C.; Tonnerre, B.; Pliska, V.; Fauchere, J. L. Free Wilson Correlation Analysis of Inhibitory Potency of a Series of Pepstatin Analogues on Plasma Renin. *J. Med. Chem.* 1987, 30, 2287-2291.
- (5) Guegan, R.; Diaz, J.; Cazaubon, C.; Beaumont, M.; Carlet, C.; Clement, J.; Demarne, H.; Mellet, M.; Richaud, J. P. Pepstatin as Novel Renin Inhibitors. *J. Med. Chem.* 1986, 29, 1152-1159.
- (6) Agarwal, N. S.; Rich, D. H. Inhibition of Cathepsin D by Substrate Analogues Containing Statine and by Analogues of Pepstatin. *J. Med. Chem.* 1986, 29, 2519-2524.
- (7) Salituro, F. G.; Agarwal, N.; Hofmann, T.; Rich, D. H. Inhibition of Aspartic Proteinases by Peptides Containing Lysine and Ornithine Side Chains in Analogues of Statine. *J. Med. Chem.* 1987, 30, 286-295.
- (8) Cumin, F.; Nisato, D.; Gagnol, J. P.; Corvol, P. A Potent Radiolabeled Human Renin Inhibitor—Enzymatic, Kinetic, and Binding Studies to Renin and Other Aspartic Proteinases. *Biochemistry* 1987, 26, 7615-7621.
- (9) Nisato, D.; Wagnon, J.; Callet, G.; Mettefeu, D.; Annens, J. L. A QSAR Study of New Pentapeptides as Renin Inhibitors Containing the Fragment Statine-Ala-Statine. *Pharmacochimistry* 1987, 10, 227-284.
- (10) Gunn, J. M.; Owens, R. A.; Liu, W. S.; Glover, G. I. Biological Activity of Aspartic Proteinase Inhibitors Related to Pepstatin. *Acta Biol. Med. Ger.* 1981, 40, 1547-1553.

Scheme 1^a



^a (a) Diisobutylaluminum hydride, toluene, -78 °C; (b) ethyl acetate is pretreated with lithium diisopropylamide in THF at -78 °C for 30 min followed by addition of aldehyde in THF, 2 h at -78 °C.

wa,^{1,2} investigators have pursued the development and biochemical study of proteinase inhibitors similar in structure to pepstatin.³⁻¹¹ The recent discovery of the aspartyl proteinase of the human immunodeficiency virus 1, (HIV-1), and its inhibition by pepstatin, has increased the search for synthetic analogues of pepstatin with increased substrate specificities.¹²⁻¹⁶ However, the therapeutic value of pepstatin is limited.

For a proteinase inhibitor to have therapeutic value, it should be chemically stable, active at low concentrations, selective for a particular proteinase in the presence of other proteinases with similar substrate specificities, and should readily penetrate cell membranes. Pepstatin satisfies the first two of these criteria. The third criterion, specificity, is satisfied only at the level of the major mechanistic class. Pepstatin, while a very effective inhibitor of aspartyl proteinases, is not very selective within this class of enzymes. Pepstatin is also known not to penetrate well cell membranes due to its size and lipophilicity.

The exact mode of inhibition of many aspartyl proteinases by pepstatin has not been well documented. However, crystal structure studies of the HIV-1 aspartyl proteinase inhibited by pepstatin show that the statine residue occupies both the P₁ and P₁' sites, acting as a dipeptide in the bound enzyme.¹⁷ Several studies have also been conducted concerning the nature of inhibition of other

- (11) Huddy, S.; Patel, G.; Heywood, G. C.; Austen, B. M.; Hermon-Taylor, J. Synthesis of Novel Semi-Synthetic Tight Binding Inhibitors of Pepsin. *Biochem. Soc. Trans.* 1989, 17, 1124-1125.
- (12) Meek, T. D.; Lambert, D. M.; Dreyer, G. B.; Carr, T. J.; Tomaszek, T. A.; Moore, M. L.; Strickler, J. E.; Debouck, C.; Hyland, L. J. Inhibition of HIV-1 protease in Infected T-lymphocytes by Synthetic Analogs. *Nature* 1990, 343, 90-92.
- (13) Tamassell, A. G.; Olsen, M. K.; Hui, J. O.; Staples, D. J.; Sawyer, T. K.; Heinrichson, R. L.; Tomich, C. S. Substrate Analog Inhibition and Active Site Titration of Purified HIV-1 Protease. *Biochemistry* 1990, 29, 264-269.
- (14) Kostka, V.; de Gruyter, B. In vitro Mutation of the Asp-25 Blocks and Pepstatin A Inhibits the Activity of Human Immunodeficiency Virus (HIV) Encoded Protease. *Proteases Retroviruses, Proc. Colloq. C. 52, 14th Int. Cong. Biochem.* 1988 1989, 103-109.
- (15) Grinde, B.; Hungnes, O.; Tjoetta, E. The Protease Inhibitor Pepstatin A Inhibits Formulation of Reverse Transcriptase in H9 Cells Infected with Human Immunodeficiency Virus 1. *AIDS Res. Hum. Retroviruses* 1989, 5, 269-274.
- (16) Krausslich, H. G.; Ingraham, R. H.; Skoog, M. T.; Wimmer, E.; Pallai, P. V.; Carter, C. A. Activity of Purified Biosynthetic Proteinase of Human Immunodeficiency Virus on Natural Substrates and Synthetic Peptides. *Proc. Natl. Acad. Sci. U.S.A.* 1989, 86, 807-811.
- (17) Fitzgerald, P. (Merck Sharp and Dohme, Rahway, NJ) Presentation: The Human Immunodeficiency Virus Protease. Three Dimensional Structure of the Inhibited and Uninhibited Enzyme. 31st Medicinal Chemistry Symposium, 1990, Buffalo, NY.

Table I. Physical Constants for Tripeptide Analogues

no.	compound	mp, °C	$[\alpha]_D^{25}$, deg (c 1.0, CH ₃ OH)	molecular formula	anal.
5a	Cbz-Val-Val-(S,S)-AHMHA ^a	201–203	–70.3	C ₂₆ H ₄₁ N ₃ O ₇	C,H,N
5b	Cbz-Val-Val-(R,S)-AHMHA	216–218	–49.5	C ₂₆ H ₄₁ N ₃ O ₇	C,H,N
6a	Cbz-Val-Val-(S,S)-AHPPA ^b	168–170	–37.9	C ₂₈ H ₃₉ N ₃ O ₇	C,H,N
6b	Cbz-Val-Val-(R,S)-AHPPA	189–191	–27.5	C ₂₈ H ₃₉ N ₃ O ₇	C,H,N
7a	Cbz-Val-Val-(S,S)-AHHPPA ^c	197–200	–73.5	C ₂₈ H ₃₉ N ₃ O ₈	C,H,N
7b	Cbz-Val-Val-(R,S)-AHHPPA	217–219	–52.3	C ₂₈ H ₃₉ N ₃ O ₈	C,H,N
8a	Cbz-Val-Val-(S,S)-AHOA ^d	218–220	–36.2	C ₂₆ H ₄₁ N ₃ O ₇	C,H,N
8b	Cbz-Val-Val-(R,S)-AHOA	232–235	–30.3	C ₂₆ H ₄₁ N ₃ O ₇	C,H,N

^aAHMHA = 4-amino-3-hydroxy-6-methylheptanoic acid. ^bAHPPA = 4-amino-3-hydroxy-5-phenylpentanoic acid. ^cAHHPPA = 4-amino-3-hydroxy-5-*p*-hydroxyphenylpentanoic acid. ^dAHOA = 4-amino-3-hydroxyoctanoic acid.

aspartyl proteinases by pepstatin and its fragments. The novel amino acid statine, (3*S*,4*S*)-4-amino-3-hydroxy-6-methylheptanoic acid (AHMHA), appears to be the major structural component responsible for the inhibitory activity of pepstatin.¹⁸ It has also been suggested that the shape of the side chain of the statine unit greatly influences the inhibitory activity. For example, Gunn et al. found peptide analogues containing a statine unit derived from L-valine to be much less active pepsin inhibitors than similar analogues containing statine units derived from L-leucine.¹⁰

In this communication we now report the preparation and determination of pepsin inhibition of eight tripeptide analogues of pepstatin. Since pepstatin is known not to penetrate cell membranes as readily as small peptides, such as leupeptin, we have synthesized tripeptide analogues of pepstatin. This was done in an effort to find a small peptide analogue with inhibitory activity comparable to that of pepstatin.

Synthesis. The preparation of the statine derivatives 4 is outlined in Scheme I. The *tert*-butyloxycarbonyl (BOC) protected amino acid methyl esters 2 were reduced directly to the aldehydes 3 with diisobutylaluminum hydride in toluene for 10 min at –78 °C. The reactions were then worked up with methylene chloride and Rochelle salt to isolate the aldehydes 3 in 65 to 90% yield. The aldehydes were then condensed with the lithium enolate of ethyl acetate at –78 °C to give the ethyl esters as mixtures of diastereomers in 70–85% yields. The lithium enolate of ethyl acetate was generated from lithium diisopropylamide (Aldrich) and ethyl acetate as a modification of Steulmann and Klostermeyer's method.¹⁹ The diastereomers were separated via high-performance liquid chromatography. Normal-phase silica gel columns were used in conjunction with SSI Model 232D gradient liquid chromatograph utilizing both UV and refractive index detection. The stereochemistry was assigned by ¹H NMR with use of the chemical shifts and coupling constants of the third and fourth carbons of the 2-oxazolidinone derivatives in comparison to similar known data.^{13,20–22}

Table II. Inhibition of Pepsin Hydrolysis of Hemoglobin by Pepstatin and Analogues

no.	compound	IC ₅₀ , μM	K _i (app), μM
1	pepstatin	0.007	4.60 × 10 ^{–11}
5a	Cbz-Val-Val-(3 <i>S</i> ,4 <i>S</i>)-AHMHA ^a	0.20	1.90 × 10 ^{–7}
5b	Cbz-Val-Val-(3 <i>R</i> ,4 <i>S</i>)-AHMHA	39.4	3.94 × 10 ^{–6}
6a	Cbz-Val-Val-(3 <i>S</i> ,4 <i>S</i>)-AHPPA ^b	0.42	4.10 × 10 ^{–7}
6b	Cbz-Val-Val-(3 <i>R</i> ,4 <i>S</i>)-AHPPA	346	3.46 × 10 ^{–4}
7a	Cbz-Val-Val-(3 <i>S</i> ,4 <i>S</i>)-AHHPPA ^c	1130	1.13 × 10 ^{–3}
7b	Cbz-Val-Val-(3 <i>R</i> ,4 <i>S</i>)-AHHPPA	1540	1.54 × 10 ^{–3}
8a	Cbz-Val-Val-(3 <i>S</i> ,4 <i>S</i>)-AHOA ^d	0.010	3.10 × 10 ^{–9}
8b	Cbz-Val-Val-(3 <i>R</i> ,4 <i>S</i>)-AHOA	17.7	1.77 × 10 ^{–6}

^aAHMHA = 4-amino-3-hydroxy-6-methylheptanoic acid.

^bAHPPA = 4-amino-3-hydroxy-5-phenylpentanoic acid.

^cAHHPPA = 4-amino-3-hydroxy-5-(*p*-hydroxyphenyl)pentanoic acid.

^dAHOA = 4-amino-3-hydroxyoctanoic acid.

Removal of the *tert*-butoxycarbonyl protecting groups was achieved with 30% trifluoroacetic acid in chloroform. The trifluoroacetate salts of the statine ethyl ester derivatives were then condensed with the mixed anhydride of carbobenzoxy-L-valyl-L-valine to give the Cbz-protected tripeptide ethyl esters in 40–70% yield. Saponification of the ethyl esters then provided the Cbz-protected tripeptide analogues (5–8) of pepstatin. The *N*-carbobenzyloxy protecting group was used on the pepstatin analogues rather than the *N*-isovaleryl protecting group because Cbz-L-valyl-L-valyl-(3*S*,4*S*)-statine has been reported to be a slightly better inhibitor of pepsin than *N*-isovaleryl-L-valyl-L-valyl-(3*S*,4*S*)-statine.²³ Physical constants for 5–8 are found in Table I.

Pepsin Assay. The potency of the synthetic pepstatin analogues was measured as inhibitors of the peptic hydrolysis of hemoglobin (Sigma), and the results are presented in Table II. Inhibition of porcine pepsin (Sigma) was measured by the following method: 0.9 mL of the inhibitor of appropriate concentration in 0.02 M KCl–HCl buffer (pH 2.0) and 1.0 mL of 0.5% hemoglobin solution were mixed and incubated for 3 min at 37 °C. Porcine pepsin, 100 μL of a 10 μg/mL solution, was added and mixed. The incubator was continued for 27 min at 37 °C. After this, 2.0 mL of 1.5 M perchloric acid was added to quench the reaction. The mixtures were allowed to warm to room temperature for 1 h. The precipitated protein was centrifuged and the absorbance measured at 280 nm. The absorbance of a blank containing no enzyme was subtracted from each reading. The inhibition of the enzyme activity was measured four times at five or more inhibitor concentrations. The average absorbance at each inhibitor concentration was utilized in calculations of the IC₅₀ values. All absorbance values were ≤0.002 standard deviations from the mean for a given inhibitor concentration; thus the relative error is minimal. The standard error for the linear regression plots was calculated and is in each case

- (18) Rich, D. H.; Sun, E. T. O.; Singh, J. Synthesis of Dideoxy-Pepstatin—Mechanism of Inhibition of Porcine Pepsin. *Biochem. Biophys. Res. Commun.* 1977, 74, 762–767.
- (19) Steulmann, R.; Klostermeyer, H. Synthesis of (3*S*,4*S*)-4-Amino-3-hydroxy-6-methylheptanoic Acid and Some of its Derivatives. *Justus Liebig's Ann. Chem.* 1975, 2245–2250.
- (20) Rich, D. H.; Sun, E. T. O.; Boparai, A. S. Synthesis of (3*S*,4*S*)-4-Amino-3-hydroxy-6-methylheptanoic Acid Derivatives. Analysis of Diastereomeric Purity. *J. Org. Chem.* 1978, 43, 3624–3626.
- (21) Morishima, H.; Takita, T.; Umezawa, H. Synthesis of 4-Amino-3-hydroxy-6-methylheptanoic Acid—Amine Component of Pepstatin. *J. Antibiot.* 1973, 26, 115–116.
- (22) Rich, D. H.; Sun, E. T. O.; Ulm, E. Synthesis of Analogues of the Carboxyl Protease Inhibitor Pepstatin. Effect of Structure of Inhibition of Pepsin and Renin. *J. Med. Chem.* 1980, 23, 27–33.

- (23) Aoyagi, T.; Morishima, H.; Nishazawa, R.; Kunimoto, S.; Takeuchi, T.; Umezawa, H.; Ikzawa, H. Biological Activity of Pepstatins, Pepstanone A, and Partial Peptides on Pepsin, Cathepsin B, and Renin. *J. Antibiot.* 1972, 25, 689–694.

less than 5%. A plot of percent inhibition versus the log of the inhibitor concentration provided a value for 50% inhibition concentration (IC_{50}). All plots are linear through the 50% inhibition value and have slopes ranging from 15 to 41. The apparent inhibition constants (K_1 (app)) were calculated as K_1 (app) = $IC_{50} - 0.5[E]_0$, where $[E]_0$ is the enzyme concentration.

Results and Conclusions

The eight Cbz-protected tripeptide analogues were compared to pepstatin as inhibitors of the pepsin hydrolysis of hemoglobin (Table II). On comparing the activities of Cbz-Val-Val-AHMHA with that of pepstatin, it was found that the length of the peptide chain alters the inhibitor activity. This agrees with data reported in the literature.^{10,22,24} Also, as expected, the stereochemistry of carbon-3, bearing the hydroxyl group, is very important to the activity of the inhibitors. The 3*S*,4*S* diastereomers was in each case shown to be a better inhibitor of the peptic hydrolysis of hemoglobin than the 3*R*,4*S* diastereomer.

Modifications made in the side chain of the novel amino acid statine, however, had the greatest effect on the pepsin inhibition. Analogues containing a statine unit derived from L-phenylalanine (6a,b) were significantly less active as pepsin inhibitors, and very little inhibition of pepsin activity is shown by the analogues containing a statine unit derived from L-tyrosine (7a,b). This is indicative of steric

restrictions within the enzyme pepsin. Interestingly, it was also found that the analogues containing a statine unit derived from L-norleucine (8a,b) displayed a better inhibition of pepsin activity than those analogues containing statine units derived from L-leucine.

In view of the preliminary success achieved in finding a small peptide with potency similar to pepstatin for pepsin inhibition, further modifications to improve hydrophilicity of aspartyl proteinase inhibitors are warranted. Further synthesis and testing of tripeptide analogues of pepstatin is currently underway.

Acknowledgment. We thank the National Institutes of Health MBRS and MARC programs for their support on this project.

Registry No. 2 (R = *i*-Bu), 63096-02-6; 2 (R = CH₂Ph), 51987-73-6; 2 (R = CH₂C₆H₄-*p*-OH), 4326-36-7; 2 (R = Bu), 87974-77-4; 3 (R = *i*-Bu), 58521-45-2; 3 (R = CH₂Ph), 72155-45-4; 3 (R = CH₂C₆H₄-*p*-OH), 134081-15-5; 3 (R = Bu), 104062-70-6; (S,S)-4 (R = *i*-Bu), 67010-43-9; (R,S)-4 (R = *i*-Bu), 67010-44-0; (S,S)-4 (R = CH₂Ph), 72155-46-5; (R,S)-4 (R = CH₂Ph), 72155-47-6; (S,S)-4 (R = CH₂C₆H₄-*p*-OH), 134081-16-6; (R,S)-4 (R = CH₂C₆H₄-*p*-OH), 134081-17-7; (S,S)-4 (R = Bu), 134081-18-8; (R,S)-4 (R = Bu), 134081-19-9; 5a, 69980-30-9; 5b, 69980-31-0; 6a, 134081-20-2; 6b, 134175-09-0; 7a, 134081-21-3; 7b, 134175-10-3; 8a, 134081-22-4; 8b, 134175-11-4; AcOEt, 141-28-6; Cbz-Val-Val-OH, 19542-54-2; pepsin, 9001-75-6.

Rose M. McConnell,* Donna Frizzell, Amanda Camp
Anissa Evans, Wanda Jones, Cathy Cagle

Department of Chemistry
University of Arkansas at Pine Bluff
Pine Bluff, Arkansas 71601

Received February 25, 1991

(24) Liu, W. S.; Smith, S. C.; Glover, G. I. Synthesis of All the Stereoisomers of Statine (4-Amino-3-hydroxy-6-methylheptanoic Acid). Inhibition of Pepsin Activity by *N*-Carbo-benzoxy-L-valyl-L-valyl-statine Derived from the Four Stereoisomers. *J. Med. Chem.* 1979, 22, 577-579.