

of an equal volume of water gave an off-white precipitate which was collected by filtration. Crystallization from ethanol yielded 64 mg (72%) of 2.

Enzyme Isolation and Assay. The thymidine kinases from HSV1- and HSV2-infected HeLa cells were isolated by the use of a thymidine 3'-(p-aminophenylphosphate) affinity column as described previously.⁵ The enzymes were assayed with limiting concentrations of [³H]thymidine as described.² Stock solutions of inhibitors in dimethyl sulfoxide were diluted into assay mixtures; control assays contained an identical concentration of the

compound solvent.

Calculations. Derivation of equations and statistical analyses were done on a IBM PC using a BASIC program written by one of the authors (J.G.).

Acknowledgment. This work was partially supported by Contract AI72644 and Grant GM21747, from the National Institutes of Health (G.W.), and by grants from Progetto AIDS-ISS and CNR Progetto finalizzato Chimica Fine (S.S.).

DNA-Directed Alkylating Agents. 5. Acridinecarboxamide Derivatives of (1,2-Diaminoethane)dichloroplatinum(II)


Ho H. Lee,[†] Brian D. Palmer,[†] Bruce C. Baguley,[†] Michael Chin,[†] W. David McFadyen,[†] Geoffrey Wickham,[‡] Deborah Thorsbourne-Palmer,[§] Laurence P. G. Wakelin,^{||} and William A. Denny^{*,†}

Cancer Research Laboratory, University of Auckland School of Medicine, Private Bag, Auckland, New Zealand, School of Science and Mathematics Education, Institute of Education, University of Melbourne, Parkville, Australia 3052, and Department of Medicinal Chemistry, Victorian College of Pharmacy, 381 Royal Parade, Parkville, Australia 3052.

Received May 21, 1991

A series of acridine-2- and -4-carboxamide-linked analogues of PtenCl₂ has been prepared and evaluated for biological activity against several tumor cell lines in vitro and in vivo. The platinum complexes were generally more cytotoxic than the corresponding ligands against wild-type P388 leukemia cells in vitro, with acridine-4-carboxamide complexes being the more effective. In contrast to cisplatin and PtenCl₂, the complexes were equally active in vitro against both wild-type and cisplatin-resistant P388 lines. The 4-carboxamide complexes showed high levels of in vivo activity (ILS >100%) against wild-type P388 using a single-dose protocol, and one compound was also significantly active in vivo in a cisplatin-resistant line, against which cisplatin and PtenCl₂ are inactive.


One focus of work on developing analogues of cisplatin (*cis*-diamminedichloroplatinum(II)) is to improve activity against cell lines which are (for a variety of reasons) resistant to cisplatin itself. Novel examples are the DACH compounds (e.g. 1-OHP; 1)¹ and the bis-cisplatin deriva-



tives (e.g. 2).² Members of these classes show almost equal cytotoxicity in cisplatin-sensitive and -resistant lines, probably (in the case of the DACH compounds) by providing DNA adducts of enhanced lethality.³

Another way of providing platinum-DNA adducts of high cytotoxicity is to "target" the platinum moiety to DNA by attachment to a suitable carrier ligand. This general concept has been discussed in detail,⁴⁻⁶ and demonstrable effects with targeted aniline mustards include altered regiospecificity and sequence-specificity of DNA

Scheme I^a



^a (i) 1,1'-Carbonyldiimidazole/DMF/50 °C; NH₂(CH₂)_nOH; (ii) MeCl/py; excess NH₂(CH₂)₂NH₂; (iii) aqueous Na₂CO₃; K₂PtCl₄/20 °C/20 h; aqueous KCl.

alkylation,^{6,7} together with increased cytotoxicity and improved in vivo antitumor activity compared with the

[†]Cancer Research Laboratory.

[‡]University of Melbourne.

[§]Victorian College of Pharmacy.


^{||}Present address: St. Luke's Cancer Research Fund, Highfield Rd., Rathgar, Dublin 6, Ireland.

(1) Mathe, G.; Kidani, Y.; Noji, M.; Moral, C.; Bourut, C.; Chenu, E. Antitumor activity of 1-OHP in mice. *Cancer Lett.* 1985, 27, 135-141.

(2) Farrell, N.; Qu, Y.; Hacker, M. P. Cytotoxicity and antitumor activity of bis(platinum) complexes. A novel class of platinum complexes active in cell lines resistant to both cisplatin and 1,2-diaminocyclohexane complexes. *J. Med. Chem.* 1990, 33, 2179-2184.

(3) Page, J. D.; Husain, I.; Sancar, A.; Chaney, S. G. Effect of the diaminocyclohexane carrier ligand on platinum adduct formation, repair and lethality. *Biochemistry* 1990, 29, 1016-1024.

(4) Denny, W. A. DNA-Intercalating agents as antitumor drugs: prospects for future design. *Anti-Cancer Drug Des.* 1989, 4, 241-263.


Table I. Physicochemical and Biological Properties of (Acridinecarboxamide)-Pt(II) Complexes and Ligands


no.	form	R	side chain	n	mp (°C)	formula	analyses	growth inhibitor:			P388/W in vivo		P388/CP in vivo	
								IC ₅₀ (μM) ^a			OD ^b	ILS ^c	OD ^b	ILS ^c
								P388/W	P388/AMSA	P33/CP				
5	A	H	2	3	235–237	C ₁₉ H ₂₂ N ₄ O·3HCl	C,H,N,Cl	3.8	–	4.9	100	NA ^d		
6	A	H	2	6	244–246	C ₂₂ H ₂₈ N ₄ O·3HCl·1/2H ₂ O	C,H,N,Cl	3.6	–	3.8	100	NA		
7	A	H	4	3	234–236	C ₁₉ H ₂₂ N ₄ O·3HClH ₂ O	C,H,N	0.74	1.6	0.59	30	64		
8	A	H	4	6	198–200	C ₂₂ H ₂₈ N ₄ O·3HCl·1/2H ₂ O	C,H,N,Cl	0.51	1.1	0.35	20	23		
9	A	NH ₂	4	3	282–284	C ₁₉ H ₂₃ N ₅ O·3HCl	C,H,N,Cl	3.1	0.87	2.6	45	45		
10	A	NH ₂	4	6	252 dec	C ₂₂ H ₂₉ N ₅ O·3HCl·8H ₂ O	C,H,N,Cl	5.2	–	2.3	65	NA		
11	B	H	2	3	261–263	C ₁₉ H ₂₂ Cl ₂ N ₄ OPt·H ₂ O	C,H,N,Pt	5.9	–	9.1	65	32		
12	B	H	2	6	266–269	C ₂₂ H ₂₈ Cl ₂ N ₄ OPt·H ₂ O	C,H,N,Pt	0.3	–	0.4	45	NA		
13	B	H	4	3	240 dec	C ₁₉ H ₂₂ Cl ₂ N ₄ OPt	C,H,N,Pt	0.06	0.43	0.4	45	103	45	25
14	B	H	4	6	206 dec	C ₂₂ H ₂₈ Cl ₂ N ₄ OPt	C,H,N	0.05	0.50	0.04	30	105	30	24
15	B	NH ₂	4	3	291 dec	C ₁₉ H ₂₃ N ₅ Cl ₂ OPt·3H ₂ O	C,H,N	0.56	0.58	1.1	20	86	20	47
16	B	NH ₂	4	6	283 dec	C ₂₂ H ₂₉ Cl ₂ N ₅ OPt·HCl·2H ₂ O	C,H,N,Cl	0.67	–	0.70	150	38		
PtCl ₂								0.30	–	1.03	13.3	70	13.3	NA
cisplatin								1.2	5.5	5.3	5.3	120	5.5	NA

^aIC₅₀: the concentration of drug (μM) to inhibit cell growth in the various cell lines by 50%, measured in 96-well cultures as described in ref 23. All measurements were done in triplicate, with a typical standard error of 10%. ^bOD: optimal dose of drug in milligrams/kilogram, administered intraperitoneally as a solution in 0.1 mL of 30% v/v ethanol/water (ligands) or 0.1 mL of 1:1:2 DMA/glycerol/water (Pt complexes), on days 1, 5, and 9 after intraperitoneal inoculation of 10⁶ P388/W leukemia cells. ^cILS: percentage increase in life span of treated animals (when treated at the optimal dose), compared with tumor-bearing control animals. Values are the mean of at least two determinations. Average life span of control animals was 11 days. ^dCompound inactive (ILS < 20) at all dose levels.

corresponding untargeted alkylators.⁵ Similar targeting of platinum species might also serve to minimize exposure to inactivating agents such as thiols, desirable since overexpression of thiols and thiol-transferring enzymes is a recognized mechanism of cellular resistance to cisplatin.^{8,9}

Examples of putative DNA-targeted platinum drugs which show improved selectivity with respect to cisplatin-resistant cell lines include the doxorubicin analogue 3^{10,11} and the 9-anilinoacridine compound 4.^{12,13} Although

Scheme II^a

^a(i) MsCl/py/20 °C/1 h; excess NH₂(CH₂)₂NH₂; (ii) (*t*-BuO)₂CO; (iii) H₂/Pd/C; (iv) phenol/NH₃/120 °C; (v) concentrated HCl/MeOH/CH₂Cl₂; (vi) aqueous Na₂CO₃; K₂PtCl₄/20 °C/20 h; aqueous KCl.

these compounds did show improved differential toxicity between wild-type and platinum-resistant cell lines, the differentials were no greater than those shown by the unplatinated ligands themselves, and it was not possible to decide the role of the 9-anilinoacridine carrier moiety. The complexes showed only minimal in vivo activity, possibly because they were very insoluble.¹² In the present paper we report studies on a further series of platinum compounds (11–16) linked to a more water-soluble acridinecarboxamide DNA-targeting carrier.

- Gourdie, T. A.; Valu, K. K.; Gravatt, G. L.; Boritzki, T. J.; Baguley, B. C.; Wakelin, L. P. G.; Wilson, W. R.; Woodgate, P. D.; Denny, W. A. DNA-directed alkylating agents. 1. Structure-activity relationships for acridine-linked aniline mustards: consequences of varying the reactivity of the mustard. *J. Med. Chem.* 1990, 33, 1177–1186.
- Kohn, K. W.; Hartley, J. A.; Mattes, W. B. Mechanisms of DNA sequence-selective alkylation of guanine N7 positions by nitrogen mustards. *Nucleic Acids Res.* 1987, 15, 10531–10549.
- Prakash, A. S.; Denny, W. A.; Gourdie, T. A.; Valu, K. K.; Woodgate, P. D.; Wakelin, L. P. G. DNA-directed alkylating ligands as potential antitumor agents: sequence specificity of alkylation by DNA-intercalating acridine-linked aniline mustards. *Biochemistry* 1990, 29, 9799–9807.
- Andrews, P. A.; Murphy, M. P.; Howell, S. B. Differential potentiation of alkylating and platinating agent cytotoxicity in human ovarian carcinoma cells by glutathione depletion. *Cancer Res.* 1985, 45, 6250–6253.
- Fram, R. J.; Woda, B. A.; Wilson, J. M.; Robichaud, N. Characterization of acquired resistance to *cis*-diamminedichloroplatinum(II) in BE human colon carcinoma cells. *Cancer Res.* 1990, 50, 72–77.
- Zunino, F.; Savi, G.; Pasini, A. Synthesis and antitumor activity of a platinum(II)-doxorubicin complex. *Cancer Chemother. Pharmacol.* 1986, 18, 180–182.
- Pasini, A. A doxorubicin-Pt(II) complex with antitumor activity. Synthesis, characterization and reactions with guanosine and DNA. *Gazz. Chim. Ital.* 1987, 117, 763–768.
- Palmer, B. D.; Lee, H. H.; Johnson, P.; Baguley, B. C.; Wickham, G.; Wakelin, L. P. G.; McFadyen, W. D.; Denny, W. A. DNA-directed alkylating agents. 2. Synthesis and biological activity of platinum complexes linked to 9-anilinoacridine. *J. Med. Chem.* 1990, 33, 3008–3014.

- Palmer, B. D.; Wickham, G.; McFadyen, W. D.; Baguley, B. C.; Denny, W. A. Synthesis, DNA binding interactions and biological activity of bis platinum(II) complexes of N,N,N',N'-tetrakis(2-aminomethyl)diamines. *Anti-Cancer Drug Des.*, in press.

Chemistry

Synthesis of the acridinecarboxamide ligands required for complexes 11–14 (Table I) was straightforward, and is shown in Scheme I. The acridine-2- and -4-carboxylic acids 17 and 20 were coupled selectively with 3-amino-propanol via the imidazoles,¹⁴ and the resulting alcohols 18 and 21 were activated with methanesulfonyl chloride and treated with excess 1,2-diaminoethane to give the desired diamines 5 and 7. Similar reactions using 6-aminohexanol provided the diamines 6 and 8 (Table I).

However, synthesis of the 9-aminoacridinecarboxamide ligands 15 and 16 required prior construction of the side chain, as shown in Scheme II. 3-Aminopropanol was N-protected with benzyloxycarbonyl groups, and the alcohol group of the resulting compound (23) was activated with methanesulfonyl chloride and reacted with excess 1,2-diaminoethane. The newly-formed amine groups of the triamine derivative 24 was then protected with Boc groups to give the differentially-protected triamine derivative 25. Selective deblocking of the N-Cbz group gave the unstable amine 26, which was reacted selectively with 9-chloroacridine-4-carbonyl chloride (27)¹⁵ to give the 9-chloro compound 28. This was converted in situ to the 9-phenoxy compound and then treated with dry ammonia at 120 °C to give the 9-aminoacridine 9, the precursor ligand for the desired platinum complexes (15). An identical sequence beginning with 6-aminohexanol gave the ligand 10, precursor of the platinum complex 16 (Table I).

The ligands 5–10 reacted readily with K_2PtCl_4 in the dark at 25 °C to form the corresponding platinum complexes 11–16. The coordination state of the platinum in the complexes was confirmed by ¹⁹⁵Pt NMR spectroscopy, with all complexes exhibiting a single signal in the range δ -2338 to -2344 ppm. This is similar to the ¹⁹⁵Pt resonance positions in the 9-anilinoacridine Pt complexes reported earlier (e.g., 4: δ -2344 ppm) and in PtenCl₂ itself (δ -2312 ppm).¹² For the 9-aminoacridinecarboxamide complexes 15 and 16 this distinguishes between alternative coordination modes via the 9-amino group, since the ¹⁹⁵Pt resonances of such compounds occur at lower fields (e.g. for *cis*-[Pt(NH₃)₂(N⁹-9-aminoacridine)Cl](NO₃), δ -2206 ppm).¹⁶

Results and Discussion

Since previous studies^{12,13} have shown that 9-anilinoacridine-linked platinum complexes (e.g., 4) are very insoluble, the compounds studied here use more water-soluble carriers. Compounds 11 and 12 place the complex off the long axis of an acridinecarboxamide intercalator (position 2), while the corresponding compounds 13 and 14 have this appended off the short axis (position 4). Compounds 15 and 16 use the more tightly DNA-binding and even more water-soluble 9-aminoacridinecarboxamide chromophore.

DNA Binding. The interaction of selected complexes with DNA has been studied in several ways. The kinetics

of the alkylation of DNA by 12 and 14 were similar to those of PtenCl₂ itself ($t_{1/2}$ = 5.5, 3.7, and 2.1 h, respectively).¹³ However, they unwound closed circular supercoiled DNA as efficiently as 9-aminoacridine, and formed interstrand DNA cross-links more effectively than PtenCl₂, as determined using nondenaturing agarose gels and linearized pBR 322 plasmid DNA¹³ (12–14 achieve complete cross-linking at a drug/base-pair ratio of 0.013, compared with a ratio of 0.038 for PtenCl₂). The sequence specificity of adduct formation of selected compounds has been studied using the *taq* DNA polymerase linear amplification method,¹⁷ and shown to be subtly different from that for PtenCl₂.

In Vitro Cytotoxicity. IC₅₀s for continuous exposure (72 h) were measured in 96-well cultures as described previously,¹⁸ and are given in Table I. The cell lines used were the drug-sensitive wild-type P388 (P388/W), a subline (P388/AMSA) resistant to the topoisomerase II agent amsacrine by its expression of a topoisomerase II isozyme,¹⁹ and a subline (P388/CP) which shows ca. 5-fold resistance to cisplatin (Table I). The unplatinated acridine-2-carboxamide carrier ligands 5 and 6 had similar IC₅₀ values (3–5 μ M) in both the wild-type and CP-resistant lines, while the isomeric 4-carboxamide ligands 7 and 8 were considerably more cytotoxic (0.4–0.7 μ M). Selected ligands also evaluated against P388/AMSA (Table I) were found to be up to 10-fold less effective, ratios typical for DNA topoisomerase II inhibitors,²⁰ and suggest that the cytotoxicity of the ligands is mediated, at least to some extent, by inhibition of topoisomerase II. With the exception of compound 11, the platinum complexes 11–16 were considerably more cytotoxic (on average 10-fold) than the corresponding ligands, but their comparative activity in the different cell lines was similar. Compounds 13 and 14 showed significantly higher cytotoxicity than PtenCl₂ itself against P388/W, but all the complexes were equally active against the wild-type and CP-resistant lines.

In Vivo Activity. The 2-carboxamide ligands 5 and 6 were not active against P388/W in vivo, but the 4-carboxamide ligands 7–10 did show moderate activity, with the C3 analogues being more active than the C6 analogues in each case. This agrees with the known SAR for acridinecarboxamides.²¹ While the complexes showed no dramatic increase in potency compared with the ligands, they did show higher activity. The preferred side-chain orientation was clearly at the 4-position, with the nature of the chromophore (9-H or 9-amino) being less critical.

- (14) Staab, H. A. Syntheses using heterocyclic amides (azolides). *Angew. Chem., Intl. Ed. Engl.* 1962, 1, 351–367.
- (15) Atwell, G. J.; Cain, B. F.; Baguley, B. C.; Finlay, G. J.; Denny, W. A. Potential Antitumor Agents. 43. Synthesis and biological activity of dibasic 9-aminoacridine-4-carboxamides: A new class of antitumor agent. *J. Med. Chem.* 1984, 27, 1481–1485.
- (16) Sundquist, W. I.; Bancroft, D. P.; Lippard, S. J. Synthesis, characterization and biological activity of *cis*-diammine-platinum(II) complexes of the CDNA intercalators 9-aminoacridine and chloroquine. *J. Am. Chem. Soc.* 1990, 112, 1590–1596.

- (17) Murray, V.; Motyka, H.; England, P. R.; Wickham, G.; Lee, H. H.; Denny, W. A.; McFadyen, W. D. An investigation of the sequence specific interaction of cisplatin and four analogues (including two acridine-tethered complexes) with DNA inside cells. Submitted to *Biochemistry*.
- (18) Finlay, G. J.; Baguley, B. C.; Wilson, W. R. Comparison of the in vitro activity of cytotoxic drugs towards human carcinoma and leukemia cell lines. *Eur. J. Cancer Clin. Oncol.* 1986, 22, 655–662.
- (19) Chung, T. D. Y.; Drake, F. H.; Tan, K. B.; Per, M.; Crooke, S. T.; Mirabelli, C. K. Characterization and immunological identification of cDNA clones encoding two human DNA topoisomerase isozymes. *Proc. Natl. Acad. Sci. U.S.A.* 1989, 86, 9431–9435.
- (20) Schneider, E.; Darkin, S. J.; Lawson, P. A.; Ching, L.-M.; Ralph, R. K.; Baguley, B. C. Cell line selectivity and DNA breakage properties of the antitumor agent N-[2-(dimethylamino)ethyl]acridine-4-carboxamide: Role of DNA topoisomerase II. *Eur. J. Cancer Clin. Oncol.* 1988, 24, 1783–1790.
- (21) Atwell, G. J.; Rewcastle, G. W.; Baguley, B. C.; Denny, W. A. Potential Antitumor Agents. 50. In vivo solid tumor activity of derivatives of N-[2-(dimethylamino)ethyl]acridine-4-carboxamide. *J. Med. Chem.* 1987, 30, 664–669.

The 4-carboxamide complexes 13 and 14 showed low in vivo activity (ILS 25%) against P388/CP (both cisplatin and PtenCl₂ are inactive), but the 9-amino derivative 15 was considerably more effective, with an ILS of 47%.

Conclusions

Previous work^{17,22} has shown that the attachment of platinum moieties (specifically PtenCl₂) to DNA-intercalating ligands does alter the nature of the interaction of the metal complex with DNA, although there is little effect on the rates of covalent adduct formation. The acridine-carboxamide complexes evaluated here showed little increase in either in vitro cytotoxicity or in vivo activity against P388/W cells compared with PtenCl₂, but did have improved activity in the P388/CP platinum-resistant line. One compound, the 9-aminoacridine-4-carboxamide complex 15, also showed significant in vivo activity against the cisplatin-resistant line.

Experimental Section

Where elemental analyses are indicated by the symbols of the elements, results were within $\pm 0.4\%$ of theoretical values. Melting points were determined on an Electrothermal apparatus using the supplied stem-corrected thermometer and are as read. ¹H NMR spectra were measured on a Bruker AM-400 spectrometer. ¹⁹⁵Pt NMR spectra were determined at 86.0 MHz for solutions (ca. 1.5×10^{-2} M) in DMF, with an external D₂O lock. Spectra were collected by using a 10- μ s pulse width and acquisition times from 0.123 to 0.4 s, with spectral widths from 11 to 33 kHz. Shifts are referenced to Na₂PtCl₆ in H₂O as 0 ppm. Platinum determinations were carried out on a Perkin-Elmer 3030 atomic absorption spectrometer.

Acridine Ligands: Preparation of *N*-[3-[(2-Aminoethyl)amino]propyl]acridine-4-carboxamide (7). Example of General Method of Scheme I. Acridine-4-carboxylic acid (20) (2.39 g, 10.1 mmol) was suspended in dry DMF (20 mL), 1,1'-carbonyldiimidazole (2.5 g, 15 mmol) was added, and the mixture was warmed to 50 °C until all solids dissolved and gas evolution ceased. The mixture was cooled to 20 °C, 3-amino-1-propanol (2 g, 30 mmol) was added, and the solution was kept at 20 °C for 45 min. Water was then added, and the solvents were removed under reduced pressure. The residue was partitioned between ice-cold 1 N Na₂CO₃ and EtOAc, and the organic layer was washed with water, dried, and evaporated to give crude *N*-(3-hydroxypropyl)acridine-4-carboxamide (21) (2.75 g, 98%), pure by TLC and suitable for direct use. A sample was recrystallized from EtOAc/petroleum ether as needles: mp 120–122 °C; ¹H NMR (CDCl₃) δ 9.05 (br s, 1 H, ArH-3), 8.93 (s, 1 H, ArH-9), 8.25–7.60 (m, 6 H, acridine protons), 3.86 (t, J = 5.8 Hz, 2 H, CH₂OH), 3.79 (t, J = 5.8 Hz, 2 H, NHCH₂), 1.95 (quintet, J_{app} = 5.8 Hz, CH₂CH₂CH₂). Anal. (C₁₇H₁₈N₂O₂) C, H, N.

The above crude alcohol 21 (2.7 g, 9.6 mmol) was dissolved in dry pyridine (50 mL) and treated with methanesulfonyl chloride (0.85 mL, 11 mmol) at 20 °C for 30 min. Water (1 mL) was then added, followed after 5 min by ethane-1,2-diamine (25 mL). The mixture was stirred for 20 h, and volatiles were removed under reduced pressure. The residue was partitioned between CH₂Cl₂ and dilute Na₂CO₃, and the organic layer was washed with water, dried, and evaporated. The residue was dissolved in 1 N HCl and extracted with CH₂Cl₂, and the aqueous layer was then evaporated to dryness under reduced pressure to give a yellow solid. Crystallization from aqueous MeOH gave the trihydrochloride salt of *N*-[3-[(2-aminoethyl)amino]propyl]acridine-4-carboxamide (7)

(2.9 g, 70% yield): mp 234–236 °C dec; ¹H NMR (CD₃SOCD₃) δ 9.5–7.75 (m, 8 H, acridine protons), 3.78 (q, J = 6 Hz [converted to t after D₂O exch], 2 H, CONHCH₂), 3.28 (br s, 4 H, CH₂NHCH₂), 3.15 (br s, 2 H, CH₂NH₂), 2.15 (m, 2 H, CH₂CH₂CH₂). Anal. in Table I.

Similar treatment of acridine-4-carboxylic acid (20) with 6-amino-1-hexanol gave *N*-(6-hydroxyhexyl)acridine-4-carboxamide (22): mp (EtOAc/CH₂Cl₂) 112–113 °C; ¹H NMR (CDCl₃) δ 9.00 (d, J = 5.4 Hz, 1 H, ArH-3), 8.90 (s, 1 H, ArH-9), 8.20–7.55 (m, 6 H, acridine protons), 3.71 (t, J = 6.9 Hz, 2 H, CH₂OH), 3.65 (t, J = 6.4 Hz, 2 H, NHCH₂), 1.9–1.5 (m, 10 H, [8 H after D₂O exchange] $-(\text{CH}_2)_4-$, NH and OH). Anal. (C₂₀H₂₂N₂O₂) C, H, N. Reaction of this with methanesulfonyl chloride followed by ethane-1,2-diamine as above gave the trihydrochloride of triamine 8: mp 198–200 °C (dec); ¹H NMR (CD₃SOCD₃) δ 9.5–7.7 (m, 8 H, acridine H), 3.60 (q, J = 6.1 Hz, 2 H, became t after D₂O exch, CONHCH₂), 3.23 (br s, 4 H, CH₂NHCH₂), 2.97 (br s, 2 H, CH₂NH₂), 1.8–1.4 (m, 8 H, $-(\text{CH}_2)_4-$). Anal. in Table I.

Similar treatment of acridine-2-carboxylic acid (17) with 3-amino-1-propanol gave *N*-(3-hydroxypropyl)acridine-2-carboxamide (18): mp (EtOAc) 150–152 °C; ¹H NMR (CD₃SOCD₃) δ 9.26 (s, 1 H, ArH-1), 8.68 (s, 1 H, ArH-9), 8.3–7.7 (m, 6 H, acridine protons), 3.56 (t, J = 6.6 Hz, 2 H, CH₂OH), 3.42 (t, J = 6.6 Hz, 2 H, NHCH₂), 1.80 (quintet, J_{app} = 6.6 Hz, 2 H, CH₂CH₂CH₂). Anal. (C₁₇H₁₈N₂O₂) C, H, N. This in turn yielded the trihydrochloride of triamine (5): mp 235–237 °C dec; ¹H NMR (CD₃SOCD₃) δ 9.82 (s, 1 H, H-9), 9.00 (s, 1 H, H-1), 8.60–7.85 (m, 6 H, acridine H), 3.50 (q, J = 6.7 Hz, became t after D₂O exch, CONHCH₂), 3.25 (br s, 4 H, became 2 \times t, J = 5.4 and 5.4 Hz after D₂O exch, NHCH₂CH₂NH₂), 3.10 (br s, 2 H, became a t, J = 7.4 Hz after D₂O exch, CH₂NHCH₂), 2.02 (m, 2 H, CH₂CH₂CH₂). Anal. in Table I.

Similar treatment of acridine-2-carboxylic acid (17) with 6-amino-1-hexanol gave *N*-(6-hydroxyhexyl)acridine-2-carboxamide (19) [mp (EtOAc/MeOH) 150.5–151.5 °C; ¹H NMR (CD₃SOCD₃) δ 9.25 (s, 1 H, ArH-1), 8.70 (s, 1 H, ArH-9), 8.4–7.6 (m, 6 H, acridine protons), 3.40 (m, 4 H, NHCH₂ and CH₂OH), 1.7–1.2 (m, 8 H, $-(\text{CH}_2)_4-$). Anal. (C₂₀H₂₂N₂O₂) C, H, N.], which in turn gave the trihydrochloride of triamine (6) [mp (aqueous MeOH) 244–246 °C (dec); ¹H NMR (CD₃SOCD₃) δ 9.83 (s, 1 H, H-1), 8.93 (s, 1 H, H-9), 8.55–7.85 (m, 6 H, acridine protons), 3.37 (q, J = 6.5 Hz, 2 H, became t on D₂O exch, CONHCH₂), 3.20 (br s, 4 H, CH₂NHCH₂), 1.75–1.35 (m, 8 H, $-(\text{CH}_2)_4-$). Anal. in Table I.]

9-Aminoacridine Ligands: Preparation of *N*-[3-[(2-aminoethyl)amino]propyl]-9-aminoacridine-4-carboxamide Trihydrochloride (9). Example of the General Method. Benzyl chloroformate (22.9 g, 19.1 mL, 0.134 mol) was added dropwise to a stirred homogeneous mixture of 3-amino-1-propanol (10.0 g, 0.134 mol) and NaHCO₃ (16.9 g, 0.201 mol) in H₂O (250 mL) at 20 °C, and the reaction was stirred for a further 1.5 h at 20 °C. Extraction with CH₂Cl₂ and usual workup gave 3-[(benzyloxycarbonyl)amino]-1-propanol (23a) (23.9 g, 86%) as a colorless solid: mp (EtOAc/petroleum ether) 49–50 °C; ¹H NMR (CDCl₃) δ 7.36 (m, 5 H, C₆H₅), 5.12 (s, 2 H, C₆H₅CH₂), 5.06 (br s, 1 H, exch with D₂O, NH), 3.67 (q, J = 6.0 Hz, 2 H, collapsing to t after D₂O, CH₂OH), 3.36 (q, J = 6.0 Hz, 2 H, collapsing to t after D₂O, NHCH₂), 2.58 (t, J = 6.0 Hz, exch with D₂O, OH), 1.71 (quintet, J = 6.0 Hz, 2 H, CH₂CH₂CH₂). Anal. (C₁₇H₁₈N₂O₃) C, H, N.

A solution of the above alcohol 23a (21.7 g, 0.104 mol) in pyridine (90 mL) was treated at 15–20 °C with methanesulfonyl chloride (12.5 mL, 1.65 mol) for 1 h. The solution was then cooled to 0 °C and water (2.4 mL) was added, followed by 1,2-diaminoethane (90 mL). After stirring at 0 °C for 2 h and then at 20 °C for 20 h, most of the solvents were removed under reduced pressure at ca. 40 °C, and the residue was partitioned between CH₂Cl₂ and aqueous 0.5 N Na₂CO₃. The organic extract was worked up as usual to give the crude triamine 24a (22.5 g, 86%) as a dark green oil, which was used immediately: ¹H NMR (CDCl₃/D₂O) δ 7.30 (m, 5 H, C₆H₅), 5.10 (s, 2 H, C₆H₅CH₂), 3.27 (t, J = 6.4 Hz, 2 H, CONHCH₂), 2.75 (t, J = 5.8 Hz, 2 H, NHCH₂CH₂NH₂), 2.66 (t, J = 6.4 Hz, 2 H, CH₂CH₂CH₂NHCH₂), 2.62 (t, J = 5.8 Hz, 2 H, CH₂NH₂), 1.66 (quintet, J = 6.4 Hz, 2 H, CH₂CH₂CH₂).

A solution of di-*tert*-butyl dicarbonate (24.4 g, 0.112 mol) was added dropwise to a solution of triamine 24a (25.5 g, 0.102 mol)

- (22) Cullinane, C.; Wickham, G.; McFadyen, W. D.; Phillips, D. R.; Palmer, B. D.; Denny, W. A. Detection of platinum-DNA intra-strand and inter-strand crosslinks by in vitro transcription: Effect of pendant intercalators. *Nucleic Acids Res.*, submitted.
- (23) Finlay, G. J.; Baguley, B. C.; Wilson, W. R. Comparison of the in vitro activity of cytotoxic drugs towards human carcinoma and leukemia cell lines. *Eur. J. Cancer Clin. Oncol.* 1986, 22, 655–662.

in dry CH_2Cl_2 (200 mL) over 70 min, and the mixture was stirred for a further 24 h at 20 °C. The mixture was then partitioned between CH_2Cl_2 and aqueous 0.5 N Na_2CO_3 , and the organic layer was worked up as usual. The resulting solid was chromatographed on SiO_2 , elution with EtOAc/petroleum ether (1:5) affording pure N,N' -bis(*tert*-butoxycarbonyl)- N -[3'-[(benzoxycarbonyl)amino]propyl]ethane-1,2-diamine (**25a**) (16.0 g, 35%) as a pale yellow solid: mp (EtOAc/petroleum ether) 62–64 °C; $^1\text{H NMR}$ (CDCl_3) δ 7.36 (m, 5 H, C_6H_5), 5.10 (s, 2 H, $\text{C}_6\text{H}_5\text{CH}_2$), 3.24 (m, 8 H, $\text{CH}_2\text{CH}_2\text{CH}_2$ and CH_2CH_2), 1.73 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.46 (s, 9 H, 3 \times CH_3), 1.42 (s, 9 H, 3 \times CH_3). Anal. ($\text{C}_{23}\text{H}_{37}\text{N}_3\text{O}_6$) C, H, N.

The above protected triamine **25a** (5.00 g, 11.1 mmol) was dissolved in MeOH/EtOAc (1:1) and hydrogenolyzed over 5% Pd/C to give the crude amine **26a**, which was immediately dissolved in a mixture of dry CH_2Cl_2 (70 mL) and Et_3N (15 mL) and added dropwise to solid 9-chloroacridine-4-carboxyl chloride (**27**) (3.94 g, 14.3 mmol, freshly prepared from 9-oxoacridin-4-carboxylic acid).¹⁵ The reaction was stirred at 20 °C for 2 h and was then diluted with CH_2Cl_2 (200 mL) and washed with aqueous 0.5 N Na_2CO_3 (200 mL). The CH_2Cl_2 layer was separated and worked up as usual to give a residue which was azeotroped with dry toluene (2 \times 50 mL) to give the crude 9-chloro amide **28a** as a brown oil. This was dissolved in dry benzene (60 mL), phenol (24 g) was added, and the mixture was heated in an open flask until the internal temperature reached 90 °C. It was then cooled to 40 °C, and anhydrous NH_3 gas was bubbled through the mixture, which was brought to 120 °C and held there for 10 min. The cooled mixture was then partitioned between CH_2Cl_2 and 2 N NaOH, and the organic layer was worked up as usual. The residue was chromatographed on SiO_2 and eluted with EtOAc/MeOH (10:1) to afford N -[3'-[N,N' -bis(*tert*-butoxycarbonyl)ethylenediamino]propyl]-9-aminoacridine-4-carboxamide (**29a**) (3.84 g, 64%) as a bright yellow solid: mp (CH_2Cl_2 /MeOH) 207–209 °C; $^1\text{H NMR}$ (CDCl_3) δ 12.60 (br s, 1 H, exch with D_2O , NH), 9.00–7.02 (m, 7 H, aromatic protons), 6.00 (br s, 3 H, exch with D_2O , NH), 3.67 (q, J = 6.0 Hz, 2 H, collapsing to t after D_2O , $\text{NHCH}_2\text{CH}_2\text{CH}_2$), 3.58–3.24 (m, 6 H, $\text{CH}_2\text{NCH}_2\text{CH}_2\text{NH}$), 2.04 (quintet, J = 6.0 Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 1.54 and 1.43 (2 s, 18 H, 6 \times CH_3). Anal. ($\text{C}_{29}\text{H}_{39}\text{N}_5\text{O}_6$) C, H, N.

This compound (3.67 g, 6.83 mmol) was dissolved in a mixture of CH_2Cl_2 /MeOH (10:1, 110 mL) and concentrated HCl (5 mL) and the reaction was stirred vigorously at 20 °C for 30 min. Solvents were removed under reduced pressure at 45 °C, and the residue was taken up in 1 N HCl and washed with CH_2Cl_2 (3 \times 30 mL). Evaporation of the aqueous layer left a bright yellow solid, which was recrystallized from CH_2Cl_2 /MeOH to afford the desired N -[3'-[(2-aminoethyl)amino]propyl]-9-aminoacridine-4-carboxamide trihydrochloride (**9**) (2.81 g, 92%): mp 282–284 °C dec; $^1\text{H NMR}$ (CD_3SOCD_3) δ 13.80 (br s, 1 H, exch with D_2O , HCl), 10.45 (br s, 2 H, exch with D_2O , 2 HCl), 9.70 (br s, 3 H, exch with D_2O , NH), 9.10–7.60 (m, 7 H, aromatic protons), 8.45 (br s, 3 H, exch with D_2O , NH), 3.52 (q, J = 7.0 Hz, 2 H, collapsing to t after D_2O , $\text{CONHCH}_2\text{CH}_2\text{CH}_2$), 3.23 (br s, $W_{1/2}$ = 5 Hz, 4 H, $\text{NHCH}_2\text{CH}_2\text{NH}_2$), 3.09 (t, J = 7.0 Hz, 2 H, $\text{CONHCH}_2\text{CH}_2\text{CH}_2$), 2.05 (quintet, J = 7.0 Hz, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_2$). Anal. in Table I.

Similar treatment of 6-aminohexanol and collection of the precipitated product by filtration gave 6-[(benzoyloxycarbonyl)amino]-1-hexanol (**23b**) in 60% yield: mp (EtOAc/petroleum ether) 79–81 °C; $^1\text{H NMR}$ (CDCl_3) δ 7.36 (m, 5 H, C_6H_5), 5.13 (br s, 1 H, exchangeable with D_2O , NH), 4.77 (br s, 1 H, exch with D_2O , OH), 3.63 (q, J = 6.4 Hz, 2 H, collapsing to t after D_2O , CH_2OH), 3.20 (q, J = 6.6 Hz, 2 H, collapsing to t after D_2O , NHCH_2), 1.62–1.32 (m, 8 H, 4 \times CH_2). Anal. ($\text{C}_{14}\text{H}_{21}\text{NO}_3$) C, H, N.

This was then treated as above to give the protected triamine **24b**, which was reacted with di-*tert*-butyl dicarbonate as above to afford N,N' -bis(*tert*-butoxycarbonyl)- N -[6-[(benzoyloxycarbonyl)amino]hexyl]ethane-1,2-diamine (**25b**) in 91% yield as a pale yellow viscous oil: $^1\text{H NMR}$ (CDCl_3) δ 7.36 (m, 5 H, C_6H_5), 5.10 (s, 2 H, $\text{C}_6\text{H}_5\text{CH}_2$), 3.22 (m, 8 H, $\text{CH}_2(\text{CH}_2)\text{CH}_2$; CH_2CH_2),

1.56–1.24 (m, 8 H, $\text{CH}_2(\text{CH}_2)_4\text{CH}_2$), 1.46 and 1.44 (2 s, 18 H, 6 \times CH_3).

The fully-protected triamine **11** was then selectively deprotected and reacted with **27** as above via the intermediates **26b** and **28b** to give N -[6-[[N,N' -bis(*tert*-butoxycarbonyl)amino]hexyl]-9-aminoacridine-4-carboxamide (**29b**) in an overall yield of 41%: mp (CH_2Cl_2 /MeOH) 198–200 °C; $^1\text{H NMR}$ (CDCl_3) δ 8.94–7.50 (m, 7 H, aromatic protons), 3.65 (br q, J = 6 Hz, 2 H, collapsing to t after D_2O , $\text{CONHCH}_2(\text{CH}_2)_5$), 3.23 (m, 6 H, $\text{CONH}(\text{CH}_2)_5\text{CH}_2\text{NCH}_2\text{CH}_2\text{NH}$), 1.86–1.20 (m, 8 H, $\text{CH}_2(\text{CH}_2)_4\text{CH}_2$), 1.44 (s, 18 H, 6 \times CH_3). Anal. ($\text{C}_{32}\text{H}_{45}\text{N}_5\text{O}_5 \cdot 0.5\text{H}_2\text{O}$) C, H, N.

Finally, deprotection of **29b** as above gave a 96% yield of N -[6-[(2-aminoethyl)amino]hexyl]-9-aminoacridine-4-carboxamide trihydrochloride (**10**), after recrystallization from MeOH/ CH_2Cl_2 : mp 251 °C dec; $^1\text{H NMR}$ (CD_3SOCD_3) δ 13.90 (s, 1 H, exch with D_2O , HCl), 10.50 (s, 1 H, exch with D_2O , HCl), 10.40 (s, 1 H, exch with D_2O , HCl), 9.55 (s, 2 H, exch with D_2O , NH), 9.45 (s, 1 H, exch with D_2O , NH), 9.05–7.60 (m, 7 H, aromatic protons), 8.40 (s, 3 H, exch with D_2O , NH), 3.44 (t, J = 7 Hz, 2 H, CONHCH_2), 3.21 (m, 4 H, $\text{NHCH}_2\text{CH}_2\text{NH}_2$), 3.00 (t, J = 7.6 Hz, 2 H, $\text{CH}_2\text{NH}(\text{CH}_2)_2\text{NH}_2$), 1.66 (m, 4 H, $\text{NHCH}_2\text{CH}_2(\text{CH}_2)_2\text{CH}_2\text{CH}_2\text{NH}$), 1.43 (m, 4 H, $\text{NH}(\text{CH}_2)_2(\text{CH}_2)_2(\text{CH}_2)_2\text{NH}$). Anal. in Table I.

Typical Platination Procedure: Synthesis of 12. The trihydrochloride **6** (48 mg, 0.1 mmol) was dissolved in water (2 mL), and 2 N aqueous Na_2CO_3 was added until the solution was just basic. A solution of K_2PtCl_4 (42 mg, 0.1 mmol) in water (1 mL) was added, and the mixture was stirred at 20 °C in the dark for 25 h. A solution of 5% aqueous KCl (10 mL) was then added, and the mixture was stirred for a further 90 min. The resulting pale yellow precipitate was collected, washed several times with deionized water, and dried to give pure N -[6-[(2-aminoethyl)amino]hexyl]acridine-2-carboxamide[dichloroplatinum(II)] (**12**) (54 mg, 86% yield): mp 266–269 °C; $^{195}\text{Pt NMR}$ (DMF) δ –2339. Anal. ($\text{C}_{22}\text{H}_{29}\text{Cl}_2\text{N}_4\text{O}_2$) in Table I.

Similar reactions gave **11** [73%; mp 261–263 °C dec; $^{195}\text{Pt NMR}$ (DMF) δ –2344. Anal. in Table I.], **13** [72%; mp 240–243 °C dec; $^{195}\text{Pt NMR}$ (DMF) δ –2339. Anal. in Table I.], and **14** [88%; mp 206 °C dec; $^{195}\text{Pt NMR}$ (DMF) δ –2339. Anal. in Table I.].

A similar procedure carried out using **9**, but with the pH of the ligand solution adjusted to 8–9 before mixing, afforded N -[3'-[(2-aminoethyl)amino]propyl]-9-aminoacridine-4-carboxamide[dichloroplatinum(II)] (**15**) (90%): mp 291 °C dec; $^{195}\text{Pt NMR}$ (DMF) δ –2339. Anal. in Table I. Likewise, ligand **10** gave complex **16** (87%) as the monohydrochloride: mp 283 °C dec; $^{195}\text{Pt NMR}$ (DMF) δ –2338. Anal. in Table I.

The platinum complexes were formulated for biological testing by suspending the free bases in dimethylacetamide and adding an equal volume of glycerol to give a homogeneous solution. Water was then added to make up the required concentration.

Acknowledgment. We thank Dr. Maruta Boyd for NMR spectra and Wendy Hodgson for preparation of the manuscript. This work was supported by the Auckland Division of the Cancer Society of New Zealand, the Medical Research Council of New Zealand, and the Anti-Cancer Council of Victoria.

Registry No. **5**, 142038-87-7; **6**, 142038-88-8; **7**, 142038-89-9; **8**, 142038-90-2; **9**, 142038-91-3; **10**, 142038-92-4; **11**, 142039-08-5; **12**, 142039-09-6; **13**, 142039-10-9; **14**, 142039-11-0; **15**, 142039-12-1; **16**, 142039-13-2; **18**, 142038-93-5; **19**, 142038-94-6; **21**, 142038-95-7; **22**, 142038-96-8; **23a**, 34637-22-4; **23b**, 17996-12-2; **24a**, 142038-97-9; **24b**, 142038-98-0; **25a**, 142038-99-1; **25b**, 142039-00-7; **26a**, 142039-01-8; **26b**, 142039-02-9; **27**, 142039-03-0; **28a**, 142039-04-1; **28b**, 142039-05-2; **29a**, 142039-06-3; **29b**, 142039-07-4; K_2PtCl_4 , 10025-99-7; ^{195}Pt , 14191-88-9; 1,1'-carbonyldiimidazole, 530-62-1; 3-aminopropanol, 156-87-6; ethylenediamine, 107-15-3; 6-aminohexanol, 4048-33-3; acridine-2-carboxylic acid, 54328-73-3; acridine-4-carboxylic acid, 31327-97-6; benzyl chloroformate, 501-53-1.