

Structure-Activity Relationships of 8-Styrylxanthines as A₂-Selective Adenosine Antagonists^{†,§}

Kenneth A. Jacobson,* Carola Gallo-Rodriguez, Neli Melman, Bilha Fischer, Michel Maillard, Andrew van Bergen, Philip J. M. van Galen, and Yishai Karton[†]

Laboratory of Bioorganic Chemistry, National Institute of Diabetes, and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892

Received November 20, 1992

A series of substituted 8-styryl derivatives of 1,3,7-alkylxanthines was synthesized as potential A₂-selective adenosine receptor antagonists, and the potency at rat brain A₁- and A₂-receptors was studied in radioligand binding experiments. At the xanthine 7-position, only small hydrophobic substituents were tolerated in receptor binding. 7-Methyl analogues were roughly 1 order of magnitude more selective for A₂ versus A₁ receptors than the corresponding 7-H analogues. 1,3-Dimethylxanthine derivatives tended to be more selective for A₂-receptors than the corresponding 1,3-diallyl, diethyl, or dipropyl derivatives. Substitutions of the phenyl ring at the 3-(monosubstituted) and 3,5-(disubstituted) positions were favored. 1,3,7-Trimethyl-8-(3-chlorostyryl)xanthine was a moderately potent (K_i vs [³H]CGS 21680 was 54 nM) and highly A₂-selective (520-fold) adenosine antagonist. 1,3,7-Trimethyl-8-[3-[(3-carboxy-1-oxopropyl)amino]styryl]xanthine was highly A₂-selective (250-fold) and of enhanced water solubility (max 19 mM). 1,3-Dipropyl-7-methyl-8-(3,5-dimethoxystyryl)xanthine was a potent (K_i = 24 nM) and very A₂-selective (110-fold) adenosine antagonist.

Introduction

Adenosine acting through the A₂-receptor subtype elicits a range of physiological responses, including the inhibition of platelet aggregation¹ dilation of blood vessels,² erythropoietin production,³ and depression of locomotor activity.⁴ The second messenger system invariably associated with the A₂-receptor is the stimulation of adenylate cyclase, while the A₁-receptor has been found to be coupled to a number of second messenger systems, including inhibition of adenylate cyclase, inhibition of calcium entry, stimulation of potassium flux, and effects on phosphoinositide metabolism.⁵

All of the adenosine agonists yet reported are derivatives of adenosine, but many classes of adenosine antagonists are known.⁵ The earliest examples of adenosine antagonists are the naturally-occurring xanthines, e.g. caffeine and theophylline, which are nonselective for adenosine receptor subtypes and have phosphodiesterase inhibitory effects as well.⁶ While thousands of ligands selective for A₁ receptors have been synthesized, it is only recently that A₂-selective agonists and antagonists have been reported.^{2,7-11} Certain 2-substituted adenosine² and adenosine-5'-carboxamide derivatives⁸ are A₂-selective agonists. Several xanthine derivatives, most notably, simple caffeine derivatives bearing a 1-propargyl group (1, Figure 1), and those doubly substituted by 7-methyl and 8-cycloalkyl




Figure 1. Xanthine and non-xanthine derivative having either high affinity or some selectivity for A₂-adenosine receptors.

groups (2),^{10,11} have been reported to be A₂-selective in binding and/or adenylate cyclase experiments. A number of non-xanthine derivatives have been found to be weakly selective for A₂-receptors, e.g. the triazoloquinazoline CGS 15943 (3),¹² the triazoloquinoxaline CP66,713 (4),¹³ and a series of imidazotriazinones, including M216765 (5).¹⁴

It would be highly desirable to have a general, high affinity A₂-antagonist probe for receptor characterization. At this time, several 8-phenylxanthines, such as XAC, 6, may serve as radioligands at striatal A₂-receptors, but this high-affinity binding is limited to certain species in which XAC is not A₁-selective.¹⁵ Shimada et al.¹¹ have reported that 1,3,7-alkylxanthine derivatives substituted with 8-styryl groups, particularly 3,4-dimethoxy and 3,4,5-trimethoxy, act as selective A₂-antagonists. In this study

* Address correspondence to Dr. Kenneth A. Jacobson, Bldg 8A, Rm B1A-17, NIDDK/NIH, Bethesda, MD 20892. Tel.: (301) 496-9024. FAX: (301) 402-0008.

[†] Permanent address: Department of Organic Chemistry, Israel Institute of Biological Research, Nes Ziona, Israel.

[‡] Abbreviations: t-Boc, *tert*-butyloxycarbonyl; CGS 21680, 4-[2-[[6-amino-9-(*N*-ethyl- β -D-ribofuranuronamidoyl]-9H-purin-2-yl)amino]ethyl]benzenepropanoic acid, DMAP, 4-(*N,N*-dimethylamino)pyridine; DMF, *N,N*-dimethylformamide; DMSO, dimethyl sulfoxide; EDAC, 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride; NECA, *N*-ethyladenosin-5'-uronamide; PIA, (*R*)-*N*-(2-phenylisopropyl)adenosine; Tris, tris(hydroxymethyl)aminomethane.

[§] Dedicated to Prof. Murray Goodman on the occasion of his 65th birthday.




Figure 2. Synthesis of 8-styrylxanthine derivatives. Reagents: (a) EDAC, DMAP, imidazole; (b) NaOH, 80 °C; alkyl bromide, 50 °C.




Figure 3. Synthesis of 8-(4-hydroxy-3,5-dimethoxystyryl)-xanthines and derivatives. Reagents: (a) C_6H_5SNa , 160 °C; (b) $R'-Br$.

we have further explored the structure-activity relationships of 8-styrylxanthines at adenosine receptors.

Results

Chemistry. 8-Styrylxanthine derivatives, substituted at the 1-, 3-, and 7-xanthine positions and at the α - and phenyl positions of the styryl moiety were synthesized according to the routes shown in Figures 2 and 3. The structures of compounds 15–56 and the affinities in radioligand binding assays at rat brain A_1 - and A_2 -receptors are given in Table I. Small alkyl substituents at the 1- and 3-positions were identical and varied from methyl to propyl. Substituents at the 7-position varied from H to 2-phenylethyl. A number of related xanthines (not 8-styryl) were prepared for comparison (Table II). The physical characteristics of the xanthine derivatives are summarized in Table III.

A *trans*-cinnamic acid, 8 (Figure 2), was condensed with a 1,3-dialkyl-5,6-diaminouracil, 7. The resulting amide, 9, was cyclized under strongly basic conditions to give the xanthine derivative 10. The intermediate cinnamic acids were obtained from commercial sources or prepared as described in the Experimental Section. After formation of the 7-H xanthine derivative 10, methylation using methyl iodide at 50 °C was carried out. The *trans* orientation of the 8-styryl group was verified for each of the derivatives based on the proton-proton coupling constants of the olefinic protons (typically ≥ 15 Hz). In the case of compound 16 (α -fluoro), the fluorine-proton coupling constant was found to be 37 Hz, consistent with the *trans* configuration.¹⁶

Synthesis of hydroxyl ring-substituted 8-styrylxanthines was attempted by the usual route (Figure 2), starting with

the 3- or 4-hydroxycinnamic acid. The intermediate amide was formed in low yield, and cyclization provided the desired xanthine in only very low yield (e.g. 18). Carrying out the sequence with hydroxyl protection in order to obtain a free hydroxyl group in the para position of the final product was attempted, but proved unsatisfactory. Acety ester and *p*-methoxybenzyl ether derivatives formed the amide intermediate 9, but the cyclization step in 4 N NaOH failed. The attempted deprotection of mono-methoxy derivatives in the series using sodium thiophenolate, trimethylsilyl iodide, or nitrogen bases at high temperature was unsuccessful. It was however possible to selectively demethylate the *p*-methoxy of 8-(3,4,5-trimethoxy)styrylxanthines, 12 (Figure 3), using sodium thiophenolate in DMF at 160 °C. The position of the free hydroxyl group (4) in 13 was determined by proton NMR. This hydroxyl group could be readily acylated or alkylated (in some cases carried out in situ following the deprotection reaction) to provide 14.

Aryl amino substituents were obtained via $Zn/HOAc$ reduction of the corresponding nitro derivative (e.g. 21) or in the case of tertiary aniline (e.g. 30a), by direct incorporation of the corresponding cinnamic acid. The N-7 position of 30a was selectively alkylated using methyl iodide at 50 °C to provide 30b. Catalytic hydrogenation of the nitrostyryl derivative 21 afforded the saturated aniline analogue 57.

Biological Evaluation. Binding assay. The analogs were tested in a radioligand binding assay for affinity at adenosine receptors in rat brain membranes. The compounds were assayed for affinity at rat A_1 cortical receptors using [3H]- N^6 -(2-phenylisopropyl)adenosine¹⁷ and at rat striatal A_2 receptors using [3H]CGS 21680 (Tables I and II).⁸ K_i values of nearly 10^{-8} M at A_2 -receptors and selectivities of hundreds of fold were achieved.

Several sites on the xanthine molecule have been varied. The N-1 and N-3 positions have been substituted with small alkyl groups (C_1 – C_3). In all cases, substituents at the 1- and 3-positions are identical. The greatest effect of elongating NMe to NPr groups at these positions was a substantial increase in A_1 -affinity, thus diminishing A_2 -selectivity. A 1,3-diethyl-7-methylxanthine, 45b, was nearly as A_2 -selective (44-fold) as the 1,3-dimethyl analogue, 44b, which was 70-fold selective. The corresponding diallyl analogue, 46 (reported previously¹¹ to be >6700 A_2 -selective), was only 13-fold selective in rat brain in this study.

The N-7 position is either unsubstituted or substituted with groups as large as 2-phenylethyl (compound 44f). Only small, hydrophobic groups (including ethyl and propargyl) at this position were tolerated in binding to either receptor. The 7-methyl analogues were found to exhibit the greatest degree of A_2 -selectivity. Figures 4 and 5 show correlations of affinity for the 7-H to 7-Me modification, which generally results in decreased A_1 -affinity and increased A_2 -affinity. In general, among 8-styrylxanthine derivatives, the 1,3,7-trimethylxanthines were A_2 -selective by factors between 10- and 500-fold, whereas the corresponding 1,3-dimethylxanthines were generally A_2 -selective by factors of only 2–6-fold. The 7-(hydroxyethyl) and phenylethyl substituents (45d and 45f) were nearly inactive, in addition to having less favorable aqueous solubility. In the 1,3-dipropyl series (Figure 5), each 7-H analogue was relatively nonselective. The selectivity of the 1,3-dipropyl-7-methyl-8-styrylxan-

thines (resulting from decreased A₁-affinity upon methylation) was highly dependent on the styryl substitution.

The effects of substitution of the 8-styryl group could be compared within the 1,3-dimethyl series and within the 1,3,7-trimethyl series. The unsubstituted styryl analogue 15a (7-H) was nonselective, but was moderately selective (41-fold) following methylation (15b). Fluorine substitution in the α -position resulted in diminished potency at both A₁- (3-fold) and A₂-receptors (7-fold). Monomethoxy substitution of the phenyl ring (compounds 17, 19, and 29) resulted in selectivity of 18–63-fold in the 7-Me series, but did not result in significant A₂-selectivity in the 7-H series. Compound 19b, the meta derivative, was the most potent and selective monomethoxy derivative, with a K_i value of 85 nM at A₂-receptors. The analogue bearing a 3-hydroxystyryl group in the 7-H series, 18, was equipotent with the methoxy compound 19a at A₂-receptors and more potent at A₁-receptors.

The A₂-potency of 1,3,7-trimethylxanthines having a variety of styryl 3-position substituents varied in the order: acetylamino > chloro, amino > fluoro, methoxy > H > trifluoromethyl > nitro. Although the 3-chloro derivative (28, K_i value of 54 nM) was slightly less potent than the 3-acetylamino derivative (23, K_i value of 39 nM, 240-fold selective) it was more selective (520-fold). It was equipotent to the amino derivative 22b, but considerably more selective. Very bulky substituents at the 3-position (urethanes 25 and 26) reduced potency at A₂-receptors roughly 20-fold, but moderate A₂-selectivity remained. A water-solubilizing 3-[(3-carboxy-1-oxopropyl)amino] group (24) resulted in decreased potency (134 nM) but high selectivity (250-fold).

For comparison to the methoxy group at the styryl 4-position, a highly electron-donating group, e.g. dimethylamino, was incorporated and resulted in greatly diminished potency at both receptors. Only the 7-Me form, 30b, displayed A₂-selectivity.

Dimethoxy substitutions at various positions of the phenyl ring were compared, and substantial differences were observed. The order of both potency and selectivity was 3,5 > 3,4 > 2,3. In the 1,3,7-trimethyl series, 3,5-dimethoxy or 3,5-difluoro substituents (33b and 34b, respectively) resulted in >200-fold selectivity.

In the 1,3-dipropyl-7-methyl series, A₂-selectivity was generally merely 5–19-fold, with only one exception (53b). The 3-chlorostyryl analogue 51b, analogous to the most selective agent in the 1,3,7-trimethyl series, was only 14-fold selective. 1,3-Dipropyl-7-methyl-8-(3,5-dimethoxystyryl)xanthine, 53b, proved to be a potent (K_i vs [³H]CGS 21680 was 24 nM) and A₂-selective (110-fold) adenosine antagonist, i.e. 5-fold more selective than the corresponding 3,4-dimethoxy analogue, 52b. Compound 52b was prepared by Shimada et al. [KF17837]¹¹ and was reported to be 190-fold selective, versus 19-fold in this study.

High selectivities were also observed among 1,3,7-trimethylxanthines that were trisubstituted on the phenyl ring. 1,3,7-Trimethyl-8-(3,4,5-trimethoxy)styrylxanthine, 44b, was 70-fold A₂-selective in binding in the rat brain (vs >5600-fold reported by Shimada et al.¹¹). The corresponding 1,3-dimethyl analogue 44a was only 6-fold A₂-selective. In general, the order of both potency and selectivity for trisubstituted phenyl substituents was 3,4,5 > 2,3,4 > 2,4,5. Among 3,4,5-substituted analogues there was considerable substitution of the 4-methoxy group tolerated at A₂-receptors. The moderately selective 3,5-

dimethoxy-4-hydroxy analogue, 35, was acylated (36) and alkylated (37, 38), resulting in enhanced A₂-selectivity and potency. The 4-acetoxy-3,5-dimethoxy analogue, 36, was 93-fold A₂-selective. Functional groups that also tended to increase water solubility, such as alkylamines (38 and 40), were included. These amino derivatives may serve as functionalized congeners²⁰ since it appears that long chain extension is possible without disrupting receptor binding. Moderately potent and selective acylated derivatives (39, 41, and 42) were prepared from the amine functionalized congeners. Butyl versus *trans*-butenylamine congeners were compared to examine the effect of altering conformational flexibility at this distal site. No major differences in potency or selectivity between butyl (e.g. 38) and butenyl (e.g. 40) analogues were found.


In an attempt to account for the discrepancy in K_i values between the present study and Shimada et al.¹¹, the effects of varying concentrations of DMSO in the assay medium were examined. DMSO was needed because of the limited aqueous solubility (in the range of 10⁻⁵ M) of most of the 8-styrylxanthines tested. To avoid precipitation associated with serial aqueous dilutions, the only point at which xanthine dissolved in DMSO was added to aqueous medium was directly into the incubation tube. We examined the effects of varying concentrations of DMSO (ranging from 0.5 to 6%) on the apparent affinity of compound 53b (Figure 6A). The apparent affinity of compound 53b at A₂ receptors was constant within the range of 0.5–6% DMSO. In addition, the total specific binding of [³H]CGS 21680 to striatal membranes was maintained, even at 6% DMSO. However, A₁-affinity appeared to be somewhat dependent on DMSO concentration (at 0.5 and 1% DMSO), and at 6% DMSO the total specific binding of [³H]PIA (data not shown) diminished to roughly 30% of its value at 1% DMSO. At the lowest concentration (0.5% DMSO), higher concentrations of the drug were required to displace [³H]PIA. This effect of increase in the apparent K_i value at ≤1% DMSO most likely relates to the xanthine precipitating from the solution, since the UV absorption (Figure 6B) does not increase in a linear fashion with the amount of xanthine added to a fixed aqueous volume. The UV absorption decreases beyond 20 μM, suggesting supersaturation.

Related, non-styrylxanthines (Table II) were tested in adenosine receptor binding for comparison to the 8-styryl derivatives. Cyclohexylcaffeine, 2, which was found to be A₂-selective in effects on adenylate cyclase¹⁰ was nonselective in binding. The saturated aniline derivative 57 was ~300-fold less potent at A₂ receptors than the corresponding styryl derivative 22b. Ring-constrained styryl analogues, 58, containing a 8-(2-benzofuran) group were synthesized. Both the 7-H and 7-Me analogues were only weak antagonists of binding at adenosine receptors (Table II).

Discussion

Potent and A₂-selective adenosine antagonists, suitable as pharmacological tools, have long been lacking. A₂-selective antagonists may also have application as therapeutic agents, e.g. in the treatment of Parkinson's disease.²⁴ The nonselective non-xanthine antagonist CGS 15943, 3, was under development as an antiasthmatic.²⁵

It was only recently that 8-styrylxanthines were reported as the first potentially useful compounds by Shimada et


Table I. Affinities of 8-Styrylxanthine Derivatives in Radioligand Binding Assays at Rat Brain A₁- and A₂-Receptors^aR₁ = R₃ and R_α = H unless noted

compd	R ₁ , R ₃	R ₇	X	K _i ^a		A ₁ /A ₂ ratio
				A ₁	A ₂	
15a	Me	H	H	654 ± 170	291 ± 40	2.3
15b	Me	Me	H	3890 ± 1150	94 ± 36	41
16 ^c	Me	H	H	2190 ± 400	2110 ± 810	1.4
17a	Me	H	2-MeO	1730 ± 420	645 ± 144	2.7
17b	Me	Me	2-MeO	4760 ± 720	267 ± 84	18
18	Me	H	3-OH	702 ± 40	303 ± 55	2.4
19a	Me	H	3-MeO	1830 ± 821	378 ± 155	4.8
19b	Me	Me	3-MeO	5430 ± 1470	84.8 ± 24.0	64
20a	Me	H	3-CF ₃	881 ± 251	343 ± 58	2.6
20b	Me	Me	3-CF ₃	3330 ± 410	134 ± 44	25
21a	Me	H	3-NO ₂	1060 ± 150	438 ± 106	2.4
21b	Me	Me	3-NO ₂	2140 ± 480	195 ± 44	11
22a	Me	H	3-NH ₂	288 ± 60	202 ± 79	1.4
22b	Me	Me	3-NH ₂	1690 ± 360	57 ± 3	30
23	Me	Me	3-(AcNH)	9470 ± 2540	39 ± 21	240
24	Me	Me	3-(HOOC(CH ₂) ₂ CONH)	35100 ± 11700	143 ± 45	250
25	Me	Me	3-(<i>t</i> -BOC-NH)	23600 ± 2500	784 ± 100	30
26	Me	Me	3-[(<i>t</i> -BOC) ₂ -N]	10800 ± 1300	740 ± 77	15
27a	Me	H	3-F	2720 ± 360	516 ± 99	5.3
27b	Me	Me	3-F	15780 ± 2860	83 ± 18	190
28	Me	Me	3-Cl	28200 ± 7000	54 ± 19	520
29a	Me	H	4-MeO	858 ± 320	472 ± 132	1.8
29b	Me	Me	4-MeO	14200 ± 3500	327 ± 75	44
30a	Me	H	4-Me ₂ N	3030 ± 300	12800	0.24
30b	Me	Me	4-Me ₂ N	5.6% ^b (3 × 10 ⁻⁵)	9270 ± 150	>1
31a	Me	H	2,3-(MeO) ₂	1600 ± 250	600 ± 204	2.7
31b	Me	Me	2,3-(MeO) ₂	5390 ± 1020	716 ± 144	7.5
32a	Me	H	3,4-(MeO) ₂	5340 ± 1440	1100 ± 250	4.8
32b	Me	Me	3,4-(MeO) ₂	13790 ± 2420	197 ± 33	70
33a	Me	H	3,5-(MeO) ₂	3044 ± 520	120 ± 36	25
33b	Me	Me	3,5-(MeO) ₂	12.5 ± 6.3% ^b (10 ⁻⁵)	75.3 ± 29.1	>200
34a	Me	H	3,5-F ₂	2330 ± 830	366 ± 77	6.4
34b	Me	Me	3,5-F ₂	14750 ± 3890	65 ± 9	230
35	Me	Me	3,5-(MeO) ₂ -4-OH	8700 ± 4100	450 ± 66	19
36	Me	Me	4-AcO-3,5-(MeO) ₂	6330 ± 1680	68 ± 22	93
37	Me	Me	4-(4-PhCH ₂ O)-3,5-(MeO) ₂	4120 ± 460	139 ± 7	30
38	Me	Me	4-(4-NH ₂ -BuO)-3,5-(MeO) ₂	6170 ± 1010	173 ± 43	36
39	Me	Me	4-[4-(<i>t</i> -BOC-NH)BuO]-3,5-(MeO) ₂	11031	265 ± 105	42
40	Me	Me	4-(4-NH ₂ - <i>trans</i> -CH ₂ CH=CHCH ₂ O)-3,5-(MeO) ₂	6280 ± 1580	228 ± 20	28
41	Me	Me	4-(4-AcNH- <i>trans</i> -CH ₂ CH=CHCH ₂ O)-3,5-(MeO) ₂	17 ± 7% ^b (10 ⁻⁵)	216 ± 40	>50
42	Me	Me	4-[4-(<i>t</i> -BOC-NH)- <i>trans</i> -CH ₂ CH=CHCH ₂ O)-3,5-(MeO) ₂	11 ± 5% ^b (10 ⁻⁵)	353 ± 62	>40
43a	Me	H	2,3,4-(MeO) ₃	26 ± 10% ^b (10 ⁻⁵)	1610 ± 260	>5
43b	Me	Me	2,3,4-(MeO) ₃	6920 ± 330	206 ± 81	34
44a	Me	H	3,4,5-(MeO) ₃	2280 ± 530	360 ± 170	6.3
44b	Me	Me	3,4,5-(MeO) ₃	9200 ± 3560	131 ± 54	70
44c	Me	Et	3,4,5-(MeO) ₃	[>100000] ^c	[18] ^c	[>5600]
44d	Me	Et	3,4,5-(MeO) ₃	6290 ± 680	882 ± 239	7.1
44e	Me	2-OHEt	3,4,5-(MeO) ₃	26 ± 9% ^b (10 ⁻⁵)	22% ^b (10 ⁻⁵)	
44f	Me	propargyl	3,4,5-(MeO) ₃	4040 ± 370	525 ± 220	7.7
45a	Et	H	3,4,5-(MeO) ₃	32 ± 9% ^b (10 ⁻⁵)	14% ^b (10 ⁻⁵)	
45b	Et	Me	3,4,5-(MeO) ₃	852 ± 277	269 ± 7	3.2
46	allyl	Me	3,4,5-(MeO) ₃	2790 ± 960	81 ± 17	34
47	Pr	H	H	1930 ± 100	131 ± 69	13
48	Pr	Me	H	[>100000] ^c	[15] ^c	[>6700]
49	Pr	Me	3-NO ₂	55 ± 28	44 ± 19	1.3
50a	Pr	H	3-NH ₂	[1800 or 22] ^d ^c	[26 or 85] ^d ^c	[69 or 0.26]
50b	Pr	Me	3-NH ₂	272 ± 68	56.2 ± 6.8	4.8
51a	Pr	H	3-F	113 ± 21	18.9 ± 5.3	6.0
51b	Pr	Me	3-F	78 ± 17	153 ± 31	0.51
51c	Pr	H	3-Cl	301 ± 64	33 ± 15	9.1
51d	Pr	Me	3-Cl	167 ± 39	216 ± 66	0.77
51e	Pr	Me	3-Cl	874 ± 222	61.3 ± 17.6	14

Table I. (Continued)

compd	R ₁ , R ₃	R ₇	X	K _i ^a		A ₁ /A ₂ ratio
				A ₁	A ₂	
52a	Pr	H	3,4-(MeO) ₂	71 ± 11 [1700] ^c	48.5 ± 8.6 [6700] ^c	1.3 [0.25]
52b ^f	Pr	Me	3,4-(MeO) ₂	577 ± 42 [1500] ^c	31.1 ± 11.8 [7.8] ^c	19 [190]
53a	Pr	H	3,5-(MeO) ₂	632 ± 152	210 ± 140	3.0
53b	Pr	Me	3,5-(MeO) ₂	2630 ± 20	24.0 ± 6.0	110
54a	Pr	H	3,5-F ₂	146 ± 25	346 ± 97	0.42
54b	Pr	Me	3,5-F ₂	382 ± 40	53 ± 15	7.2
55a	Pr	H	2,3,4-(MeO) ₃	97 ± 19	64.0 ± 15.6	1.5
55b	Pr	Me	2,3,4-(MeO) ₃	379 ± 128	68.5 ± 12.6	5.5
56a	Pr	H	2,4,5-(MeO) ₃	143 ± 19	323 ± 74	0.44
56b	Pr	Me	2,4,5-(MeO) ₃	689 ± 239	327 ± 52	2.1

^a Expressed in nM (single determination or mean ± SEM for three or more determinations) vs [³H]PIA (1 nM) at rat A₁-receptors and vs [³H]CGS 21680 (5 nM) at rat striatal A₂-receptors. ^b Percent displacement of specific binding at the concentration indicated in parentheses. ^c Values in brackets are from ref 11 and represent K_i values vs [³H]NECA in rat striatum and vs [³H]CHA in guinea pig brain, unless noted. ^d Affinities at both A₁- and A₂-receptors measured in rat brain (Erickson et al.¹⁹). ^e R_α = F / KF17837.

Table II. Affinities of Related Xanthine Derivatives in Radioligand Binding Assays at Rat Brain A₁- and A₂-Receptors^a

compd	R ₇	R ₈	K _i ^a		A ₁ /A ₂ ratio
			A ₁	A ₂	
2	Me	cyclohexyl	[28000] ^b	17100	1.6
57	Me	2-(3-aminophenyl)ethyl	15% ^c (10 ⁻⁵)	18000	
58a	H	7-methoxybenzofuran-2-yl	1770 ± 70	3900 ± 940	0.5
58b	Me	7-methoxybenzofuran-2-yl		4740	

^a Expressed in nM (single determination or mean ± SEM for three or more determinations) vs [³H]PIA (1 nM) at rat A₁-receptors and vs [³H]CGS 21680 (5 nM) at rat striatal A₂-receptors. ^b Reference 10. ^c Percent displacement of specific binding at the concentration indicated in parentheses.

al.¹¹ These authors found that 8-styryl derivatives of 1,3-dimethylxanthines were the most selective for A₂-receptors (selectivities greater than 5000-fold were reported), but the affinities of the corresponding 1,3-propyl analogues at both subtypes were greater (the most potent compound having a K_i value of 7.8 nM at A₂ receptors).

The selectivity factors in the present study were generally much less than those reported by Shimada et al.¹¹ The principal reason may be that A₁-affinity in this study was measured in the same species as A₂-affinity (rat), whereas Shimada et al. measured A₁ affinity in guinea pig brain and A₂ affinity in rat brain. The species dependence of affinity of alkylxanthines at both A₁ and A_{2a} receptors is well-documented.^{21,22} Invariably, the affinity of xanthines at A₁-receptors is higher in the rat than in the guinea pig, but the interspecies affinity ratios have been found to vary from only 2-fold for theophylline to as much as 20-fold for 8-phenyltheophylline.²¹ Indeed, the reported variation in A₁-affinities for 8-styrylxanthines in different species differ even more: up to 33-fold (e.g. compound 47 has an A₁-affinity in rat of 55 nM versus 1800 nM in guinea pig;¹¹ Erickson et al.¹⁹ have determined a K_i value at rat A₁-receptors of 22 nM). Thus, comparing guinea pig A₁-values to rat A₂-affinities results in artificially high selectivity ratios, and we therefore feel strongly that affinities should be compared within one species in order to yield useful figures.

In addition, some unexplained and substantial differences (e.g. compound 50a) were observed between K_i values

versus [³H]CGS 21680 in this study and versus [³H]NECA in the study of Shimada et al.¹¹ (both having been measured in rat striatal membranes).

Another potential reason for discrepancies with previous results in binding assays was the amount of DMSO present. Shimada et al.¹¹ utilized approximately 1% DMSO in the assay medium, and we used 2%. We demonstrated (Figure 6) that at 0.5% DMSO a 1,3-dipropyl-7-methylxanthine derivative, 53b, did not remain dissolved in aqueous solution at concentrations greater than 10 μM. This would affect in particular A₁-displacement curves for many compounds in this study, for which data points beyond xanthine concentrations of 10 μM are required. Thus, the addition of insufficient DMSO to the medium (or serial aqueous dilutions) might tend to overestimate the selectivity of the A₂-selective xanthines, but would not be expected to alter the apparent affinity at A₂ receptors (Figure 6A).

Functional studies, for example to characterize the antagonism of adenosine agonist-elicited increases in cyclic AMP, will be an important component of future work with these xanthine derivatives.

The most significant findings in this study are as follows: (1) the position of styryl substitution (meta favored) is a determinant of potency and selectivity (compare 17b, 19b, and 29b), (2) increasing the size of small alkyl groups at the 1- and 3-xanthine position (e.g. 45b versus 44b) increases potency at both receptors and decreases A₂-selectivity, and (3) A₂-selectivity and moderate affinity are maintained with long chain extension from the para position of the styryl ring (e.g. 41). It would seem that this position of the 8-styryl group when bound to the receptor is located in a relatively insensitive region. This finding is not unexpected given previous similar observations for para-substituted 8-phenylxanthines.²⁰ A₂-Selectivities of thousands of fold reported previously¹¹ were not observed in this study, although the selectivities of up to 520-fold (compound 28), promise to be useful in physiological studies. A₂-Antagonists of particular interest are compounds 23, 24, 27b, 28, 33b, and 34b (A₂-selectivity of 200-fold or greater); compounds 23, 28, 49, 50b, 52b, 53b, and 54b (A₂-affinity 50 nM or less); and compounds 22b, 38, and 40 (amine functionalized congeners). Compound 24 also has enhanced water solubility; the maximal solubility in a 0.1 M potassium phosphate solution at pH 7.4 was 19 mM.

Table III. Characterization of Xanthine Derivatives

compd	% yield	mp (°C)	formula	analysis	compd	% yield	mp (°C)	formula	analysis
15a	51	>280	C ₁₅ H ₁₄ N ₄ O ₂ ·1/2H ₂ O	H,N;C ^b	38	90 ^e	207–212	C ₂₂ H ₂₈ N ₅ O ₅	c
15b	81	220–222	C ₁₆ H ₁₆ N ₄ O ₂ ·1/4H ₂ O	C,H;N ^b	39	15 ^d	184.5–186.5	C ₂₇ H ₃₇ N ₅ O ₇ ·1/2H ₂ O	C,H,N ^c
16	57	>280	C ₁₅ H ₁₃ N ₄ O ₂ F	C,H,N	40	18	200–206	C ₂₂ H ₂₇ N ₅ O ₅	H,N;C ^{b,c}
17a	31	>300	C ₁₆ H ₁₆ N ₄ O ₃ ·2/5H ₂ O	C,H,N	41	7	229–232 dec	C ₂₄ H ₂₈ N ₅ O ₆	C,H,N ^c
17b	74	238–240	C ₁₇ H ₁₈ N ₄ O ₃	C,H,N	42	75	192–195	C ₂₇ H ₃₅ N ₅ O ₇	C,H,N ^c
18	3	>300	C ₁₅ H ₁₄ N ₄ O ₃	C,H,N ^c	43a	72	165	C ₁₈ H ₂₀ N ₄ O ₅	C,H;N ^b
19a	65	>280	C ₁₆ H ₁₆ N ₄ O ₃	C,H,N	43b	43	189–193	C ₁₉ H ₂₂ N ₄ O ₅ ·1/2H ₂ O	C,H;N ^b
19b	61	212–215	C ₁₇ H ₁₈ N ₄ O ₃ ·1/2H ₂ O	C,H;N ^b	44a	63	>280	C ₁₈ H ₂₀ N ₄ O ₅	C,H,N
20a	55	>300	C ₁₆ H ₁₃ N ₄ O ₂ F ₃	C,H,N,F	44b	82	245–247	C ₁₉ H ₂₂ N ₄ O ₅	C,H,N
20b	84	232–236	C ₁₇ H ₁₅ N ₄ O ₂ F ₃ ·1/2H ₂ O	C,H,N	44c	84	225–229	C ₂₀ H ₂₄ N ₄ O ₅	C,H,N
21a	56	>300	C ₁₅ H ₁₃ N ₅ O ₄	C,H,N	44d	70	251–254	C ₂₀ H ₂₄ N ₄ O ₂	C,H,N
21b	84	306–308	C ₁₆ H ₁₅ N ₅ O ₄	C,H,N	44e	79	235–237	C ₂₁ H ₂₂ N ₄ O ₅	C,H,N
22a	85	>300	C ₁₅ H ₁₅ N ₅ O ₂ ·1/2H ₂ O	C,H,N	44f	71	215–218	C ₂₆ H ₂₈ N ₄ O ₅	C,H,N
22b	92	222–224	C ₁₆ H ₁₇ N ₅ O ₂ ·0.85H ₂ O	C,H,N ^c	45a	20 ^a	286–289	C ₂₀ H ₂₄ N ₄ O ₅ ·1/4H ₂ O	C,H,N ^c
23	77	>300	C ₁₈ H ₁₉ N ₅ O ₃ ·3/5H ₂ O	C,H,N ^c	45b	64	207–210	C ₂₁ H ₂₆ N ₄ O ₅	C,H,N ^c
24	78	>300	C ₂₀ H ₂₁ N ₅ O ₅ ·0.7H ₂ O	C,H,N ^c	47	52 ^a	257–260	C ₁₉ H ₂₂ N ₄ O ₂	C,H,N
25	59	>300	C ₂₁ H ₂₅ N ₅ O ₄ ·1/2H ₂ O	C,H,N ^{b,c}	48	91	215–217	C ₂₀ H ₂₃ N ₅ O ₄	C,H,N
26	27	175–177	C ₂₆ H ₃₃ N ₅ O ₆ ·2/5H ₂ O	C,H,N ^c	49	92	145–148	C ₂₀ H ₂₅ N ₅ O ₂ ·3/4H ₂ O	C,H,N
27a	87	>310	C ₁₅ H ₁₃ N ₄ O ₂ F _{1/2} H ₂ O	C,H;N ^{b,c}	50a	61	264–265	C ₁₉ H ₂₁ N ₄ O ₂ F	C,H,N
27b	75	208–209	C ₁₆ H ₁₅ N ₄ O ₂ F	C,H,N	50b	83	155–157	C ₂₀ H ₂₃ N ₄ O ₂ F _{1/4} H ₂ O	C,H,N ^c
28	10	205	C ₁₆ H ₁₅ N ₄ O ₂ Cl	C,H,N	51a	18 ^a	257–259	C ₁₉ H ₂₁ N ₄ O ₂ Cl	C,H,N
29a	4	>320	C ₁₆ H ₁₆ N ₄ O ₃	C,H,N	51b	67	164–166	C ₂₀ H ₂₃ N ₄ O ₂ Cl	C,H,N ^c
29b	55	220–222	C ₁₇ H ₁₈ N ₄ O ₃	C,H,N ^c	52a	48	250–253	C ₂₁ H ₂₆ N ₄ O ₄ ·1/4H ₂ O	C,H,N
30a	43	>230	C ₁₇ H ₁₉ N ₅ O ₂	c	52b	78	164–164	C ₂₂ H ₂₈ N ₄ O ₄ ·3/4H ₂ O	C,H,N ^c
30b	29	>230	C ₁₈ H ₂₁ N ₅ O ₂	C,H;N ^{b,c}	53a	100	150–152	C ₂₁ H ₂₆ N ₄ O ₄ ·2/5H ₂ O	C,H,N ^c
31a	32 ^a	299–301	C ₁₇ H ₁₈ N ₄ O ₄	C,H,N	53b	59	166–167	C ₂₂ H ₂₈ N ₄ O ₄	C,H,N ^c
31b	49	233.5–235	C ₁₈ H ₂₀ N ₄ O ₄ ·1/2H ₂ O	C,H,N	54a	78	275–278	C ₁₉ H ₂₀ N ₄ O ₂ F ₂ ·3/4H ₂ O	C,H,N
32a	4	>295	C ₁₇ H ₁₈ N ₄ O ₄	C,H,N ^c	54b	85	161–163	C ₂₀ H ₂₂ N ₄ O ₂ F ₂ ·0.9H ₂ O	C,H,N
32b	63	230–232	C ₁₈ H ₂₀ N ₄ O ₄ ·1/2H ₂ O	C,H,N	55a	32	241–244	C ₂₂ H ₂₈ N ₄ O ₅	C,H,N
33a	18	>320	C ₁₇ H ₁₈ N ₄ O ₄	C,H,N	55b	88	107.5–109	C ₂₅ H ₃₀ N ₄ O ₅	C,H,N
33b	63	228–230	C ₁₈ H ₂₀ N ₄ O ₄	C,H,N	56a	11	252–254	C ₂₂ H ₂₈ N ₄ O ₅	C,H,N
34a	76	>310	C ₁₅ H ₁₂ N ₄ O ₂ F ₂	C,H,N	56b	82	193–194	C ₂₃ H ₃₀ N ₄ O ₅ ·1/2H ₂ O	C,H,N
34b	87	238–239	C ₁₆ H ₁₄ N ₄ O ₂ F ₂ ·3/4H ₂ O	C,H,N	57	67	158–160	C ₁₆ H ₁₉ N ₅ O ₂	C,H,N ^c
35	54	269 dec	C ₂₀ H ₂₂ N ₄ O ₆	C,H,N ^c	58a	78	>280	C ₁₆ H ₁₄ N ₄ O ₄ ·1/2H ₂ O	C,H,N
36	57	274–279	C ₁₈ H ₂₀ N ₄ O ₅ ·1/4H ₂ O	C,H,N ^c	58b	99	273–275	C ₁₇ H ₁₆ N ₄ O ₄	C,H,N
37	75	190–194	C ₂₅ H ₂₆ N ₄ O ₅ ·1/2H ₂ O	C,H,N					

^a Yield calculated from 1,3-dialkyl-6-amino-5-nitrosouracil. ^b Analyses: % N found (calcd) 15b, 17.60 (18.62); 19b, 15.86 (16.71); 25, 16.00 (16.66); 27a, 17.58 (18.11); 30b, 18.15 (20.63); 43a, 14.06 (15.06); 43b, 13.37 (14.17); % C found (calcd) 15a, 62.42 (61.85); 40, 58.66 (59.28). ^c Accurate mass, measured (ppm from calculated), in EI mode, unless noted: 18, 298.1055 (–3.7); 22b, 311.1373 (5.6); 23, 253.1483 (–1.4); 24, 411.1556 (3.2); 25, 411.1894 (–3.1); 26, 511.2450 (3.7); 27a, 300.1018 (2.3); 29b, 326.1371 (–2.4); 30a, 325.1537 (–0.5); 30b, 339.1688 (–4.1); 32a, 342.1326 (–0.6); 35, 372.1436 (0.7); 36, 414.1543 (0.9); 38 (FAB), 444.2255 (0.8); 39, 543.2684 (–1.7); 40, 441.2001 (–2.5); 41, 483.2131 (2.7); 42, 541.2544 (1.4); 45a (FAB), 401.1812 (–1.3); 45b, 414.1898 (–1.3); 50b, 370.1795 (–2.7); 51b, 386.1492 (–4.5); 52b, 412.2110 (–0.1); 53a, 398.1937 (–4.3); 53b, 412.2093 (–4.3); 57, 313.1521 (–5.7). ^d From compound 44b. ^e From compound 39.




Figure 4. Correlation of affinity at adenosine receptors for 7-H versus 7-Me analogues of 1,3-dimethyl-8-styrylxanthine derivatives. Inhibition constants in nM given for A₁-receptors (□) and A₂-receptors (●).

A goal of this study is to identify adenosine antagonists that may be utilized as radioligands. Compound 53b is of sufficiently high affinity and A₂-selectivity to be useful in binding assays when radiolabeled. An additional objective is to design functionalized congeners for eventual




Figure 5. Correlation of affinity at adenosine receptors for 7-H versus 7-Me analogues of 1,3-dipropyl-8-styrylxanthine derivatives. Inhibition constants in nM given for A₁-receptors (○) and A₂ receptors (●).

covalent labeling of the receptor, as was carried out for A₂ receptors using an agonist ligand²³ and for A₁-receptors.²⁰ The arylamino derivatives 22b and 49 and the alkylamino derivatives 38 and 40 are chemically functionalized for further reaction leading to A₂-selective conjugates.




Figure 6. (A) Dependence of observed IC₅₀ values on DMSO concentration in competitive radioligand binding of 1,3-dipropyl-8-(3,5-dimethoxystyryl)xanthine, **53b**, at A₁- and A₂-receptors. IC₅₀ values are expressed in nM (mean ± SEM for three or more determinations) vs [³H]PIA (1 nM) at rat A₁-receptors (□) and vs [³H]CGS21680 (5 nM) at rat striatal A₂-receptors (○). (B) UV absorption of water solutions following addition of compound **53b** dissolved in DMSO (theoretical final concentration assuming complete dissolution given on abscissa). The final concentration of DMSO was 0.5% in each case. The peak absorption occurred at 345 nm, with a molar extinction coefficient (ε) of 13 200.

Experimental Section

Chemistry. New compounds were characterized (and resonances assigned) by 300-MHz proton nuclear magnetic resonance spectroscopy using a Varian GEMINI-300 FT-NMR spectrometer. Unless noted, chemical shifts are expressed as ppm downfield from tetramethylsilane. Synthetic intermediates were characterized by chemical ionization mass spectrometry (NH₃) and xanthine derivatives by fast atom bombardment mass spectrometry (positive ions in a glycerol matrix) on a JEOL SX102 mass spectrometer. In the EI mode accurate mass was determined using a VG7070F mass spectrometer. C, H, and N analyses were carried out by Atlantic Microlabs (Norcross, GA), and ±0.4% was acceptable. All xanthine derivatives were judged to be homogeneous using thin-layer chromatography following final purification. 7-Methoxy-2-benzofurancarboxylic acid, *trans*-cinnamic acid, and the following derivatives thereof were purchased from Aldrich (St. Louis, MO): α-fluoro, 2-methoxy, 3,4-dimethoxy, 3,5-difluoro, and 3,5-dimethoxy. 3- and 4-methoxy derivatives of *trans*-cinnamic acid were purchased from Fluka (Ronkonoma, NY). The following derivatives of *trans*-cinnamic acid were purchased from Lancaster (Windham, NH): 2,3-dimethoxy, 3,4,5-trimethoxy, 2,3,4-trimethoxy, 2,4,5-trimethoxy, and 3-fluoro. The following derivatives of *trans*-cinnamic acid were from Janssen Chimica (Geel, Belgium): 3-trifluoromethyl, 3-chloro, and 3-nitro. 2-Chloroadenosine was purchased from Research Biochemicals, Inc. (Natick, MA). Compound **46** was the gift of Dr. Ray Olsson (University of Southern Florida, Tampa,

FL). 8-Cyclohexylcaffeine, **2**, was the gift of Dr. John W. Daly (NIH). Analytical TLC plates and silica gel (230–400 mesh) were purchased from VWR (Bridgeport, NJ).

Synthesis of Xanthine Derivatives. General Procedure A: Reaction of Substituted Cinnamic Acid Derivatives with 5,6-Diamino-1,3-dimethyluracil To Yield Amide **9.** The substituted cinnamic acid (1 equiv) was dissolved in a minimum volume of DMF containing 1,3-dialkyl-5,6-diaminouracil (1.5 equiv). 1-[3-(Dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (1 equiv) was added, followed by a catalytic amount (0.05 equiv) of 4-(*N,N*-dimethylamino)pyridine and 0.05 equiv of imidazole. The mixture was stirred at room temperature for 3 h, and saturated sodium chloride solution was added (for 1,3-dipropyl derivatives water was used here), to form a precipitate or amorphous insoluble fraction. The insoluble residue was filtered and dissolved in 4 N aqueous sodium hydroxide containing sufficient methanol to obtain a clear solution.

The mixture was heated at 60 °C for 2 h or until the complete disappearance of starting material, as judged using TLC (silica plate, CHCl₃/CH₃OH/HOAc, 85:10:5, v/v). The mixture was cooled and acidified to pH = 1 with 6 N aqueous hydrochloric acid solution.

The precipitate was washed with water, dried, and further purified using a preparative silica plate (85–95% CHCl₃/5–15% methanol/1–5% HOAc). This procedure was used to make compounds **17a**, **29a**, **32a**, **33a**, etc.

General Procedure B: Methylation of N-7 8-Styrylxanthines. An 8-styrylxanthine derivative (1 equiv) was dissolved in a minimum of DMF. Excess finely powdered anhydrous potassium carbonate was added and the solution was left in an ultrasonic bath for 10 min. Methyl iodide (5 equiv) was added. The mixture was stirred at 60 °C for 30 min or until the complete disappearance of starting material, as judged using TLC (silica, chloroform/methanol/acetic acid, 95:4:1, v/v). The reaction mixture was cooled, and excess concentrated aqueous ammonia solution was added. The precipitate was washed with water, dried in vacuo, and further purified either by crystallization or by chromatography on a preparative thin-layer plate (85–95% chloroform/5–15% methanol/1–5% acetic acid). This procedure was used to make compounds **17b**, **29b**, **32b**, **33b**, etc.

1,3-Dimethyl-8-(2-methoxystyryl)xanthine (17a). Compound **17a** was made from 2-methoxycinnamic acid according to general procedure A and triturated with hot methanol: Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 3.27 (s, 3 H, N₃-CH₃), 3.35 (s, 3 H, N₁-CH₃), 3.5 (s, 3 H, OCH₃), 3.9 (s, 3 H, N₇-CH₃), 7.1 (d, 1 H, *J* = 18 Hz), 7.0–7.2 (m, 2 H), 7.4 (m, 1 H), 7.7 (d, 1 H, *J* = 8 Hz), 7.8 (d, 1 H, *J* = 18 Hz). MS (CI/NH₃): *m/e* 313 (MH⁺, base) 281, 117.

1,3,7-Trimethyl-8-(2-methoxystyryl)xanthine (17b). Compound **17b** was made from **17a** according to general procedure B. Mp: 238–240 °C. ¹H NMR DMSO-*d*₆: δ 3.24 (s, 3 H, N₃-CH₃), 3.48 (s, 3 H, N₁-CH₃), 3.90 (s, 3 H, OCH₃), 4.06 (s, 3 H, N₇-CH₃), 7.0–7.14 (m, 2 H), 7.34 (d, 1 H, *J* = 16 Hz), 7.4 (m, 1 H), 7.9 (d, 1 H, *J* = 8 Hz), 8.0 (d, 1 H, *J* = 16 Hz). MS (CI/NH₃): *m/e* 327 (MH⁺) base peak.

1,3-Dimethyl-8-[3-(trifluoromethyl)styryl]xanthine (20a). Compound **20a** was made from 3-(trifluoromethyl)cinnamic acid according to general procedure A. Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 3.26 (s, 3 H, NCH₃), 3.48 (s, 3 H, NCH₃), 7.19 (d, 1 H, *J* = 16 Hz), 7.64 (t, 1 H, *J* = 8 Hz), 7.70 (d, 1 H, *J* = 7 Hz), 7.72 (d, 1 H, *J* = 16 Hz), 7.94 (d, 1 H, *J* = 8 Hz), 7.96 (s, 1 H). MS (CI): *m/e* 350 (base), 329, 292.

1,3,7-Trimethyl-8-[3-(trifluoromethyl)styryl]xanthine (20b). Compound **20b** was made from **20a** according to general procedure B. Mp: 232–236 °C. ¹H NMR DMSO-*d*₆: δ 3.25 (s, 3 H, NCH₃), 3.49 (s, 3 H, NCH₃), 4.09 (s, 3 H, N₇-CH₃), 7.58 (d, 1 H, *J* = 16 Hz), 7.67 (t, 1 H, *J* = 8 Hz), 7.72 (d, 1 H, *J* = 8 Hz), 7.78 (d, 1 H, *J* = 16 Hz), 8.09 (d, 1 H, *J* = 7 Hz), 8.26 (s, 1 H). MS (EI): *m/e* 364.

1,3-Dimethyl-8-(3-nitrostyryl)xanthine (21a). Compound **21a** was made from 3-nitrocinnamic acid according to general procedure A (temperature raised to 80 °C for 3 h, recrystallized from methanol). Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 3.25 (s, 3 H, NCH₃), 3.48 (s, 3 H, NCH₃), 7.22 (d, 1 H, *J* = 16 Hz), 7.70 (t, 1 H, *J* = 8 Hz), 7.76 (d, 1 H, *J* = 16 Hz), 8.10 (d, 1 H, *J* = 8

H_z), 8.18 (d, 1 H, *J* = 8 Hz), 8.41 (s, 1 H). MS (EI): *m/e* 327 (base), 310, 280.

1,3,7-Trimethyl-8-(3-nitrostyryl)xanthine (21b). Compound 21b was made from 21a according to general procedure B. Mp: 306–308 °C. ¹H NMR DMSO-*d*₆: δ 3.23 (s, 3 H, NCH₃), 3.47 (s, 3 H, NCH₃), 4.08 (s, 3 H, N₇-CH₃), 7.63 (d, 1 H, *J* = 16 Hz), 7.71 (t, 1 H, *J* = 8 Hz), 7.80 (d, 1 H, *J* = 16 Hz), 8.18 (d, 1 H, *J* = 8 Hz), 8.23 (d, 1 H, *J* = 8 Hz), 8.70 (s, 1 H). MS (EI): *m/e* 341 (base), 294.

1,3-Dimethyl-8-(3-aminostyryl)xanthine (22a). Compound 22a was made from 21a by reducing with Zn/acetic acid for 3 h. Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 3.24 (s, 3 H, NCH₃), 3.46 (s, 3 H, NCH₃), 5.19 (s, 2 H, NH₂), 6.56 (d, 1 H, *J* = 8 Hz), 6.74 (d, 1 H, *J* = 8 Hz), 6.76 (s, 1 H), 6.84 (d, 1 H, *J* = 16 Hz), 7.05 (t, 1 H, *J* = 8 Hz), 7.49 (d, 1 H, *J* = 16 Hz). MS (CI/NH₃): *m/e* 315 (M + NH₄⁺), 298 (MH⁺, base).

1,3,7-Trimethyl-8-(3-aminostyryl)xanthine (22b). Compound 22b was made from 21b using Zn/acetic acid as reducing agent for 3 h. Mp: 222–224 °C. ¹H NMR DMSO-*d*₆: δ 3.22 (s, 3 H, N-CH₃), 3.46 (s, 3 H, NCH₃), 4.00 (s, 3 H N₇-CH₃), 5.14 (s, 2 H, NH₂), 6.58 (d, 1 H, *J* = 8 Hz, H-4), 6.87 (s, 1 H, H-2), 6.92 (d, 1 H, *J* = 8 Hz, H-6), 7.07 (t, 1 H, *J* = 8 Hz, H-5), 7.14 (d, 1 H, *J* = 16 Hz), 7.51 (d, 1 H, *J* = 16 Hz). MS (CI/NH₃): *m/e* 312 (MH⁺).

1,3,7-Trimethyl-8-[3-(acetylamino)styryl]xanthine (23). Compound 23 was made from 22b with acetic anhydride in DMF and DMAP for 1 h. Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 2.06 (s, 3 H, COCH₃), 3.23 (s, 3 H, NCH₃), 3.47 (s, 3 H, NCH₃), 4.03 (s, 3 H, N₇-CH₃), 7.24 (d, 1 H, *J* = 16 Hz), 7.34 (t, 1 H, *J* = 8 Hz), 7.50 (t, 1 H, *J* = 8 Hz), 7.54 (d, 1 H, *J* = 8 Hz), 7.61 (d, 1 H, *J* = 16 Hz), 7.86 (s, 1 H). MS (CI/NH₃): *m/e* 354 (MH⁺).

1,3,7-Trimethyl-8-[3-(3-carboxy-1-oxopropyl)amino]styryl]xanthine (24). Compound 24 was made from 22b with succinic anhydride in DMF and DMAP. Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 2.28 (t, 2 H, *J* = 7 Hz), 2.43 (t, 2 H, *J* = 7 Hz), 3.23 (s, 3 H, NCH₃), 3.47 (s, 3 H, NCH₃), 4.03 (s, 3 H, N₇-CH₃), 7.24 (d, 1 H, *J* = 16 Hz), 7.32 (t, 1 H, *J* = 8 Hz), 7.45 (d, 1 H, *J* = 8 Hz), 7.54 (d, 1 H, *J* = 8 Hz), 7.61 (d, 1 H, *J* = 16 Hz), 7.82 (s, 1 H). MS (CI/NH₃): *m/e* 394 (M - OH), 312, 209 (base). UV: λ_{max} (methanol) 349 nm, log ε = 4.48. The maximal aqueous solubility following dissolution in K₂HPO₄ (0.1 M) was determined to be 19 mM.

1,3,7-Trimethyl-8-[3-[(*tert*-butyloxy)carbonyl]amino]styryl]xanthine (25). Compound 25 was made from 22b with di-*tert*-butyl dicarbonate and DMAP in DMF. Mp: >300 °C. ¹H NMR DMSO-*d*₆: δ 1.40 (s, 9 H, CH₃COO), 3.17 (s, 3 H, NCH₃), 3.41 (s, 3 H, NCH₃), 3.89 (s, 3 H, N₇-CH₃), 7.23 (d, 1 H, *J* = 16 Hz), 7.33 (d, 1 H, *J* = 8 Hz), 7.51 (t, 1 H, *J* = 8 Hz), 7.57 (s, 1 H), 7.67 (d, 1 H, *J* = 16 Hz), 7.75 (d, 1 H, *J* = 8 Hz). MS (CI/NH₃): *m/e* 414 (M - CH₃ + NH₄⁺, base), 338, 314, 312.

1,3,7-Trimethyl-8-[3-bis[(*tert*-butyloxy)carbonyl]amino]styryl]xanthine (26). Compound 26 was made from 22b with di-*tert*-butyl dicarbonate and DMAP in DMF. Mp: 175–177 °C. ¹H NMR DMSO-*d*₆: δ 1.39 (s, 18 H, CH₃COO), 3.23 (s, 3 H, NCH₃), 3.46 (s, 3 H, NCH₃), 4.03 (s, 3 H, N₇-CH₃), 7.17 (d, 1 H, *J* = 8 Hz), 7.42 (t, 1 H, *J* = 8 Hz), 7.43 (d, 1 H, *J* = 16 Hz), 7.67 (d, 1 H, *J* = 16 Hz), 7.69 (d, 1 H, *J* = 8 Hz), 7.74 (s, 1 H). MS (CI/NH₃): *m/e* 514 (M - CH₃ + NH₄⁺), 414 (base).

1,3-Dimethyl-8-(4-methoxystyryl)xanthine (29a). Compound 29a was made from 4-methoxycinnamic acid according to general procedure A. Mp: >320 °C. ¹H NMR DMSO-*d*₆: δ 3.24 (s, 3 H, N₃-CH₃), 3.46 (s, 3 H, N₇-CH₃), 3.78 (s, 3 H, OCH₃), 6.85 (d, 1 H, *J* = 16 Hz), 7.0 (d, 2 H, *J* = 8 Hz), 7.55 (d, 2 H, *J* = 8 Hz), 7.6 (d, 1 H, *J* = 16 Hz). MS (CI/NH₃): *m/e* 313 (MH⁺, base) 172.

1,3,7-Trimethyl-8-(4-methoxystyryl)xanthine (29b). Compound 29b was made from 29a according to general procedure B. Mp: >320 °C. ¹H NMR DMSO-*d*₆: δ 3.22 (s, 3 H, N₃-CH₃), 3.45 (s, 3 H, N₇-CH₃), 3.8 (s, 3 H, OCH₃), 4.0 (s, 3 H, N₇-CH₃), 7.0 (d, 1 H, *J* = 8 Hz), 7.2 (d, 1 H, *J* = 16 Hz), 7.66 (d, 1 H, *J* = 16 Hz), 7.72 (d, 1 H, *J* = 8 Hz). MS (CI/NH₃): *m/e* 327 (MH⁺, base) 205.

1,3-Dimethyl-8-[4-(dimethylamino)styryl]xanthine (30a). A solution of 4-(dimethylamino)cinnamic acid (0.1 g, 0.52 mmol), 1-hydroxybenzotriazole (0.14 g, 1.04 mmol), and EDAC (0.19 g, 1.04 mmol) in DMF (1 mL) was sonicated for 1 h. 1,3-Dimethyl-

5,6-diaminouracil (0.088 g, 0.52 mmol) was added, and the mixture was heated for 3 h at 80 °C. The dark red solution was cooled to room temperature, and the product was obtained as a deep yellow precipitate (0.045 g). An additional crop was obtained by cooling the mother liquor in an ice bath and adding 10 volumes of brine (combined yield 38%). ¹H NMR CD₃OD: δ 7.54 (d, 1 H, *J* = 15.5 Hz), 7.45 (d, 2 H, 8.8 Hz), 6.74 (d, 2 H, *J* = 8.8 Hz), 6.56 (d, 1 H, *J* = 15.5 Hz), 3.42, 3.27 (s, 3 H, CH₃), 3.00 (s, 6 H, N(CH₃)₂). MS (CI): *m/e* 344 (MH⁺).

The above amide (0.045 g, 0.13 mmol) was suspended in methanol (1 mL), and 4 N NaOH (1 mL) was added. The resulting solution was stirred at 80–90 °C for 1.5 h. HCl (18%) was added carefully to the ice-cooled reaction solution to pH = 7–8. A yellow precipitate was obtained (0.018 g, 43%). ¹H NMR DMSO-*d*₆: δ 7.54 (d, 1 H, *J* = 16 Hz), 7.44 (d, 2 H, *J* = 8.5 Hz), 6.74 (d, 2 H, *J* = 16 Hz), 6.74 (d, 2 H, *J* = 16 Hz), 3.47, 3.25 (s, 3 H, CH₃), 2.97 (s, 6 H, N(CH₃)₂). MS (CI): *m/e* 326 (MH⁺).

1,3-Dimethyl-8-(2,3-dimethoxystyryl)xanthine (31a). Compound 31a was made from 2,3-dimethoxycinnamic acid according to general procedure A (recrystallized from methanol). Mp: 299–301 °C. ¹H NMR DMSO-*d*₆: δ 3.25 (s, 3 H, N₃-CH₃), 3.47 (s, 3 H, NCH₃), 3.78 (s, 3 H, OCH₃), 3.82 (s, 3 H, OCH₃), 7.05 (d, 1 H, *J* = 17 Hz), 7.05 (dd, 1 H, *J* = 2, 8 Hz), 7.11 (t, 1 H, *J* = 8 Hz), 7.26 (dd, 1 H, *J* = 2, 8 Hz), 7.84 (d, 1 H, *J* = 17 Hz) MS (CI/NH₃): *m/e* 360 (M + NH₄⁺), 343 (base peak).

1,3,7-Trimethyl-8-(2,3-dimethoxystyryl)xanthine (31b). Compound 31b was made from 31a according to the general procedure B. Mp: 233–235 °C. ¹H NMR DMSO-*d*₆: δ 3.23 (s, 3 H, NCH₃), 3.47 (s, 3 H, NCH₃), 3.78 (s, 3 H, OCH₃), 3.83 (s, 3 H, OCH₃), 4.02 (s, 3 H, N₇-CH₃), 7.06 (d, 1 H, *J* = 8 Hz), 7.10 (t, 1 H, *J* = 8 Hz), 7.32 (d, 1 H, *J* = 16 Hz), 7.51 (d, 1 H, *J* = 8 Hz), 7.90 (d, 1 H, *J* = 16 Hz). MS (EI): *m/e* 356 (base), 325.

1,3-Dimethyl-8-(3,4-dimethoxystyryl)xanthine (32a). Compound 32a was made from 3,4-dimethoxycinnamic acid according to general procedure A. Mp: >320 °C. ¹H NMR DMSO-*d*₆: δ 3.25 (s, 3 H, N₃-CH₃), 3.46 (s, 3 H, N₁-CH₃), 3.78 (s, 3 H, OCH₃), 3.82 (s, 3 H, OCH₃), 6.96 (d, 1 H, *J* = 16 Hz), 6.98 (d, 1 H, *J* = 8 Hz), 7.14 (d, 1 H, *J* = 8 Hz), 7.25 (s, 1 H). MS (CI/NH₃): *m/e* 343 (MH⁺), 172 (base peak).

1,3,7-Trimethyl-8-(3,4-dimethoxystyryl)xanthine (32b). Compound 32b was made from 32a according to general procedure B. Mp: 230–232 °C. ¹H NMR DMSO-*d*₆: δ 3.29 (s, 3 H, N₃-CH₃), 3.52 (s, 3 H, N₇-CH₃), 3.85 (s, 3 H, OCH₃), 3.9 (s, 3 H, OCH₃), 4.09 (s, 3 H, N₇-CH₃), 7.05 (d, 1 H, *J* = 8 Hz), 7.25 (d, 1 H, *J* = 16 Hz), 7.30 (d, 1 H, *J* = 8 Hz), 7.48 (s, 1 H), 7.66 (d, 1 H, *J* = 16 Hz). MS (CI): *m/e* 357 (MH⁺ base), 209.

1,3-Dimethyl-8-(3,5-dimethoxystyryl)xanthine (33a). Compound 33a was made from 3,5-dimethoxycinnamic acid according to general procedure A. Mp: >320 °C. ¹H NMR DMSO-*d*₆: δ 3.24 (s, 3 H, N₃-CH₃), 3.46 (s, 3 H, N₁-CH₃), 3.78 (s, 6 H, OCH₃), 6.5 (s, 1 H), 6.78 (s, 2 H), 7.02 (d, 1 H, *J* = 16 Hz), 7.54 (d, 1 H, *J* = 16 Hz). MS (CI): *m/e* 343 (MH⁺ base), 166, 136.

1,3,7-Trimethyl-8-(3,5-dimethoxystyryl)xanthine (33b). Compound 33b was made from 33a according to general procedure B. Mp: 228–230 °C. ¹H NMR DMSO-*d*₆: δ 3.22 (s, 3 H, N₃-CH₃), 3.45 (s, 3 H, N₁-CH₃), 3.79 (s, 6 H, OCH₃), 4.04 (s, 3 H, N₇-CH₃), 6.5 (s, 1 H), 6.97 (s, 2 H), 7.32 (d, 1 H, *J* = 16 Hz), 7.58 (d, 1 H, *J* = 16 Hz).

General Procedure C: Demethylation Followed by O-Alkylation of 8-(3,5-Dimethoxy-4-hydroxystyryl)xanthines. 1,3,7-Trialkyl-8-(3,4,5-trimethoxystyryl)xanthine (1 equiv) was dissolved in a minimum of DMF, and 1.5 equiv of sodium thiophenoxide was added. The solution was heated to 150–160 °C for 20 min or until judged complete using TLC. Ether was added, and the resulting orange precipitate was isolated by filtration. This solid was dissolved in DMF, and water and 6 N HCl were added until pH 2 to give 35 as a yellow precipitate. The appropriate halide (2 equiv for monohalide and 8 equiv for dihalide) was added to a solution of 35 in DMF, followed by finely powdered, anhydrous K₂CO₃. The solution was left in an ultrasonic bath for 15 min and further heated at 50–80 °C for 2 h or until judged complete using TLC. The reaction mixture was cooled and extracted with petroleum ether. The crude product was precipitated by water (for product of reaction with monohalides) or reacted further (for dihalides) with concentrated aqueous ammonia and chromatographed on preparative TLC

using 90–95% chloroform/5–10% methanol/1% aqueous ammonia. This procedure was used to make compounds 37–40.

The 4-hydroxy intermediate 35 was also recrystallized (MeOH/H₂O). ¹H NMR DMSO-*d*₆: δ 3.23 and 3.46 (each s, 3 H, N₁- and N₃-CH₃); 3.83 (s, 6 H, 3,5-(OCH₃)), 4.03 (s, 3 H, N₇-CH₃), 7.08 (s, 2 H, Ar), 7.18 and 7.61 (each d, 1 H, C=C, *J* = 15.7 Hz), 8.82 (br s, 1 H, ArOH).

1,3,7-Trimethyl-8-[3,5-dimethoxy-4-(benzyloxy)styryl]xanthine (37). Compound 37 was made from benzyloxy bromide according to general procedure C. Mp: 190–195 °C. ¹H NMR CDCl₃: δ 3.42 (s, 3 H, N₃CH₃), 3.63 (s, 3 H, N₅CH₃), 3.89 (s, 6 H, OCH₃), 5.06 (s, 2 H, OCH₂), 6.8 (s, 2 H), 6.78 (d, 1 H, *J* = 16 Hz), 7.3–7.5 (m, 5 H), 7.7 (d, 1 H, *J* = 16 Hz). MS (CI): *m/e* 463 (MH⁺ base), 375, 357.

1,3,7-Trimethyl-8-[3,5-dimethoxy-4-[(4-aminobutyl)oxy]styryl]xanthine (38). Compound 38 was made from 1,4-dibromobutane according to general procedure C. MS (CI): *m/e* 444 (MH⁺ base), 373, 359.

1,3,7-Trimethyl-8-[3,5-dimethoxy-4-[[4-[(*tert*-butyloxy)carbonyl]amino]butyl]oxy]styryl]xanthine (39). Compound 39 was made from 38 using di-*tert*-butyl dicarbonate in CHCl₃ (30 min). The chloroform was removed under a stream of N₂, and the crude product was purified using a preparative plate (silica, ethyl acetate/petroleum ether, 70:30). ¹H NMR CDCl₃: δ 1.41 (s, 9 H, CH₃), 1.6–1.8 (m, 4 H, CH₂), 3.2 (m, 2 H, CH₂NH), 4.0 (m, 2 H, OCH₂), 3.39 (s, 3 H, N₃-CH₃), 3.6 (s, 2 H, N₇-CH₃), 3.88 (s, 6 H, OCH₃), 4.05 (s, 3 H, N₇-CH₃), 6.74 (s, 2 H), 6.75 (d, 1 H, *J* = 16 Hz), 7.7 (d, 1 H, *J* = 16 Hz). MS (CI): *m/e* 544 (MH⁺ base) 44, 359.

1,3,7-Trimethyl-8-[3,5-dimethoxy-4-[(4-amino-*trans*-butenyl)oxy]styryl]xanthine (40). Compound 40 was made from 1,4-dibromo-*trans*-2-butene according to general procedure C. ¹H NMR CDCl₃: δ 3.41 (s, 3 H, N₃-CH₃), 3.63 (s, 3 H, N₁-CH₃), 3.91 (s, 6 H, OCH₃), 4.06 (s, 3 H, N₇-CH₃), 4.43 (s, 2 H, CH₂NH₂), 5.94 (s, 2 H, OCH₂), 6.78 (s, 2 H), 6.79 (d, 1 H, *J* = 16 Hz). MS (CI): *m/e* 442 (MH⁺ base) 373, 357, 124.

7-Ethyl-1,3-trimethyl-8-(3,4,5-trimethoxystyryl)xanthine (44c). Compound 44c was made from compound 44a according to general procedure B, except that ethyl iodide was used instead of methyl iodide. ¹H NMR DMSO-*d*₆: δ 1.34 (t, 3 H, CH₃Et, *J* = 7 Hz), 3.25 and 3.47 (each s, 3 H, NCH₃), 3.70 (s, 4 H, 4-OCH₃), 3.86 (s, 6 H, 3,5-(OCH₃)), 4.54 (q, 2 H, N₇-CH₂), 7.13 (s, 2 H, Ar), 7.30 and 7.68 (each d, 1 H, C=C, *J* = 16 Hz).

1,3-Dipropyl-7-methyl-8-styrylxanthine (47). 5-Amino-6-nitroso-1,3-dipropyluracil^{19,20} was suspended in DMF (10 mmol/100 mL) and hydrogenated over 5% Pd/C at 40 psi overnight. The clear solution was filtered through Celite and could be stored at –20 °C.

trans-Cinnamic acid (0.47 g) and EDAC (0.65 g) were added to 2.1 mmol of the above solution and stirred for 4 h. An additional 0.3 g of EDAC was added. After an additional 2 h, half-saturated NaCl solution was added and the mixture was extracted with ethyl acetate (6×). The organic layer was dried over Na₂SO₄ and evaporated to an oil, which was used without further purification.

The above oil was dissolved in methanol (30 mL) and treated with 4 N NaOH (20 mL). After refluxing for 15 min, the mixture was cooled, ice was added, and it was acidified using 6 N HCl. A precipitate formed and was recovered by filtration. The NMR and MS were consistent with the assigned structure of 47. Recrystallized from DMF/water.

ϵ_{342} for 47 in methanol (λ_{max}) was 35 100. A smaller absorption peak was at 265 nm.

1,3,7-Trimethyl-8-[2-(3-aminophenyl)ethyl]xanthine (57). Compound 54 was made from 21b with H₂/Pd at 50 psi in DMF for 3 h. Mp: 158–160 °C. ¹H NMR DMSO-*d*₆: δ 2.82 (t, 2 H, *J* = 8 Hz), 2.96 (t, 2 H, *J* = 8 Hz), 3.20 (s, 3 H, NCH₃), 3.42 (s, 3 H, NCH₃), 3.69 (s, 3 H, N₇-CH₃), 4.95 (s, 2 H, NH₂), 6.34–6.39 (3 H, H-2, H-4, H-6), 6.90 (t, 1 H, *J* = 8 Hz, H-5). MS (CI/NH₃): *m/e* 314 (MH⁺).

Biological Methods. Receptor Binding. Rat cerebral cortical membranes and striatal membranes were prepared^{12,13} and treated with adenosine deaminase (2 units/mL) for 30 min at 37 °C prior to storage at –70 °C. Solid samples of the adenosine derivatives were dissolved in DMSO and stored in the dark at –20 °C. The stock solutions were diluted with DMSO to a concentration of ≤0.1 mM prior to adding to the aqueous medium.

The final concentration of DMSO in the assay medium was generally 2%.

Inhibition of binding of 1 nM [³H]-N⁶-(2-phenylisopropyl)-adenosine (Du Pont NEN, Boston, MA) to A₁-receptors in rat cerebral cortex membranes was measured as described.¹⁷ Membranes (~100 μg of protein/tube) were incubated for 1.5 h at 37 °C in a total volume of 0.5 mL of 50 mM Tris HCl, at pH 7.4. Test drugs were dissolved in DMSO and added in 10-μL aliquots, resulting in a final DMSO concentration of 2%. Bound and free radioligand were separated by addition of 3 mL of a buffer containing 50 mM Tris HCl at pH 7.4 at 5 °C followed by vacuum filtration using a Brandel cell harvester (Brandel, Gaithersburg, MD) and a Whatman GF/B glass-fiber filter with additional washes totaling 9 mL of buffer. Nonspecific binding was determined with 10 μM 2-chloroadenosine.

Inhibition of binding of 5 nM [³H]CGS 21680 was carried out as follows. Membranes (~80 μg of protein/tube, prepared according to ref 8) were incubated for 1 h at 25 °C in a total volume of 0.5 mL of 50 mM Tris HCl containing 10 mM MgCl₂ at pH 7.4. Test drugs were dissolved in DMSO and added in 10-μL aliquots, resulting in a final DMSO concentration of 2%. Nonspecific binding was defined using 20 μM 2-chloroadenosine. Filtration was carried out using a Brandel cell harvester, as above, using Tris HCl/MgCl₂ as the washing buffer.

At least six different concentrations spanning 3 orders of magnitude, adjusted appropriately for the IC₅₀ of each compound, were used. IC₅₀ values, computer-generated using a nonlinear regression formula on the GraphPAD program (Institute for Scientific Information), were converted to apparent K_i values using K_D values^{12,13} of 1.0 and 14 nM for [³H]PIA and [³H]CGS 21680 binding, respectively, and the Cheng–Prusoff equation.¹⁸

Acknowledgment. N.M., C.G., and A.v.B. thank the Cystic Fibrosis Foundation for financial support. M.M. thanks Cortex Pharmaceuticals, Inc., Irvine, CA, for financial support.

References

- (1) Lohse, M. J.; Elger, B.; Lindenborn-Fotinos, J.; Klotz, K. N.; Schwabe, U. Separation of solubilized A₂ adenosine receptors of human platelets from non-receptor [³H]NECA binding sites by gel filtration. *Naunyn Schmiedeberg's Arch. Pharmacol.* 1988, 337, 64–68.
- (2) Ueda, M.; Thompson, R. D.; Arroyo, L. H.; Olsson, R. A. 2-Aralkoxyadenosines: Potent and selective agonists at the coronary artery A₂ adenosine receptor. *J. Med. Chem.* 1991, 34, 1340–1344.
- (3) Ueno, M.; Brookins, J.; Beckman, B.; Fisher, J. W. A₁ and A₂ adenosine receptor regulation of erythropoietin production. *Life Sci.* 1988, 43, 229–237.
- (4) Nikodijevic O.; Sarges, R.; Daly, J. W.; Jacobson, K. A. Characterization of A₁- and A₂-adenosine receptor mediated components in locomotor depression elicited by adenosine agonists. *J. Pharmacol. Exp. Ther.* 1991, 259, 286–294.
- (5) Van Galen, P. J. M.; Stiles, G. L.; Michaels, G.; Jacobson, K. A. Adenosine A₁ and A₂ receptors: Structure-function relationships. *Med. Res. Rev.* 1992, 12, 423–471.
- (6) Choi, O. H.; Shamim, M. T.; Padgett, W. L.; Daly, J. W. Caffeine and theophylline analogues: correlation of behavioral effects with activity as adenosine receptor antagonists and as phosphodiesterase inhibitors. *Life Sci.* 1988, 43, 387–398.
- (7) Seale, T.; Abl, K. A.; Shamim, M. T.; Carney, J. M.; Daly, J. W. 3,7-Dimethyl-1-propargylxanthine: A potent and selective *in vivo* antagonist of adenosine analogs. *Life Sci.* 1988, 43, 1671–1684.
- (8) Jarvis, M. F.; Schutz, R.; Hutchison, A. J.; Do, E.; Sills, M. A.; Williams, M. [³H]CGS 21680, an A₂ selective adenosine receptor agonist directly labels A₂ receptors in rat brain tissue. *J. Pharmacol. Exp. Ther.* 1989, 251, 888–893.
- (9) Abiru, T.; Miyashita, T.; Watanabe, Y.; Yamaguchi, T.; Machida, H.; Matsuda, A. Nucleosides and nucleotides. 107. 2-(Cycloalkylalkynyl)adenosines: Adenosine A₂ receptor agonists with potent antihypertensive effects. *J. Med. Chem.* 1992, 35, 2253–2260.
- (10) Shamim, M. T.; Ukena, D.; Padgett, W. L.; Daly, J. D. Effects of 8-phenyl and 8-cyclohexyl substituents on the activity of mono-, di-, and trisubstituted alkylxanthines with substitution at the 1-, 3-, and 7-positions. *J. Med. Chem.* 1989, 32, 1231–1237.
- (11) Shimada, J.; Suzuki, F.; Nonaka, H.; Ishii, A.; Ichikawa, S. (E)-1,3-Dialkyl-7-methyl-8-(3,4,5-trimethoxystyryl)xanthines: Potent and selective adenosine A₂ antagonists. *J. Med. Chem.* 1992, 35, 2342–2345.

- (12) Francis, J. E.; Cash W. D.; Psychoyos, S.; Ghai, G.; Wenk, P.; Friedmann, R. C.; Atkins, C.; Warren, V.; Furness, P.; Hyun, J. L.; et al. Structure-activity profile of a series of novel triazoloquinazoline adenosine antagonists. *J. Med. Chem.* 1988, *31*, 1014-1020.
- (13) Sarges, R.; Howard, H. R.; Browne, R. G.; Lebel, L. A.; Seymour, P. A.; Koe, B. K. 4-Amino[1,2,4]triazolo[4,3-a]quinoxalines. A novel class of potent adenosine receptor antagonists and potential rapid-onset antidepressants. *J. Med. Chem.* 1990, *33*, 2240-2254.
- (14) James, R.; Stutchbury, N. C. J. Imidazo[1,5-a]triazin-4-ones: The use of electrostatic potentials in the design of adenosine antagonists. Abstract presented at the 203rd American Chemical Society Meeting, San Francisco, 1992.
- (15) Ji, X.-D.; Stiles, G. L.; van Galen, P. J. M.; Jacobson, K. A. Characterization of human striatal A₂-adenosine receptors using radioligand binding and photoaffinity labeling. *J. Receptor Res.* 1992, *12*, 149-169.
- (16) Becker, E. D. *High Resolution NMR: Theory and Chemical Applications*; Academic Press: New York, 1980.
- (17) Schwabe, U.; Trost, T. Characterization of adenosine receptors in rat brain by (-)[³H]N⁶-phenylisopropyladenosine. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 1980, *313*, 179-187.
- (18) Cheng, Y. C.; Prusoff, W. H. Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 percent inhibition (IC₅₀) of an enzyme reaction. *Biochem. Pharmacol.* 1973, *22*, 3099-3108.
- (19) Erickson, R. H.; Hiner, R. N.; Feeney, S. W.; Blake, P. R.; Rzeszutarski, W. J.; Hicks, R. P.; Costello, D. G.; Abreu, M. E. 1,3,8- Trisubstituted xanthines. Effects of substitution pattern upon adenosine receptor A1/A2 affinity. *J. Med. Chem.* 1991, *34*, 1431-1435.
- (20) Jacobson, K. A.; Barone, S.; Kammula, U.; Stiles, G. L. Electrophilic derivatives of purines as irreversible inhibitors of A1-adenosine receptors. *J. Med. Chem.* 1989, *32*, 1043-1051.
- (21) Ukena, D.; Jacobson, K. A.; Padgett, W.; Ayala, C.; Shamim, M. T.; Kirk, K. L.; Daly, J. W. Species differences in structure-activity relationships of adenosine agonists and xanthine antagonists at brain A1 adenosine receptors. *FEBS Lett.* 1986, *209*, 122-128.
- (22) Stone, G. A.; Jarvis, M. F.; Sills, M. A.; Weeks, B.; Snowhill, E. W.; Williams, M. Species differences in high-affinity adenosine A₂ binding sites in striatal membranes from mammalian brain. *Drug Dev. Res.* 1988, *15*, 31-46.
- (23) Barrington, W. W.; Jacobson, K. A.; Hutchison, A. J.; Williams, M.; Stiles, G. L. Identification of the A₂ adenosine receptor binding subunit by photoaffinity crosslinking. *Proc. Natl. Acad. Sci. U.S.A.* 1989, *86*, 6572-6576.
- (24) Ferré, S.; Fuxe, K.; von Euler, G.; Johansson, B.; Fredholm, B. B. Adenosine-dopamine interactions in the brain. *Neurosci.* 1992, *51*, 501-512.
- (25) Jacobson, K. A.; van Galen, P. J. M.; Williams, M. Adenosine receptors: Pharmacology, structure activity relationships and therapeutic potential. *J. Med. Chem.* 1992, *35*, 407-422.