

Synthesis and Opioid Activity of Dynorphin A-(1-13)NH₂ Analogues Containing *cis*- and *trans*-4-Aminocyclohexanecarboxylic Acid[†]

Kristin R. Snyder, Thomas F. Murray, Gary E. DeLander, and Jane V. Aldrich*

Oregon State University, College of Pharmacy, Corvallis, Oregon 97331

Received October 16, 1992

It has been proposed that the "message" sequence of dynorphin A (Dyn A) exists in an extended conformation in aqueous solution (Schiller, P. W. *Int. J. Pept. Protein Res.* 1983, 21, 307-312). Molecular modeling suggested that *trans*-4-aminocyclohexanecarboxylic acid (*trans*-ACCA) might function as a conformationally constrained replacement for Gly²-Gly³ of Dyn A in such an extended conformation. ACCA was synthesized by catalytic hydrogenation of *p*-aminobenzoic acid, and the *cis* and *trans* isomers were separated by fractional recrystallization. Analogues of Dyn A-(1-13)-NH₂ containing *cis*- and *trans*-ACCA were prepared by solid-phase peptide synthesis using the Fmoc chemical protocol. Results from radioligand binding assays indicated that the peptides have modest affinity for κ opioid receptors (K_i 's = 9.1 and 13.4 nM for [*cis*-ACCA²⁻³]- and [*trans*-ACCA²⁻³]Dyn A-(1-13)NH₂, respectively) and modest κ -receptor selectivity (K_i ratio ($\kappa/\mu/\delta$) = 1/13/210 and 1/21/103, respectively). [*cis*-ACCA²⁻³]- and [*trans*-ACCA²⁻³]Dyn A-(1-13)-NH₂ are the first reported Dyn A analogues constrained in the "message" sequence that are selective for κ receptors. The *cis*-ACCA analogue showed very weak opioid activity (IC₅₀ = 4.0 μ M) in the guinea pig ileum.

Introduction and Rationale

Dynorphin A (Dyn A), a 17-amino acid peptide, has been postulated to be an endogenous κ opioid receptor ligand.¹ Dyn A shares with other mammalian opioid peptides a common N-terminal tetrapeptide "message" sequence which is important for opioid activity, while containing a unique C-terminal "address" sequence which imparts selectivity for κ opioid receptors.² The shortened Dyn A-(1-13), with the sequence Tyr-Gly-Gly-Phe-Leu-Arg-Arg-Ile-Arg-Pro-Lys-Leu-Lys, accounts for essentially all of the biological activity of Dyn A in the guinea pig ileum (GPI) assay.³

Since Dyn A is a linear peptide, it can adopt a number of possible conformations in solution, with the conformation dependent on the environment. In aqueous solution results from several spectral techniques, circular dichroism,^{4,5} IR spectroscopy,^{6,7} Raman spectroscopy,⁸ and NMR,^{7,9,10} are consistent with a random coil and/or extended β -strand conformation for Dyn A-(1-13). Fluorescent energy transfer experiments with [Trp⁴]Dyn A-(1-13) in dilute aqueous solution suggested that the N-terminal portion of Dyn A is in an extended conformation.¹¹ On the planar surface of a neutral lipid membrane,⁶ or in the presence of sodium dodecyl sulfate,⁴ however, Dyn A-(1-13) appears to adopt an α -helical structure. Binding to an anionic phospholipid⁹ or the addition of cerebroside sulfate⁵ does not induce the helical structure in the peptide.

The question remains of whether a conformation observed in solution reflects the conformation Dyn A

adopts at opioid receptors. Preparing conformationally constrained derivatives is another way to examine the biologically active conformations of a peptide. The two reported analogues of Dyn A constrained in the "message" sequence, cyclo[D-Cys²,Cys⁵]Dyn A-(1-13)¹² and cyclo-[D-Orn²,Asp⁵]Dyn A-(1-8),¹³ show high potency in the GPI, but also exhibit high μ -receptor affinity. The other reported cyclic analogues of Dyn A, cyclized via either a lactam¹³ or a disulfide,¹⁴ are constrained in the C-terminal "address" portion of the peptide.

Since the N-terminal "message" sequence appears to be important for the opioid activity of Dyn A, we incorporated conformational constraints into this region of the peptide. Tyr¹ and Phe⁴ residues are important for opioid activity and potency,² so conformational constraints involving Gly² and/or Gly³, which would affect the relative orientation of the aromatic rings at positions 1 and 4, were examined. Since several of the conformational studies described above, particularly Schiller's fluorescent energy transfer experiments,¹¹ suggested an extended conformation for the N-terminus of Dyn A, a conformational constraint consistent with this proposed conformation was chosen for preparation.

Results and Discussion


Design Rationale and Synthesis. Molecular modeling with the AMBER^{15,16} program was used to examine possible conformational constraints for incorporation into positions 2 and 3 in Dyn A-(1-13)NH₂. These studies suggested that *trans*-4-aminocyclohexanecarboxylic acid (*trans*-ACCA) might replace Gly²-Gly³ in an extended conformation. The calculated nitrogen-carbonyl carbon distance was 5.70 Å for *trans*-ACCA in the diequatorial conformation vs 6.12 Å between the nitrogen of Gly² and the carbonyl carbon of Gly³ when the peptide was in an extended conformation. The ACCA dipeptide replacement is equivalent to constraining ψ_2 and ϕ_3 while still allowing free rotation around ϕ_2 and ψ_3 .

ACCA was synthesized by hydrogenation of *p*-aminobenzoic acid with PtO₂ as a catalyst¹⁷ (Scheme I), which

* Author to whom correspondence should be addressed.

[†] Abbreviations used for amino acids follow the rules of the IUPAC-IUB Commission of Biochemical Nomenclature outlined in *J. Biol. Chem.* (1972, 247, 977-983). Amino acids are in the L-configuration. Additional abbreviations used are as follows: ACCA, 4-aminocyclohexanecarboxylic acid; Aib, α -aminoisobutyric acid; DAMGO, [D-Ala²,MePhe⁴,glyol]-enkephalin; DPDPE, [D-Pen²,D-Pen³]enkephalin; Dyn A, dynorphin A; FAB-MS, fast atom bombardment mass spectrometry; Fmoc, 9-fluorenylmethoxycarbonyl; Fmoc-ON-Su, N-(9-fluorenylmethoxycarbonyl)-succinimide; GPI, guinea pig ileum; TFA, trifluoroacetic acid; THF, tetrahydrofuran.

distance was 5.70 Å for *trans*-ACCA in the diequatorial conformation vs 6.12 Å between the nitrogen of Gly² and the carbonyl carbon of Gly³ when the peptide was in an extended conformation. The ACCA dipeptide replacement is equivalent to constraining ψ_2 and ϕ_3 while still allowing free rotation around ϕ_2 and ψ_3 .




ACCA was synthesized by hydrogenation of *p*-aminobenzoic acid with PtO₂ as a catalyst¹⁷ (Scheme I), which yielded a mixture of *cis* and *trans* isomers (approx ratio of 2.5/1 *cis/trans*). Separation by fractional recrystallization¹⁸ from EtOH yielded the *cis* isomer, which was pure by ¹H NMR. Subsequent recrystallization from EtOH/ether yielded the *trans* isomer, but ¹H NMR indicated that the *trans* isomer contained from 10–15% up to 35% of the *cis* isomer, depending on the batch. Following fractional recrystallization, each isomer was converted separately to its Fmoc (9-fluorenylmethoxycarbonyl) derivative. Fmoc protection of these branched amino acids proved to be difficult, and literature procedures¹⁹ had to be modified²⁰ to obtain the desired product (see Experimental Section). HPLC verified that Fmoc-*cis*-ACCA contained only the *cis* isomer. In the case of Fmoc-*trans*-ACCA, the contaminating *cis* isomer (*t*_R = 29.3 min) was not well resolved from the desired Fmoc-*trans*-ACCA (*t*_R = 29.7 min) by HPLC, so this mixture was used in the synthesis of the *trans*-ACCA analogue of Dyn A-(1-13)-NH₂.

Both the *cis* and *trans* isomers were incorporated separately into Dyn A-(1-13)NH₂ (Scheme I). The peptides were prepared as amides because of the enhanced metabolic stability of Dyn A-(1-13) amide vs its corresponding acid.²¹ The peptides were synthesized on a PAL resin²² using Fmoc-protected amino acids by procedures described previously.²³ The peptides were deprotected and cleaved from the PAL resin using trifluoroacetic acid (TFA) and purified by preparative reverse-phase HPLC. The purification of [*cis*-ACCA²⁻³]Dyn A-(1-13)NH₂ was straightforward, while the purification of the *trans* analogue proved difficult due to the necessity of separating it from the contaminating *cis* analogue. This resulted in a very low yield of pure [*trans*-ACCA²⁻³]Dyn A-(1-13)-NH₂.

Opioid Receptor Binding Affinities and Opioid Activity. The peptides were evaluated for opioid receptor affinity at κ receptors by measuring the inhibition of binding of [³H]bremazocine to guinea pig cerebellar membranes, and for μ and δ receptor affinities in rat forebrain membranes using [³H]DAMGO ([D-Ala²-MePhe⁴,glyol]enkephalin) and [³H]DPDPE ([D-Pen²,D-Pen⁵]enkephalin), respectively (Table I).

Both [*cis*-ACCA²⁻³]- and [*trans*-ACCA²⁻³]Dyn A-(1-13)NH₂ bound to κ opioid receptors with modest affinity (*K*_i = 9–13 nM), with the *cis*-ACCA analogue exhibiting slightly greater affinity than the *trans* isomer. The affinity of these analogues for κ receptors is 1/60 to 1/90 that of the

Scheme I



parent peptide Dyn A-(1-13)NH₂. Introduction of either isomer of ACCA into Dyn A-(1-13)NH₂ causes an even larger decrease in affinity for μ receptors, resulting in *K*_i's greater than 100 nM. Therefore, both of these analogues are κ selective, with [*trans*-ACCA²⁻³]Dyn A-(1-13)NH₂ showing slightly better κ selectivity (κ/μ ratio = 1/21) than [*cis*-ACCA²⁻³]Dyn A-(1-13)NH₂ (κ/μ ratio = 1/13). Both of these peptides had very little affinity for δ receptors (*K*_i's > 1000 nM).

Both the *cis*- and *trans*-ACCA Dyn A analogues show binding affinity similar to the Dyn A-(1-13)NH₂ analogues containing an L-amino acid at position 2 reported by Story et al.,²⁴ all of which had *K*_i's in the 2–20 nM range. The ACCA-substituted peptides, however, have better κ vs μ selectivity than any of these 2-substituted dynorphin analogues. [Aib²]Dyn A-(1-13),²⁵ which incorporates an α,α -disubstituted amino acid into position 2 similar to the disubstitution α to the amine of ACCA, surprisingly has lower affinity for κ receptors (*K*_i = 50 nM). [Aib²]Dyn A-(1-13) has 10–30-fold higher affinity for μ receptors (*K*_i = 10 nM) than [*cis*- and [*trans*-ACCA²⁻³]Dyn A-(1-13)-NH₂; thus, the Aib²-substituted peptide is μ selective while the ACCA-substituted peptides are κ selective.

[*cis*-ACCA²⁻³]Dyn A-(1-13)NH₂ was evaluated for opioid activity in the guinea pig ileum (GPI) assay. Its potency (IC₅₀ = 4.09 μ M, 95% confidence = 2.88–5.80 μ M) was much lower than the parent Dyn A-(1-13)NH₂ (IC₅₀ = 0.24 nM, 95% confidence = 0.21–0.29 nM), but this analogue exhibited a full dose-response curve and naloxone antagonized its effects (data now shown). The low GPI activity of [*cis*-ACCA²⁻³]Dyn A-(1-13)NH₂ parallels the results obtained for the analogues containing L-amino acids in position 2.²⁴ There was insufficient *trans* compound to allow for its testing in the GPI.

Conclusions

[*cis*- and *trans*-ACCA²⁻³]Dyn A-(1-13)NH₂ are the first reported Dyn A analogues conformationally constrained in the "message" sequence that are selective for κ opioid receptors. The *cis*- and *trans*-ACCA²⁻³-substituted pep-

Table I. Opioid Receptor Binding Affinities of Dyn A-(1-13)NH₂ and [ACCA]Dyn A-(1-13)NH₂ Analogues

analogue	K _i (nM)			ratio κ/μ/δ ^a
	[³ H]bremazocine (κ)	[³ H]DAMGO (μ)	[³ H]DPDPE (δ)	
Dyn A-(1-13)NH ₂	0.146 ± 0.004	0.194 ± 0.002	3.88 ± 0.09	1/1.3/27
[<i>cis</i> -ACCA ²⁻³]Dyn A-(1-13)NH ₂	9.10 ± 0.92	117 ± 2	1910 ± 65	1/13/210
[<i>trans</i> -ACCA ²⁻³]Dyn A-(1-13)NH ₂	13.4 ± 0.6	275 ± 2	1370 ± 53	1/21/103

^a Ratio of K_i/K_i/K_i.

and selectivity between the κ-selective [*cis*-ACCA²⁻³] and [*trans*-ACCA²⁻³]Dyn A-(1-13)NH₂ and the μ-selective Dyn A analogue [Aib²]Dyn A-(1-13)²⁵ could be due to differences in conformation, the larger size of ACCA vs Aib, or the substitution at the position corresponding to the C_α of Gly³ in the ACCA-substituted peptides.

The good discrimination of κ vs μ receptors exhibited by the *cis*- and *trans*-ACCA derivatives of Dyn A-(1-13)-NH₂ is encouraging and suggests that further modification at positions 2 and/or 3 might lead to Dyn A analogues with higher κ receptor affinity and opioid potency.

Experimental Section

Materials. The reagents and instrumentation used were those described previously.²³ Elemental analysis was performed by MWH Laboratories, Phoenix, AZ. NMR spectroscopy was done on a 400-MHz Bruker NMR in the Department of Chemistry at Oregon State University.

Hydrogenation of *p*-Aminobenzoic Acid to ACCA.¹⁷ *p*-Aminobenzoic acid (Aldrich, 1.056 g, 7.71 mmol) in 30% EtOH (100 mL) was hydrogenated under 50 psi of H₂, with PtO₂ (278 mg) as a catalyst, at room temperature overnight. This was repeated six times and the solutions combined. After removing the platinum by filtration, the filtrate was concentrated in vacuo until crystals appeared. The crystals were dissolved in H₂O and boiled, and the solution was treated with charcoal. ACCA was precipitated by the addition of 55 mL of EtOH/ether (10/1), followed by additional ether until no more crystals appeared. This yielded 4.916 g (70.7%) of ACCA as a mixture of *cis* and *trans* isomers; TLC: *R*_f (C₁₈, reverse phase, MeOH/H₂O, 60/40) 0.33.

Separation of *Cis* and *Trans* Isomers of ACCA.¹⁸ The crystals were dissolved in water, and EtOH was added to crystallize *cis*-ACCA. The crystals were collected, and the process was repeated three times to obtain additional *cis* isomer. The remaining filtrate was concentrated and 55 mL of EtOH/ether (10/1) added to crystallize the *trans*-ACCA. The crystals were collected, and the process was repeated three times with the addition of excess ether. The isomers were then recrystallized from H₂O/EtOH (*cis*) or H₂O/EtOH/ether (*trans*) to yield 3.125 g (44.9%) of *cis*-ACCA and 1.259 g (18.1%) of *trans*-ACCA. Data for *cis*-ACCA follow. Mp: 258–264 °C. FAB-MS: *m/z* 144 (M + 1). ¹H NMR (*d*-TFA): δ 1.76–2.34 (m, 8 H, -CH₂CH₂-), 2.85 (m, 1 H, CHCO₂H), 3.53 (m, 1 H, CHNH₂). Anal. (C₇H₁₃NO₂·0.4H₂O): C, H, N. HCl salt mp: 207–208 °C (lit. mp:¹⁸ 217 °C). Data for *trans*-ACCA follow. Mp: 262–267 °C. FAB-MS: *m/z* 144 (M + 1). ¹H NMR (*d*-TFA): δ 1.60–1.70 (m, 4 H, -CH₂-), 2.33 (m, 4 H, -CH₂-), 2.52 (m, 1 H, CHCO₂H), 3.46 (m, 1 H, CHNH₂). Anal. (C₇H₁₃NO₂·0.4H₂O): C, H, N. HCl salt mp: 272–273 °C (lit. mp:¹⁸ 273 °C).

Fmoc-*cis* and Fmoc-*trans*-ACCA.²⁰ *cis*-ACCA (393 mg, 2.73 mmol) and NaOH (109 mg, 2.73 mmol) were dissolved in H₂O (1.97 mL). A suspension of Fmoc-ON-Su (Bachem, 737 mg, 2.18 mmol) in THF (1.97 mL) was added to the ACCA solution, followed by additional H₂O and THF (1.97 mL of each). After 5 min NaHCO₃ (229 mg, 2.73 mmol) and THF (1.97 mL) were added; all reactants went into solution after 25 min. The reaction was stirred until TLC (CHCl₃/MeOH/AcOH, 89/10/1) indicated that all of the Fmoc-ON-Su had reacted (overnight). The solution was then acidified with excess 1.5 N HCl and the product, which precipitated out, dissolved in EtOAc. The HCl layer was extracted with additional EtOAc. The combined EtOAc extracts were then washed with 1.5 N HCl (2×), H₂O (2×), and saturated NaCl and dried over anhydrous MgSO₄ or Na₂SO₄. The solvent was

evaporated in vacuo to give a thick oil. Dissolution in EtOAc, precipitation with hexane, and evaporation of solvent yielded 737 mg (92.6%) of Fmoc-*cis*-ACCA as a glassy solid. Mp: 128–130 °C. FAB-MS: *m/z* 366 (M + 1). HPLC²⁶ (0 to 75% B over 50 min, 1.5 mL/min): *t*_R = 29.3 min. ¹H NMR (*d*₆-DMSO): δ 1.40–1.58 (m, 6 H, -CH₂CH₂-), 1.85 (m, 2 H, -CH₂-), 2.38 (br s, 1 H, CHCO₂H), 3.41 (m, 1 H, -CHNH-), 4.22 (m, 3 H, Fmoc-CHCH₂O-), 7.29 (d, 1 H, -NH-), 7.31 (t, 2 H, Ar), 7.40 (t, 2 H, Ar), 7.69 (d, 2 H, Ar), 7.87 (d, 2 H, Ar). Anal. (C₂₂H₂₃NO₄): C, H, N.

Protection of the *trans* isomer (429 mg, 3 mmol) as described for *cis*-ACCA yielded 760 mg (86.7%) of Fmoc-*trans*-ACCA. FAB-MS: *m/z* 366 (M + 1). HPLC²⁶ (0 to 75% B over 50 min, 1.5 mL/min): *t*_R = 29.7 min (Fmoc-*trans*-ACCA, 66%), *t*_R = 29.3 min (Fmoc-*cis*-ACCA, 33%). Data for an analytical sample recrystallized from THF/H₂O follow. Mp: 220–227 °C. ¹H NMR (*d*₆-DMSO): δ 1.19–1.35 (m, 4 H, -CH₂CH₂-), 1.85–1.94 (m, 4 H, -CH₂CH₂-), 2.08 (br t, 1 H, CHCO₂H), 3.18 (m, 1 H, -CHNH-), 4.20 (m, 1 H, Fmoc-CHCH₂O-), 4.26 (m, 2 H, Fmoc-CHCH₂O-), 7.22 (d, 1 H, -NH-), 7.32 (t, 2 H, Ar), 7.40 (t, 2 H, Ar), 7.67 (d, 2 H, Ar), 7.87 (d, 2 H, Ar). Anal. (C₂₂H₂₃NO₄): C, H, N.

Peptide Synthesis Using Fmoc-Protected Amino Acids. Both peptides were synthesized, cleaved from the resin, and purified by procedures described previously.²³ Before removing the Fmoc group from ACCA, the entire coupling sequence for that amino acid was repeated to ensure complete coupling. Following cleavage from the resin the peptides were purified by preparative reverse-phase HPLC²⁷ (0 to 50% B over 75 min, 20 mL/min for the *cis* analogue, and 15 to 45% B over 60 min, 20 mL/min for the *trans* analogue). Data for [*cis*-ACCA²⁻³]Dyn A-(1-13)NH₂ follow. Yield: 30 mg (24%). FAB-MS: *m/z* 1614 (M + 1). HPLC²⁶ (0 to 75% B over 50 min, 1.5 mL/min): *t*_R = 19.5 min. Amino acid analysis: Tyr (1) 0.93, Phe (1) 0.99, Leu (2) 2.02, Arg (3) 3.15, Ile (1) 0.97, Pro (1) 0.97, Lys (2) 2.01. Data for [*trans*-ACCA²⁻³]Dyn A-(1-13)NH₂ follow. Yield: 4.4 mg (3.3%). FAB-MS: *m/z* 1614 (M + 1). HPLC²⁶ (15 to 35% B over 40 min, 1.5 mL/min): *t*_R = 14.4 min. Amino acid analysis: Tyr (1) 0.79, Phe (1) 1.05, Leu (2) 2.02, Arg (3) 3.12, Ile (1) 0.99, Pro (1) 1.02, Lys (2) 2.02.

Binding and Guinea Pig Ileum Assays. Guinea pig cerebellar membranes and rat forebrain membranes were prepared and binding assays performed as previously described,²⁴ except that 100 nM DAMGO was included in the incubation mixtures of [³H]bremazocine with guinea pig cerebellar membranes (κ binding assays). Guinea pig ileum assays were performed as previously described.²⁴ In experiments involving naloxone, the antagonist was added to the tissue bath 10 min prior to determination of the agonist dose-response curve.

Acknowledgment. This research was supported by a grant from the National Institute on Drug Abuse (R01 DA05195). We would like to acknowledge the technical assistance of Valerie Caldwell for binding assays and Martin Knittle for guinea pig ileum assays. We would also like to thank Dwight Weller for use of the VAX workstation and AMBER software and for his assistance in learning and using AMBER.

References

- Chavkin, C.; James, I. F.; Goldstein, A. Dynorphin Is a Specific Endogenous Ligand of the κ Opioid Receptor. *Science* 1982, 215, 413–415.
- Chavkin, C.; Goldstein, A. Specific Receptor for the Opioid Peptide Dynorphin: Structure-Activity Relationships. *Proc. Natl. Acad. Sci. U.S.A.* 1981, 78, 6543–6547.

- (3) Goldstein, A.; Tachibana, S.; Lowney, L. I.; Hunkapiller, M.; Hood, L. Dynorphin-(1-13), an Extraordinarily Potent Opioid Peptide. *Proc. Natl. Acad. Sci. U.S.A.* 1979, 76, 6666-6670.
- (4) Maroun, R.; Mattice, W. L. Solution Conformations of the Pituitary Opioid Peptide Dynorphin-(1-13). *Biochem. Biophys. Res. Commun.* 1981, 103, 442-446.
- (5) Wu, C.-S. C.; Lee, N. M.; Loh, H. H.; Yang, J. T. Competitive Binding of Dynorphin-(1-13) and β -Endorphin to Cerebroside Sulfate in Solution. *J. Biol. Chem.* 1986, 261, 3687-3691.
- (6) Erne, D.; Sargent, D. F.; Schwyzer, R. Preferred Conformation, Orientation, and Accumulation of Dynorphin A-(1-13)-tridecapeptide on the Surface of Neutral Lipid Membranes. *Biochemistry* 1985, 24, 4261-4263.
- (7) Renugopalakrishnan, V.; Rapaka, R. S.; Huang, S.-G.; Moore, S.; Hutson, T. B. Dynorphin A (1-13) Peptide NH Groups are Solvent Exposed; FT-IR and 500 MHz ^1H NMR Spectroscopic Evidence. *Biochem. Biophys. Res. Commun.* 1988, 151, 1220-1225.
- (8) Rapaka, R. S.; Renugopalakrishnan, V.; Collette, T. W.; Dobbs, J. C.; Carreira, L. A.; Bhatnagar, R. S. Conformational Features of Dynorphin A-(1-13). *Laser Raman Spectroscopic Studies. Int. J. Pept. Protein Res.* 1987, 30, 284-287.
- (9) Lancaster, C. R. D.; Mishra, P. K.; Hughes, D. W.; St.-Pierre, S. A.; Bothner-By, A. A.; Epand, R. M. Mimicking the Membrane-Mediated Conformation of Dynorphin A-(1-13)-Peptide: Circular Dichroism and Nuclear Magnetic Resonance Studies in Methanolic Solution. *Biochemistry* 1991, 30, 4715-4726.
- (10) Zhou, N.; Gibbons, W. A. A ^1H Nuclear Magnetic Resonance Study of the Opioid Peptide Dynorphin-(1-13) in Aqueous Solution. *J. Chem. Soc. Perkin Trans. 2* 1986, 637-644.
- (11) Schiller, P. W. Fluorescence Study on the Solution Conformation of Dynorphin in Comparison to Enkephalin. *Int. J. Pept. Protein Res.* 1983, 21, 307-312.
- (12) Schiller, P. W.; Eggmann, B.; Nguyen, T. M.-D. Comparative Structure-Function Studies with Analogs of Dynorphin-(1-13) and [Leu⁵]Enkephalin. *Life Sci.* 1982, 31, 1777-1780.
- (13) Schiller, P. W.; Nguyen, T. M.-D.; Lemieux, C. Synthesis and Opioid Activity Profiles of Cyclic Dynorphin Analogs. *Tetrahedron* 1988, 44, 733-743.
- (14) Kawasaki, A. M.; Knapp, R. J.; Kramer, T. H.; Wire, W. S.; Vasquez, O. S.; Yamamura, H. I.; Burks, T. F.; Hruby, V. J. Design and Synthesis of Highly Potent and Selective Cyclic Dynorphin A Analogues. *J. Med. Chem.* 1990, 33, 1874-1879.
- (15) Weiner, P. K.; Kollman, P. A. AMBER: Assisted Model Building with Energy Refinement. A General Program for Modeling Molecules and Their Interactions. *J. Comput. Chem.* 1981, 2, 287-303.
- (16) Weiner, S. J.; Kollman, P. A.; Nguyen, D. T.; Case, D. A. An All Atom Force Field for Simulations of Proteins and Nucleic Acids. *J. Comput. Chem.* 1986, 7, 230-252.
- (17) Skaric, V.; Kovacevic, M.; Skaric, D. Synthesis of Peptides Containing *cis*- or *trans*-3- or 4-Aminocyclohexanecarboxylic Acid Residues. *J. Chem. Soc. Perkin Trans. 1* 1976, 1199-1201.
- (18) Ferber, E.; Bruckner, H. Uber das Isochinuclidin. *Chem. Ber.* 1943, 76, 1019-1027.
- (19) Milton, R. C. deL.; Becker, E.; Milton, S. C. F.; Baxter, J. E. J.; Elsworth, J. F. Improved Purities for Fmoc-amino Acids from Fmoc-ONSu. *Int. J. Pept. Protein Res.* 1987, 30, 431-432.
- (20) Anderson, H., Synthetech, Inc., Albany, OR, personal communication, 1991.
- (21) Leslie, F. M.; Goldstein, A. Degradation of Dynorphin-(1-13) by Membrane-Bound Rat Brain Enzymes. *Neuropeptides* 1982, 2, 185-196.
- (22) Albericio, F.; Barany, G. An Acid-Labile Anchoring Linkage for Solid-Phase Synthesis of C-Terminal Peptide Amides Under Mild Conditions. *Int. J. Pept. Protein Res.* 1987, 30, 206-216.
- (23) Snyder, K. R.; Story, S. C.; Heidt, M. E.; Murray, T. F.; DeLander, G. E.; Aldrich, J. V. Effect of Modification of the Basic Residues of Dynorphin A-(1-13) Amide on κ Opioid Receptor Selectivity and Opioid Activity. *J. Med. Chem.* 1992, 35, 4330-4333.
- (24) Story, S. C.; Murray, T. F.; DeLander, G. E.; Aldrich, J. V. Synthesis and Opioid Activity of 2-Substituted Dynorphin A-(1-13) Amide Analogues. *Int. J. Pept. Protein Res.* 1992, 40, 89-96.
- (25) Lemaire, S.; Parent, P.; Lapierre, C.; Michelot, R. N,N-Diallylated Analogs of Dynorphin A-(1-13) as Potent Antagonists for the Endogenous Peptide and Selective κ Opioid Analgesics. In *The International Narcotics Research Conference (INRC) '89, Progress in Clinical and Biological Research*; Quirion, R., Jhamandas, K., Gianoulakis, C., Eds.; Alan R. Liss, Inc.: New York, 1990; Vol. 328, pp 45-48.
- (26) Vydac 214TP54 column, solvent A = 0.1% TFA in H₂O, solvent B = 0.1% TFA in AcCN.
- (27) Protein Plus preparative column, solvents A and B are as above.