

The Importance of Micelle-Bound States for the Bioactivities of Bifunctional Peptide Derivatives for δ/μ Opioid Receptor Agonists and Neurokinin 1 Receptor Antagonists

Takashi Yamamoto,[†] Padma Nair,[†] Neil E. Jacobsen,[†] Peg Davis,[‡] Shou-wu Ma,[‡] Edita Navratilova,[‡] Sharif Moye,[‡] Josephine Lai,[‡] Henry I. Yamamura,^{‡,▽} Todd W. Vanderah,[‡] Frank Porreca,[‡] and Victor J. Hruby^{*,†}

Department of Chemistry, University of Arizona, 1306 East University Boulevard, Tucson, Arizona 85721, Department of Pharmacology, University of Arizona, 1501 North Campbell Avenue, Tucson, Arizona 85724

Received April 4, 2008

To provide new insight into the determining factors of membrane-bound peptide conformation that might play an important role in peptide–receptor docking and further biological behaviors, the dodecylphosphocholine (DPC) micelle-bound conformations of bifunctional peptide derivatives of δ -preferring opioid agonists and NK1 antagonists (**1**: Tyr-D-Ala-Gly-Phe-Met-Pro-Leu-Trp-O-3,5-Bzl(CF₃)₂; **2**: Tyr-D-Ala-Gly-Phe-Met-Pro-Leu-Trp-NH-3,5-Bzl(CF₃)₂; **3**: Tyr-D-Ala-Gly-Phe-Met-Pro-Leu-Trp-NH-Bzl) were determined based on 2D NMR studies. Although the differences in the primary sequence were limited to the C-terminus, the obtained NMR conformations were unexpectedly different for each compound. Moreover, their biological activities showed different trends in direct relation to the compound-specific conformations in DPC micelles. The important result is that not only were the NK1 antagonist activities different (the pharmacophore located at the C-terminus) but the opioid agonist activities (this pharmacophore was at the structurally preserved N-terminus) also were shifted, suggesting that a general conformational change in the bioactive state was induced due to relatively small and limited structural modifications.

Introduction

The importance of peptide and protein conformation has become increasingly appreciated recently, as their biological importance has been widely recognized. One notable example is protein misfolding, which has been implicated in a large number of diseases such as protein folding disorders (PFD).^{1–3} In the diseases known as amyloidoses,³ large quantities of misfolded proteins or peptides undergo aggregation, resulting in destroying brain cells and other tissues. These diseases, such as Alzheimer's disease and Parkinson's disease, have high rates of pathogenesis and give rise to the huge social problems all over the world. The Alzheimer's β -amyloid protein (A β P) is the well-known offending substance whose conformational change to the β -sheet oligomer and subsequent aggregation was shown to enhance its neurotoxicity.^{3–5} Another important misfolded protein is the cellular prion protein PrP^C, whose secondary structure was changed into a β -rich conformation (PrP^{Sc}) to cause the prion diseases.⁶ Thus, many scientific efforts have been made to seek the origin of such misfoldings.^{7–10} However, the full picture of their mechanisms is still largely unknown.

The conformation of small and linear peptides, such as the endogenous opioid peptide enkephalin¹¹ and its analogues,^{12,13} have been thought to have "random" conformations in aqueous solution due to their high flexibility.¹⁴ However, in the presence of trifluoroethanol (TFE) or membrane-mimicking surroundings, they could have structured conformations because their secondary structural elements can be stabilized by the environmental effects.^{12,13,15,16} Therefore, the circumstances surrounding such

a peptide should be important especially for their conformational definitions.¹⁷

Many G-protein coupled receptors (GPCRs), which are the typical membrane-bound proteins, have their ligand binding sites in or near the lipophilic trans-membrane (TM) domains.^{18–21} Because the docking event of such a receptor and a ligand must take place near the membrane, ligand–membrane interactions should be important and this research topic has been occasionally explored for decades.^{22,23} In fact, it has been pointed out that the membrane promotes ligand–receptor docking^{24–27} and that the ligand adoption into membrane followed by membrane-catalyzed 2D search should be more efficient than the three-dimensional ligand–receptor binding through solvent space.^{12,13,23,24,28} Hence, understanding ligand–membrane interactions and membrane-bound structures of ligands is indispensable for further insight into their diverse biological behaviors.^{12,13,28}

The opioid and NK1^a receptors are both GPCRs which have significant importance in pain signal transmission, and their

^a Abbreviations: Abbreviations used for amino acids and designation of peptides follow the rules of the IUPAC-IUB Commission of Biochemical Nomenclature in *J. Biol. Chem.* **1972**, *247*, 977–983. The following additional abbreviations are used: AcOH, acetic acid; Boc, *tert*-butoxycarbonyl; BuOH, butanol; Bzl, benzyl; BSA, bovine serum albumin; CCK, cholecystokinin; Cl-HOBT, 1-hydroxy-6-chlorobenzotriazole; CHO, Chinese hamster ovary; DMEM, Dulbecco's modified Eagle's medium; DCM, dichloromethane; DIEA, diisopropylethylamine; DMF, *N,N*-dimethylformamide; DMSO, dimethylsulfoxide; DOR, δ opioid receptor; DPC, Dodecylphosphocholine; DPDPE, c[D-Pen²,D-Pen⁵]-enkephalin; DQF-COSY, double quantum filtered correlation spectroscopy; DAMGO, [D-Ala²,NMePhe⁴,Gly⁵-ol]-enkephalin; ESI, electrospray ionization; Fmoc, fluorenylmethoxycarbonyl; GPI, guinea pig isolated ileum; HCTU, 1H-benzotriazolium-1-[bis(dimethylamino)methylene]-5-chloro-hexafluorophosphate-(1-),3-oxide; HEK, human embryonic kidney; HRMS, high-resolution mass spectroscopy; LMMP, longitudinal muscle/myenteric plexus; MOR, μ opioid receptor; MVD, mouse vas deferens; NK1, neurokinin-1; NOESY, nuclear Overhauser enhancement; rmsd, root-mean-square deviation; RP-HPLC, reverse phase high performance liquid chromatography; SAR, structure–activity relationships; SEM, mean standard error; SPPS, solid phase peptide synthesis; TFA, trifluoroacetic acid; TOCSY, total correlation spectroscopy; Trp-NH-3,5-Bzl(CF₃)₂, 3',5'-(bistrifluoromethyl)-benzyl amide of tryptophan; Trp-NH-Bzl, benzyl amide of tryptophan; Trp-O-3,5-Bzl(CF₃)₂, 3',5'-(bistrifluoromethyl)-benzyl ester of tryptophan.

* To whom correspondence should be addressed. Phone: (520)-621-6332. Fax: (520)-621-8407. E-mail: hruby@email.arizona.edu.

[†] Department of Chemistry, University of Arizona.

[‡] Department of Pharmacology, University of Arizona.

[▽] Dedicated to the memory of Professor Henry "Hank" I. Yamamura our dear friend, colleague and teacher who passed away peacefully on the evening of September 4, 2008.




Figure 1. Sequence of the bifunctional ligands synthesized and examined.

ligand binding sites are found in or near the TM domains.^{20,21} The opioid receptors are classified into three subtypes (μ , δ , and κ) and have endogenous peptides as their ligands such as endomorphin, enkephalin, and dynorphin.^{20,29,30} It is well-known that the 11-amino acid peptide substance P acts as an excitatory and pronociceptive neurotransmitter of pain signaling through the neurokinin-1 NK1 receptor.^{31–37} Recently, combining the agonist effect at the opioid receptors together with the blocking of signals through NK1 receptors has been reported to have several merits for the analgesic effects such as enhanced potency and the prevention of opioid-induced tolerance.^{31–37} Thus, we have been developing in one molecule bifunctional peptide derivatives that act as opioid agonists and NK1 antagonists, with extensive discussions on their structure–activity relationships (SAR), which were based on their primary sequences.^{38–40} However, their three-dimensional conformations in membrane and compound–membrane interactions should also be important for further understanding the biological activities of ligands both in vitro and in vivo.

In the present article, we report our new findings that small and local structural modifications induced large conformational changes in membrane-mimicking dodecylphosphocholine (DPC) micelles,^{41–46} together with a discussion of their relevance to the biological activities. The starting point of our bifunctional design was the successful C-terminal modifications on TY001 (Tyr¹-D-Ala²-Gly³-Phe⁴-Pro⁵-Leu⁶-Trp⁷-O-3',5'-Bzl(CF₃)₂) to yield several potent bifunctional derivatives with better affinities at the μ opioid receptors rather than at the δ receptor.⁴⁰ Because the selective agonists at the δ opioid receptor have analgesic activity with fewer adverse effects, but less potency than μ -selective drugs,^{47–49} a δ -selective opioid agonist with enhanced analgesic activity, which also has NK1 receptor antagonist activity, provides a novel way to find a potent drug candidate for prolonged pain control. Thus, **1** (TY005: Tyr¹-D-Ala²-Gly³-Phe⁴-Met⁵-Pro⁶-Leu⁷-Trp⁸-O-3',5'-Bzl(CF₃)₂), which has a Met⁵ as the key amino acid residue for the δ -opioid selectivity and has been shown to have potent analgesic activities in vivo,^{38–40} was chosen as the current lead compound to find novel δ -preferring bifunctional ligands with strong analgesic potency. The C-terminus of **1** was modified to give two novel derivatives **2** (TY027: Tyr¹-D-Ala²-Gly³-Phe⁴-Met⁵-Pro⁶-Leu⁷-Trp⁸-NH-3',5'-Bzl(CF₃)₂) and **3** (TY025: Tyr¹-D-Ala²-Gly³-Phe⁴-Met⁵-Pro⁶-Leu⁷-Trp⁸-NH-Bzl) (Figure 1).

Our main intention for these structural modifications was to seek induced conformational changes in the presence of lipid media and to examine the influences of such changes on manipulating bioactivities. Therefore, the NMR conformations of **1–3** were determined in aqueous solution with membrane-mimicking perdeuterated dodecylphosphocholine (DPC) micelles. Fluorescence experiments as well as additional NMR studies using paramagnetic agents also were performed to determine the difference in membrane–compound interactions. Moreover, the biological activities of **2** and **3** were extensively evaluated using radioligand binding assays, GTP γ S binding assays, and tissue-based functional experiments using guinea

pig ileum (GPI) and mouse isolated vas deferens (MVD) tissues.^{38,40,50,51}

Results and Discussion

Peptide Synthesis. The peptide derivatives **1–3** (Figure 1) were synthesized using previously published methods.⁴⁰ Briefly, Boc-Tyr(tBu)-D-Ala-Gly-Phe-Met-Pro-Leu-Trp(Boc)-OH was synthesized using *N*^α-Fmoc chemistry with HBTU as the coupling reagent on a 2-chlorotrityl resin, followed by esterification or amidation of the protected C-terminal intermediate. The esterification was performed employing cesium carbonate to form the cesium salt of the protected peptide to react with 3',5'-bis(trifluoromethyl)benzyl bromide, and standard EDC/Cl-HOBt coupling chemistry with two equivalents of reactant amine was used for the amidation. Final deprotection with the cleavage cocktail (82.5% v/v TFA, 5% water, 5% thioanisole, 2.5% 1,2-ethanedithiol, and 5% phenol to quench the highly stabilized carbocations released from permanent protecting groups) gave the final crude peptides, which were purified by RP-HPLC (>99%) and characterized by analytical HPLC, ¹H NMR, HRMS, and TLC. The assignments of NMR resonances are available in the Supporting Information.

Secondary Structure Analysis Based on Assigned ¹H NMR. NMR experiments were performed to obtain structural information in DPC micelles as previously reported.^{45,46} Two-dimensional NMR studies including TOCSY, DQF-COSY, and NOESY, were performed on all three bifunctional peptide derivatives **1–3** in pH 4.5 buffer (45 mM CD₃CO₂Na/HCl, 1 mM NaN₃, 90% H₂O/10% D₂O) with 40-fold perdeuterated DPC micelles. At concentrations above the critical micelle point, DPC forms micelles with an aggregate number of 50–90, corresponding to one or two peptide molecules per micelle.⁵²

The obtained NOESY data showed the reasonably good quality as seen in Figure 2. A total of 136, 155, and 184 nonredundant NOE restraints were used for **1**, **2**, and **3**, respectively, based on the NOESY cross-peak volumes including sequential (50, 63, and 72, respectively), medium-range (2–4 residues; 31, 36, and 46, respectively), and long-range (1, 0, and 3, respectively) restraints. The distribution of these restraints along the peptide chain is shown in Figure 3D. Only one dihedral angle restraint was used: the Leu φ angle in **3**. The total numbers of restraints were 136, 155, and 185, respectively (15.1, 17.2, and 20.4 per residue). The large numbers of NOEs per residue for small and linear peptides suggest that the peptide derivatives exist in well-defined conformations in the DPC micelles.

The interresidual NOE connectivities and the ³*J*_{HN–H α} coupling constants of all of the peptide derivatives are illustrated in Figure 3. The C-terminal benzyl moiety of **2** and **3** are represented as residue 9. The ³*J*_{HN–H α} coupling constants for all residues in all three peptide derivatives were within the range of 6–8 Hz except for Leu⁷ in **3**. This is most likely due to conformational averaging of the peptides in solution.⁵³ The observed NOE patterns, including *d*_{NN}(*i*, *i* + 1), *d* _{α N}(*i*, *i* + 1) and some medium-range (*i*, *i* + 2 or 3) connectivities, suggest the possibility of β -turn structures around residues 1–4 in all three peptide derivatives as well as around residues 5–8 in **2**.⁵⁴ A few longer-range *d* _{α N}(*i*, *i* + 3) and *d* _{α N}(*i*, *i* + 4) connectivities found in **3** indicate the existence of a helical structure in this molecule, consistent with its H α CSI pattern (Figure 3C).⁵⁴

Structural Calculations. The 20 structures with the lowest total energies after rMD refinement were used to represent the structure of the peptide derivatives in DPC micelles. Throughout the ¹H NMR studies, only one major rotamer was found for




Figure 2. Fingerprint (H^N-H^{α}) region of the NOESY spectrum of (A) **1**, (B) **2**, and (C) **3** in DPC micelles. Intraresidue H^N-H^{α} NOE cross-peaks are labeled with residue numbers, and arrows indicate the connectivity path from *N*-terminal to *C*-terminal. X9 represents the cross-peaks derived from the corresponding *C*-terminal H^N and benzyl protons.




Figure 3. Diagram of H^N-H^{α} coupling constants, NOE connectivities, and H^{α} chemical shift index (CSI) for (A) **1**, (B) **2**, and (C) **3**. The H^{α} CSI was calculated using the random-coil values reported by Andersen et al.^{86,87} The residual interresidue NOE distance restraints of **1** (left), **2** (middle), and **3** (right) (D). Each column shows the sequential ($i, i+1$; open), medium-range ($i, i+2-4$; hatched) and long-range restraints ($i, i+>4$; filled), respectively. The residue Bzl or 9 stands for the respective *C*-terminal moieties.

peptides **1–3** and the populations of minor rotamers were all negligible. The Met⁵–Pro⁶ bond of the major rotamers were fixed in the *trans* configuration based on the observations of $^5H^{\alpha}$ to Pro⁶ H^{δ} sequential NOEs together with the absence of sequential $^5H^{\alpha}$ – $^6H^{\alpha}$ NOEs in the structural calculations of **1–3**. On the basis of the observations, *cis*-isomers of peptides **1–3** have negligible population in the lipid-mimicking DPC micelles and thus considered to have negligible effects on the biological behaviors. Because the experimental NOEs and $^3J_{HN-H^{\alpha}}$ were the averaged values in the NMR experiment time course, the

obtained low-energy structures did not correspond to a single conformation under these experimental conditions. However, the ensemble of an appropriate number of obtained structures provides sufficient structural figures based on the “averaged” or “dynamic” view of the conformations. Thus, statistics for the 20 best structures were performed as shown in Table 1. The average restraint violation energies were low (2.48, 2.95, and 1.13 kcal mol⁻¹ for **1**, **2**, and **3**, respectively), with average maximum NOE distance violations of 0.17, 0.11, and 0.11 Å with no dihedral angle violations. The 19 structures were aligned

Table 1. Structural Statistics

	compd 1		compd 2		compd 3	
	final 20 structures	most stable structure	final 20 structures	most stable structure	final 20 structures	most stable structure
rmsd from NOE dist restraints (\AA) ^a	0.025 ± 0.004	0.028	0.027 ± 0.004	0.016	0.016 ± 0.001	0.016
rmsd from backbone φ angle restraints (deg) ^b	c	c	c	c	0.0 ± 0.0	0.0
NOE dist restraints violations						
>0.01 \AA	13.9 ± 2.5	11	14.6 ± 1.5	14	14.2 ± 1.1	15
>0.1 \AA	2.5 ± 1.4	3	3.9 ± 1.2	3	1.0 ± 0.0	1
max dist violations (\AA)	0.17 ± 0.04	0.22	0.16 ± 0.02	0.13	0.11 ± 0.00	0.12
dihedral backbone angle violations						
>0.1°	c	c	c	c	0 ± 0	0
>1°	c	c	c	c	0 ± 0	0
max dihedral violations (deg)	c	c	c	c	0 ± 0	0
rms deviation from ideal geometry ^d						
bond length (\AA) ^e	0.0061 ± 0.0004	0.0063	0.0052 ± 0.0002	0.0052	0.0035 ± 0.00004	0.0035
bond valence angles (deg) ^f	2.14 ± 0.11	2.16	1.78 ± 0.05	1.72	1.25 ± 0.003	1.25
out-of-plane angles (deg) ^g	3.57 ± 0.63	3.25	2.73 ± 0.40	2.90	1.54 ± 0.09	1.45
AMBER energies (kcal mol ⁻¹)			9.08			
restraint ^h	2.48 ± 0.67	2.78	2.95 ± 0.58	2.41	1.13 ± 0.07	1.20
bond stretching	2.07 ± 0.22	2.20	1.42 ± 0.08	1.40	1.68 ± 0.02	1.70
bond angles	19.28 ± 1.84	19.49	12.8 ± 0.8	11.89	14.28 ± 0.22	14.00
dihedral angles	12.24 ± 1.9	9.54	9.57 ± 1.61	7.99	14.52 ± 0.38	14.53
planarity	1.63 ± 1.11	1.15	0.74 ± 0.33	0.61	0.19 ± 0.03	0.17
van der Waals ⁱ	-11.65 ± 3.1	-12.61	-12.23 ± 1.4	-13.80	-17.4 ± 0.75	-18.41
electrostatic ^j	-9.6 ± 0.91	-9.93	-11.5 ± 0.68	-11.84	-9.59 ± 0.41	-9.82
total	13.25 ± 2.12	9.08	-0.01 ± 2.25	-4.44	3.14 ± 0.78	1.61
atomic rmsd (\AA): final 19 structures vs most stable structure						
	compd 1		compd 2		compd 3	
	backbone atoms (N, C $^{\alpha}$, C')	all non-hydrogen atoms	backbone atoms (N, C $^{\alpha}$, C')	all non-hydrogen atoms	backbone atoms (N, C $^{\alpha}$, C')	all non-hydrogen atoms
calculated on whole molecule	1.80 ± 0.47	2.72 ± 0.92	1.14 ± 0.43	2.09 ± 0.64	0.19 ± 0.20	0.84 ± 0.28
calculated only on 1–4 res.	1.11 ± 0.54	2.49 ± 1.12	1.05 ± 0.63	2.16 ± 0.98	0.14 ± 0.30	0.32 ± 0.66
calculated only on 5–8 res. and C-terminus	0.75 ± 0.26	1.82 ± 0.90	0.45 ± 0.38	1.02 ± 0.25	0.04 ± 0.01	0.76 ± 0.42

^a The total number of NOE restraints were 136 for **1**, 155 for **2**, and 184 for **3**, respectively. ^b Two backbone φ angle restraints were applied only on **3**. ^c No restraints used. ^d Derived from the rMD calculations using the AMBER force field in DISCOVER. ^e The number of bond length were 160 for **1**, 161 for **2**, and 155 for **3**, respectively. ^f The number of bond valence angles were 285 for **1**, 287 for **2**, and 275 for **3**, respectively. ^g The number of out-of-plane angles were 36 for **1**, 36 for **2**, and 37 for **3**, respectively. ^h Calculated with force constants of 25 kcal mol⁻¹ \AA^{-2} and 100 kcal mol⁻¹ rad⁻² for the NOE distance and dihedral angle restraints, respectively. ⁱ Calculated with the Lennard-Jones potential using the AMBER force field and a 12 \AA cutoff. ^j Calculated with a distance-dependent dielectric constant ($\epsilon = 4r$).

with the most stable structure using all backbone atoms (Figure 4A), only the backbone atoms of residues 1–4 (Figure 4B), or 5–8 (Figure 4C). The backbone rmsd's of the 19 structures with respect to the most stable structure were 1.80, 1.14, and 0.19 \AA for **1**, **2**, and **3**, respectively, for all residues. The rmsd values are significantly decreased if alignment is carried out only on the backbone atoms of residues 5–8 (**1**: 0.75; **2**: 0.45; **3**: 0.04), indicating that the C-terminal half is much better defined by the NMR restraints than the N-terminal half (residues 1–4). This may be due to greater flexibility in the N-terminal portion. The decrease in rmsd going from a flexible ester (**1**) to a more rigid amide (**2**) linkage at the C-terminus was expected, but the much larger decrease resulting from removal of two trifluoromethyl groups of **2** (**3**) was surprising.

In Met-Enkephalin (Tyr¹-Gly²-Gly³-Phe⁴-Met⁵-OH) and Leu-Enkephalin (Tyr¹-Gly²-Gly³-Phe⁴-Leu⁵-OH), which form the basis for the design of the N-terminal portion of peptide sequence of **1–3**, a β -turn structure was often found between Tyr¹ and Phe⁴ by several methods including X-ray crystallography and NMR spectroscopy in environments which mimic the membrane bilayer.^{55,56} In the case of peptide derivatives **1** and **2**, a distance of less than 7 \AA between the C $_{\alpha}$ of D-Ala² and the C $_{\alpha}$ of Met⁵ was observed in 15 of the best 20 structures for **1**, and in all 20 structures for **2**, whereas the β -turn between Tyr¹ and Phe⁴ were found in only 5 and 3 structures in the best 20 structures of **1** and **2**, respectively (Table 2).⁵⁷ This result implied the existence of an alternative β -turn in the residues

2–5 for both **1** and **2**, and all of them were classified as type IV ("distorted") β -turn by their backbone dihedral angles. A second β -turn structure was found from Pro⁶ to the C-terminal benzyl moiety (residue 9) in which the distance between the C $_{\alpha}$ (Pro⁶) and the benzylic carbon (CH₂) of the C-terminus was less than 7 \AA in 17 and 19 of the best 20 structures for **1** and **2**, respectively (Table 2). For peptide derivative **2**, 6 of 19 turns found in **2** were classified as type I β -turns, and a hydrogen bond between the H^N of residue 9 and the carbonyl oxygen atom in Pro⁶ was observed in 9 of the 19 (Table 3). The C-terminal ester (**1**) showed only distorted β -turns in this region, with no hydrogen bonds, consistent with the larger backbone rmsd observed for C-terminal half (0.75 \AA vs 0.45 \AA for **2**). These implied that the secondary structure elements were consistent with the observed NOE connectivities (Figure 3).

Comparing the tandem β -turn structures of **1** and **2**, the peptide with no trifluoromethyl groups in the C-terminus (**3**) showed significantly different structural properties. First, the backbone of **3** had a well-defined helical structure, consistent with the NOE connectivities and CSI values (Figure 4). It is noteworthy that not only the C-terminal half-of **3**, but also its N-terminus was found to be quite structured (Table 1). Thus, removal of two trifluoromethyl groups from the C-terminus led to a drastic conformational change for the whole molecules from tandem β -turn structures to a helical structure (Figure 4).

This well-defined structure of **3** was also confirmed by the Ramachandran plot and angular order parameters (Figure 5).⁵⁸




Figure 4. Ensembles of the best 20 calculated structures in 40-fold DPC micelle/pH 4.5 buffer for (A) **1**, (B) **2**, and (C) **3** with the lowest restraint energy, aligned on backbone atoms of residues (a) 1–8, (b) 1–4, and (c) 5–8, from *N*-terminal (up in the left image) to *C*-terminal (down). Only backbone atoms were illustrated in (a) and (b) for easier comparison, and the most stable conformers (c) are shown with all non-hydrogen atoms.

Table 2. Number of Structures with Less than 7 Å Distance between α Carbons of *i*th and (*i* + 3)th Residues^a

	Tyr ¹ -Phe ⁴ residue	D-Ala ² -Met ⁵ residue	Gly ³ -Pro ⁶ residue	Met ⁵ -Trp ⁸ residue	Pro ⁶ -Bzl ⁹ residue
1	5	15	0	0	17
2	3	20	0	0	19
3^b	0	2	20	20	18

^a Out of the best 20 calculated structures. ^b Helical structure was found in which no β -turn structures should not be defined according to the original definition.⁵⁷ Bzl stands for the cross-peaks derived from the corresponding aromatic protons of benzyl moiety (residue 9).

Table 3. Observed Hydrogen Bonds^a

molecule	no. ^b	donor	acceptor	distance (Å) ^c	angle (deg) ^d
1	14	Leu ⁷ H ^N	Met ⁵ O	1.91 ± 0.07	141.1 ± 6.0
2	9	Bzl ⁹ H ^{Ne}	Pro ⁶ O	2.16 ± 0.11	158.5 ± 1.9
	7	Gly ³ H ^N	Tyr ¹ O	2.05 ± 0.11	137.8 ± 8.1
	5	Trp ⁸ H ^N	Met ⁵ O	2.04 ± 0.02	132.3 ± 1.1
3		no hydrogen bond observed			

^a The hydrogen bonds which were observed in more than five structures were listed. ^b The number of structures of the final 20 for which the listed hydrogen bond is observed. ^c The distance is the mean proton–oxygen distance (\pm SD) in the structures for which a hydrogen bond is observed.

^d The angle is the mean N–H...O angle (\pm SD) in the structures for which a hydrogen bond is observed. ^e Amide proton of *C*-terminal benzyl moiety.

In the Ramachandran plot of **3**, only seven clear spots, corresponding to the respective residues 2–8, were found. Among them, only Gly³ has positive φ angles in all of its 20 best structures. On the other hand, the corresponding Ramachandran plots for **1** and **2** showed more scattered views together with positive φ angles for Gly³ (10 structures in **1** and 3 structures in **2**), Phe⁴ (3 structures in **1**), Met⁵ (13 structures in **1** and all 20 structures in **2**), and Leu⁷ (1 structures in **1** and 6 structures in **2**) in the seven L-amino acids and some of D-Ala² (3 structures in **1** and 3 structures in **2**) have negative φ angles. It is interesting that Met⁵ of **1** and **2**, located between two β -turns, had frequent positive φ angles. As for the angular order parameters, both the parameters for φ and ψ angles in **3** were close to 1 in all the residues, whereas **1** and **2** had smaller values in some residues, implying a better-defined structure for **3** than

for **1** and **2**, especially in the *N*-terminus where no structural modification was made.

On the basis of the NMR structural analysis, it is clear that the limited modifications at the *C*-terminal moiety gave rise to several changes in the conformations of the peptide derivatives **1–3** in the presence of membrane-like DPC micelles. The change of ester (**1**) into amide (**2**) resulted in a better-defined conformation especially in the *C*-terminal portion of the peptide, and a more rigid β -turn structural element with intramolecular hydrogen-bond was found for residues 6–9 of **2**. The removal of the trifluoromethyl groups from the *C*-terminus of **3** induced much larger changes in the three-dimensional structures, which changed from a tandem β -turn (**1** and **2**) to a structured helical conformation, with the smallest rmsd values for alignment at residues 5–8 (Table 1), implying a large influence of the trifluoromethyl groups on the conformations of the entire molecules in the lipid environment. Because the structural modifications were limited only to the *C*-terminus, it is not clear how the simple electrostatic or steric changes of these modifications affects the conformations of the entire molecule. Because the molar ratios of **1–3** were low compared to DPC, they presumably form complexes with lipid molecules and thus induce perturbations in the interaction between the compound and micelles due to their unique *C*-terminus. It might be suggested that the highly lipophilic and electronegative trifluoromethyl groups induced such a perturbation if these trifluoromethyl groups interact with the core of micelles, which mostly consists of lipophilic and electrostatically neutral hydrocarbon chains. The ester-to-amide substitution, in which the oxygen atom (hydrogen-bond acceptor) was replaced to the nitrogen with a proton to be a hydrogen-bond donor, also can introduce some changes in the interactive mode between these compounds and membrane-like micelles. Therefore, the interaction of compound with DPC micelles might be compound-specific, which is consistent with the different three-dimensional conformations for each compound. To confirm these implications, further NMR experiments using paramagnetic agents as




Figure 5. The D-Ala² (crosses), Gly³ (open circle), and Met⁵ with positive ϕ angles (circled) were indicated in the Ramachandran ϕ, ψ plots for (A) **1**, (B) **2**, and (C) **3** for residues 2–7 of 20 final structures. Angular order parameters for ϕ (D) and ψ (E) angles calculated from the 20 final structures for **1** (open circles), **2** (filled squares), and **3** (crosses). For calculating the ψ angles of Trp⁸, noncarbonyl oxygen atoms of the C-terminal ester (**1**) and nitrogen atoms of C-terminal amide (**2** and **3**) were used instead of N ($i + 3$), respectively.

Table 4. Solubility and Lipophilicity of Peptide Derivatives

no.	lipophilicity		solubility ^c ($\mu\text{g/mL}$)
	$\log D_{7.4}$ ^a	AlogP ^b	
1	>4.0	5.74	<0.2
2	>4.0	5.45	<0.2
3	3.6	3.97	1.1

^a Logarithm of octanol/saline distribution coefficient in 0.05 N HEPES buffer in 0.1 N NaCl solution. ^b Calculated with ALOGPS 2.1 software. See refs 83, 84. ^c Solubility in 0.05 N HEPES buffer in 0.1 N NaCl solution.

well as fluorescence experiments were performed to estimate such differences in the membrane-compound interactions.

Fluorescence Study. It is well-known that the intrinsic fluorescence spectrum of the indole ring in tryptophan shifts to shorter wavelength (“blue-shifted”) as the polarity of the solvent surrounding the tryptophan residue decreases, and this blue-shift is a good index to monitor the lipophilicity of the environment close to the tryptophan.^{59,60} The fluorescence of Trp⁸ in **1–3** was measured in the presence and absence of DPC micelles in order to examine the interaction between the peptides and membrane-like micelles. The fluorescence spectra in DPC micelles were compared to the spectra in the EtOH–buffer solution (EtOH: pH 7.4 HEPES buffer = 1:1), which was chosen as the standard because the solubilities of the peptide derivatives **1–3** in aqueous media were too low to run the experiment (Table 4).⁶¹ The emission spectra were obtained by excitation at 290 nm to avoid excitation of the tyrosine residue.

Obvious blue-shifts of fluorescence spectra from the standard solution were observed in all three peptide derivatives (6 nm, **1**; 10 nm, **2**; 10 nm, **3**, respectively, Figure 6), implying that the indole ring in Trp⁸ at the C-terminal of all peptides was buried inside the micelles and that the peptides have strong interactions with the lipophilic core of the micelles, at least the C-terminus.

Because poorly water-soluble peptides **1–3** were easily dissolved at millimolar concentration in the presence of micelles, strong interactions between the peptide derivatives **1–3** and micelles are also suggested (Table 4). Among compounds **1–3**, **1** showed a smaller blue-shift than that of **2** and **3**, suggesting that the surroundings near the indole ring of Trp⁸ in C-terminal

ester (**1**) were less lipophilic than was the case for the C-terminal amides (**2** and **3**), both of which had better-defined structures in DPC micelles than **1** (Table 1). Therefore, the lipophilicity near Trp⁸ was mostly influenced by ester-amide substitution rather than the removal of trifluoromethyl groups, and the lipophilic environment at the C-terminus might have some responsibility for the structured conformations of the peptide derivatives.

Paramagnetic Broadening Studies on ¹H NMR. To obtain further information about the interaction between peptide derivatives **1–3** and DPC micelles, we used a nitroxyl spin-labeled 5-doxyloleic acid (5-DOXYL), and Mn²⁺ ions (MnCl₂) to induce selective broadening of NMR resonances close to the paramagnetic probes.^{12,62} The cross-peaks of protons exposed to an aqueous exterior are broadened or disappear due to the paramagnetic effect of Mn²⁺, while cross-peaks of protons located inside the micelles and close to the phosphate groups of DPC are broadened by the free radical on the doxyl group, which is bound to carbon 5 of the stearic acid. The paramagnetic effects of these agents on the peptide resonances were studied by comparing TOCSY spectra in the presence and absence of the paramagnetic agents, and all peaks were classified into three categories according to their sensitivities to the paramagnetic agents: missed by 5-DOXYL only (in the micelle, but not deeply buried), missed by both Mn²⁺ and 5-DOXYL (the proton is at or near the surface of micelles) and preserved by either agent (deeply buried in the micelle) (Figure 7 and Supporting Information). In fact, none of the ¹H resonances were missed by Mn²⁺ only, indicating a strong association with the micelle for all three derivatives.

For all three peptide derivatives **1–3**, nearly all of the H^N related cross-peaks were categorized as sensitive to both Mn²⁺ and 5-DOXYL, implying that the peptide backbones are located at or near the surface of micelles. On the other hand, most of the side chain resonances were missed only by 5-DOXYL or were nonsensitive to either agent. Thus, generally, the backbones of **1–3** lie close to the surface with their side chains buried in the micelles. However, there is one notable exception to this general observation: the H^N resonances of Met⁵ in **1–3** were affected only by 5-DOXYL, implying that the protons are not




Figure 6. Fluorescence spectra of (A) **1**, (B) **2**, (C) **3**: the spectra were recorded at 500 $\mu\text{g/mL}$ at 290 nm excitation in 40-fold DPC micelle/pH 7.4 HEPES buffer (solid line) or EtOH:pH 7.4 HEPES buffer = 1:1 solution (broken line).




Figure 7. Typical example of the paramagnetic effects on TOCSY spectra. The aromatic region of **3** with DPC micelles (left column), with 200 μM Mn^{2+} (middle) and 5-DOXYL stearic acid (right). Preserved resonances (labeled) are in a phase not missed by the phase-specific radical probe (Mn^{2+} or DOXYL). Spectra were compared from the same noise level. The full spectra for **1–3** are available in the Supporting Information.

Table 5. Binding Affinities of Bifunctional Peptides at δ/μ Opioid Receptors and NK1 Receptors

no.	hDOR ^a , [³ H]DPPPE ^b		rMOR ^a , [³ H]DAMGO ^c			hNK1 ^d , [³ H]substance P ^e		rNK1 ^d , [³ H]substance P ^f		$K_i(\text{hNK1})/K_i(\text{rNK1})$
	LogIC ₅₀ ^g	K_i (nM)	LogIC ₅₀ ^g	K_i (nM)	$K_i(\mu)/K_i(\delta)$	LogIC ₅₀ ^g	K_i (nM)	LogIC ₅₀ ^g	K_i (nM)	
1 ^h	-8.2 ± 0.06	2.8	-7.1 ± 0.11	36	13	-9.9 ± 0.25	0.084	-9.0 ± 0.10	0.29	3.4
2	-8.8 ± 0.07	0.66	-7.4 ± 0.05	16	24	-10.9 ± 0.10	0.0065	-7.6 ± 0.05	7.3	1100
3	-9.1 ± 0.09	0.44	-8.4 ± 0.03	1.8	4.1	-8.4 ± 0.42	3.20	-5.6 ± 0.06	700	220
biphalin ⁱ		2.6		1.4	0.54					
4						-8.8 ± 0.02	0.73	-6.4 ± 0.12	130	180

^a Competition analyses were carried out using membrane preparations from transfected HN9.10 cells that constitutively expressed the δ and μ opioid receptors, respectively. ^b $K_d = 0.45 \pm 0.1$ nM. ^c $K_d = 0.50 \pm 0.1$ nM. ^d Competition analyses were carried out using membrane preparations from transfected CHO cells that constitutively expressed rat or human NK1 receptors. ^e $K_d = 0.40 \pm 0.17$ nM. ^f $K_d = 0.16 \pm 0.03$ nM. ^g The LogIC₅₀ ± standard errors are expressed as logarithmic values determined from the nonlinear regression analysis of data collected from two independent experiments performed in duplicate (four independent experimental values per drug concentration). The K_i values are calculated using the Cheng and Prusoff equation to correct for the concentration of the radioligand used in the assay. ^h See ref 38. ⁱ See ref 85.

exposed to the surface of the micelles. Moreover, the side-chain cross-peaks of Met⁵ in only **1** were missed by either paramagnetic agent, indicating that Met⁵ side chain of **1** was exposed to the surface of the micelle, whereas those of **2** and **3** were sensitive only to 5-DOXYL, indicating a different orientation in the side-chain of Met⁵.

The sensitivity of side-chain protons to broadening by 5-DOXYL provides insight into their portioning in depth in the micelles. The cross-peaks of two different aromatic protons (*para* and *ortho*) in the *C*-terminal benzyl moiety were eliminated in **1** and **2** by 5-DOXYL, but the resonance of **3** was preserved in spite of the hydrophobic trifluoromethyl group in **1** and **2**. On the other hand, the cross-peaks of the aromatic protons of Trp⁸ were unaffected in **1** but were sensitive to 5-DOXYL for **2** and **3**. Therefore, the *C*-terminal benzyl moiety (**1** and **2**) and Trp⁸ (**2** and **3**) appear to be located close to the phosphate moiety of DPC micelles, but the Trp⁸ of **1** and the *C*-terminus of **3** were rather deeply buried into the micelles.

The important implication from these paramagnetic experiments was that all the compounds (**1–3**) showed different orientations at each *C*-terminus in DPC micelles. Thus, both of the modifications, the ester-to-amide substitution and the removal of trifluoromethyl groups, were shown to change the

interaction between compound and membrane-like micelles, as suggested from the NMR conformations of **1–3**.

Biological Activities. Finally, to evaluate the biological influences of conformations in the presence of membrane-mimicking micelles and compound-micelles interactions, the bioactivities of peptide derivatives **2** and **3** were evaluated using well-established assay systems (Tables 5 and 7; the details of assay systems are found in the Experimental Section and references herein).^{38,40,50,51}

Compared to the subnanomolar-level affinities of the *C*-terminal ester derivative **1** ($K_i = 0.29$ nM), the *C*-terminal amide derivative **2** was 25 times less potent but still showed binding affinities in the nanomolar range for the rNK1 receptor ($K_i = 7.3$ nM) (Table 5). The binding affinity at the rNK1 of **3**, which had no trifluoromethyl group in the *C*-terminal benzyl moiety, was decreased to a $K_i = 700$ nM. However, in the functional assay using the GPI to examine their antagonist activities, the K_e value for **3** (10 nM) was similar to that for **2** (9.9 nM) and better than that for **1** (25 nM) (Table 7). Several factors, such as pharmacokinetic differences in the different assay systems, might be responsible for this inconsistency, but one good explanation should be provided by the known species difference between rat and guinea pig NK1 receptors.⁶³ It is

Table 6. Opioid Agonist Functional Activities in [³⁵S]GTP γ S Binding Assays

no.	hDOR ^a			rMOR ^a		
	Log EC ₅₀ ^b	EC ₅₀ (nM)	E _{max} (%) ^c	Log EC ₅₀ ^b	EC ₅₀ (nM)	E _{max} (%) ^c
1 ^d	-8.5 ± 0.21	2.9	48	-7.5 ± 0.09	32	46
2	-8.1 ± 0.11	8.6	58	-8.2 ± 0.17	7.0	55
3	-8.6 ± 0.13	2.6	52	-7.7 ± 0.18	21	47
biphalin	-9.0 ± 0.17	1.1	83			
DPDPE	-8.8 ± 0.25	1.6	69			
DAMGO				-7.4 ± 0.19	37	150

^a Expressed from HN9.10 cell. ^b The log EC₅₀ ± standard error are logarithmic values determined from the nonlinear regression analysis of data collected from two independent experiments performed in duplicate (four independent experimental values per drug concentration). ^c Net total bound/basal binding × 100. ^d See ref 38.

Table 7. Functional Assay Result for Bifunctional Peptide Ligands at Opioid and Substance P Receptors

no.	opioid agonist		substance P GPI
	MVD (δ), IC ₅₀ (nM) ^a	GPI (μ), IC ₅₀ (nM) ^a	antagonist, Ke (nM) ^b
1 ^c	22 ± 1.2	360 ± 130	25 ± 8.8
2	15 ± 2.0	490 ± 29	10 ± 2.1
3	4.8 ± 0.35	61 ± 9.6	9.9 ± 2.8
biphalin	2.7 ± 1.5	8.8 ± 0.3	
4			250 ± 87

^a Concentration at 50% inhibition of muscle contraction at electrically stimulated isolated tissues. ^b Inhibitory activity against the substance P induced muscle contraction in the presence of 1 μ M naloxone. Ke: concentration of antagonist needed to inhibit substance P to half-its activity. ^c See ref 38.

well-known that the human NK1 receptor has higher homology to the guinea pig NK1 receptor than to the rat or mouse NK1, and some NK1 antagonists have a large species difference.⁶³ In fact, **3** showed a 220 times better K_i value at the hNK1 receptor (3.2 nM) compared to the value at the rNK1 receptor. A smaller difference was found in **1**, whose K_i value for the hNK1 was 0.084 nM (3.4-fold species difference). Surprisingly, **2**, which is the C-terminal benzyl amide with two trifluoromethyl groups, showed the largest difference between the affinities at the rNK1 and at the hNK1 (1100-fold) and the K_i value for the hNK1 receptor was 6.5 μ M, suggesting that this analogue would be very potent in humans.

Therefore, substitution of the C-terminal ester for an amide gave rise to increased species difference at the human and rat NK1 receptors. This substitution changed hydrogen-bonding interactions and planarity of the C-terminus, which might induce a conformational change in the lipidic surroundings as found in the NMR conformations (Figure 4). The summation of these changes could explain the activity shifts at the NK1 receptors. The existence of two trifluoromethyl groups at the C-terminus plays an important role in the affinities for both rNK1 and hNK1 receptors. Because the 3',5'-dimethyl-substituted analogue of Ac-Trp-O-3',5'-Bzl(CF₃)₂ (**4**, L-732,138)²¹ was reported to have 16-fold reduced affinity for the human NK1 receptor as compared to **4**,²¹ one good explanation of the influence by trifluoromethyl introduction was that a more electronegative phenyl ring at the C-terminus might be preferred by the NK1 receptors.⁶⁴ Another possibility is that the induced conformational changes to a well-defined helical structure due to the trifluoromethyl groups interfered with the binding of the compound at the NK1 receptor.

For the δ and μ opioid agonist activities, **2** showed 24-fold δ selectivity, as expected from the existence of Met⁵,³⁸ with 4-fold higher affinity at the hDOR (K_i = 0.66 nM) and two times higher affinity at the rMOR (K_i = 16 nM) than **1**. It is interesting that a small modification at the C-terminus, which is far from the opioid agonist pharmacophore, can influence the opioid activities so much (Table 5). The elimination of two

trifluoromethyl groups led to further increased affinity at both the DOR (K_i = 0.44 nM) and the rMOR (K_i = 1.8 nM) with 4.1-fold δ -selectivity. The binding affinity results correlated well with the in vitro GTP γ S binding assays and the functional assays using GPI and MVD tissues (Table 6 and 7).

Here, **2** showed potent activity in the MVD assay with the best δ selectivity (IC₅₀ = 15 nM in MVD and 490 nM in GPI). The IC₅₀ value of **3** in the GPI assay (IC₅₀ = 61 nM) was a large increase from those of **1** and **2**, with the best IC₅₀ value in the MVD assay (4.8 nM). Therefore, **3** was found to be a bifunctional peptide derivative possessing potent agonist activities for both δ and μ opioid receptors together with a nanomolar level hNK1 antagonist activity. On the other hand, **2** was characterized as a very potent hNK1 antagonist with potent and δ selective opioid agonist activities, which also has nanomolar level affinity at the rNK1 receptor. Thus, **2** can be considered as a promising candidate for a potent analgesic drug and it is currently being tested in several animal models using rats and will be published as a separate biological paper. Though both **2** and **3** have different biological profiles, they are expected to be potent analgesics for pain control in humans.

It should be noted that **2**, which had the better-defined conformation in DPC micelles than **1**, showed increased opioid activities, although the primary sequences for opioid agonist pharmacophore were exactly the same for both of **1** and **2** (Figure 1). Moreover, **3**, which also had the same sequence for opioid pharmacophore at the N-terminus, showed much improved opioid activities versus **1** and **2**. As discussed above, **3** had a structured helical conformation in DPC micelles, whereas **1** and **2** showed β -rich conformations. Moreover, the interactions between the compounds and the membranes were different for compounds **1**–**3**.

Consequently, these results show that even small and limited structural modification in the ligand could induce relatively large conformational changes from a β -structure to a helical conformation in membrane-mimicking DPC micelles, together with significantly shifted biological activities. In addition, the reported data suggests that the removal of trifluoromethyl groups from the C-terminus induced much larger changes in the three-dimensional conformations than the substitution of the ester to amide, implying a large influence of lipophilic and electronegative trifluoromethyl groups on the conformations of whole molecules in a lipid media. The importance of this article is that such a small and local structural modification in the C-terminus might affect on the bioactive states not only at the C-terminus (NK1 pharmacophore) but also at the N-terminal (opioid pharmacophore, which was preserved for all three derivatives), and the observed differences of the conformations in the membrane-like micelles could indicate their relevance to the changes in their bioactive form. These findings provide significant information regarding the nature of peptide drug–membrane interactions as well as the specific driving force to

use conformations to manipulate bioactivities, both of which can be a guide to new aspects of modern medicinal chemistry to understand the folding of bioactive peptides or proteins.

Experimental Section

Materials. All amino acid derivatives and coupling reagents were purchased from EMD Biosciences (Madison, WI), Bachem (Torrance, CA), SynPep (Dublin, CA), and Chem Impex International (Wood Dale, IL). 2-Chlorotrityl resin was acquired from Iris Biotech GmbH (Marktredwitz, Germany). Perdeuterated DPC was purchased from C/D/N Isotopes (Quebec, Canada). ACS grade organic solvents were purchased from VWR Scientific (West Chester, PA), and other reagents were obtained from Sigma-Aldrich (St. Louis, MO) and used as obtained. The polypropylene reaction vessels (syringes with frits) were purchased from Torviq (Niles, MI). Myo-[2-³H(N)]-inositol, [tyrosyl-3,5-³H(N)] D-Ala²-MePhe⁴-Glyol⁵-enkephalin (DAMGO), [tyrosyl-2,6-³H(N)]-[2-D-penicillamine, 5-D-penicillamine] enkephalin (DPDPE), [³H]-substance P, and [³⁵S]-guanosine 5'-(γ -thio) triphosphate were purchased from Perkin-Elmer (Wellesley, MA). Bovine serum albumin (BSA), protease inhibitors, Tris, and other buffer reagents were obtained from Sigma (St. Louis, MO). Culture medium (MEM, DMEM, and IMDM), penicillin/streptomycin, and fetal calf serum (FCS) were purchased from Invitrogen (Carlsbad, CA).

Boc-Tyr(tBu)-D-Ala-Gly-Phe-Met-Pro-Leu-Trp(Boc)-OH. The peptide was synthesized manually by the *N*^α-Fmoc solid-phase methodology using HBTU as the coupling reagents as previously reported.⁴⁰ 2-Chlorotrityl resin (2.0 g, 1.56 mmol/g) was placed into a 50 mL polypropylene syringe with the frit on the bottom and swollen in DMF (20 mL) for 1 h. The resin was washed with DMF (3 × 15 mL) and then with DCM (3 × 15 mL). Fmoc-Trp(Boc)-OH (1.2 equiv) was dissolved in 30 mL of DCM, and then DIEA (5 equiv) was added. The reaction mixture was transferred into the syringe with the resin then shaken for 2 h. The resin was washed three times with DMF (15 mL) and three times with DCM (15 mL) and then with DMF (3 × 15 mL). The Fmoc protecting group was removed by 20% piperidine in DMF (1 × 2 min and 1 × 20 min). The deprotected resin was washed with DMF (3 × 15 mL), DCM (3 × 15 mL), and then with DMF (3 × 15 mL). Fmoc-Leu-OH (3 equiv) and HBTU (2.9 equiv) were dissolved in 30 mL of DMF, then DIEA (6 equiv) was added. The coupling mixture was transferred into the syringe with the resin, then shaken for 2 h. All other amino acids, Pro, Met, Phe, Gly, D-Ala, and Tyr, were consecutively coupled using the procedures described above, using the TNBS test (all the amino acids except for Met) or chloranil test (only for Met) to check the extent of coupling. In case of a positive test result, the coupling was repeated until a negative test result was obtained. The resulting batch of the resin-bound protected Boc-Tyr(tBu)-D-Ala-Gly-Phe-Met-Pro-Leu-Trp(Boc) was carefully washed with DMF (3 × 15 mL), DCM (3 × 15 mL), DMF (3 × 15 mL), and DCM (3 × 15 mL) and dried under reduced pressure. The peptide was cleaved off the solid support with 1% v/v TFA in DCM (30 mL) for 30 min, and most of the organic solvent was removed under reduced pressure. The obtained crude peptides were precipitated out by the addition of chilled petroleum ether (45 mL) to give a white precipitate. The suspensions were centrifuged for 20 min at 7000 rpm, and then the liquid was decanted off. The crude peptides were washed with petroleum ether (2 × 50 mL), and after the final centrifugation, the peptides were dried under vacuum (2 h) to obtain the title compound (2.89 g, 74.8%). The purity of the final products (93.6%) was checked by analytical RP-HPLC using a Hewlett-Packard 1100 system (230 nm) on a reverse phase column (Waters NOVA-Pak C-18 column, 3.9 mm × 150 mm, 5 μ m, 60Å). The peptide was eluted with a linear gradient of aqueous CH₃CN/0.1% CF₃CO₂H (10–90% in 40 min) at a flow rate of 1.0 mL/min. The crude peptide was used for next reactions without further purification. ¹H NMR (DMSO-*d*₆): 0.79 (3H, d, *J* = 6.6 Hz), 0.84 (3H, d, *J* = 6.6 Hz), 1.10 (3H, d, *J* = 6.6 Hz), 1.24 (9H, s), 1.26 (9H, s), 1.38–1.41 (2H, m), 1.61 (10H, s), 1.70–1.77 (3H, m), 1.80–1.94

(3H, m), 2.01 (3H, s), 2.42–2.47 (2H, m), 2.67 (1H, d-d, *J* = 3.6, 13.2 Hz), 2.75 (1H, d-d, *J* = 9.6, 13.8 Hz), 2.87 (1H, d-d, *J* = 4.2, 13.2 Hz), 2.94 (1H, d-d, 2.4, 13.2 Hz), 3.04 (1H, d-d, 8.4, 15.0 Hz), 3.14 (1H, d-d, 4.8, 15.0 Hz), 3.48–3.53 (1H, m), 3.54–3.63 (2H, m), 3.68 (1H, d-d, *J* = 5.4, 16.8 Hz), 4.15 (1H, d-d, *J* = 6.0, 12.0 Hz), 4.20–4.30 (2H, m), 4.32 (1H, d-d, 3.6, 7.8 Hz), 4.48–4.55 (2H, m), 4.58 (1H, *J* = 7.2, 14.4 Hz), 6.83 (2H, d, *J* = 7.8 Hz), 6.92 (1H, d, *J* = 8.4 Hz), 7.12 (2H, d, *J* = 7.8 Hz), 7.14–7.24 (7H, m), 7.30 (1H, t, *J* = 7.8 Hz), 7.47 (1H, s), 7.87–7.92 (2H, m), 7.98–8.08 (4H, m), 8.30 (1H, d, *J* = 7.2 Hz). MS (ESI): 1262 (M + Na)⁺.

H-Tyr-D-Ala-Gly-Phe-Met-Pro-Leu-Trp-O-3,5-Bzl(CF₃)₂·TFA (1). Boc-Tyr(tBu)-D-Ala-Gly-Phe-Met-Pro-Leu-Trp(Boc)-OH (2.0 g, 1.61 mmol) and 3,5-bis(trifluoromethyl)benzyl bromide (1.24 g, 4.02 mmol) were dissolved in DMF (8 mL). Cesium carbonate (1.05 g, 3.22 mmol) was added to the solution at r.t. After stirring for 2 h, saturated aqueous sodium bicarbonate (200 mL) was added to the solution and extracted with ethyl acetate (200 mL) three times. The combined organic phases were washed with 5% aqueous citrate and saturated aqueous sodium chloride (200 mL each) and dried over sodium sulfate. The solvent was evaporated off, and the crude peptide was precipitated in cold petroleum ether (45 mL) and centrifuged two times and dried under reduced pressure. The obtained protected peptide was treated with 82.5% v/v TFA, 5% water, 5% thioanisole, 2.5% 1,2-ethanedithiol, and 5% phenol (10 mL, 1 h). The crude peptide was precipitated out by the addition of chilled diethyl ether (45 mL) to give white precipitates. The suspension was centrifuged for 20 min at 7000 rpm, then the liquid was decanted. The crude peptides were washed with diethyl ether (2 × 45 mL), and after the final centrifugation, the peptides were dried under vacuum (2 h). The resulting white residues (2.48 g, quantitative) were dissolved in a 3:1 mixture of acetonitrile and distilled water (5 mL) and the insoluble impurities were removed by passing the solutions through syringe filters (Gelman Laboratory, Ann Arbor, MI, Acrodisc 13 mm syringe filter with 0.45 μ M PTFE membrane). Final purification was accomplished by preparative RP-HPLC and then lyophilized.

H-Tyr-D-Ala-Gly-Phe-Met-Pro-Leu-Trp-NH-3,5-Bzl(CF₃)₂·TFA (2). Boc-Tyr(tBu)-D-Ala-Gly-Phe-Met-Pro-Leu-Trp(Boc)-OH (3.0 g, 2.42 mmol) and Cl-HOBt (428 mg, 2.66 mmol) were dissolved in DMF (10 mL). 3,5-Bistrifluoromethylbenzyl amine (1.17 g, 4.84 mmol) and EDC (508 mg, 2.66 mmol) were added to the solution at r.t. and stirred until the starting material was not detected in TLC and then saturated aqueous sodium bicarbonate (250 mL) was added. The reaction mixture was extracted with ethyl acetate (250 mL) three times. The combined organic phases were washed with 5% aqueous citrate and saturated aqueous sodium chloride (250 mL each) and then dried over sodium sulfate. The solvent was evaporated, and the crude peptide was precipitated in cold petroleum ether (45 mL). The product was twice dispersed in cold petroleum ether, centrifuged and decanted, and then dried under reduced pressure. The obtained protected peptide was treated with 82.5% v/v TFA, 5% water, 5% thioanisole, 2.5% 1,2-ethanedithiol, and 5% phenol (1.5 mL, 1 h). The crude peptide was precipitated out by the addition of chilled diethyl ether (45 mL) to give a white precipitate. The resulting peptide suspension was centrifuged for 20 min at 7000 rpm, and the liquid was decanted. The crude peptide was washed with diethyl ether (2 × 45 mL), and after a final centrifugation, the peptides were dried under vacuum (2 h). The resulting white residue (3.42 g, quantitative) was dissolved in a 3:1 mixture of acetonitrile and distilled water (5 mL), and the insoluble impurities were removed by passing the solutions through syringe filters (Gelman Laboratory, Acrodisc 13 mm syringe filter with 0.45 μ M PTFE membrane). Final purification was accomplished by preparative RP-HPLC. The pure title compound was obtained after lyophilization.

H-Tyr-D-Ala-Gly-Phe-Met-Pro-Leu-Trp-NH-Bzl·TFA (3). The title peptide was prepared using the same method as described for H-Tyr-D-Ala-Gly-Phe-Pro-Leu-Trp-NH-3,5-Bzl(CF₃)₂·TFA (2). The crude peptide was obtained quantitatively.

Characterization of Peptides. Preparative RP-HPLC was performed on Waters Delta Prep 4000 with Waters XTerra C-18 column (19 mm \times 250 mm, 10 μ m, a linear gradient of 33–53% or 40–60% acetonitrile/0.1% TFA at a flow rate of 15.0 mL/min). The purified peptides were characterized by HRMS, TLC, analytical HPLC, and ^1H -1D-NMR. Sequential assignment of proton resonances was achieved by 2D-TOCSY NMR experiments.^{65–69} High-resolution MS were taken in the positive ion mode using ESI methods at the University of Arizona Mass Spectrometry Facility. TLC was performed on aluminum sheets coated with a 0.2 mm layer of silica gel 60 F₂₅₄ Merck using the following solvent systems: (1) CHCl_3 :MeOH:AcOH = 90:10:3, (2) EtOAc:*n*-BuOH:water:AcOH = 5:3:1:1, and (3) *n*-BuOH:water:AcOH = 4:1:1. TLC chromatograms were visualized by UV light and by ninhydrin spray followed by heating (hot plate). Analytical HPLC was performed on a Hewlett-Packard 1100 or Hewlett-Packard 1090m with Waters NOVA-Pak C-18 column (3.9 mm \times 150 mm, 5 μ m, 60 Å) or Vydac 218TP104 C-18 column (4.6 mm \times 250 mm, 10 μ m, 300 Å). ^1H -1D-NMR spectra were obtained on Bruker DRX-500 or DRX-600 spectrometer. 2D-TOCSY NMR spectra were performed on a Bruker DRX-600 spectrometer equipped with a 5 mm Nalorac triple-resonance single-axis gradient probe. The NMR experiments were conducted in DMSO-*d*₆ solution at 298 K. Spectra were referenced to residual solvent protons as 2.49 ppm. The processing of NMR data was performed with the XwinNmr software (Bruker BioSpin, Fremont, CA). In the TOCSY experiments, the TPPI mode⁷⁰ with MLEV-17 Mixing Sequence⁶⁵ were used with a mixing time of 62.2 ms, at a spin-lock field of 8.33 kHz. TOCSY spectra were acquired with 2k complex pairs in t_2 and 750 or 1024 FIDs using a 90°-shifted sine-squared window function in both dimensions and zero filling to a final matrix size of 2048 (F2) \times 1024 (F1) points.

NMR Spectroscopy in DPC Amphipathic Media.^{45,46} All NMR spectra were recorded on a Bruker DRX600 600 MHz spectrometer with 5 mm Nalorac triple-resonance single-axis gradient probe. The peptide concentration for the NMR experiments varied from 3.5 to 4 mM. The samples were prepared by dissolving the peptide in 0.5 mL of 45 mM sodium acetate-*d*₃ buffer (pH 4.5) containing 40 equiv of dodecylphosphocholine-*d*₃₈ and 1 mM sodium azide (90% H₂O/10% D₂O), followed by sonication for 5 min. Two-dimensional double quantum filtered correlation (DQF-COSY), nuclear Overhauser effect⁷¹ (NOESY), and total correlation spectra⁶⁶ (TOCSY) were acquired using standard pulse sequences and processed using XwinNmr and Felix 2000 (Accelrys Inc., San Diego, CA). The mixing time for NOESY spectra was 450 ms. All 2D spectra were acquired in the TPPI mode with 2k or 1k complex data points in t_2 and 750 real data points in t_1 , and the spectral processing used 90°-shifted sine bell window functions in both dimensions. For suppressing the H₂O signal, the 3–9–19 WATERGATE pulse sequence was used.⁷² Experiments were conducted at 310 K and referenced to the H₂O shift (4.631 ppm). Coupling constants ($^3J_{\text{NH-H}\alpha}$) were measured from 2D DQF-COSY spectra by analysis of the fingerprint region. The matrix rows of each of the upper and lower halves of a cross-peak were summed to give an antiphase 1D spectrum, which was fitted using a 5-parameter Levenberg–Marquardt nonlinear least-squares protocol⁷³ to a general antiphase doublet. The analysis yielded two independent determinations of the J coupling and line width for each cross-peak, one from the upper half and one from the lower half, and the one with the better fitted curve was used for structure calculations. Cross-peak volumes for determination of distance restraints were measured using the Felix 2000 software. In the radical experiment using Mn²⁺, a stock solution of 5 mM MnCl₂ was prepared and added to the sample to achieve a total concentration of 200 μ M in Mn²⁺. The DPC micelles with 5-DOXYL stearic acid were prepared as the same procedure with about 1 mg mL⁻¹ of 5-DOXYL stearic acid but sonicating for 30 min.

Conformational Structure Determination. The methods used for structure calculations have been described previously.^{45,46} Throughout the ^1H NMR studies, only one major rotamer for **1–3** was found and the populations of minor rotamers were all

negligible. The Met⁵–Pro⁶ bond of the major rotamers were fixed in the trans configuration based on the observations of $^5\text{H}^\alpha$ to Pro⁶ H ^{δ} sequential NOEs together with the absence of sequential $^5\text{H}^\alpha$ – $^6\text{H}^\alpha$ NOEs in the structural calculations of **1–3**. The volumes of the assigned cross-peaks in the 2D NOESY spectrum were converted into upper distance bounds of 3.0, 3.8, 4.8, or 5.8 Å. For overlapping cross-peaks, the distance categories was increased by one or two levels depending on the qualitative estimate of the extent of overlap. Pseudoatoms were created for nonstereospecifically assigned methylene protons with a correction of 1.0 Å applied to their upper bound distances. In addition to the distance constraints, φ dihedral angle constraints derived from $^3J_{\text{HN-H}\alpha}$ coupling constants were set to between –90 and 40° for $^3J_{\text{HN-H}\alpha} < 6$ Hz and to between –150 and –90° for $^3J_{\text{HN-H}\alpha} > 8$ Hz. Dihedral angle constraints of $180^\circ \pm 5^\circ$ for peptide bonds (ω) were also used to maintain the planarity of these bonds.

Simulated annealing molecular dynamics analysis was done for all the peptides to obtain an ensemble of NMR structures using the NOE-derived distance constraints and dihedral angle (φ) constraints using the DGII⁷⁴ program within the software package Insight II 2000 (Accelrys Inc., San Diego, CA). Solvent was not explicitly included in the calculations. All the embedded structures successfully passed the simulated annealing step and were minimized using the consistent valency force field (CVFF) (Accelrys Inc.). The 50 structures with the lowest penalty function were further refined by two rounds of restrained molecular dynamics (rMD) using the all-atom AMBER force field with additional parameters for fluorine atom^{75–78} using the standalone DISCOVER ver. 2.98 program (Accelrys Inc.). A 12.0 Å cutoff for nonbonded interactions and a distance-dependent dielectric constant (4 r) were used. All amide bonds were constrained to trans conformation by a 100 kcal mol⁻¹ rad⁻² energy penalty. The distance constraints and dihedral angles (φ) constraints were applied with a force constant of 25 kcal mol⁻¹ Å⁻² and 100 kcal mol⁻¹ rad⁻² were applied, respectively. After 100 steps of steepest descents minimization and 1000 steps of conjugate gradient minimization on the initial structures, an rMD equilibration at 500 K was performed for 1.5 ps, during which a scale factor of 0.1 was applied to the experimental restraint force constants. During the next 2 ps, full values of the experimental restraint force constants were applied. A further 1 ps rMD simulation was run at 500 K, and the system was then cooled to 0 K over 3 ps. After another 1 ps at 0 K, 100 cycles of steepest descents and 2000 steps of conjugate gradient minimization were performed. The final 20 structures with the lowest energies were used for the analysis. All calculations were performed on a Silicon Graphics Octane computer.

Fluorescence Emission Spectra. The Fluorescence spectra were recorded on a Cary Eclipse fluorescence spectrometer (Varian, Darmstadt, Germany). Emission spectra were obtained by excitation at 290 nm and recorded in the wavelength range of 310–420 nm with continuous stirring at 25 °C. A scan step was 1 nm and scan speed was 120 nm min⁻¹. Excitation and emission bandwidths were set at 5 and 10 nm, respectively. The peptide concentration was 500 μ M in HEPES buffer (10 mM HEPES, 150 mM NaCl, 1 mM Na₃, 0.1 mM EDTA, pH = 7.40) with 40-fold DPC or standard solution (EtOH:HEPES buffer = 1:1).⁶¹ At least two scans were accumulated and averaged for each spectrum.

Octanol/Saline Distribution (logD_{7.4}).³⁸ HEPES buffer (0.05 M HEPES buffer in 0.1 M NaCl, pH 7.4, 500 μ L) was added to 2 mg of peptide and mixed with 500 μ L of 1-octanol. The sample was shaken at r.t. for 12 h to allow equilibrating. The sample was centrifuged at 6500 rpm in a VanGuard V6500 (GlaxoSmithKline, Research Triangle Park, NC) for 15 min. The layers were separated and each layer was centrifuged once again. The peptide concentrations in the obtained layers were analyzed by HPLC (30–70% of acetonitrile containing 0.1% TFA within 20 min and up to 95% within additional 5 min, 1 mL/min, 230 nm, Vydac 218TP104 C-18 column). The logarithm of 1-octanol/saline distribution (logD_{7.4}) was calculated as the ratio of peptide concentration in the 1-octanol and saline phases.

Solubility. HEPES buffer (0.05 M HEPES buffer in 0.1 M NaCl, pH 7.4, 500 μ L) was added to 1 mg of peptide. The sample was vortexed for 30 s, sonicated for 5 min, shaken at r.t. for 2 h, and then allowed to be stayed overnight to equilibration. The sample was filtrated with an Acrodisc Syringe Filter (13 mm, 0.45 μ m pore, PTFE membrane, Pall Life Sciences, East Hills, NY). The peptide concentration of the obtained filtrate was analyzed by HPLC (30–70% of acetonitrile containing 0.1% TFA within 20 min and up to 95% within an additional 5 min, 1 mL/min, 230 nm, Vydac 218TP104 C-18 column).

Cell Lines. For opioid receptors, the cDNA for the human δ opioid receptor (DOR) was a gift from Dr. Brigitte Kieffer (IGBMC, Illkirch, France). The cDNA for the rat μ opioid receptor (MOR) was a gift from Dr. Huda Akil (University of Michigan, MI). Stable expression of the rat MOR (pCDNA3) and the human DOR (pCDNA3) in the neuroblastoma cell line HN9.10 were achieved with the respective cDNAs by calcium phosphate precipitation followed by clonal selection in neomycin. Expression of the respective receptors was initially verified and the level of expression periodically monitored by radioligand saturation analysis (see below). All cells were maintained at 37 $^{\circ}$ C, with a 95% air/5% CO₂ humidified atmosphere in a Forma Scientific incubator in Dulbecco's modified Eagle's medium (DMEM) with 10% bovine serum albumin (BSA) and 100 U mL penicillin/100 μ g mL streptomycin. For NK1 receptor, the hNK1/CHO and rNK1/CHO cell lines were obtained from Dr. James Krause (University of Washington Medical School, St. Louis, MI). Expression of the receptor was verified as previously described by Krause et al.⁷⁹ All cells were maintained at a 37 $^{\circ}$ C, 95% air and 5% CO₂, humidified atmosphere, in a Forma Scientific incubator in Ham's F12 with 2.5 mM HEPES, 10% fetal bovine serum, and 100 U mL penicillin/100 μ g mL streptomycin/500 μ g mL Geneticin.

Radioligand Labeled Binding Assays. For opioid receptors, crude membranes were prepared as previously described⁸⁰ from the transfected cells that express the MOR or the DOR. The protein concentration of the membrane preparations was determined by the Lowry method, and the membranes were stored at -80 $^{\circ}$ C until use. Membranes were resuspended in assay buffer (50 mM Tris, pH 7.4, containing 50 μ g/mL bacitracin, 30 μ M bestatin, 10 μ M captopril, 100 μ M phenylmethylsulfonyl fluoride (PMSF), 1 mg/mL BSA). For saturation analysis, six concentrations of tritiated [D-Ala², NMePhe⁴, Gly⁵-ol]-enkephalin (³H]DAMGO) (0.02–6 nM, 47.2 Ci/mmol), or six concentrations of tritiated [D-Pen², D-Pen⁵]-enkephalin (³H]DPDPE) (0.1–10 nM, 44 Ci/mmol) were each mixed with 200 μ g of membranes from MOR or DOR expressing cells, respectively, in a final volume of 1 mL. For competition analysis, 10 concentrations of a test compound were each incubated with 50 μ g of membranes from MOR or DOR expressing cells and the K_d concentration of [³H]DAMGO (1.0 nM, 50 Ci/mmol), or of [³H]DPDPE (1.0 nM, 44 Ci/mmol), respectively. Naloxone at 10 μ M was used to define the nonspecific binding of the radioligands in all assays. All assays were carried out in duplicates and were repeated. The samples were incubated in a shaking water bath at 25 $^{\circ}$ C for 3 h, followed by rapid filtration through Whatman grade GF/B filter paper (Gaithersburg, MD) presoaked in 1% polyethyleneimine, washed 4 times each with 2 mL of cold saline, and the radioactivity determined by liquid scintillation counting (Beckman LS5000 TD).

For the NK1 receptors, competition binding assays for the NK1 receptors were carried out on crude membranes prepared from transfected CHO cells expressing the human or rat NK1 receptor, respectively. Ten concentrations of a test compound were each incubated with 50–100 μ g of membrane homogenate and 0.3–0.4 nM [³H] substance P (135 Ci/mmol, Perkin-Elmer) in 1 mL final volume of assay buffer (50 mM Tris, pH 7.4, containing 5 mM MgCl₂, 50 μ g/mL bacitracin, 30 μ M bestatin, 10 μ M captopril, 100 μ M phenylmethylsulfonyl fluoride (PMSF) at 25 $^{\circ}$ C for 20 min. Substance P at 10 μ M was used to define the nonspecific binding. Membrane concentrations used in the assay were within the tissue

linearity range. The [³H] substance P concentration was selected based on the saturation binding experiments, which showed a high affinity binding with K_d values of 0.40 ± 0.17 for hNK1 and 0.16 ± 0.03 nM for rNK1, respectively. The incubation times correspond to the binding equilibrium as determined from the kinetics experiments. The incubation was stopped by rapid filtration through a GF/B glass filter (Brandel Inc.) presoaked in 0.5% polyethyleneimine, followed by four washes with 2 mL of ice-cold saline using a Brandel Harvester apparatus. The filter-bound radioactivity was measured by liquid scintillation counting (Beckman LS 6000SC). Log IC₅₀ values were determined with standard errors from nonlinear regression analysis of at least two independent experiments performed in duplicates using GraphPad Prism 4 software (GraphPad, San Diego, CA).

For competition analysis using [³H]DAMGO, [³H]DPDPE or [³H]substance P values were calculated from the IC₅₀ by the Cheng and Prusoff equation.

[³⁵S]GTP γ S Binding Assay. The method was carried out as previously described.⁸⁰ Membrane preparation (10 μ g) to a final volume of 1 mL of incubation mix (50 mM Hepes, pH 7.4, 1 mM EDTA, 5 mM MgCl₂, 30 μ M GDP, 1 mM dithiothreitol, 100 mM NaCl, 0.1 mM PMSF, 0.1% BSA, 0.1 nM [³⁵S]GTP γ S) was added along with various concentrations, in duplicates or triplicates, of the test drug and incubated for 90 min at 30 $^{\circ}$ C in a shaking water bath. Reactions were terminated by rapid filtration through Whatman GF/B filters (presoaked in water), followed by four washes with 4 mL of ice-cold wash buffer (50 mM Tris, 5 mM MgCl₂, 100 mM NaCl, pH 7.4). The radioactivity was determined by liquid scintillation counting as above. Basal level of [³⁵S]GTP γ S binding was defined as the amount bound in the absence of any test drug. Nonspecific binding was determined in the presence of 10 μ M unlabeled GTP γ S. Total binding was defined as the amount of radioactivity bound in the presence of the test drug. The effect of the drug at each concentration on [³⁵S]GTP γ S binding was calculated as a percentage by the following equation: [total bound-basal]/[basal-nonspecific] \times 100. Data were expressed as log EC₅₀ \pm standard error from at least two independent experiments performed in duplicate analyzed by nonlinear regression analysis using GraphPad Prism4.

Guinea Pig Isolated Ileum Assay. The in vitro tissue bioassays were performed as described previously.³⁸ Male Hartley guinea pigs under ether anesthesia were sacrificed by decapitation and a nonterminal portion of the ileum removed and the longitudinal muscle with myenteric plexus (LMMP) was carefully separated from the circular muscle as described previously.⁸¹ These tissues were tied to gold chains with suture silk and mounted between platinum wire electrodes in 20 mL organ baths at a tension of 1 g and bathed in oxygenated (95:5 O₂:CO₂) Krebs's bicarbonate buffer at 37 $^{\circ}$ C and stimulated electrically (0.1 Hz, 0.4 ms duration) at supramaximal voltage. Following an equilibration period, compounds were added cumulatively to the bath in volumes of 14–60 μ L until maximum inhibition was reached. A baseline for PL-017 was constructed to determine tissue integrity and allow calculation of antagonist activity before opioid analogue testing began. If no agonist activity was observed at 1 μ M, a repeat PL-017 dose–response curve was constructed to test for antagonist qualities.

All substance P antagonist compound testing was performed in the presence of 1 μ M naloxone to block opioid effects on unstimulated tissue. Two minutes after naloxone was added to the bath, the test compound was added. Four minutes after naloxone was added, the test dose of substance P was added to the bath, the peak height was noted, and the tissues were washed. Agonist activity of the analogue was also observed during this period. Testing stopped at 1 mM concentrations of the test compound.

Mouse Isolated Vas Deferens (MVD) Assay. The in vitro tissue bioassay was performed as described previously.³⁸ Male ICR mice under ether anesthesia were sacrificed by cervical dislocation and the vasa deferentia removed. Tissues were tied to gold chains with suture silk and mounted between platinum wire electrodes in 20 mL organ baths at a tension of 0.5 g and bathed

in oxygenated (95:5 O₂:CO₂) magnesium free Krebs's buffer at 37 °C and stimulated electrically (0.1 Hz, single pulses, 2.0 ms duration) at supramaximal voltage as previously described.⁸² Following an equilibration period, compounds were added to the bath cumulatively in volumes of 14–60 μ L until maximum inhibition was reached. Response to an IC₅₀ dose of DPDPE (10 nM) was measured to determine tissue integrity before compound testing.

Analysis of the GPI and MVD assays. For opioid data analysis, percentage inhibition was calculated using the average tissue contraction height for 1 min preceding the addition of the agonist divided by the contraction height 3 min after exposure to the dose of agonist. IC₅₀ values represent the mean of not less than 4 tissues. IC₅₀ and E_{max} estimates were determined by computerized nonlinear least-squares analysis (the pharmacological statistics package FlashCalc: Dr. Michael Ossipov, University of Arizona, Tucson, AZ). For substance P data analysis, the height of the maximum peak produced during the control substance P dose–response curve was used as a 100% response and other values calculated as a percentage. Ke values represent the mean of not less than 4 tissues. Ke estimates were determined by computerized nonlinear least-squares analysis (FlashCalc).

Acknowledgment. The work was supported by a grant from the USDHS, National Institute on Drug Abuse, DA-13449. We thank Dr. Guangxin Lin and Dr. Jinfa Ying for kind assistance and advice with the NMR measurements and structure calculations, Dr. Robin Polt for fruitful discussion, especially about the paramagnetic NMR experiments, Dr. John J. Osterhout for use of the fluorescence spectrometer, Magdalena Kaczmarek for culturing cells, and the University of Arizona Mass Spectrometry Facility for the mass spectra measurements. We express appreciation to Margie Colie and Brigid Blazek for assistance with the manuscript.

Supporting Information Available: The ¹H NMR, HPLC, and MS data of the peptide derivatives 1–3 are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Dobson, C. M. Protein folding and misfolding. *Nature* **2003**, *426*, 884–890.
- (2) Stefani, M.; Dobson, C. M. Protein aggregation and aggregate toxicity: new insights into protein folding, misfolding diseases and biological evolution. *J. Mol. Med.* **2003**, *81*, 678–699.
- (3) Selkoe, D. J. Folding proteins in fatal ways. *Nature* **2003**, *426*, 900–904.
- (4) Barrow, C. J.; Zagorski, M. G. Solution structures of beta peptide and its constituent fragments: relation to amyloid deposition. *Science* **1991**, *253*, 179–182.
- (5) Fraser, P. E.; Nguyen, J. T.; Surewicz, W. K.; Kirschner, D. A. pH-Dependent structural transitions of Alzheimer amyloid peptides. *Biophys. J.* **1991**, *60*, 1190–1201.
- (6) Prusiner, S. B. Prions. *Proc. Natl. Acad. Sci. U.S.A.* **1998**, *95*, 13363–13383.
- (7) Trzesniewska, K.; Brzyska, M.; Elbaum, D. Neurodegenerative aspects of protein aggregation. *Acta Neurobiol. Exp.* **2004**, *64*, 41–52.
- (8) Reiss, C.; Lesnik, T.; Parvez, H.; Parvez, S.; Ehrlich, R. Conformational toxicity and sporadic conformational diseases. *Toxicology* **2000**, *153*, 115–121.
- (9) Vendruscolo, M.; Zurdo, J.; MacPhee, C. E.; Dobson, C. M. Protein folding and misfolding: a paradigm of self-assembly and regulation in complex biological systems. *Philos. Trans. R. Soc. London, Ser. A* **2003**, *361*, 1205–1222.
- (10) Lopez de la Paz, M.; de Mori, G. M.; Serrano, L.; Colombo, G. Sequence dependence of amyloid fibril formation: insights from molecular dynamics simulations. *J. Mol. Biol.* **2005**, *349*, 583–596.
- (11) Hughes, J.; Smith, T. W.; Kosterlitz, H. W.; Fothergill, L. A.; Morgan, B. A.; Morris, H. R. Identification of two related pentapeptides from the brain with potent opiate agonist activity. *Nature* **1975**, *258*, 577–580.
- (12) Palian, M. M.; Boguslavsky, V. I.; O'Brien, D. F.; Polt, R. Glycopeptide–membrane interactions: glycosyl enkephalin analogues adopt turn conformations by NMR and CD in amphipathic media. *J. Am. Chem. Soc.* **2003**, *125*, 5823–5831.
- (13) Dhanasekaran, M.; Palian, M. M.; Alves, I.; Yeomans, L.; Keyari, C. M.; Davis, P.; Bilsky, E. J.; Eggleton, R. D.; Yamamura, H. I.; Jacobsen, N. E.; Tollin, G.; Hruby, V. J.; Porreca, F.; Polt, R. Glycopeptides related to beta-endorphin adopt helical amphipathic conformations in the presence of lipid bilayers. *J. Am. Chem. Soc.* **2005**, *127*, 5435–5448.
- (14) Spadaccini, R.; Temussi, P. A. Natural peptide analgesics: the role of solution conformation. *Cell. Mol. Life Sci.* **2001**, *58*, 1572–1582.
- (15) Rudolph-Bohner, S.; Quarzago, D.; Czisch, M.; Ragnarsson, U.; Moroder, L. Conformational preferences of Leu-enkephalin in reverse micelles as membrane-mimicking environment. *Biopolymers* **1997**, *41*, 591–606.
- (16) Kallick, D. A.; Tessmer, M. R.; Watts, C. R.; Li, C. Y. The use of dodecylphosphocholine micelles in solution NMR. *J. Magn. Reson., Ser. B* **1995**, *109*, 60–65.
- (17) Kameda, T.; Takada, S. Secondary structure provides a template for the folding of nearby polypeptides. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 17765–17770.
- (18) Berthold, M.; Bartfai, T. Modes of peptide binding in G protein-coupled receptors. *Neurochem. Res.* **1997**, *22*, 1023–1031.
- (19) Noeske, T. G., A.; Parsons, C. G.; Weila, T. Allosteric modulation of family 3 GPCRs. *QSAR Comb. Sci.* **2006**, *25*, 134–146.
- (20) Eguchi, M. Recent advances in selective opioid receptor agonists and antagonists. *Med. Res. Rev.* **2004**, *24*, 182–212.
- (21) Cascieri, M. A.; Macleod, A. M.; Underwood, D.; Shiao, L. L.; Ber, E.; Sadowski, S.; Yu, H.; Merchant, K. J.; Swain, C. J.; Strader, C. D.; Fong, T. M. Characterization of the interaction of N-acyl-L-tryptophan benzyl ester neurokinin antagonists with the human neurokinin-1 receptor. *J. Biol. Chem.* **1994**, *269*, 6587–6591.
- (22) D'Alagni, M.; Delfini, M.; Di Nola, A.; Eisenberg, M.; Paci, M.; Roda, L. G.; Veglia, G. Conformational study of [Met⁵]enkephalin-Arg-Phe in the presence of phosphatidylserine vesicles. *Eur. J. Biochem.* **1996**, *240*, 540–549.
- (23) Deber, C. M.; Behnam, B. A. Role of membrane lipids in peptide hormone function: binding of enkephalins to micelles. *Proc. Natl. Acad. Sci. U.S.A.* **1984**, *81*, 61–65.
- (24) Sargent, D. F.; Schwyzer, R. Membrane lipid phase as catalyst for peptide–receptor interactions. *Proc. Natl. Acad. Sci. U.S.A.* **1986**, *83*, 5774–5778.
- (25) Wienk, H. L.; Wechselberger, R. W.; Czisch, M.; de Kruijff, B. Structure, dynamics, and insertion of a chloroplast targeting peptide in mixed micelles. *Biochemistry* **2000**, *39*, 8219–8227.
- (26) Wienk, H. L.; Wechselberger, R. W.; Czisch, M.; de Kruijff, B. Structure, dynamics, and insertion of a chloroplast targeting peptide in mixed micelles. *Biochemistry* **2000**, *39*, 8219–8227.
- (27) Bryson, E. A.; Rankin, S. E.; Carey, M.; Watts, A.; Pinheiro, T. J. Folding of apocytochrome c in lipid micelles: formation of α helix precedes membrane insertion. *Biochemistry* **1999**, *38*, 9758–9767.
- (28) Saffman, P. G.; Delbruck, M. Brownian motion in biological membranes. *Proc. Natl. Acad. Sci. U.S.A.* **1975**, *72*, 3111–3113.
- (29) Janecka, A.; Fichna, J.; Janecki, T. Opioid receptors and their ligands. *Curr. Top. Med. Chem.* **2004**, *4*, 1–17.
- (30) Gentilucci, L.; Tolomelli, A. Recent advances in the investigation of the bioactive conformation of peptides active at the micro-opioid receptor. Conformational analysis of endomorphins. *Curr. Top. Med. Chem.* **2004**, *4*, 105–121.
- (31) King, T.; Gardell, L. R.; Wang, R.; Vardanyan, A.; Ossipov, M. H.; Malan, T. P., Jr.; Vanderah, T. W.; Hunt, S. P.; Hruby, V. J.; Lai, J.; Porreca, F. Role of NK-1 neurotransmission in opioid-induced hyperalgesia. *Pain* **2005**, *116*, 276–288.
- (32) Ripley, T. L.; Gadd, C. A.; De Felipe, C.; Hunt, S. P.; Stephens, D. N. Lack of self-administration and behavioural sensitisation to morphine, but not cocaine, in mice lacking NK1 receptors. *Neuropharmacology* **2002**, *43*, 1258–1268.
- (33) King, T.; Ossipov, M. H.; Vanderah, T. W.; Porreca, F.; Lai, J. Is paradoxical pain induced by sustained opioid exposure an underlying mechanism of opioid antinociceptive tolerance? *Neurosignals* **2005**, *14*, 194–205.
- (34) Ma, W.; Zheng, W. H.; Kar, S.; Quirion, R. Morphine treatment induced calcitonin gene-related peptide and substance P increases in cultured dorsal root ganglion neurons. *Neuroscience* **2000**, *99*, 529–539.
- (35) Powell, K. J.; Quirion, R.; Jhamandas, K. Inhibition of neurokinin-1-substance P receptor and prostanoid activity prevents and reverses the development of morphine tolerance in vivo and the morphine-induced increase in CGRP expression in cultured dorsal root ganglion neurons. *Eur. J. Neurosci.* **2003**, *18*, 1572–1583.
- (36) Misterek, K.; Maszczyńska, I.; Dorociak, A.; Gumulka, S. W.; Carr, D. B.; Szyfelbein, S. K.; Lipkowski, A. W. Spinal co-administration of peptide substance P antagonist increases antinociceptive effect of the opioid peptide buprenorphine. *Life Sci.* **1994**, *54*, 939–944.
- (37) Gu, G.; Kondo, I.; Hua, X. Y.; Yaksh, T. L. Resting and evoked spinal substance P release during chronic intrathecal morphine infusion:

- parallels with tolerance and dependence. *J. Pharmacol. Exp. Ther.* **2005**, *314*, 1362–1369.
- (38) Yamamoto, T.; Nair, P.; Davis, P.; Ma, S. W.; Navratilova, E.; Moye, M.; Tumati, S.; Vanderah, T. W.; Lai, J.; Porreca, F.; Yamamura, H. I.; Hruby, V. J. Design, Synthesis and Biological Evaluation of Novel Bifunctional C-terminal Modified Peptides for δ/μ Opioid Receptor Agonists and Neurokinin-1 Receptor Antagonists. *J. Med. Chem.* **2007**, *50*, 2779–2786.
- (39) Yamamoto, T.; Nair, P.; Davis, P.; Ma, S. W.; Moye, S.; Largent, T.; Vanderah, T. W.; Lai, J.; Porreca, F.; Yamamura, H. I.; Hruby, V. J. Design, Structure-Activity Relationships and Biological Evaluation of Novel Bifunctional C-terminal Modified Peptides for δ/μ Opioid Receptor Agonists and Neurokinin-1 Receptor Antagonists. In 232nd ACS National Meeting, San Francisco, CA, 2006, pp MED1-7.
- (40) Yamamoto, T.; Nair, P.; Vagner, J.; Davis, P.; Ma, S. W.; Navratilova, E.; Moye, M.; Tumati, S.; Vanderah, T. W.; Lai, J.; Porreca, F.; Yamamura, H. I.; Hruby, V. J. A Structure-Activity Relationship Study and Combinatorial Synthetic Approach of C-Terminal Modified Bifunctional Peptides That Are δ/μ Opioid Receptor Agonists and Neurokinin 1 Receptor Antagonists. *J. Med. Chem.* **2008**, *51*, 1369–1376.
- (41) Braun, W.; Wider, G.; Lee, K. H.; Wuthrich, K. Conformation of glucagon in a lipid-water interphase by ^1H nuclear magnetic resonance. *J. Mol. Biol.* **1983**, *169*, 921–948.
- (42) Thornton, K.; Gorenstein, D. G. Structure of glucagon-like peptide (7-36) amide in a dodecylphosphocholine micelle as determined by 2D NMR. *Biochemistry* **1994**, *33*, 3532–3539.
- (43) Arora, A.; Abildgaard, F.; Bushweller, J. H.; Tamm, L. K. Structure of outer membrane protein A transmembrane domain by NMR spectroscopy. *Nat. Struct. Biol.* **2001**, *8*, 334–338.
- (44) Karslake, C.; Piotto, M. E.; Pak, Y. K.; Weiner, H.; Gorenstein, D. G. 2D NMR and structural model for a mitochondrial signal peptide bound to a micelle. *Biochemistry* **1990**, *29*, 9872–9878.
- (45) Ying, J.; Ahn, J. M.; Jacobsen, N. E.; Brown, M. F.; Hruby, V. J. NMR solution structure of the glucagon antagonist [desHis1, desPhe6, Glu9]glucagon amide in the presence of perdeuterated dodecylphosphocholine micelles. *Biochemistry* **2003**, *42*, 2825–2835.
- (46) Jacobsen, N. E.; Abadi, N.; Sliwkowski, M. X.; Reilly, D.; Skelton, N. J.; Fairbrother, W. J. High-resolution solution structure of the EGF-like domain of heregulin- α . *Biochemistry* **1996**, *35*, 3402–3417.
- (47) Porreca, F.; Heyman, J. S.; Mosberg, H. I.; Omnaas, J. R.; Vaught, J. L. Role of μ and δ receptors in the supraspinal and spinal analgesic effects of [D-Pen², D-Pen⁵]enkephalin in the mouse. *J. Pharmacol. Exp. Ther.* **1987**, *241*, 393–400.
- (48) Porreca, F.; Mosberg, H. I.; Hurst, R.; Hruby, V. J.; Burks, T. F. Roles of μ , δ and κ opioid receptors in spinal and supraspinal mediation of gastrointestinal transit effects and hot-plate analgesia in the mouse. *J. Pharmacol. Exp. Ther.* **1984**, *230*, 341–348.
- (49) Audjgier, Y.; Mazarguil, H.; Gout, R.; Cros, J. Structure-activity relationships of enkephalin analogs at opiate and enkephalin receptors: correlation with analgesia. *Eur. J. Pharmacol.* **1980**, *63*, 35–46.
- (50) Agnes, R. S.; Lee, Y. S.; Davis, P.; Ma, S. W.; Badghisi, H.; Porreca, F.; Lai, J.; Hruby, V. J. Structure-activity relationships of bifunctional peptides based on overlapping pharmacophores at opioid and cholecystokinin receptors. *J. Med. Chem.* **2006**, *49*, 2868–2875.
- (51) Lee, Y. S.; Agnes, R. S.; Badghisi, H.; Davis, P.; Ma, S. W.; Lai, J.; Porreca, F.; Hruby, V. J. Design and synthesis of novel hydrazide-linked bifunctional peptides as δ/μ opioid receptor agonists and CCK-1/CCK-2 receptor antagonists. *J. Med. Chem.* **2006**, *49*, 1773–1780.
- (52) Lazaridis, T.; Mallik, B.; Chen, Y. Implicit solvent simulations of DPC micelle formation. *J. Phys. Chem. B* **2005**, *109*, 15098–15106.
- (53) Smith, L. J.; Bolin, K. A.; Schwalbe, H.; MacArthur, M. W.; Thornton, J. M.; Dobson, C. M. Analysis of main chain torsion angles in proteins: prediction of NMR coupling constants for native and random coil conformations. *J. Mol. Biol.* **1996**, *255*, 494–506.
- (54) Wagner, G.; Neuhaus, D.; Worgotter, E.; Vasak, M.; Kagi, J. H.; Wuthrich, K. Nuclear magnetic resonance identification of “half-turn” and 3(10)-helix secondary structure in rabbit liver metallothionein-2. *J. Mol. Biol.* **1986**, *187*, 131–135.
- (55) Smith, D.; Griffin, J. F. Conformation of [Leu⁵]enkephalin from X-ray diffraction: features important for recognition at opiate receptor. *Science* **1978**, *199*, 1214–1216.
- (56) Graham, W. H.; Carter, E. S., II; Hicks, R. P. Conformational analysis of Met-enkephalin in both aqueous solution and in the presence of sodium dodecyl sulfate micelles using multidimensional NMR and molecular modeling. *Biopolymers* **1992**, *32*, 1755–1764.
- (57) Wilmot, C. M.; Thornton, J. M. Analysis and prediction of the different types of beta-turn in proteins. *J. Mol. Biol.* **1988**, *203*, 221–232.
- (58) Hyberts, S. G.; Goldberg, M. S.; Havel, T. F.; Wagner, G. The solution structure of eglin c based on measurements of many NOEs and coupling constants and its comparison with X-ray structures. *Protein Sci.* **1992**, *1*, 736–751.
- (59) Beechem, J. M.; Brand, L. Time-resolved fluorescence of proteins. *Annu. Rev. Biochem.* **1985**, *54*, 43–71.
- (60) Vivian, J. T.; Callis, P. R. Mechanisms of tryptophan fluorescence shifts in proteins. *Biophys. J.* **2001**, *80*, 2093–2109.
- (61) Raguz, M.; Brnjac-Kraljevic, J. Resolved fluorescence emission spectra of PRODAN in ethanol/buffer solvents. *J. Chem. Inf. Model* **2005**, *45*, 1636–1340.
- (62) Neidigh, J. W.; Fesinmeyer, R. M.; Prickett, K. S.; Andersen, N. H. Exendin-4 and glucagon-like-peptide-1: NMR structural comparisons in the solution and micelle-associated states. *Biochemistry* **2001**, *40*, 13188–13200.
- (63) Datar, P.; Srivastava, S.; Coutinho, E.; Govil, G. Substance P: structure, function, and therapeutics. *Curr. Top. Med. Chem.* **2004**, *4*, 75–103.
- (64) Lewis, R. T.; Macleod, A. M.; Merchant, K. J.; Kelleher, F.; Sanderson, I.; Herbert, R. H.; Cascieri, M. A.; Sadowski, S.; Ball, R. G.; Hoogsteen, K. Tryptophan-derived NK1 antagonists: conformationally constrained heterocyclic bioisosteres of the ester linkage. *J. Med. Chem.* **1995**, *38*, 923–933.
- (65) Braunschweiler, L.; Ernst, R. R. Coherence Transfer by Isotropic Mixing: Application of Proton Correlation Spectroscopy. *J. Magn. Reson.* **1983**, *53*, 521–528.
- (66) Davis, D. G.; Bax, A. Assignment of complex proton NMR spectra via two-dimensional homonuclear Hartmann-Hahn spectroscopy. *J. Am. Chem. Soc.* **1985**, *107*, 2820–2821.
- (67) Subramanian, S.; Bax, A. Generation of pure phase NMR subspectra for measurement of homonuclear coupling constants. *J. Magn. Reson.* **1987**, *71*, 325–330.
- (68) Rance, M. Improved techniques for homonuclear rotating-frame and isotropic mixing experiments. *J. Magn. Reson.* **1987**, *74*, 557–564.
- (69) Bax, A.; Davis, D. G. MLEV-17-based two-dimensional homonuclear magnetization transfer spectroscopy. *J. Magn. Reson.* **1985**, *65*, 355–360.
- (70) Marion, D. W.; Wuthrich, K. Application of phase sensitive two-dimensional correlated spectroscopy (COSY) for measurements of ^1H - ^1H spin-spin coupling constants in proteins. *Biochem. Biophys. Res. Commun.* **1983**, *113*, 967–974.
- (71) Kumar, A.; Ernst, R. R.; Wuthrich, K. A two-dimensional nuclear Overhauser enhancement (2D NOE) experiment for the elucidation of complete proton-proton cross-relaxation networks in biological macromolecules. *Biochem. Biophys. Res. Commun.* **1980**, *95*, 1–6.
- (72) Piotto, M.; Saudek, V.; Sklenář, V. Gradient-tailored excitation for single-quantum NMR spectroscopy of aqueous solutions. *J. Biomol. NMR* **1992**, *2*, 661–665.
- (73) Press, W. H.; Vetterling, W. T.; Teukolsky, S. A. Numerical Recipes in C. In *The Art of Scientific Computing* Cambridge University Press: New York, 1988.
- (74) Havel, T. F. An evaluation of computational strategies for use in the determination of protein structure from distance constraints obtained by nuclear magnetic resonance. *Prog. Biophys. Mol. Biol.* **1991**, *56*, 43–78.
- (75) Weiner, S. J.; Kollman, P. A.; Case, D. A.; Singh, U. C.; Ghio, C.; Alagona, G. S.; Profeta, J.; Weiner, P. A New Force Field for Molecular Mechanical Simulation of Nucleic Acids and Proteins. *J. Am. Chem. Soc.* **1984**, *106*, 765–784.
- (76) Weiner, S. J.; Kollman, P. A.; Case, D. A. An all atom force field for simulations of proteins and nucleic acids. *J. Comput. Chem.* **1986**, *7*, 230–252.
- (77) Gough, C. A.; DeBolt, S. E.; Kollman, P. A. Derivation of fluorine and hydrogen atom parameters using liquid simulations. *J. Comput. Chem.* **1992**, *13*, 963–970.
- (78) Fluorine Parameters. <http://amber.scripps.edu/Questions/fluorine.html>. Accessed February 2007.
- (79) Boyd, N. D.; Kage, R.; Dumas, J. J.; Krause, J. E.; Leeman, S. E. The peptide binding site of the substance P (NK-1) receptor localized by a photoreactive analogue of substance P: presence of a disulfide bond. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 433–737.
- (80) Lorenzen, A.; Fuss, M.; Vogt, H.; Schwabe, U. Measurement of guanine nucleotide-binding protein activation by A1 adenosine receptor agonists in bovine brain membranes: stimulation of guanosine-5'-O-(3-[35S]thio) triphosphate binding. *Mol. Pharmacol.* **1993**, *44*, 115–123.
- (81) Porreca, F.; Burks, T. F. Affinity of normorphine for its pharmacologic receptor in the naive and morphine-tolerant guinea pig isolated ileum. *J. Pharmacol. Exp. Ther.* **1983**, *225*, 688–693.
- (82) Porreca, F.; LoPresti, D.; Ward, S. J. Opioid agonist affinity in the guinea pig ileum and mouse vas deferens. *Eur. J. Pharmacol.* **1990**, *179*, 129–139.

- (83) Tetko, I. V.; Gasteiger, J.; Todeschini, R.; Mauri, A.; Livingstone, D.; Ertl, P.; Palyulin, V. A.; Radchenko, E. V.; Zefirov, N. S.; Makarenko, A. S.; Tanchuk, V. Y.; Prokopenko, V. V. Virtual computational chemistry laboratory: design and description. *J. Comput.-Aided Mol. Des.* **2005**, *19*, 453–63.
- (84) VCCLAB. Virtual Computational Chemistry Laboratory. <http://www.vcclab.org>. Accessed February 2007.
- (85) Lipkowski, A. W.; Misicka, A.; Davis, P.; Stropova, D.; Janders, J.; Lachwa, M.; Porreca, F.; Yamamura, H. I.; Hruby, V. J. Biological activity of fragments and analogues of the potent dimeric opioid peptide, biphalin. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2763–2766.
- (86) Wishart, D. S.; Sykes, B. D.; Richards, F. M. The Chemical Shift Index: A Fast and Simple Method for the Assignment of Protein Secondary Structure through NMR Spectroscopy. *Biochemistry* **1992**, *31*, 1647–1651.
- (87) Andersen, N. H.; Neidigh, J. W.; Harris, S. W.; Lee, G. M.; Liu, Z. H.; Tong, H. Extracting Information from the Temperature Gradients of Polypeptide NH Chemical Shifts. 1. The Importance of Conformational Averaging. *J. Am. Chem. Soc.* **1997**, *119*, 8547–8561.

JM800389V