



# Erratum to ‘The corrosion of Alloy 690 in high-temperature aqueous media – thermodynamic considerations’ [J. Nucl. Mater. 294 (2001) 141–147] <sup>☆</sup>

R.J. Lemire <sup>\*</sup>, G.A. McRae

*AECL, Chalk River Laboratories, Chalk River, Ont., Canada K0J 1J0*

Unfortunately, in the final printed version of the paper, in Table 2 the final column indicating the Gibbs energies for formation of metals at 573 K is incorrect: the subscript ‘298’ should be ‘573’. Thus, the table should have appeared as follows.

Table 2

Gibbs energies of formation for metals in Alloy 690 with respect to the metals in their standard states at 298.15 K

Metal	$\Delta_f G_{298}^0$ (kJ mol <sup>-1</sup> ) (pure fcc)	$\Delta_f G_{298}^0$ (kJ mol <sup>-1</sup> ) (in Alloy 690)	$\Delta_f G_{573}^0$ (kJ mol <sup>-1</sup> ) (in Alloy 690)
Cr	10.65	12.52	-1.34
Ni	0.0	-4.94	-16.84
Fe	5.4	-6.08	-24.38

The publisher apologises for any confusion caused by this mistake.

<sup>☆</sup> PII of original article S0022-3115(01)00468-8.

<sup>\*</sup> Corresponding author. Tel.: +1-613 584 3311x3116; fax: +1-613 584 1220.

*E-mail address:* lemire@aecl.ca (R.J. Lemire).