



Application of the IEAF-2001 activation data library to activation analyses of the IFMIF high flux test module

U. Fischer ^{a,*}, P.P.H. Wilson ^b, D. Leichtle ^a, S.P. Simakov ^a, U.v. Möllendorff ^a,
A. Konobeev ^c, Yu. Korovin ^c, P. Pereslavytsev ^c, I. Schmuck ^a

^a *Forschungszentrum Karlsruhe, Association FZK-Euratom, Institut für Reaktorsicherheit, Postfach 3640, 76021 Karlsruhe, Germany*

^b *University of Wisconsin-Madison, 1500 Engineering Drive, Madison, WI 53706, USA*

^c *Institute of Nuclear Power Engineering, Studgorodok 1, 249020 Obninsk, Russia*

Abstract

A complete activation data library IEAF-2001 (intermediate energy activation file) has been developed in standard ENDF-6 format with neutron-induced activation cross sections for 679 target nuclides from $Z = 1$ (hydrogen) to $Z = 84$ (polonium) and incident neutron energies up to 150 MeV. Using the NJOY processing code, an IEAF-2001 working library has been prepared in a 256 energy group structure for enabling activation analyses of the International Fusion Material Irradiation Facility (IFMIF) D–Li neutron source. This library was applied to the activation analysis of the IFMIF high flux test module using the recent Analytical and Laplacian Adaptive Radioactivity Analysis activation code which is capable of handling the variety of reaction channels open in the energy domain above 20 MeV. The IEAF-2001 activation library was thus shown to be suitable for activation analyses in fusion technology and intermediate energy applications such as the IFMIF D–Li neutron source.

© 2002 Elsevier Science B.V. All rights reserved.

1. Introduction

The International Fusion Material Irradiation Facility (IFMIF) [1] will provide an intense high-energy neutron field produced by 40 MeV deuterons impinging on a thick lithium target. The resulting neutron source spectrum will extend up to $\cong 55$ MeV neutron energy. Numerous reaction channels are open in this energy region giving rise to a variety of activation and transmutation reactions which do not exist in the traditional energy domain below 20 MeV.

To perform activation calculations for the IFMIF D–Li neutron source requires (i) a complete activation data library comprising all target nuclides that may be present in the IFMIF facility and taking into account all activation and transmutation reactions that may occur

over the whole neutron energy range from 55 MeV down to thermal energy, (ii) an activation code capable of handling an arbitrary number for activation reaction channels. A suitable activation data library, the Intermediate Energy Activation File (IEAF-2001) [2] has been developed by a collaboration of Forschungszentrum Karlsruhe and the Institute of Nuclear Power Engineering, Obninsk, as part of the IFMIF project. The activation code ALARA (Analytical and Laplacian Adaptive Radioactivity Analysis), previously developed at the University of Wisconsin, Madison as an advanced computational tool for simulating induced activation in nuclear facilities [3], has the ability to handle the IEAF-2001 activation cross section data in a straightforward way.

In the following a short description is given of the IEAF-2001 library and the main features of the ALARA activation code. The application of the IEAF-2001 data to the activation analysis of the IFMIF high flux test module (HFTM) is presented including a description of the methodological approach applied. Results of the

* Corresponding author. Tel.: +49-7247 82 3407; fax: +49-7247 82 3718.

E-mail address: ulrich.fischer@irs.fzk.de (U. Fischer).

activation analysis for the HFTM with the low activation (LA) steel EUROFER-97 as structural material are presented and discussed.

2. Intermediate Energy Activation File IEF-2001

The Intermediate Energy Activation File IEF-2001 [2] has been developed by a collaboration of Forschungszentrum Karlsruhe and the Institute of Nuclear Power Engineering, Obninsk, with the objective to provide the nuclear data base required for the activation analysis of the IFMIF D–Li neutron source. The IEF-2001 library includes 679 (stable and unstable) target nuclides from $Z = 1$ (hydrogen) to 84 (polonium) with neutron induced reaction reactions from 10^{-5} eV to 150 MeV incident neutron energy.

The European Activation File EAF-99 [4] served as basis for the activation cross-section data below 20 MeV neutron energy. Threshold reaction cross-sections were evaluated on the basis of geometry dependent hybrid exciton and evaporation models and were adjusted to the EAF-99 cross-sections at 20 MeV. For these calculations, a modified version of the ALICE code [5] was used that takes into account pre-equilibrium emissions of d, t, ^3He , and α -clusters and γ -rays [6]. A new computational approach based on diffraction theory and a modified intra-nuclear cascade model [7] was employed to evaluate the reaction cross sections for the light nuclei up to $Z = 12$.

The IEF-2001 data library has been prepared in standard ENDF-6 data format making use of the MT = 5 (neutron, anything) option with the excitation functions stored in MF = 3 and the product nuclide vectors in MF = 6 (LAW = 0). An MF = 2 resonance parameter skeleton has been formally added to allow the processing of the activation library with the NJOY code [8].

A 256 group library has been generated with NJOY/GROUPR on the basis of the IEF-2001 point data to enable activation calculations for the IFMIF D–Li neutron source. The group data were generated for a temperature of 300 K assuming a flat weighting spectrum for the group cross section averaging. The final multi-group data set is a GENDF-formatted (group-wise ENDF) working library as provided by the GROUPR-module of NJOY and can be used by any activation code capable of handling an arbitrary number of reaction channels.

The 256 energy group structure consists of the 175 VITAMIN-J groups below 20 MeV and 81 groups above. Constant energy bins of 1 MeV and 2 MeV were used for the energy domains 20–50 and 50–150 MeV, respectively. This group structure is suitable for applications both in fusion technology and intermediate energy application such as irradiation simulations for the IFMIF D–Li neutron source.

3. ALARA activation code

ALARA is a next generation activation code designed for accuracy, speed and usability, and validated for use in fusion activation calculations [9]. Its basic features include multi-point simulations in a variety of geometries, user defined accuracy, and the tracking of stable isotope accumulation including light ions. The advanced features include the exact modelling of arbitrary hierarchical irradiation schedules and the ability to perform ‘reverse’ calculations. Most important for this application, ALARA does not rely on a table of fixed reaction types and reaction channels. The reactions are defined entirely by the library, requiring only a list of resultant isotopes and cross-sections for the production of each. This makes it ideal for extension to intermediate and high energy activation calculations.

4. IFMIF high flux test module

IFMIF provides a total neutron production of $\cong 1.1 \times 10^{17} \text{ s}^{-1}$ by bombarding a flowing liquid Lithium target with two 125 mA deuteron beams of 40 MeV. This results in a maximum neutron flux density of $\cong 7.5 \times 10^{14} \text{ cm}^{-2} \text{ s}^{-1}$ at the front surface of the HFTM which is placed downstream the beam adjacent to the Lithium target. The HFTM is a steel container of 20 cm width, 5 cm height and 5 cm depth housing 27 helium cooled rigs which hold the material specimens enclosed in capsules. The specimens, made of the LA steel Eurofer [10], will be subjected to a displacement damage accumulation of 20–50 dpa (iron) per full power year to simulate the radiation load of the highest loaded structural material components of a future fusion power reactor.

5. Neutron transport and activation calculations

The geometrical model of the HFTM consists of a rectangular Eurofer steel block with a mass density of 6.24 g/cm^3 and the proper dimensions divided into small cubic segments of size $0.5 \times 0.5 \times 0.5 \text{ cm}^3$. Neutron flux spectra are calculated with the newly developed McDeLicious code [11] in 256 energy groups for each of the 1000 segments of one HFTM quadrant. McDeLicious is a further enhancement of the McDeLi code, based on a built-in semi-empirical D–Li reaction model [12], with the ability to sample the generation of D–Li source neutrons from evaluated $\text{d} + {}^6\text{Li}$ cross-section data. Both McDeLi and McDeLicious make use of the MCNP Monte Carlo code [13] for performing the neutron and photon transport calculation. Two 125 mA deuteron beams are assumed in the McDeLicious calculation impinging on the lithium target at an angle of

10° with respect to the horizontal centre plane, see Fig. 1. The neutron flux spectra were normalised to a beam current of 2×125 mA. Fig. 2 compares the neutron flux spectra as calculated for the HFTM with McDeLicious and McDeLi. Note that only the McDeLicious approach is capable of representing the high energy part of the neutron spectrum above 40 MeV which is due to exothermic $\text{Li}(d, xn)$ reactions. It is essential to take into account this high energy component as it may affect the activation characteristics of the materials under irradiation.

ALARA activation calculations are performed for the Eurofer LA steel in each of the 1000 spatial segments taking proper account of the associated neutron flux spectrum. A continuous irradiation over a period of five full power years is assumed for the activation calculations with a truncation tolerance of 10^{-5} . The elemental composition of the Eurofer LA steel (Table 1) has been adopted from the specification for Eurofer 97 [10]. Yield data of manufactured Eurofer 97 largely agree with the specified elemental compositions.




Fig. 1. Schematic HFTM model with deuteron beams and lithium target.




Fig. 2. Neutron spectra calculated with McDeLi and McDeLicious for the IFMIF HFTM.

Table 1

Elemental composition of the Eurofer LA steel [10] assumed for the activation calculations

Element	wt%
Fe	88.983
O	0.010
C	0.105
Si	0.050
Mn	0.400
P	0.005
S	0.005
Cr	9.000
Ni	0.005
Mo	0.005
V	0.200
Nb	0.001
B	0.001
N	0.030
Al	0.010
Co	0.005
Cu	0.005
Ta	0.070
Ti	0.010
W	1.100

6. HFTM activation and afterheat

The specific activity and the afterheat power density of the HFTM are displayed in Figs. 3 and 4, respectively, as function of the cooling time. These data have been averaged over the HFTM volume. The total activity inventory and afterheat power as calculated for the HFTM with a volume of 500 cm^3 and a mass of 3.12 kg steel is given in Table 2. At shutdown, the HFTM afterheat power amounts to 3.5% of the direct nuclear heating at operation. One day after shutdown, this value




Fig. 3. Specific activity of HFTM as function of the cooling time.




Fig. 4. Afterheat power density of HFTM as function of the cooling time.

decreases to less than 1%. Only very few radio-nuclides are significantly contributing to the activity and the decay heat such as Mn-54, Mn-56 and Fe-55 in the time range up to a few years after irradiation and H-3, C-14 afterwards. Both Mn-54,-Mn-56, Fe-55 and H-3 are primarily produced through (n, p), (n, 2n) and (n, t) activation reactions on the natural iron isotopes whereas C-14 is an activation product of N-14. As a result of the various (n, t)-reactions, there is a total tritium generation of some 4 mg per full power year in the HFTM. Threshold reactions in the high energy range above 20 MeV incident neutron energy do not significantly contribute to the activity and the decay heat of the HFTM. Likewise, the HFTM activation is not affected by reaction products of the heavy Eurofer constituents W and Ta.

7. Conclusion and outlook

The newly developed activation data library IEAF-2001 has been tested through the application to the activation analysis of the IFMIF HFTM using the recent ALARA activation code. The IEAF-2001 activation library was thus shown to be suitable for activation analyses in fusion technology and intermediate energy applications such as the IFMIF D–Li neutron source.

The HFTM activation analysis revealed that threshold reactions in the high energy range above 20 MeV incident neutron energy do not significantly contribute to the activity and the decay heat. Likewise, the HFTM activation is not affected by reaction products of the heavy Eurofer constituents W and Ta.

The IEAF-2001 library is thus ready for distribution via the NEA data bank [2]. The IEAF-2001 cross-section data can be used with any activation code capable of handling an arbitrary number of reaction channels. Testing of the IEAF data against integral experiments is considered an important next step to qualify and further improve the IEAF library.

Acknowledgements

This work has been performed in the framework of the nuclear fusion programme of Forschungszentrum Karlsruhe and is supported by the European Union within the European Fusion Technology Programme.

References

- [1] M. Martone (Ed.), IFMIF International Fusion Materials Irradiation Facility, Conceptual Design Activity, Final Report, IFMIF CDA TEAM, ENEA Frascati, Report ENEA-RT/ERG/FUS/96-11, December 1996.
- [2] Yu. Korovin, A.Yu. Konobeev, P.E. Pereslavl'tsev, U. Fischer, U.v. Möllendorff, in: International Conference on Nuclear Data for Science and Technology (ND2001), 7–12 October, 2001, Tsukuba, Japan.
- [3] P.P.H. Wilson, D.L. Henderson, ALARA: Analytical and Laplacian Adaptive Radioactivity Analysis, vol. I, Technical Manual, University of Wisconsin, Report UWFDMM 1080, January 1998.
- [4] J.-Ch. Sublet, J. Kopecky, R.A. Forrest, The European Activation File: EAF-99 cross section library, UKAEA Fusion, Report UKAEA FUS 408, 1998.
- [5] M. Blann, ALICE 87 pre-compound nuclear model code, Report IAEA-NDS-93 REV.O, 1988.
- [6] A.Yu. Konobeyev, Yu.A. Korovin, P.E. Pereslavl'tsev, Code ALICE/ASH for the calculation of excitation functions, energy and angular distributions of emitted particles in nuclear reactions, INPE Report, Obninsk, February 1997.
- [7] A.Yu. Konobeyev, Yu.A. Korovin, P.E. Pereslavl'tsev, U. Fischer, U.v. Möllendorff, Nucl. Sci. Eng. 139 (2001) 1.

Table 2
Total afterheat power and activity inventory of the HFTM

	Shut down	Cooling time							
		1 h	1 d	30 d	1 y	5 y	10 y	100 y	1000 y
Activity (10^{14} Bq)	22	20	16	15	10	3.3	0.94	2.8×10^{-4}	7.2×10^{-6}
Afterheat power (W)	244	172	51	41	19	1.0	0.10	7.1×10^{-5}	4.6×10^{-6}

- [8] R.E. MacFarlane, D.W. Muir, R.M. Boicourt, The NJOY Nuclear Data Processing System, vol. I: User's Manual, Los Alamos National Laboratory, September 1993, LA-12740-M.
- [9] P.P.H. Wilson, H. Tsige-Tamirat, et al., *Fus. Technol.* 34 (1998).
- [10] R. Lindau, Forschungszentrum Karlsruhe, personal communication, 1997 and 2001.
- [11] S.P. Simakov et al., these Proceedings.
- [12] P.P.H. Wilson, Neutronics of the IFMIF Neutron Source: Development and Analysis, Forschungszentrum Karlsruhe, Report FZKA 6218, January 1999.
- [13] F. Briesmeister (Ed.), MCNP – A General Monte Carlo N-Particle Transport Code, Version 4C, LA-13709-M, April 2000.