


PRELIMINARY COMMUNICATION

Reformatsky-type reactions of α -functionally substituted organotin compounds

The chemical reactivity of the Sn- \dot{C} bond in compounds $R_3Sn\dot{C}H_2X$ (X, functional group with -M effect such as CN or COOEt) and $R_3Sn\dot{C}H_2CH_2X$ (defined as α - and β -functionally substituted, respectively) differs considerably¹ with respect to both nucleophilic reactions such as alkaline hydrolysis² or attack by lithium aluminium hydride³ or Grignard reagents³ and electrophilic reactions such as cleavage by halogens⁴. Ionic cleavage of the Sn- \dot{C} bond in compounds $R_3Sn\dot{C}H_2X$ is facilitated by the ability of the group X to stabilize negative charge on the adjacent carbon atom (*cf.* ref. 2), *e.g.*:




A similar resonance stabilization of an incipient carbanion is not possible for compounds $R_3SnCH_2\dot{C}H_2X$ and, therefore, attack is on X rather than on Sn- \dot{C} ^{2,3}.

The extreme susceptibility towards electrophilic attack (*e.g.* by halogen⁴) of compounds R_3SnCH_2X suggested to us that interaction with organic substrates containing strongly electrophilic carbon might enable their synthetic application as carbon-functional chain extension reagents. One isolated example of such a reaction has been reported in the literature, *i.e.* the addition of triethylstannylacetone to benzaldehyde⁵.

Quite recently the generation of carbon-carbon bonds via organotin compounds has received attention. Both triethyl(phenylethynyl)tin⁶ and triethylallyltin⁷ have been added to a few aldehydes and ketones. α -Functionally substituted organotin compounds undergo condensation with both allyl and benzyl bromide⁸. All these reactions have in common that the group transferred from tin to carbon may give rise to a resonance-stabilized carbanion.

In the present communication we wish to report that α -functionally substituted organotin nitriles, ketones, esters and amides readily add across the carbonyl group of a variety of aldehydes and ketones with quantitative formation of the corresponding β -triorganostannoxy-substituted carbon-functional derivatives:




Representative examples of such reactions have been listed in Table 1.

Since the Sn-O bond of the adducts obtained is easily hydrolyzed under acidic conditions an attractive new route to β -hydroxynitriles, -ketones, -esters and -amides has become available, *e.g.*:

TABLE 1


ADDITION OF α -FUNCTIONALLY SUBSTITUTED ORGANOTINS TO ALDEHYDES AND KETONES

Reaction ^a	Product ^b	Conditions for complete reaction
I	Bu ₃ SnCH ₂ CN + C ₆ F ₅ CHO Bu ₃ SnOCH(C ₆ F ₅)CH ₂ CN B.p., 146–156°/0.1 mm; n_D^{20} , 1.4813	16 h, room temp.
II	Bu ₃ SnCH ₂ CN + CCl ₃ CHO Bu ₃ SnOCH(CCl ₃)CH ₂ CN B.p., 130–133°/0.08 mm; n_D^{20} , 1.4995	16 h, room temp.
III	Et ₃ SnCH ₂ COMe + C ₆ F ₅ CHO Et ₃ SnOCH(C ₆ F ₅)CH ₂ COMe	Exothermic at room temp.
IV	Et ₃ SnCH ₂ COMe + C ₆ H ₅ CHO ^c Et ₃ SnOCH(C ₆ H ₅)CH ₂ COMe	Exothermic at room temp.
V	Et ₃ SnCH ₂ COMe + furfural Et ₃ SnOCH(C ₄ H ₃ O)CH ₂ COMe	Exothermic at room temp.
VI	Et ₃ SnCH ₂ COMe + <i>n</i> -PrCHO Et ₃ SnOCH(<i>n</i> -Pr)CH ₂ COMe	Exothermic at room temp.
VII	Et ₃ SnCH ₂ COMe + C ₆ H ₅ COCF ₃ Et ₃ SnOC(C ₆ H ₅)(CF ₃)CH ₂ COMe	Exothermic at room temp.
VIII	Et ₃ SnCH ₂ COMe + ClCH ₂ COCH ₂ Cl Et ₃ SnOC(CH ₂ Cl) ₂ CH ₂ COMe	Exothermic at room temp.
IX	Et ₃ SnCH ₂ COMe + cyclohexanone C ₆ H ₁₀ (Et ₃ SnO)(MeCOCH ₂) <i>gem</i>	48 h, room temp.
X	Et ₃ SnCH ₂ COOEt + C ₆ F ₅ CHO Et ₃ SnOCH(C ₆ F ₅)CH ₂ COOEt B.p., 107–108°/0.07 mm; n_D^{20} , 1.4751	16 h, room temp.
XI	Et ₃ SnCH ₂ COOEt + C ₆ H ₅ CHO Et ₃ SnOCH(C ₆ H ₅)CH ₂ COOEt	8 h/50°, ZnCl ₂ cat.
XII	Et ₃ SnCH ₂ COOEt + CCl ₃ CHO Et ₃ SnOCH(CCl ₃)CH ₂ COOEt B.p., 108–114°/0.25 mm; n_D^{20} , 1.4973	Exothermic at room temp.
XIII	Et ₃ SnCH ₂ COOEt + C ₆ H ₅ COCF ₃ Et ₃ SnOC(C ₆ H ₅)(CF ₃)CH ₂ COOEt	5 h/60°, ZnCl ₂ cat.
XIV	Et ₃ SnCH ₂ CONEt ₂ + C ₆ F ₅ CHO Et ₃ SnOCH(C ₆ F ₅)CH ₂ CONEt ₂	Exothermic at room temp.
XV	Et ₃ SnCH ₂ CONEt ₂ + furfural Et ₃ SnOCH(C ₄ H ₃ O)CH ₂ CONEt ₂	4 h/60°
XVI	Et ₃ SnCH ₂ CONEt ₂ + CCl ₃ CHO Et ₃ SnOCH(CCl ₃)CH ₂ CONEt ₂	Exothermic at room temp.

^a Reactions were carried out by mixing equimolar amounts of the reactants under N₂ in the absence of a solvent.^b All compounds have been characterized by NMR spectroscopy. ^c Cf. ref. 5; the reaction proceeds equally well with *o*- and *p*-chloro-, *m*- and *p*-nitrobenzaldehyde.

As appears from Table 1 adduct-formation runs to completion under particularly mild conditions if an electron-withdrawing group is attached to the carbonyl group. If necessary the reaction can be catalyzed by Lewis acids (reactions XI and XIII, Table 1; cf. ref. 7) suggesting once more that electrophilic attack of the carbonyl carbon on the Sn–C bond is the rate-determining step in these reactions (cf. ref. 9).


The 1 : 1 adduct XII in the presence of excess of chloral affords a 1 : 2 adduct (cf. comparable reactions with trialkyltin hydrides⁹ and alkoxides¹⁰):



Although IR spectroscopy is useful, the progress of the additions with aldehydes is more conveniently followed by observing the very characteristic changes in the PMR spectrum. Due to the presence of a centre of asymmetry the spectra of the aldehyde adducts reveal an ABX pattern with eight lines at high field and four lines at low field.

The products listed in Table I are high-boiling liquids which cannot be distilled without some decomposition taking place. Since as appears from the PMR spectra, the products formed are better than 98 % pure no further attempts at purification were made.

In order to avoid olefin-forming elimination malonic acid⁷ or oxalic acid is preferred over ethereal hydrochloric acid⁶ for converting the adducts into the corresponding β -hydroxy derivatives, e.g.:



The direct introduction of carbon-functional groups has remained an important problem in organic synthesis (functionally substituted halides are not amenable to Grignard-formation; the Reformatsky-reaction which is often attended by side reactions is largely limited to introduction of the ester function). Therefore, we wish to draw attention to this particularly facile organometallic method of carbon-functional chain extension, the scope of which is the subject of further investigation at our Institute.

Details of this research will be published in the full paper.

Acknowledgement

Financial support by the International Tin Research Council (Dr. E. S. HEDGES), London, is gratefully acknowledged. Thanks are due to Mr. F. VERBEEK for skilful experimental assistance. The authors are indebted to Dr. W. DRENTH, Dr. A. J. LEUSINK and Mr. J. W. MARSMAN for discussions.

*Institute for Organic Chemistry TNO
Utrecht (The Netherlands)*

J. G. NOLTES
H. M. J. C. CREEMERS
G. J. M. VAN DER KERK

- 1 J. G. NOLTES AND G. J. M. VAN DER KERK, *Functionally Substituted Organotin Compounds*, Tin Research Institute, Greenford, 1958.
- 2 G. J. M. VAN DER KERK AND J. G. NOLTES, *J. Appl. Chem.*, 9 (1959) 113.
- 3 G. J. M. VAN DER KERK AND J. G. NOLTES, *J. Appl. Chem.*, 9 (1959) 176.
- 4 G. J. M. VAN DER KERK AND J. G. NOLTES, *J. Appl. Chem.*, 9 (1959) 179.
- 5 S. V. PONOMAREV AND I. F. LUTSENKO, *Zh. Obshch. Khim.*, 34 (1964) 3450.
- 6 R. G. MIRSHOV AND V. M. VLASOV, *Zh. Obshch. Khim.*, 36 (1966) 562.
- 7 K. KÖNIG AND W. P. NEUMANN, *Tetrahedron Letters* (1967) 495.
- 8 M. PEREYRE, G. COLIN AND J. VALADE, *C. R. Acad. Sci, Paris, Ser. C*, 264 (1967) 1204.
- 9 A. J. LEUSINK, Ph.D.Thesis, Utrecht, 1966.
- 10 A. G. DAVIES AND W. R. SYMES, *J. Chem. Soc. (C)*, (1967) 1009.

(Received October 24th, 1967)