


product. The product of butadiene-1,1,4,4- d_4 dimerization was confirmed by its NMR spectrum together with other evidence to be 3-methyl- d_3 -hepta-1,4,6-triene-1,1,4,7,7- d_5 . The NMR spectrum of the deuterio-methylheptatriene as compared with that of 3-methylhepta-1,4,6-triene indicated the absence of the doublet (τ 8.90) assigned to the methyl protons, the change of the sextet (τ 7.15) into the doublet assigned to H* and the simplification of the multiplets of vinyl protons. The result suggests that the hydrogen transfer in the linear dimerization of butadiene by the cobalt complex catalyst occurs only at 1- and 4-positions of butadiene.



Department of Industrial Chemistry,
University of Tokyo,
Hongo, Tokyo (Japan)
Tokyo Institute of Technology,
Research Laboratory of Resources
Utilization, Meguro, Tokyo (Japan)

TARO SAITO
YASUZO UCHIDA
AKIRA MISONO
AKIO YAMAMOTO
KAZUHIKO MORIFUJI
SAKUJI IKEDA


- 1 S. OHTSUKA, T. TAKETOMI AND T. KIKUCHI, *J. Chem. Soc. Japan, Ind. Chem. Sect.*, 66 (1963) 1094.
- 2 T. SAITO, T. OHNO, Y. UCHIDA AND A. MISONO, *J. Chem. Soc. Japan, Ind. Chem. Sect.*, 66 (1963) 1099.
- 3 S. TANAKA, K. MABUCHI AND N. SHIMAZAKI, *J. Org. Chem.*, 29 (1964) 1626.
- 4 S. OHTSUKA, T. KIKUCHI, Y. TANAKA, F. IMAIZUMI AND K. MORI, *The Chemistry of High Polymers (Japan)*, 22 (1965) 24.
- 5 G. N. SCHRAUZER AND J. KOHNLE, *Chem. Ber.*, 97 (1964) 3056.
- 6 D. DOLPHIN AND A. W. JOHNSON, *Chem. Commun.*, (1965) 494.
- 7 H. YAMAZAKI AND N. HAGIHARA, *Bull. Chem. Soc. Japan*, 38 (1965) 2212.
- 8 R. B. KING, *Inorg. Chem.*, 5 (1966) 82.
- 9 A. YAMAMOTO, K. MORIFUJI, S. IKEDA, T. SAITO, Y. UCHIDA AND A. MISONO, *J. Am. Chem. Soc.*, 87 (1965) 4652.

Received June 27th, 1966

J. Organometal. Chem., 6 (1966) 572-573


Halogenated organotin compounds as CX₂ transfer agents

We have expended considerable effort on studies of the CX₂ transfer reactions of phenyl(trihalomethyl)mercury compounds¹, and it seemed of interest to evaluate trihalomethyl compounds of other metals as dihalocarbene sources. The earlier work of Clark and Willis² showed trimethyl(trifluoromethyl)tin to be a source of CF₂ at 140-150°. This suggested that other trihalomethyltin compounds should undergo thermal α -elimination of trimethyltin halide to give dihalocarbenes, and we chose to commence our investigation of this aspect of halomethyl metal chemistry with a study of trimethyl(trichloromethyl)tin and trimethyl(bromodichloromethyl)tin. The preparation of the former was accomplished in 62 % yield using the reaction of trimethyltin chloride with trichloromethyl lithium³⁻⁵ (eqn. 1). A route to (CH₃)₃SnCCl₂Br was provided




J. Organometal. Chem., 6 (1966) 573-576

The somewhat difficult purification of these trihalomethyltin compounds and their facile hydrolysis prompted us to investigate the possible use of organotin trihaloacetates as CX_2 transfer agents. The thermal decarboxylation of some organotin esters



is known to occur, e.g., $(C_4H_9)_3SnO_2CCH_2CN \longrightarrow (C_4H_9)_3SnCH_2CN^8$ and $(C_4H_9)_3SnO_2CH \longrightarrow (C_4H_9)_3SnH^9$, to mention two recent examples, and thus it seemed possible that decarboxylation and CX_2 transfer could be effected by heating an $R_3SnO_2CCX_3$ compound in the presence of an olefin. This procedure, if successful, would have the advantage of using stable, easily prepared and purified organotin starting materials. Such a route to *gem*-dihalocyclopropanes could be realized, but the product yields were only moderate due to competing side reactions.

It was found that triphenyltin trichloroacetate was stable when heated in refluxing cyclohexene ($\sim 80^\circ$), but that decarboxylation and CCl_2 -transfer took place when this tin ester was heated in cyclooctene at reflux ($\sim 140^\circ$) under nitrogen. Gas evolution occurred, and subsequent analysis of the reaction mixture using GLPC showed that 9,9-dichlorobicyclo[6.1.0]nonane had been formed in 56% yield (eqn. 4).



Benzene (39%) also was present, and this suggests that a competing radical decomposition of the tin ester had occurred. Reaction of cyclooctene and the tin ester in 5:1 ratio in a solvent (chlorobenzene or diglyme, at reflux) gave only slightly diminished yields of the dichlorocyclopropane product. Very similar results were obtained with trimethyltin trichloroacetate. The reaction of triphenyltin tribromoacetate with cyclooctene at reflux for 3 h gave 9,9-dibromobicyclo[6.1.0]nonane in 36% yield. Attempts to effect CF_2 transfer to olefins with triphenyltin trifluoroacetate or chlorodifluoroacetate were unsuccessful. Studies are in progress which hopefully will reveal whether R_3SnCCl_3 and R_3SnCBr_3 are intermediates in these reactions or whether decarboxylation and CX_2 transfer to olefin occur simultaneously in a concerted fashion.

It is readily apparent that phenyl(trihalomethyl)mercury compounds are to be preferred as CX_2 transfer agents. They are easily prepared and purified, are stable toward hydrolysis and their somewhat diminished stability allows rapid transfer of CX_2 to olefins and other substrates to be carried out at 80° .

Details of this work will be presented at a later date. We publish our present findings at this time in view of the current interest in trihalomethyl derivatives of the Group IV elements, notably the previously reported $(C_4H_9)_3SnCCl_3$ ¹⁰, $(C_6H_5)_3Pb-CCl_3$ ^{11,12} and $(C_6H_5)_3PbCBr_3$ ¹².

Acknowledgements

The authors are grateful to the Directorate of Chemical Sciences, U.S. Air Force Office of Scientific Research, for generous support of this work and to M&T Chemicals, Inc. for gifts of chemicals. This investigation was supported in part by Public Health Fellowship 5-F1-GM-23,742-02 (to F.M.A.).

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139, (U.S.A.)

DIETMAR SEYFERTH^a
FRANK M. ARMBRECHT JR.^b
BELA PROKAI^c
RONALD J. CROSS^d

- 1 D. SEYFERTH, in *Proceedings of the Robert A. Welch Foundation Conferences on Chemical Research. IX. Organometallic Compounds*, Robert A. Welch Foundation, Houston, Texas, 1966, in press (review).
- 2 H. C. CLARK AND C. J. WILLIS, *J. Am. Chem. Soc.*, **82** (1960) 1888.
- 3 W. T. MILLER, JR. AND D. M. WHALEN, *J. Am. Chem. Soc.*, **86** (1964) 2089.
- 4 G. KÖBRICH, K. FLORY AND W. DRISCHEL, *Angew. Chem.*, **76** (1964) 536.
- 5 D. F. HOEG, D. I. LUSK AND A. L. CRUMBLISS, *J. Am. Chem. Soc.*, **87** (1965) 4147.
- 6 M. E. GORDON, K. V. DARRAGH AND D. SEYFERTH, *J. Am. Chem. Soc.*, **88** (1966) 1831.
- 7 D. SEYFERTH AND J. Y.-P. MUI, unpublished work.
- 8 G. J. M. VAN DER KERK AND J. G. A. LUIJTEN, *J. Appl. Chem.*, **6** (1956) 93.
- 9 M. OHARA AND R. OKAWARA, *J. Organometal. Chem.*, **3** (1965) 484.
- 10 A. G. DAVIES AND W. R. SYMES, *J. Organometal. Chem.*, **5** (1966) 394.
- 11 L. C. WILLEMSSENS AND G. J. M. VAN DER KERK, *Investigations in the Field of Organolead Chemistry*, International Lead Zinc Research Organization, Inc., New York, 1965, pp. 42, 98.
- 12 A. G. DAVIES AND R. J. PUDDEPHATT, *J. Organometal. Chem.*, **5** (1966) 590.

Received July 19th, 1966

-
- ^a Alfred P. Sloan Foundation Fellow, 1962-1966.
 - ^b National Institutes of Health Predoctoral Fellow.
 - ^c Postdoctoral Research Associate, 1964-1966.
 - ^d Postdoctoral Research Associate, 1965-1966.

J. Organometal. Chem., **6** (1966) 573-576