

RELATIVE ELECTRON-RELEASING EFFECTS OF TRIMETHYL-SILYL, -GERMYL, AND -STANNYL GROUPS

R. W. BOTT, C. EABORN AND D. R. M. WALTON

*Department of Chemistry, University of Leicester,
and**The Chemical Laboratory, University of Sussex, Brighton, (Great Britain)*

(Received December 2nd, 1963)

INTRODUCTION

Because of the uncertainty about the relative electronegativities of the Group IVB elements¹, the rates of cleavage by aqueous-methanolic perchloric acid of the aryl-silicon bonds were measured for *p*-Me₃M-CH₂-C₆H₄-SiMe₃ compounds in which M = Si and Ge, and it was found that the compound containing germanium was more reactive than that containing silicon, indicating that the trimethylgermyl group releases electrons more strongly than the trimethylsilyl group². (Aspects of the infrared spectra of Me₃M-O-CO-Me compounds indicate the same order³.) Information could not be obtained about the trimethylstannyl group because with the compound *p*-Me₃Sn-CH₂-C₆H₄-SiMe₃ the benzyl-tin bond was cleaved more rapidly than the aryl-silicon bond. To remedy this omission, we have measured the rates of cleavage of the aryl-tin bonds of the compounds *p*-Me₃M-CH₂-C₆H₄-SnMe₃ in which M = Si, Ge, and Sn; the benzyl-tin bond is stable at the low acid concentrations involved.

RESULTS AND DISCUSSION

The results are shown in the table as pseudo first-order rate constants, *k*₁, and as relative rates; the strength of the acid shown is that of the aqueous acid before mixing.

TABLE
CLEAVAGE OF Me₃M-CH₂-C₆H₄-SnMe₃ COMPOUNDS BY A MIXTURE OF METHANOL
(5 vol.) AND AQUEOUS PERCHLORIC ACID (2 vol.) AT 30.25°





M	10 ² ·HClO ₄ ⁻ (M)	10 ² · <i>k</i> ₁ (min ⁻¹)	Rel. Rate	M	10 ² ·HClO ₄ ⁻ (M)	10 ² · <i>k</i> ₁ (min ⁻¹)	Rel. Rate
Si	2.10	56	1.00	Si	1.07	34	1.00
Ge	2.10	76	1.36	Ge	1.07	46	1.35
Sn	2.10	180	3.21	Sn	1.07	109	3.21

The order of reactivity is (M =) Sn > Ge > Si, and so the order of electron release to the reaction centre is Me₃Sn > Me₃Ge > Me₃Si. The difference between the effects of the trimethylgermyl and -silyl groups is smaller than in cleavage of

Me₃M-CH₂-C₆H₄-SiMe₃ compounds², in keeping with the generally smaller spread of rates in protodesannylation than in protodesilylation (*e.g.*, $f_p^{Me} = 5.4$, in the former⁴, and 21 in the latter⁵ reaction).

It was previously assumed that the magnitude of the electron release to the aromatic ring by a Me₃M-CH₂ group was a measure of the inductive release of electrons by the Me₃M group to a saturated carbon atom². It is not, however, unambiguously so, since there is a possibility of some interaction between the p_π electrons of the ring and the d -orbitals of the metal atom, the overall electron release towards the ring representing the difference between inductive supply of electrons and return of them by the p_π - d_π bonding. Such bonding would be analogous to the d_π - p_π bonding suggested to occur between silicon and a halogen or oxygen atom on a carbon atom adjacent to the silicon⁶. If there is interaction between the metal atom and the p_π -orbital of the 1-carbon atom of the ring it must, of course, be small, since the Me₃M-CH₂ groups are strongly electron releasing groups (markedly more so than the methyl group itself), and in our opinion its variation from metal to metal is unlikely to reverse the trend set by the inductive release of electrons, which we thus believe to increase in the order Me₃Si < Me₃Ge < Me₃Sn*.

(*p*-Trimethylstannylbenzyl)trimethylstannane was made from the Grignard reagent obtained in tetrahydrofuran from (*p*-bromobenzyl)trimethylstannane (which was itself made from trimethylstannyl lithium⁸ and *p*-bromobenzyl bromide in the same solvent). Attempts to make it from (*p*-chlorobenzyl)trimethylstannane were unsuccessful, and some of them confirmed the ease with which the benzyl-tin bond is cleaved by organolithium reagents (*cf.* ref. 9) or lithium. For example, the chloride reacted readily with lithium in tetrahydrofuran, and when the resulting mixture was hydrolysed, *p*-chlorotoluene was obtained in good yield, but when bromotrimethylstannane was added before the hydrolysis, hexamethyldistannane was obtained in good yield and most of the (*p*-chlorobenzyl)trimethylstannane was recovered. The reactions involved appear to be those in equations (1)-(4).



Again, when (*p*-chlorobenzyl)trimethylstannane was treated with *n*-butyllithium in ether and the mixture hydrolysed, *p*-chlorotoluene and *n*-butyltrimethylstannane were obtained in approximately equimolar proportions.

Such cleavages account for the failures² to prepare (*p*-carboxybenzyl)trimethylstannane from the *p*-chloro-compound via the *p*-lithio-compound.

* It is not known with certainty how the ease of d_π - p_π bonding varies for tetrahedrally-bonded silicon, germanium, and tin, but there is evidence that it falls from silicon to germanium in the anions derived from Me₃M-Ph compounds⁷.

EXPERIMENTAL

*Trimethyl(*p*-trimethylstannylbenzyl)germane*

Bromotrimethylstannane (5.4 g, 0.022 mole) was refluxed for 2 h, with the organolithium reagent from (*p*-chlorobenzyl)trimethylgermane¹⁰ (5.4 g, 0.022 mole) and lithium (c.70 g, 0.10 g-atom) in ether (200 ml). Treatment with saturated aqueous ammonium chloride, followed by separation, drying, and fractionation of the ethereal layer gave trimethyl(*p*-trimethylstannylbenzyl)germane (1.2 g, 15%), b.p. 114–115°/2.4 mm, n_D^{25} 1.5278. (Found: C, 41.9; H, 6.7. C₁₃H₂₄GeSn calcd.: C, 42.0; H, 6.5%) and unchanged (*p*-chlorobenzyl)trimethylgermane (2.8 g, 52%).

*(*p*-Bromobenzyl)trimethylstannane*

A little (ca. 2 ml) of a solution of bromotrimethylstannane (30 g, 0.123 mole) in tetrahydrofuran (50 ml) was added to lithium shot (2.1 g, 0.30 g-atom) in the same solvent (80 ml), the mixture becoming warm and dark red. The remaining bromotrimethylstannane solution was added with stirring during 30 min, the temperature of the mixture being kept at about –10°. After a further 10 min stirring, the solution was filtered under nitrogen through glass wool. The yield of trimethylstannylolithium was 73% (by titration).

p-Bromobenzyl bromide (5.0 g, 0.020 mole) in tetrahydrofuran (80 ml) was added during 10 min to a solution of trimethylstannylolithium (0.022 mole) in the same solvent (40 ml). The mixture was warmed on a water bath for 10 min then treated with aqueous ammonium chloride. Ether extraction, followed by washing, drying (Na₂SO₄), and fractionation of the ethereal layer gave (*p*-bromobenzyl)trimethylstannane (6.0 g, 89%), b.p. 99°/1.5 mm (lit.² b.p. 126°/6 mm), n_D^{25} 1.5770.

*Trimethyl(*p*-trimethylstannylbenzyl)stannane*

(*p*-Bromobenzyl)trimethylstannane (16.7 g, 0.05 mole) was brought into reaction with magnesium turnings (1.94 g, 0.08 g-atom) in tetrahydrofuran (40 ml) containing ethyl bromide (ca. 0.5 ml). Bromotrimethylstannane (14.6 g, 0.060 mole) in tetrahydrofuran (20 ml) was added, and the mixture was refluxed for 15 min, then cooled and worked up in the usual way to give trimethyl(*p*-trimethylstannylbenzyl)stannane (6.7 g, 64%), b.p. 98–100°/0.04 mm, n_D^{25} 1.5550. (Found: C, 37.7; H, 5.8. C₁₃H₂₄Sn₂ calcd.: C, 37.4; H, 5.8%.)

*Attempted preparation of trimethyl(*p*-trimethylstannylbenzyl)stannane from (*p*-chlorobenzyl)trimethylstannane*

(i) Most of the *p*-chlorobenzyl chloride was recovered from attempts to couple it with bromotrimethylstannane via a Wurtz-Fittig reaction with sodium in boiling light petroleum (b.p. 60–80°) or toluene, or via a Grignard reagent in tetrahydrofuran.

(ii) An exothermic reaction occurred when (*p*-chlorobenzyl)trimethylstannane (72 g, 0.25 mole) was added to lithium shot (4.1 g, 0.60 g-atom) in tetrahydrofuran (150 ml) at 0°. The mixture was halved; one half was treated with aqueous ammonium chloride, and the usual working-up (involving ether extraction) gave *p*-chlorotoluene (12.7 g, 0.11 mole, 88%). The other half was boiled for 10 min with bromotrimethylstannane before the hydrolysis, and the usual working-up gave hexamethyldistannane (33 g, 0.10 mole, 80%), and (*p*-chlorobenzyl)trimethylstannane (34.6 g, 0.12 mole, 96%).

(iii) A mixture of *n*-butyllithium (0.10 mole) and (*p*-chlorobenzyl)trimethylstannane (28.8 g, 0.10 mole) in ether (130 ml) was stirred for 2 h at room temperature. The usual working-up gave unchanged (*p*-chlorobenzyl)trimethylstannane (8.2 g, 28%) and material (16 g) of b.p. 150–164°, which was shown by vapour-phase chromatography to be an approximately equimolar mixture of *p*-chlorotoluene and *n*-butyltrimethylstannane.

Rate studies

A methanolic solution (5 ml) of the $p\text{-Me}_3\text{M-CH}_2\text{-C}_6\text{H}_4\text{-SnMe}_3$ compound [$10^4 \times$ concn.: (M =) Si, 13; Ge, 12; Sn, 7 M] was mixed with 2 ml of aqueous perchloric acid of the strength specified in the Table. Rate measurements were determined spectrophotometrically at $30.25^\circ \pm 0.02^\circ$ by the method previously described^{4,5}, "infinity" values being measured on the reaction mixture after 10 half-lives. The wave-lengths employed were (M =) Si, 278.5; Ge, 256; Sn, 272 m μ .

Products

With trimethyl(*p*-trimethylstannylbenzyl)stannane there was a slight possibility that cleavage of the benzyl-Sn might accompany cleavage of the aryl-Sn bond, but the ultra-violet spectrum of the reaction mixture after 10 half-lives was identical with that of a solution of benzyltrimethylstannane. A small-scale preparative cleavage was carried out with stronger acid than used for rate studies. A solution of trimethyl(*p*-trimethylstannylbenzyl)stannane (5.0 g) in methanol (80 ml) was mixed with 1.0 M aqueous perchloric acid (20 ml); the mixture was warmed on a water bath for 12 min, and then added to an excess of saturated aqueous ammonium chloride. Ether extraction, followed by washing, drying (Na_2SO_4) and fractionation of the extract gave benzyltrimethylstannane (3.4 g, 71%), b.p. 96°/1.1 mm, n_D^{25} 1.5420. No toluene or trimethyl-*p*-tolylstannane was detected.

ACKNOWLEDGEMENTS

This work was supported by the Office of Aerospace Research, United States Air Force, through its European Office. We thank Dr. L. SPIALTER of the Aeronautical Research Laboratory, Wright Air Development Center, for his interest and encouragement, and the Royal Society for the loan of apparatus.

SUMMARY

The rates of cleavage of the aryl-tin bond of $p\text{-Me}_3\text{M-CH}_2\text{-C}_6\text{H}_4\text{-SnMe}_3$ compounds by aqueous methanolic perchloric acid increases in the order (M =) Si < Ge < Sn, which means that electron release by the Me_3M groups also increases in this order.

REFERENCES

- 1 A. L. ALLRED AND E. G. ROCHOW, *J. Inorg. Nucl. Chem.*, 5 (1958) 264, 269; 20 (1961) 167; A. L. ALLRED, *J. Inorg. Nucl. Chem.*, 17 (1961) 215; R. S. DRAGO, *J. Inorg. Nucl. Chem.*, 15 (1960) 237.
- 2 R. W. BOTT, C. EABORN, K. C. PANDE AND T. W. SWADDLE, *J. Chem. Soc.*, (1962) 1217.
- 3 T. N. SRIVASTAVA AND M. ONYSZCHUK, *Can. J. Chem.*, 41 (1963) 1244.

- 4 C. EABORN AND J. A. WATERS, *J. Chem. Soc.*, (1961) 542.
- 5 C. EABORN, *J. Chem. Soc.*, (1956) 4858.
- 6 A. G. BROOK, M. A. QUIGLEY, G. J. D. PEDDLE, N. V. SCHWARTZ AND C. M. WARNER, *J. Am. Chem. Soc.*, 82 (1960) 5102; O. M. STEWARD AND O. R. PIERCE, *J. Am. Chem. Soc.*, 83 (1961) 4932.
- 7 J. A. BEDFORD, J. R. BOLTON, A. CARRINGTON AND R. H. PRINCE, *Trans. Faraday Soc.*, 59 (1963) 53.
- 8 C. TAMBORSKI, F. E. FORD, W. L. LERN, G. J. MOORE AND E. J. SOLOSKI, *J. Org. Chem.*, 27 (1962) 619; C. TAMBORSKI, F. E. FORD AND E. J. SOLOSKI, *J. Org. Chem.*, 28 (1963) 237.
- 9 H. GILMAN AND S. D. ROSENBERG, *J. Org. Chem.*, 24 (1959) 2063.

J. Organometal. Chem., 2 (1964) 154-158