

Although quite stable, all these solvato complexes are decomposed by halides: KBr pellets of II for example show a  $\nu(\text{N}-\text{H})$  at  $3200 \text{ cm}^{-1}$  (characteristic of compounds I [5]) instead of  $3340 \text{ cm}^{-1}$ .

Further studies of reactions 1 and 2 with a range of chelating C—N ligands are in progress.

## References

- 1 P.K.F. Chin and F.R. Hartley, Inorg. Chem., 15 (1976) 982.
- 2 C. White, S.J. Thompson and P.M. Maitles, Chem. Commun., (1976) 409.
- 3 H.C. Clark and C.R. Jablonski, Inorg. Chem., 13 (1974) 2213.
- 4 H.C. Clark and H. Kurosawa, J. Organometal. Chem., 36 (1972) 399.
- 5 J. Dehand, M. Pfeffer and M. Zinsius, Inorg. Chim. Acta, 13 (1975) 229.
- 6 L.J. Bellamy, The Infrared Spectra of Complex Molecules, 2nd edition, Methuen, London, 1958.
- 7 H. Suhr, Organische Chemie in Einzeldarstellungen, Vol. 8, Anwendung der kernmagnetischen Resonanz in der organischen Chemie, Springer Verlag, Berlin, 1965.

## Erratum

---

*J. Organometal. Chem.*, Vol. 108, No. 1 (March 16th, 1976)

Page 9, Table 2

The  $^{11}\text{B}$  chemical shift values for  $\text{Cl}_2\text{BC}_2\text{F}_3$  and  $\text{BrB}(\text{C}_2\text{F}_3)_2$  should read -49.4.