

Preliminary communication

MANGANESE(I) CARBONYLS WITH σ -BONDED ALKYNYL LIGANDS. THE CRYSTAL STRUCTURE OF $[\text{Co}_2(\text{CO})_6][\text{PhC}\equiv\text{CMn}(\text{CO})_4\{\text{P}(\text{C}_6\text{H}_{11})_3\}]$

G. A. CARRIEDO, V. RIERA*,

Departamento de Química Inorgánica, Universidad de Oviedo, Oviedo (Spain)

D. MIGUEL,

Departamento de Química Inorgánica, Universidad de Valladolid, Valladolid (Spain)

A. M. MANOTTI LANFREDI and A. TIRIPICCHIO*



Istituto di Chimica Generale ed Inorganica, Università di Parma, Centro di Studio per la Strutturistica Diffattometrica del C.N.R., Via M. D'Azeglio 85, I-43100 Parma (Italy)

(Received April 24th, 1984)

Summary

The preparation of some σ -alkynylmanganese(I) carbonyl complexes is described. The crystal structure of the compound $[\text{Co}_2(\text{CO})_6][\text{PhC}\equiv\text{CMn}(\text{CO})_4(\text{PCy}_3)]$ (Cy = cyclohexyl) has been established by X-ray diffraction.

Octahedral manganese(I) carbonyl derivatives containing σ -bonded alkynyl ligands have been little studied, and as far as we are aware, only the compound $\text{PhC}\equiv\text{CMn}(\text{CO})_5$ [1] and the somehow related acetylide complex $\text{BrMn}(\text{CO})_4\text{C}\equiv\text{CPh}_3$ [2] have been reported. We have found that various new types of σ -alkynyl complexes of manganese(I) can be prepared from halocarbonyls $\text{BrMn}(\text{CO})_n\text{L}_{5-n}$ or perchlorato complexes $\text{O}_3\text{ClOMn}(\text{CO})_n\text{L}_{5-n}$ by means of the general reactions 1 and 2; the ligands are phosphines, diphosphines or phosphites; R = Ph or Bu^t.



(L₂ = dppe or bipy; R = H or Ph)

The products are fairly stable crystalline solids and, in contrast to the inertness of the "parent" $\text{PhC}\equiv\text{CMn}(\text{CO})_5$ [1], can be used as precursors for other

TABLE 1
ANALYTICAL, INFRARED, AND MELTING POINT DATA FOR SOME σ -ALKYNYL COMPOUNDS

Compound	Analysis (Found (calcd.) (%))		IR frequencies σ (cm^{-1})		m.p. ($^{\circ}\text{C}$)	Reaction
	C	H	(C—C)	(C—O)		
I <i>cis</i> -[Mn(CO) ₄ (PCy ₃) ₂ (C≡CPh)]	65.77 (65.69)	6.91 (6.98)	2106w	2074m; 1983s,br;1948s	162	1
II <i>fac</i> -[Mn(CO) ₃ (dppe)(C≡CPh)]	69.20 (69.60)	4.48 (4.58)	2097w	2019s,1943s,1920s	168	1
III <i>fac</i> -[Mn(CO) ₃ (dppe)(C≡CH)]	65.94 (66.20)	4.36 (4.48)	—	2024s,1942s,1918s	166	2
IV <i>mer</i> -[Mn(CO) ₃ (PCy ₃) ₂ (C≡CPh)]	69.76 (70.48)	9.13 (8.93)	2093w	2009w,1913s,1885m	175	3a
V [Mn(CO)(dppe) ₂ (C≡CPh)]	74.65 (74.69)	5.76 (5.44)	2056w	1832m	145	3b

^a In Cl₂CH₂.

derivatives, by thermally- or UV-promoted replacement of CO by other ligands L such as phosphines or isocyanides, as shown in equations 3a and 3b.





Table 1 lists analytical, infrared and melting point data for some representative compounds made by reactions 1, 2 and 3.

The acetylenic-like nature of compounds $\text{PhC}\equiv\text{CMn}(\text{CO})_n\text{L}_{5-n}$ was clearly demonstrated by treating *cis*- $\text{PhC}\equiv\text{CMn}(\text{CO})_4(\text{PCy}_3)$ with $\text{Co}_2(\text{CO})_8$ in hexane at room temperature (reaction 4). The product $[\text{Co}_2(\text{CO})_6][\text{PhC}\equiv\text{CMn}(\text{CO})_4(\text{PCy}_3)]$ (VI) could be isolated as dark green crystals (m.p. 126°C dec.) ($\bar{\nu}(\text{CO})$ - at 2082w, 2057w, 2035s, 2014m, 1998m, 1984vw, 1968w and 1958w cm^{-1} in hexane. The structure of VI was determined by an X-ray analysis; suitable crystals being obtained from a $\text{CH}_2\text{Cl}_2/\text{EtOH}$ mixture.




Fig. 1. A view of the compound $[\text{Co}_2(\text{CO})_6][\text{PhC}\equiv\text{CMn}(\text{CO})_4(\text{PCy}_3)]$. Important bond distances (Å) and angles (°): Co(1)—Co(2) 2.469(3), Co(1)—C(12) 2.002(10), Co(2)—C(12) 1.971(11), Co(1)—C(11) 2.053(9), Co(2)—C(11) 2.045(9), Mn—P 2.362(3), Mn—C(11) 2.063(7), C(11)—C(12), 1.313(11); P—Mn—C(11) 178.8(3), Mn—C(11)—C(12) 144.5(8), C(11)—C(12)—C(13) 143.3(9).

Crystal data: $C_{36}H_{38}Co_2MnO_{10}P \cdot \frac{1}{2}C_2H_5OH$, $M = 1715.01$, monoclinic, space group $P2_1/c$, a 13.545(5), b 18.691(7), c 16.944(9) Å, β 110.89(3)°, V 4008(3) Å³, $Z = 4$, D_c 1.421 g cm⁻³, $F(000)$ 1764, $\mu(Mo-K_\alpha)$ 12.06 cm⁻¹. The intensities of 6370 independent reflections were collected on a Siemens AED single-crystal diffractometer (with θ in the range $3 \leq \theta \leq 24^\circ$); niobium-filtered $Mo-K_\alpha$ radiation and the $\theta/2\theta$ scan technique were used. The structure was solved by Patterson and Fourier methods and refined by full-matrix least-squares on the basis of 3402 observed reflections (having $I \geq 2\sigma(I)$) to an R value of 0.061. The atomic coordinates for the non hydrogen atoms, the calculated coordinates for the hydrogen atoms, and the thermal parameters are available on request from authors or from the Director of the Cambridge Crystallographic Data Centre, University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW.

The structure of VI is shown in Fig. 1 together with the more significant bond distances and angles. The structure was mainly as expected, and is similar to that proposed for the known $[Co_2(CO)_6][PhC \equiv CFe(\eta-C_5H_5)(CO)L]$ ($L = CO, PMe_2Ph$) [3], although no X-ray structures have been reported for these species.

In VI the acetylide ligand is σ -bonded to the manganese atom and π -bonded to both cobalt atoms. The Mn—C(11) bond distance (2.063(7) Å) is much longer than in $BrMn(CO)_4C \equiv CPPh_3$ (1.981(14) Å) [4] where the acetylide is only σ -bonded to the Mn atom. The interaction with the Co atoms also results in a lengthening of the acetylenic C—C bond (C(11)—C(12) 1.313(11) Å), which is very close to that of the standard ethylenic double bond length (1.337 Å), and in a marked deviation from linearity (Mn—C(11)—C(12) 144.5(8)°). In $BrMn(CO)_4C \equiv CPPh_3$ the acetylenic C—C bond length is 1.216(14) Å and the Mn—C—C angle is very close to 180° (176.3°) [4].

An unexpected feature of VI is that the PCy₃ ligand is *trans* to the acetylide ligand, whereas it was *cis* to this ligand in the starting complex. This indicates that in reaction 4 there has been a rearrangement of the $Mn(CO)_4(PCy_3)$ fragment.

The reactions of σ -alkynyl and their derivatives with some nucleophiles and electrophiles and with other metal carbonyls are now under investigation.

Acknowledgment. We thank the Spanish C.A.C.Y.T. (Project No. 1184/81) for financial support.

References

- 1 N.A. Ustynyuk, V.N. Vinogradova and P.V. Petrovskii, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 3 (1982) 680.
- 2 W.C. Kaska, D.K. Mitchell, R.F. Reichelderfer and W.D. Korte, *J. Am. Chem. Soc.*, 96 (1974) 2847.
- 3 K. Yasufuku and H. Yamazaki, *Bull. Chem. Soc. Jap.*, 45 (1972) 2664.
- 4 S.Z. Goldberg, E.N. Duesler and K. Raymond, *Inorg. Chem.*, 11 (1972) 1397.