





Corrigendum

Carbon dioxide fixation by lanthanide complexes; by M.N. Bochkarev, E.A. Fedorova, Yu.F. Radkov, S.Ya. Khorshev, G.S. Kalinnikov and G.A. Razuvayev (*J. Organomet. Chem.*, 258 (1984) C29—C33)

Page C30, equations 3 and 4 should read:



Page C31, paragraph 2 should read:

Tri-t-butoxy-praseodymium and -neodymium add three moles of CO₂ per lanthanide atom in hexane. In this case, according to the IR spectrum (1580, 1370 and 1250 cm⁻¹, Fig. 1) the insertion of CO₂ into the O—Ln bond occurs with formation of carbonate derivatives (t-BuOCO₂)₃Ln.

Page C31, the legend to Fig. 1 should read:

Fig. 1. IR spectra of the lanthanide complexes: (a) $[(\text{Me}_3\text{Si})_2\text{N}]_3\text{Pr} \cdot 2\text{CO}_2$, (b) $[(\text{Me}_3\text{SiCH}_2)_3\text{Sn}]_3\text{Pr} \cdot 2\text{CO}_2 \cdot \text{DME}$ and $[(\text{Me}_3\text{Si})_2\text{N}]_2\text{PrSn}(\text{CH}_2\text{SiMe}_3)_3 \cdot 2\text{CO}_2 \cdot \text{DME}$, (c) $(\text{Me}_3\text{SiCH}_2)_3\text{SnPr}(\text{OH})_2 \cdot 2\text{CO}_2$, (d) $(\text{t-BuOCO}_2)_3\text{Ln}$.

Erratum

Role of the cation in the reaction of Co(CO)₄⁻ with RuCl₃ · xH₂O. Synthesis and Molecular Structure of the Ruthenium Cluster [N(PPh₃)₂]₂[Ru₄(μ-Cl)₄ · (μ-CO)₂(CO)₈]; by P. Braunstein, J. Rosé, Y. Dusausoy and J.P. Mangeot (*J. Organomet. Chem.*, 256 (1983) 125—134)

Page 126, lines 31-33 should read:

The formation of [PPN][RuCo₃(CO)₁₂], from the reaction of [PPN]-[Co(CO)₄], was in an unexpectedly low yield, in view of the previous results obtained with Na[Co(CO)₄] [9,10]. Unfortunately this complex is somewhat difficult to separate