

## Corrigenda

---

**Palladium assisted organic reactions. VII. The preparations of cyclopalladated primary and secondary benzylamines;** by P.W. Clark and S.F. Dyke (*J. Organomet. Chem.*, 281 (1985) 389–396).

Recently in reporting the preparation of a number of cyclopalladated primary and secondary benzylamines, and providing extensive  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectral data to establish structures of our products, we stated that such complexes had not previously been made. Our attention has been drawn [1] to a note by S. Baba and S. Kawaguchi [2] in which the parent compound, benzylamine was reported to undergo cyclopalladation with  $\text{Pd}(\text{acac})_2$ . (Some IR spectral evidence was quoted. The cyclopalladation failed under the same conditions when *N*-methyl- or *N,N*-dimethyl-benzylamines were used.)

- 1 We are grateful to Dr. R. O'Neil, CIBA-Geigy Industrial Chemicals, Manchester (Great Britain) for drawing attention to our oversight.
  - 2 S. Baba and S. Kawaguchi, *Inorg. Nucl. Chem. Lett.*, 11 (1975) 415.
- 

**Oxymetallation. XIX. Preparation of the first oxymercurials derived from cyclopropane;** by A.J. Bloodworth and C.J. Cooksey (*J. Organomet. Chem.*, 295 (1985) 131–135).

Page 135,  $\delta(\text{Hg})$  for compounds 1, 2, and 3 should read 1019, 958, and 997, respectively.