

## JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 378 (1989)

---

### SUBJECT INDEX

#### Aluminium

Trimethylaluminium; trapping of the dimethylaluminium radical in laser-powdered homogeneous pyrolysis of (G.A. Atiya, A.S. Grady, S.A. Jackson, N. Parker, D.K. Russell), (378) 307

#### Antimony

(Me<sub>3</sub>Ge)<sub>2</sub>Sb-Sb(GeMe<sub>3</sub>)<sub>2</sub> and related distibanes, crystal structures and thermal properties of (S. Roller, M. Dräger, H.J. Breunig, M. Ates, S. Güleç), (378) 327

Tertiary stibines, reduction of some organic compounds by (Y.-Z. Huang, C. Chen, F. Zhu, Y. Liao), (378) 147

#### Arsenic

Bidentate P,P or P,As ligands (L-L) in HRu<sub>3</sub>(C≡CBu<sup>l</sup>)(CO)<sub>7</sub>(L-L), chelating and bridging behaviour of (E. Sappa, G. Predieri, A. Tiripicchio, C. Vignali), (378) 109

#### Boron

Diethyl- and diisopropylamino-trifluoromethylboranes, synthesis reactions with HF, HCl, HBr and H<sub>2</sub>O of; crystal structure of (CF<sub>3</sub>)<sub>2</sub>BN(i-Pr)<sub>2</sub> revealing a very short B-N bond (D.J. Brauer, H. Bürger, F. Dörrenbach, G. Pawelke, W. Weuter), (378) 125

Triferrocenylborane and its amine adducts (T. López, A. Campero), (378) 91

#### Calcium

M(C<sub>5</sub>H<sub>4</sub>CH<sub>3</sub>)<sub>2</sub> · CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> (M = Ca, Yb), investigations of (A. Hammel, W. Schwarz, J. Weidlein), (378) 347

#### Chromium

(π<sup>6</sup>-C<sub>6</sub>H<sub>4</sub>R<sub>3</sub>)(CO)<sub>3</sub>Cr (R = Me, Et), photo-induced rearrangement with acetylenes of; molecular structure of (C<sub>6</sub>H<sub>3</sub>Me<sub>3</sub>)(CO)Cr(μ-CO){μ-C<sub>2</sub>[C(OEt)<sub>2</sub>H]<sub>2</sub>}Cr(CO)<sub>4</sub> (H.G. Alt, H.E. Engelhardt, R.D. Rogers, S.T. Abu-Orabi), (378) 33

[Cr(CO)<sub>5</sub>I], induced photoredox reaction by ligand to metal charge transfer excitation of (H. Kunkely, A. Vogler), (378) C15

Pentacarbonylhalogenaminophosphine complexes (CO)<sub>5</sub>MPRXNR'<sub>2</sub>, formation of (K. Diemert, W. Kuchen, D. Lorenzen), (378) 17

Thio- and seleno-carbene-chromium, -tungsten and -manganese complexes, formation of (R. Aumann, J. Schröder), (378) 57

Tricarbonyl-chromium, -molybdenum and -tungsten complexes of bis(2-diphenylphosphinoethyl)-amine, formation and characterization of (J. Ellermann, M. Moll, N. Will), (378) 73

Tri(methylene)methane, rearrangements to 1,3-diene ligands to chromium and iron complexes of (R. Aumann, B. Trentmann), (378) 171

#### Cobalt

Mono- and bis-metallocene calix[4]arene hydrophobic receptor molecules, synthesis, structure and electrochemical properties of (P.D. Beer, A.D. Keefe, M.G.B. Drew), (378) 437

## Copper

- [Cu<sub>3</sub>(μ<sub>3</sub>-η<sup>1</sup>-C≡CPh)(μ-dppm)<sub>3</sub>][BF<sub>4</sub>]<sub>2</sub>, synthesis and crystal structure of (M.P. Gamasa, J. Gimeno, E. Lastra, A. Aguirre, S. Garcia-Granda), (378) C11
- 4,4-Dimethyl-1-phenylpenta-1,2-diene (1-Ph-3-t-Bu-allene), chiral induction in the synthesis catalyzed by chiral phosphine complexes of palladium of (W. de Graaf, J. Boersma, G. van Koten, C.J. Elsevier), (378) 115
- Heterobinuclear complexes [WM(CO)<sub>3</sub>(PPh<sub>3</sub>)<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)] (M = Rh, Cu), <sup>13</sup>C and <sup>31</sup>P NMR studies on geometrical non-rigidity in (W.E. Lindsell, P.J. Tomb), (378) 245
- Nitrosoarene complexes of rhodium(III), iridium(III), copper(I) and mercury(II), mode of bonding in (G. Vasapollo, C.F. Nobile, A. Sacco, B.G. Gowenlock, L. Sabbatini, C. Malitesta, P. Zambonin), (378) 239

## Germanium

- Bis(triphenylgermane)tetrakis(tetrahydrofuran)ytterbium, synthesis and structure of (M.N. Bochkarev, I.M. Penyagina, L.N. Zakharov, Yu.F. Rad'kov, E.A. Fedorova, S.Ya. Khorshev, Yu.T. Struchkov), (378) 363
- (Me<sub>3</sub>Ge)<sub>2</sub>Sb-Sb(GeMe<sub>3</sub>)<sub>2</sub> and related distibanes, crystal structures and thermal properties of (S. Roller, M. Dräger, H.J. Breunig, M. Ates, S. Güleç), (378) 327

## Gold

- <sup>195</sup>Au and <sup>14</sup>C-labeled antineoplastic gold(I) phosphine complexes, synthesis of (K.T. Ganes, J.R. Heys, S.W. Landvatter, J. Statler), (378) 293

## Hafnium

- (η<sup>4</sup>-Butadiene)-zirconocene, and -hafnocene with 2-methyl-2-nitrosopropane of (G. Erker, M.G. Humphrey), (378) 163

## Iridium

- [M(COD)(μ-L)]<sub>2</sub> (M = Rh, Ir; L = substituted hydroxypyridinate), variable temperature <sup>1</sup>H NMR studies of (G.S. Rodman, K.R. Mann), (378) 255
- Nitrosoarene complexes of rhodium(III), iridium(III), copper(I) and mercury(II), mode of bonding in (G. Vasapollo, C.F. Nobile, A. Sacco, B.G. Gowenlock, L. Sabbatini, C. Malitesta, P. Zambonin), (378) 239

## Iron

- Carbon disulphide and carbon dioxide adducts (η-C<sub>5</sub>H<sub>5</sub>)(CO)<sub>2</sub>Fe-CX<sub>2</sub><sup>-</sup>, reactions with organoiron electrophiles of (M.E. Giuseppetti-Dery, B.E. Landrum, J.L. Shibley, A.R. Cutler), (378) 421
- Chiral substituted ferrocenes, stereo- and enantio-selectivity in catalytic hydrogenolysis to give cyclopentanes of (V.I. Sokolov, L.L. Troitskaya, N.S. Khrushcheva, O.A. Reutov), (378) 227
- S-(1-Ferrocenylethyl)thioglycolic acid, kinetics of the synthesis of (D. Scutaru, L. Tătaru, I. Mazișu, B. Scutaru, T. Lixandru, C. Simionescu), (378) 99
- Mono- and bis-metallocene calix[4]arene hydrophobic receptor molecules, synthesis, structure and electrochemical properties of (P.D. Beer, A.D. Keefe, M.G.B. Drew), (378) 437
- Solid state polymerization of diacetylenes; crystal and molecular structure of 1,4-bis(1'-ethylferrocenyl)buta-1,3-diyne (V.E. Shklover, Yu.E. Ovchinnikov, I.A. Zamaev, Yu.T. Struchkov, V.V. Dement'ev, T.M. Frunze, B.A. Antipova), (378) 235
- Triferrocenylborane and its amine adducts (T. López, A. Campero), (378) 91
- Tri(methylene)methane, rearrangements to 1,3-diene ligands to chromium and iron complexes of (R. Aumann, B. Trentmann), (378) 171
- Trimethylplatinum(IV) halide complexes, [PtXMe<sub>3</sub>{(C<sub>5</sub>H<sub>4</sub>ECH<sub>3</sub>)<sub>2</sub>Fe}] (E = S, Se; X = Cl, Br, I), syntheses, structures and stereodynamics of (E.W. Abel, N.J. Jong, K.G. Orell, A.G. Osborne, V. Šik), (378) 473

## Lithium

Lithium and zirconocene(IV)-phosphido complexes of *ortho*-phenylenebisphosphido ligands, 1,2-(PH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> (E. Hey), (378) 375

## Magnesium

4,4-Dimethyl-1-phenylpenta-1,2-diene (1-Ph-3-*t*-Bu-allene), chiral induction in the synthesis catalyzed by chiral phosphine complexes of palladium of (W. de Graaf, J. Boersma, G. van Koten, C.J. Elsevier), (378) 115

## Manganese

Dicarbonylcyclopentadienylmanganese fragments, reactions with propiolic aldehyde acetals of (C. Löwe, H.-U. Hund, H. Berke), (378) 211

[Mn(CO)<sub>5</sub>Br] and Re[CO]<sub>5</sub>Br], Pd-catalysed carbonyl ligand substitution reactions with isocyanides of (N.J. Coville, P. Johnston, A.E. Leins, A.J. Markwell), (378) 401

Thio- and seleno-carbene-chromium, -tungsten and -manganese complexes, formation of (R. Aumann, J. Schröder), (378) 57

## Mercury

Dibenzylmercury radical cations, ESR study of (C.J. Rhodes, H.A. Agirbas), (378) 303

Nitrosoarene complexes of rhodium(III), iridium(III), copper(I) and mercury(II), mode of bonding in (G. Vasapollo, C.F. Nobile, A. Sacco, B.G. Gowenlock, L. Sabbatini, C. Malatesta, P. Zambonin), (378) 239

## Metalloenes

( $\eta^4$ -Butadiene)-zirconocene, and -hafnocene with 2-methyl-2-nitrosopropane of (G. Erker, M.G. Humphrey), (378) 163

Chiral substituted ferrocenes, stereo- and enantio-selectivity in catalytic hydrogenolysis to give cyclopentanes of (V.I. Sokolov, L.L. Troitskaya, N.S. Khrushcheva, O.A. Reutov), (378) 227

Lithium and zirconocene(IV)-phosphido complexes of *ortho*-phenylenebisphosphido ligands, 1,2-(PH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> (E. Hey), (378) 375

Mono- and bis-metallocene calix[4]arene hydrophobic receptor molecules, synthesis, structure and electrochemical properties of (P.D. Beer, A.D. Keefe, M.G.B. Drew), (378) 437

Solid state polymerization of diacetylenes; crystal and molecular structure of 1,4-bis(1'-ethylferrocenyl)buta-1,3-diyne (V.E. Shklover, Yu.E. Ovchinnikov, I.A. Zamaev, Yu.T. Struchkov, V.V. Dement'ev, T.M. Frunze, B.A. Antipova), (378) 235

Titanocene chiral derivatives accessible from substituted dihydropentalene and azulene precursors (P. Burger, H.-U. Hund, K. Evertz, H.-H. Brintzinger), (378) 153

Triferrocenylborane and its amine adducts (T. López, A. Campero), (378) 91

Trimethylplatinum(IV) halide complexes, [PtXMe<sub>3</sub>{(C<sub>5</sub>H<sub>4</sub>ECH<sub>3</sub>)<sub>2</sub>Fe}] (E = S, Se; X = Cl, Br, I), syntheses, structures and stereodynamics of (E.W. Abel, N.J. Jong, K.G. Orell, A.G. Osborne, V. Šik), (378) 473

## Molybdenum

Molybdenum  $d^6$ -ML<sub>5</sub> fragments of hydrido derivatives, investigation of C-H activation by (L. Dahlenburg, B. Pietsch), (378) 199

Molybdenum(II) and tungsten(II) polyisonitrile substituted half-sandwich complexes (A.C. Filippou, W. Grünleitner), (378) 387

Pentacarbonylhalogenaminophosphine complexes (CO)<sub>5</sub>MPRXNR'<sub>2</sub>, formation of (K. Diemert, W. Kuchen, D. Lorenzen), (378) 17

Transformation of phenylacetylene into *ortho*-metallated phenylvinylidene cluster having a Re<sub>2</sub>Mo<sub>2</sub> core; molecular structure of (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Mo<sub>2</sub>Re<sub>2</sub>( $\mu_4$ -C=C(H)C<sub>6</sub>H<sub>4</sub>)( $\mu$ -H)( $\mu$ -CO)(CO)<sub>7</sub> (A.D. Shaposhnikova, G.L. Kamalov, R.A. Stadnichenko, A.A. Pasynskii, I.L. Eremenko, Yu.T. Struchkov, A.I. Yanovsky, P.V., Petrovskii), (378) 67

Tricarbonyl-chromium, -molybdenum and -tungsten complexes of bis(2-diphenylphosphinoethyl)-amine, formation and characterization of (J. Ellermann, M. Moll, N. Will), (378) 73

## Nickel

Heteronuclear di- and tri-metal  $\mu$ -carbyne compounds from the thiocarbyne  $[\text{HB}(\text{pz})_3](\text{CO})_2\text{W}\equiv\text{CSMe}$ , synthesis of (R.A. Doyle, R.J. Angelici, F.G.A. Stone), (378) 81

Nickel(O) induced CC coupling of fluoro-substituted alkenes with phenyl isocyanate (H. Hoberg, D. Guhl), (378) 279

## Osmium

Tetranuclear mixed-metal hydride complexes,  $\text{LWM}_3(\text{CO})_{11}(\mu\text{-H})_3$ , (L =  $\text{C}_5\text{H}_5$ ,  $\text{C}_5\text{Me}_5$  and M = Ru, Os), synthesis and characterization of (Y. Chi, C.-Y. Cheng, S.-L. Wang), (378) 45

## Palladium

4,4-Dimethyl-1-phenylpenta-1,2-diene (1-Ph-3-t-Bu-allene), chiral induction in the synthesis catalyzed by chiral phosphine complexes of palladium of (W. de Graaf, J. Boersma, G. van Koten, C.J. Elsevier), (378) 115

Metal complexes of C-nitroso compounds, correlation of NO stretching frequencies with coordination modes in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (378) 493

$[\text{Mn}(\text{CO})_5\text{Br}]$  and  $[\text{Re}(\text{CO})_5\text{Br}]$ , Pd-catalysed carbonyl ligand substitution reactions with isocyanides of (N.J. Coville, P. Johnston, A.E. Leins, A.J. Markwell), (378) 401

Platinum and palladium  $\alpha$ -ketoacyl complexes, reactivity toward olefins and acetylenes of (W.M. Vetter, A. Sen), (378) 485

## Phosphorus

Acrylonitrile, insertion reactions with hydridocarbonylbis(triphenylphosphine)ruthenium(II) carboxylates of (S.S. Deshpande, S. Gopinathan, C. Gopinathan), (378) 103

$^{195}\text{Au}$  and  $^{14}\text{C}$ -labeled antineoplastic gold(I) phosphine complexes, synthesis of (K.T. Garnes, J.R. Heys, S.W. Landvatter, J. Statler), (378) 293

Bidentate P,P or P,As ligands (L-L) in  $\text{HRu}_3(\text{C}\equiv\text{CBu}^1)(\text{CO})_7(\text{L-L})$ , chelating and bridging behaviour of (E. Sappa, G. Predieri, A. Tiripicchio, C. Vignali), (378) 109

$[\text{Cu}_3(\mu_3\text{-}\eta^1\text{-C}\equiv\text{CPh})(\mu\text{-dppm})_3][\text{BF}_4]_2$ , synthesis and crystal structure of (M.P. Gamasa, J. Gimeno, E. Lastra, A. Aguirre, S. Garcia-Granda), (378) C11

Dicarbonyl( $\eta^5$ -cyclopentadienyl)[ $\eta^2$ -aryl(methylthio)carbene]tungsten tetrafluoroborate, reaction with tri-methylphosphine with formation of an  $\eta^3$ -thio-ketene intermediate of (N. Ullrich, H. Keller, C. Stegmair, F.R. Kreissl), (378) C19

4,4-Dimethyl-1-phenylpenta-1,2-diene (1-Ph-3-t-Bu-allene), chiral induction in the synthesis catalyzed by chiral phosphine complexes of palladium of (W. de Graaf, J. Boersma, G. van Koten, C.J. Elsevier), (378) 115

Lithium and zirconocene(IV)-phosphido complexes of *ortho*-phenylenebisphosphido ligands, 1,2-(PH) $_2\text{C}_6\text{H}_4$  (E. Hey), (378) 375

Molybdenum  $d^6$ -ML $_5$  fragments of hydrido derivatives, investigation of C-H activation by (L. Dahlenburg, B. Pietsch), (378) 199

Nickel(O) induced CC coupling of fluoro-substituted alkenes with phenyl isocyanate (H. Hoberg, D. Guhl), (378) 279

Pentacarbonylhalogenaminophosphine complexes  $(\text{CO})_5\text{MPRXNR}'_2$ , formation of (K. Diemert, W. Kuchen, D. Lorenzen), (378) 17

Tricarbonyl-chromium, -molybdenum and -tungsten complexes of bis(2-diphenylphosphinoethyl)amine, formation and characterization of (J. Ellermann, M. Moll, N. Will), (378) 73

## Platinum

Heteronuclear di- and tri-metal  $\mu$ -carbyne compounds from the thiocarbyne  $[\text{HB}(\text{pz})_3](\text{CO})_2\text{W}\equiv\text{CSMe}$ , synthesis of (R.A. Doyle, R.J. Angelici, F.G.A. Stone), (378) 81

Metal complexes of C-nitroso compounds, correlation of NO stretching frequencies with coordination modes in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (378) 493

Platinum and palladium  $\alpha$ -ketoacyl complexes, reactivity toward olefins and acetylenes of (W.M. Vetter, A. Sen), (378) 485

Trimethylplatinum(IV) halide complexes,  $[\text{PtXMMe}_3\{(\text{C}_5\text{H}_4\text{ECH}_3)_2\text{Fe}\}]$  (E = S, Se; X = Cl, Br, I), syntheses, structures and stereodynamics of (E.W. Abel, N.J. Jong, K.G. Orell, A.G. Osborne, V. Šik), (378) 473

## Rhenium

[Mn(CO)<sub>5</sub>Br] and Re[CO]<sub>5</sub>Br, Pd-catalysed carbonyl ligand substitution reactions with isocyanides of (N.J. Coville, P. Johnston, A.E. Leins, A.J. Markwell), (378) 401

## Rhodium

Dihydrobis(triphenylphosphine)rhodium(III) complexes of aromatic N-donor carboxylates, <sup>15</sup>N NMR spectra of (L. Carlton, M.-P. Belciug), (378) 469

Heterobinuclear complexes [WM(CO)<sub>3</sub>(PPh<sub>3</sub>)<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)] (M = Rh, Cu), <sup>13</sup>C and <sup>31</sup>P NMR studies on geometrical non-rigidity in (W.E. Lindsell, P.J. Tomb), (378) 245

Heteronuclear di- and tri-metal μ-carbyne compounds from the thiocarbyne [HB(pz)<sub>3</sub>](CO)<sub>2</sub>W≡CSMe, synthesis of (R.A. Doyle, R.J. Angelici, F.G.A. Stone), (378) 81

Hydrocarboxylation of isoprene catalysed by iodocarbonylrhodium derivatives (L. Garlaschelli, M. Marchionna, M.C. Iapalucci, G. Longoni), (378) 457

[M(COD)(μ-L)]<sub>2</sub> (M = Rh, Ir; L = substituted hydroxypyridinate), variable temperature <sup>1</sup>H NMR studies of (G.S. Rodman, K.R. Mann), (378) 255

Nitrosoarene complexes of rhodium(III), iridium(III), copper(I) and mercury(II), mode of bonding in (G. Vasapollo, C.F. Nobile, A. Sacco, B.G. Gowenlock, L. Sabbatini, C. Malitesta, P. Zambonin), (378) 239

## Ruthenium

Acrylonitrile, insertion reactions with hydridocarbonylbis(triphenylphosphine)ruthenium(II) carboxylates of (S.S. Deshpande, S. Gopinathan, C. Gopinathan), (378) 103

Bidentate P,P or P,As ligands (L-L) in HRu<sub>3</sub>(C≡CBu<sup>t</sup>)(CO)<sub>7</sub>(L-L), chelating and bridging behaviour of (E. Sappa, G. Predieri, A. Tiripicchio, C. Vignali), (378) 109

[η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>RuOMe]<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>Ru)<sub>3</sub>(μ<sub>3</sub>-OMe)<sub>2</sub>](PF<sub>6</sub>), comparison of molecular structures of (U. Kölle, J. Kossakowski, R. Boese), (378) 449

Tetranuclear mixed-metal hydride complexes, LW<sub>3</sub>(CO)<sub>11</sub>(μ-H)<sub>3</sub>, (L = C<sub>5</sub>H<sub>5</sub>, C<sub>5</sub>Me<sub>5</sub> and M = Ru, Os), synthesis and characterization of (Y. Chi, C.-Y. Cheng, S.-L. Wang), (378) 45

## Selenium

Thio- and seleno-carbene-chromium, -tungsten and -manganese complexes, formation of (R. Aumann, J. Schröder), (378) 57

Trimethylplatinum(IV) halide complexes, [PtXMMe<sub>3</sub>{(C<sub>5</sub>H<sub>4</sub>ECH<sub>3</sub>)<sub>2</sub>Fe}] (E = S, Se; X = Cl, Br, I), syntheses, structures and stereodynamics of (E.W. Abel, N.J. Jong, K.G. Orell, A.G. Osborne, V. Šik), (378) 473

## Silicon

Chloroplatinic acid catalysed cyclization of silanes bearing pendant acetylenic groups (M.G. Steinmetz, B.S. Udayakumar), (378) 1

Isosteric isobutyl(isopropoxy)silanethiols, preparation and spectroscopic properties of (J. Pickies, W. Wojnowski), (378) 317

(Me<sub>3</sub>Ge)<sub>2</sub>Sb-Sb(GeMe<sub>3</sub>)<sub>2</sub> and related distibanes, crystal structures and thermal properties of (S. Roller, M. Dräger, H.J. Breunig, M. Ates, S. Güleç), (378) 327

Phenylsilanes formation of silyltriflates from (W. Uhlig, A. Tzschach), (378) C1

## Sulphur

Carbon disulphide and carbon dioxide adducts (η-C<sub>5</sub>H<sub>5</sub>)(CO)<sub>2</sub>Fe-CX<sub>2</sub><sup>-</sup>, reactions with organoiron electrophiles of (M.E. Giuseppetti-Dery, B.E. Landrum, J.L. Shibley, A.R. Cutler), (378) 421

Dicarbonyl(η<sup>5</sup>-cyclopentadienyl)[η<sup>2</sup>-aryl(methylthio)carbene]tungsten tetrafluoroborate, reaction with tri-methylphosphine with formation of an η<sup>3</sup>-thio-ketene intermediate of (N. Ullrich, H. Keller, C. Stegmair, F.R. Kreissl), (378) C19

Dimethyltin dichloride-dibenzylsulphoxide, crystal structure of (S.W. Ng, A.L. Rheingold), (378) 339

Heteronuclear di- and tri-metal μ-carbyne compounds from the thiocarbyne [HB(pz)<sub>3</sub>](CO)<sub>2</sub>W≡CSMe, synthesis of (R.A. Doyle, R.J. Angelici, F.G.A. Stone), (378) 81

Isosteric isobutyl(isopropoxy)silanethiols, preparation and spectroscopic properties of (J. Pickies, W. Wojnowski), (378) 317

Thio- and seleno-carbene-chromium, -tungsten and -manganese complexes, formation of (R. Aumann, J. Schröder), (378) 57

Trimethylplatinum(IV) halide complexes, [PtXMe<sub>3</sub>{(C<sub>5</sub>H<sub>4</sub>ECH<sub>3</sub>)<sub>2</sub>Fe}] (E = S, Se; X = Cl, Br, I), syntheses, structures and stereodynamics of (E.W. Abel, N.J. Jong, K.G. Orell, A.G. Osborne, V. Šik), (378) 473

### Thallium

Titanocene chiral derivatives accessible from substituted dihydropentalene and azulene precursors (P. Burger, H.-U. Hund, K. Evertz, H.-H. Brintzinger), (378) 153

### Tin

Dimethyltin dichloride-dibenzylsulphoxide, crystal structure of (S.W. Ng, A.L. Rheingold), (378) 339

(Me<sub>3</sub>Ge)<sub>2</sub>Sb-Sb(GeMe<sub>3</sub>)<sub>2</sub> and related distibanes, crystal structures and thermal properties of (S. Roller, M. Dräger, H.J. Breunig, M. Ates, S. Güleç), (378) 327

Metal complexes of C-nitroso compounds, correlation of NO stretching frequencies with coordination modes in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (378) 493

Tris(pyrazolyl)borates, preparations of (G.G. Lobbia, F. Bonati, P. Cecchi, A. Cingolani, A. Lorenzotti), (378) 139

### Titanium

( $\eta^4$ -Butadiene)-zirconocene, and -hafnocene with 2-methyl-2-nitrosopropane of (G. Erker, M.G. Humphrey), (378) 163

### Tungsten

Dicarbonyl( $\eta^5$ -cyclopentadienyl)[ $\eta^2$ -aryl(methylthio)carbene]tungsten tetrafluoroborate, reaction with tri-methylphosphine with formation of an  $\eta^3$ -thio-ketene intermediate of (N. Ullrich, H. Keller, C. Stegmair, F.R. Kreissl), (378) C19

Heteronuclear di- and tri-metal  $\mu$ -carbyne compounds from the thiocarbyne [HB(pz)<sub>3</sub>](CO)<sub>2</sub>W $\equiv$ CSMe, synthesis of (R.A. Doyle, R.J. Angelici, F.G.A. Stone), (378) 81

Molybdenum(II) and tungsten(II) polyisonitrile substituted half-sandwich complexes (A.C. Filippou, W. Grünleitner), (378) 387

Pentacarbonylhalogenaminophosphine complexes (CO)<sub>5</sub>MPRXNR'<sub>2</sub>, formation of (K. Diemert, W. Kuchen, D. Lorenzen), (378) 17

Tetranuclear mixed-metal hydride complexes, LWM<sub>3</sub>(CO)<sub>11</sub>( $\mu$ -H)<sub>3</sub>, (L = C<sub>5</sub>H<sub>5</sub>, C<sub>5</sub>Me<sub>5</sub> and M = Ru, Os), synthesis and characterization of (Y. Chi, C.-Y. Cheng, S.-L. Wang), (378) 45

Thio- and seleno-carbene-chromium, -tungsten and -manganese complexes, formation of (R. Aumann, J. Schröder), (378) 57

Thio- and seleno-ketene imines from thio- and seleno-carbene complexes of chromium, tungsten, manganese and isocyanides (R. Aumann, J. Schröder, C. Krüger, R. Goddard), (378) 185

Tricarbonyl-chromium, -molybdenum and -tungsten complexes of bis(2-diphenylphosphinoethyl)-amine, formation and characterization of (J. Ellermann, M. Moll, N. Will), (378) 73

### Ytterbium

Bis(triphenylgermane)tetrakis(tetrahydrofuran)ytterbium, synthesis and structure of (M.N. Bochkarev, I.M. Penyagine, L.N. Zakharov, Yu.F. Rad'kov, E.A. Fedorova, S.Ya. Khorshev, Yu.T. Struchkov), (378) 363

M(C<sub>5</sub>H<sub>4</sub>CH<sub>3</sub>)<sub>2</sub> · CH<sub>3</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub> (M = Ca, Yb), investigations of (A. Hammel, W. Schwarz, J. Weidlein), (378) 347

### Zinc

4,4-Dimethyl-1-phenylpenta-1,2-diene (1-Ph-3-t-Bu-allene), chiral induction in the synthesis catalyzed by chiral phosphine complexes of palladium of (W. de Graaf, J. Boersma, G. van Koten, C.J. Elsevier), (378) 115

### Zirconium

Lithium and zirconocene(IV)-phosphido complexes of *ortho*-phenylenebisphosphido ligands, 1,2-(PH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub> (E. Hey), (378) 375

**Book reviews**

Advances in metal carbene chemistry, NATO ASI Series, Series C: Mathematical and physical sciences, Volume 269; edited by U. Schubert (P.A. Chaloner), (378) C9

Chemical reactions on polymers, ACS Symposium Series Volume 364, edited by J.L. Benham and J.F. Kinstle (P.A. Chaloner), (378) C7