
JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 367 (1989)

SUBJECT INDEX**Aluminium**

(C₅H₅)₂ZrCl(AlCl₄), crystal and molecular structure and unusual hydrocarbon solubility properties of; relevance to Ziegler-Natta polymerization chemistry (M.V. Gaudet, M.J. Zaworotko, T.S. Cameron, A. Linden), (367) 267

Boron

Boron hydride insertion into Zr-C bond of dimeric (η²-formaldehyde)zirconocene (G. Erker, U. Hoffmann, R. Zwettler, G. Krüger), (367) C15

(1,1-Dichloroalkyl)lithium, reactions with boronic acid of; facile synthesis of tertiary carbinols (B. Junchai, Z. Weike, D. Hongxun), (367) C9

2,4,6,8,9,10-Hexachloro-2,4,6,8,9,10-hexaboroadamantane (W. Haubold, W. Keller, G. Sawitzki), (367) 19
Methyl 4-tri-n-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hyarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Chromium

η⁶-Arenetricarbonylchromium(0) π-complexes, failing nucleophilic substitution of hydrogen in (S. Ostrowski, M. Mąkosza), (367) 95

Benchrotrenic alcohols, asymmetric synthesis with baker's yeast of (J. Gillois, G. Jaouen, D. Buisson, R. Azerad), (367) 85

Group 6 metal tetracarbonyl complexes, *cis*-[M(CO)₄{(C₅H₄ECH₃)₂Fe}] (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1'-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Tricarbonyl(boat *p*-xylene)chromium complexes, ab initio calculations of the charge densities in (I. Ono, S. Mita, S. Kondo, N. Mori), (367) 81

Cobalt

(η⁵-C₅Me₅)₂CoRh(μ-CO)₂, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197

Dicarbonylcyclopentadienylmethylthiolatoiron(II), reactions with monocynoacetylene of; crystal structures of [FeC(O)C(CN)=CH(SCH₃)(η⁵-C₅H₅)(CO)] and of [μ-η²-Fe(C≡CCN)(η⁵-C₅H₅)(CO)₂][Co(CO)₃]₂ (R. Kergoat, M.M. Kubicki, L.C. Gomes de Lima, H. Scordia, J.E. Guerschais, P. L'Haridon), (367) 143

Tetrakis- and hexakis-(trimethylsilyl)cobaltocenium ions, ring rotation hindrance in (J. Okuda), (367) C1

Trimethylphosphinecobalt(I) complexes: reaction with cyclopentatriene and cyclooctatetraene; crystal structure of [Co(η⁴-C₇H₈)(PMe₃)₃]BPh₄ (L.C. Ananias de Carvalho, M. Dartiguenave, Y. Dartiguenave), (367) 187

Copper

Cu^I-macrocyclic tridentate amine systems, reactions with CO, C₂H₄ and cyclic olefins of (P. Chaudhuri, K. Oder), (367) 249

Germanium

Tetrakis(2,6-diisopropylphenyl)digermene, synthesis of and its reductive cleavage to the corresponding digermenyllithium(DME) complex (J. Park, S.A. Batcheller, S. Masamune), (367) 39

Iron

1,6-Bis(triphenylmethyl)biferrocenium triiodide, electron transfer in (T.-Y. Dong, T.-L. Hsu), (367) 313
Chalcogens, contrasting roles in synthesis of mixed metal complexes $(\text{CO})_6\text{Fe}_2(\mu_3\text{-E})_2\text{M}(\text{PPh}_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

$[(\text{C}_5\text{H}_4\text{Me})\text{FeCO}]_3\text{NO}$ and $[(\text{C}_5\text{Me}_5)\text{FeCO}]_3\text{NO}$, synthesis and structure of (J. Müller, I. Sonn, T. Akhnoukh), (367) 133

Dicarbonylcyclopentadienylmethylthiolatoiron(II), reactions with monocyanoacetylene of; crystal structures of $[\text{FeC}(\text{O})\text{C}(\text{CN})=\text{CH}(\text{SCH}_3)(\eta^5\text{-C}_5\text{H}_5)(\text{CO})]$ and of $[\mu\text{-}\eta^2\text{-Fe}(\text{C}=\text{CCN})(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2][\text{Co}(\text{CO})_3]_2$ (R. Kergoat, M.M. Kubicki, L.C. Gomes de Lima, H. Scordia, J.E. Guerschais, P. L'Haridon), (367) 143

β -Keto phosphines derived from ferrocene: syntheses and structures of $[\text{Ph}_2\text{PCH}_2\text{C}(\text{O})(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{L}^1)]$ and *trans*- $[\text{PdCl}_2\text{L}_2]$ (P. Braunstein, T.M. Gomes Carneiro, D. Matt, F. Balegroune, D. Grandjean), (367) 117

Ruthenium dinuclear cluster with N- and O-containing bridges, reactions with CO_2 of (F. Neumann, G. Süss-Fink), (367) 175

Lithium

(1,1-Dichloroalkyl)lithium, reactions with boronic acid of; facile synthesis of tertiary carbinols (B. Junchai, Z. Weike, D. Hongxun), (367) C9

$\text{Et}_3\text{SiCF}=\text{CFM}$ (M = Li, ZnBr), preparation and reactivity of (P. Martinet, R. Sauvêtre, J.-F. Normant), (367) 1

Tetrakis(2,6-diisopropylphenyl)digermene, synthesis of and its reductive cleavage to the corresponding digermenyllithium(DME) complex (J. Park, S.A. Batcheller, S. Masamune), (367) 39

Magnesium

Grignard reactions of 1-phenylethyl chloride with aldehydes and ketones, reactivity of benzyl Grignard reagent (C. Bernardon), (367) 11

Mercury

Methyl 4-tri-n-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Metalloenes

1,6-Bis(triphenylmethyl)biferrocenium triiodide, electron transfer in (T.-Y. Dong, T.-L. Hsu), (367) 313
Boron hydride insertion into Zr-C bond of dimeric $(\eta^2\text{-formaldehyde})\text{zirconocene}$ (G. Erker, U. Hoffmann, R. Zwettler, G. Krüger), (367) C15

Dicyclopentadienyltitanium dihalides, electronic structure as indicated by UV photoelectron spectroscopy of (T. Vondrák, K. Mach, V. Varga), (367) 69

Dicyclopentadienyltricarboxyldirrhodium, search for ^{13}C equilibrium isotope effects in (S. Berger, M. Ochs), (367) 343

Group 6 metal tetracarbonyl complexes, *cis*- $[\text{M}(\text{CO})_4\{(\text{C}_5\text{H}_4\text{ECH}_3)_2\text{Fe}\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1'-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

β -Keto phosphines derived from ferrocene: syntheses and structures of $[\text{Ph}_2\text{PCH}_2\text{C}(\text{O})(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{L}^1)]$ and *trans*- $[\text{PdCl}_2\text{L}_2]$ (P. Braunstein, T.M. Gomes Carneiro, D. Matt, F. Balegroune, D. Grandjean), (367) 117

Palladium(0)-mediated reactions of $\text{BrZnCH}(\text{CH}_3)\text{COO}^t\text{Bu}$ with vinyl and aryl triflates, influence of catalyst on (F. Orsini, F. Pelizzoni, L.M. Vallarino), (367) 375

Tetrakis- and hexakis-(trimethylsilyl)cobaltocenium ions, ring rotation hindrance in (J. Okuda), (367) C1

Molybdenum

- Group 6 metal tetracarbonyl complexes, *cis*-[M(CO)₄((C₅H₄ECH₃)₂Fe)] (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1'-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275
- Ligand-centred coupling of alkynyl radical cations [Mo(C≡CR)(Ph₂PCH₂CH₂PPh₂)(η-C₇H₇)]⁺ (R = Ph or Buⁿ): crystal structure of [Mo₂(Ph₂PCH₂CH₂PPh₂)₂(η-C₇H₇)₂(μ-C₄R₂)] [PF₆]₂ (R = Ph) (R.L. Beddoes, C. Bitcon, A. Ricalton, M.W. Whiteley), (367) C21
- Molybdenum(II) and tungsten(II) cationic seven-coordinate complexes, synthesis of (P.K. Baker, M. Bamber, G.W. Rogers), (367) 101

Nickel

- 2,2-Bipyridine-nickel complexes as catalysts for electrosynthesis of symmetrical ketones from organic halides (L. Garnier, Y. Rollin, J. Périchon), (367) 347
- Chalcogens, contrasting roles in synthesis of mixed metal complexes (CO)₆Fe₂(μ₃-E)₂M(PPh₃)₂ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243
- Nickel complexes containing phosphorus(III) ligands and their related Ziegler catalysts, electrochemical studies of (A.A. Pozdeeva, U.M. Dzhemilev, N.R. Popod'ko, R.I. Khusnutdinov, S.I. Zhdanov, G.A. Tolstikov), (367) 205
- Oligomerisation of ethene with catalysts exhibiting enzyme-like activities (S.J. Brown, A.F. Masters), (367) 371
- Organometallic complexes, extent and direction of optical induction (T. Bartik, I. Gerdes, P. Heimbach, H.-G. Schulte), (367) 359
- (Tmeda)Ni(H₂C=CHCOOCH₃)₂, synthesis and structure of (W. Kaschube, K.R. Pörschke, K. Seevogel, C. Krüger), (367) 233

Osmium

- Aromatic olefinic ruthenium(0) and osmium(0) complexes, C₆Me₆Ru(CO)(C₂H₃R) and C₆H₆Os(L)(C₂H₄), direct reductive synthesis of (H. Werner, K. Roder), (367) 339
- Hydrido(methyl)osmium compounds, C₆H₆OsH(CH₃)(PR₃), synthesis, properties and reactivity of (K. Roder, H. Werner), (367) 321

Palladium

- Chalcogens, contrasting roles in synthesis of mixed metal complexes (CO)₆Fe₂(μ₃-E)₂M(PPh₃)₂ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243
- Methyl 4-tri-*n*-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259
- Montmorillonitesilylaminepalladium(II) chloride, selective hydrodebromination of aryl bromides by (Y.V. Subba Rao, K. Mukkanti, B.M. Choudary), (367) C29
- Oxidative addition–reductive elimination reactions; crystal structure of Me₂(*p*-BrC₆H₄CH₂)Pd(phen)Br (A.J. Canty, A.A. Watson, B.W. Skelton, A.H. White), (367) C25
- Palladium(0)-mediated reactions of BrZnCH(CH₃)COO^tBu with vinyl and aryl triflates, influence of catalyst on (F. Orsini, F. Pelizzoni, L.M. Vallarino), (367) 375

Phosphorus

- Aromatic olefinic ruthenium(0) and osmium(0) complexes, C₆Me₆Ru(CO)(C₂H₃R) and C₆H₆Os(L)(C₂H₄), direct reductive synthesis of (H. Werner, K. Roder), (367) 339
- t*-Butyl-substituted tin–phosphor compounds, preparation and structure of (D. Hänssgen, H. Aldenhoven, M. Nieger), (367) 47
- Hexahydrido(η⁵-pentamethylcyclopentadienyl)rhenium(VII), (η⁵-C₅Me₅)ReH₆, reactivity of (W.A. Herrmann, H.G. Theiler, E. Herdtweck, P. Kiprof), (367) 291
- Hydrido(methyl)osmium compounds, C₆H₆OsH(CH₃)(PR₃), synthesis, properties and reactivity of (K. Roder, H. Werner), (367) 321
- β-Keto phosphines derived from ferrocene; syntheses and structures of [Ph₂PCH₂C(O)(η⁵-C₅H₄)Fe(η⁵-C₅H₅)](L¹) and *trans*-[PdCl₂L₂]¹ (P. Braunstein, T.M. Gomes Carneiro, D. Matt, F. Balegrone, D. Grandjean), (367) 117

- Ligand-centred coupling of alkynyl radical cations $[\text{Mo}(\text{C}\equiv\text{CR})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)(\eta\text{-C}_7\text{H}_7)]^+$ (R = Ph or Buⁿ): crystal structure of $[\text{Mo}_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2(\eta\text{-C}_7\text{H}_7)_2(\mu\text{-C}_4\text{R}_2)][\text{PF}_6]_2$ (R = Ph) (R.L. Beddoes, C. Bitcon, A. Ricalton, M.W. Whiteley), (367) C21
- Molybdenum(II) and tungsten(II) cationic seven-coordinate complexes, synthesis of (P.K. Baker, M. Bamber, G.W. Rogers), (367) 101
- Nickel complexes containing phosphorus(III) ligands and their related Ziegler catalysts, electrochemical studies of (A.A. Pozdeeva, U.M. Dzhemilev, N.R. Popod'ko, R.I. Khusnutdinov, S.I. Zhdanov, G.A. Tolstikov), (367) 205
- Organometallic complexes, extent and direction of optical induction (T. Bartik, I. Gerdes, P. Heimbach, H.-G. Schulte), (367) 359
- Palladium complexes as catalyst in reduction of aromatic acyl fluorides (R. Braden, T. Himmler), (367) C12
- Ruthenium dinuclear cluster with N- and O-containing bridges, reactions with CO₂ of (F. Neumann, G. Süss-Fink), (367) 175
- Thiolate-amine complexes $[\text{Ru}(\text{CO})_2(\text{bmae})]$ and $[\text{Ru}(\text{CO})_2(\text{}^{\text{bu}}\text{bmae})]$ ($\text{bmae}^{2-} = 1,2\text{-bis}(2\text{-mercaptoanilino})\text{ethane}(-2)$; $\text{}^{\text{bu}}\text{bmae} = 1,2\text{-bis}(2\text{-mercapto-3,5-di-}t\text{-butylalino})\text{ethane}(-2)$), UV photolysis of (D. Sellmann, O. Käßler, F. Knoch), (367) 161
- Trimethylphosphinecobalt(I) complexes: reaction with cyclopentatriene and cyclooctatetraene; crystal structure of $[\text{Co}(\eta^4\text{-C}_7\text{H}_8)(\text{PMe}_3)_3]\text{BPh}_4$ (L.C. Ananias de Carvalho, M. Dartiguenave, Y. Dartiguenave), (367) 187

Platinum

- Chalcogens, contrasting roles in synthesis of mixed metal complexes $(\text{CO})_6\text{Fe}_2(\mu_3\text{-E})_2\text{M}(\text{PPh}_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

Rhenium

- Four isomeric (α -diimine) $\text{Re}(\text{CO})_3\text{Hal}$ complexes, synthesis, electrochemistry and emission spectroscopy in solution of (W. Kaim, H.E.A. Kramer, C. Vogler, J. Rieker), (367) 107
- Hexahydrido(η^5 -pentamethylcyclopentadienyl)rhenium(VII), $(\eta^5\text{-C}_5\text{Me}_5)\text{ReH}_6$, reactivity of (W.A. Herrmann, H.G. Theiler, E. Herdtweck, P. Kiprof), (367) 291

Rhodium

- $(\eta^5\text{-C}_5\text{Me}_5)_2\text{CoRh}(\mu\text{-CO})_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197
- Dicyclopentadienyltricarbonyldirrhodium, search for ¹³C equilibrium isotope effects in (S. Berger, M. Ochs), (367) 343

Ruthenium

- Aromatic olefinic ruthenium(0) and osmium(0) complexes, $\text{C}_6\text{Me}_6\text{Ru}(\text{CO})(\text{C}_2\text{H}_3\text{R})$ and $\text{C}_6\text{H}_6\text{Os}(\text{L})(\text{C}_2\text{H}_4)$, direct reductive synthesis of (H. Werner, K. Roder), (367) 339
- Ruthenium dinuclear cluster with N- and O-containing bridges, reactions with CO₂ of (F. Neumann, G. Süss-Fink), (367) 175
- Thiolate-amine complexes $[\text{Ru}(\text{CO})_2(\text{bmae})]$ and $[\text{Ru}(\text{CO})_2(\text{}^{\text{bu}}\text{bmae})]$ ($\text{bmae}^{2-} = 1,2\text{-bis}(2\text{-mercaptoanilino})\text{ethane}(-2)$; $\text{}^{\text{bu}}\text{bmae} = 1,2\text{-bis}(2\text{-mercapto-3,5-di-}t\text{-butylalino})\text{ethane}(-2)$), UV photolysis of (D. Sellmann, O. Käßler, F. Knoch), (367) 161

Selenium

- Chalcogens, contrasting roles in synthesis of mixed metal complexes $(\text{CO})_6\text{Fe}_2(\mu_3\text{-E})_2\text{M}(\text{PPh}_3)_2$ (E = Te, Se, S; M = Ni, Pd, Pt) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243
- $(\eta^5\text{-C}_5\text{Me}_5)_2\text{CoRh}(\mu\text{-CO})_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197
- Group 6 metal tetracarbonyl complexes, *cis*- $[\text{M}(\text{CO})_4\{(\text{C}_5\text{H}_4\text{ECH}_3)_2\text{Fe}\}]$ (M = Cr, Mo, W; E = S or Se); crystal structure of 1,1'-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Silicon

$\text{Et}_3\text{SiCF}=\text{CFM}$ ($\text{M} = \text{Li}, \text{ZnBr}$), preparation and reactivity of (P. Martinet, R. Sauvêtre, J.-F. Normant), (367) 1

Palladium complexes as catalyst in reduction of aromatic acyl fluorides (R. Braden, T. Himmler), (367) C12

Silanol silylation by a chlorosilane in the presence of neutral nucleophiles, mechanism of (S. Rubinsztajn, M. Cypryk, J. Chojnowski), (367) 27

Tetrakis- and hexakis-(trimethylsilyl)cobaltocenium ions, ring rotation hindrance in (J. Okuda), (367) C1

Sulphur

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(\text{CO})_6\text{Fe}_2(\mu_3\text{-E})_2\text{M}(\text{PPh}_3)_2$ ($\text{E} = \text{Te}, \text{Se}, \text{S}; \text{M} = \text{Ni}, \text{Pd}, \text{Pt}$) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

$(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO})_2$, reaction with sulphoxides of (T.L. Chen, A. Shaver, T.H. Chan), (367) C5

$(\eta^5\text{-C}_5\text{Me}_5)_2\text{CoRh}(\mu\text{-CO})_2$, reaction with sulphur and selenium of (H. Brunner, N. Janietz, J. Wachter, B. Nuber, M.L. Ziegler), (367) 197

Dicarbonylcyclopentadienylmethylthiolatoiron(II), reactions with monocynoacetylene of; crystal structures of $[\text{FeC}(\text{O})\text{C}(\text{CN})=\text{CH}(\text{SCH}_3)(\eta^5\text{-C}_5\text{H}_5)(\text{CO})]$ and of $[\mu\text{-}\eta^2\text{-Fe}(\text{C}=\text{CCN})(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2][\text{Co}(\text{CO})_3]_2$ (R. Kergoat, M.M. Kubicki, L.C. Gomes de Lima, H. Scordia, J.E. Guerschais, P. L'Haridon), (367) 143

Group 6 metal tetracarbonyl complexes, *cis*- $[\text{M}(\text{CO})_4\{(\text{C}_5\text{H}_4\text{ECH}_3)_2\text{Fe}\}]$ ($\text{M} = \text{Cr}, \text{Mo}, \text{W}; \text{E} = \text{S}$ or Se); crystal structure of 1,1'-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Thiolate-amine complexes $[\text{Ru}(\text{CO})_2(\text{bmae})]$ and $[\text{Ru}(\text{CO})_2(\text{}^{\text{bu}}\text{bmae})]$ ($\text{bmae}^{2-} = 1,2\text{-bis}(2\text{-mercaptoanilino})\text{ethane}(-2)$; $\text{}^{\text{bu}}\text{bmae} = 1,2\text{-bis}(2\text{-mercapto-3,5-di-t-butylalanino})\text{ethane}(-2)$). UV photolysis of (D. Sellmann, O. Käppler, F. Knoch), (367) 161

Tellurium

Chalcogens, contrasting roles in synthesis of mixed metal complexes $(\text{CO})_6\text{Fe}_2(\mu_3\text{-E})_2\text{M}(\text{PPh}_3)_2$ ($\text{E} = \text{Te}, \text{Se}, \text{S}; \text{M} = \text{Ni}, \text{Pd}, \text{Pt}$) of (P. Mathur, I.J. Mavunkal, V. Rugmini), (367) 243

Tin

Bis[8-(dimethylamino)1-naphthyl-*C,N*]tin(II) and its $\text{W}(\text{CO})_5$ adduct, synthesis and characterization of (J.T.B.H. Jastrzebski, P.A. van der Schaaf, J. Boersma, G. van Koten, D. Heijdenrijk, K. Goubitz, D.J.A. de Ridder), (367) 55

t-Butyl-substituted tin-phosphor compounds, preparation and structure of (D. Hänssgen, H. Aldenhoven, M. Nieger), (367) 47

Methyl 4-tri-*n*-butylstannylbenzoate, *P*-(methoxycarbonyl)phenylmercuric chloride and *P*-(methoxycarbonyl)phenylboronic acid, synthesis and iodination of (M.D. Hylarides, D.S. Wilbur, S.W. Hadley, A.R. Fritzberg), (367) 259

Titanium

Bis(cyclopentylcyclopentadienyl)titanium dichloride and bis(cyclohexylcyclopentadienyl)titanium dichloride, crystal structures of (M. Kapon, G.M. Reisner), (367) 77

$(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{CO})_2$, reaction with sulphoxides of (T.L. Chen, A. Shaver, T.H. Chan), (367) C5

Dicyclopentadienyltitanium dihalides, electronic structure as indicated by UV photoelectron spectroscopy of (T. Vondrák, K. Mach, V. Varga), (367) 69

Tungsten

Bis[8-(dimethylamino)1-naphthyl-*C,N*]tin(II) and its $\text{W}(\text{CO})_5$ adduct, synthesis and characterization of (J.T.B.H. Jastrzebski, P.A. van der Schaaf, J. Boersma, G. van Koten, D. Heijdenrijk, K. Goubitz, D.J.A. de Ridder), (367) 55

Group 6 metal tetracarbonyl complexes, *cis*- $[\text{M}(\text{CO})_4\{(\text{C}_5\text{H}_4\text{ECH}_3)_2\text{Fe}\}]$ ($\text{M} = \text{Cr}, \text{Mo}, \text{W}; \text{E} = \text{S}$ or Se); crystal structure of 1,1'-bis(methylseleno)ferrocene tetracarbonyltungsten (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik, P.A. Bates, M.B. Hursthouse), (367) 275

Molybdenum(II) and tungsten(II) cationic seven-coordinate complexes, synthesis of (P.K. Baker, M. Bamber, G.W. Rogers), (367) 101

Vanadium

Cyclopentadienylalkynevanadium(I) CO-free complexes, facile synthesis and reactivity of (B. Hessen, J.H. Teuben), (367) C18

Zinc

$\text{Et}_3\text{SiCF}=\text{CFM}$ ($M = \text{Li}, \text{ZnBr}$), preparation and reactivity of (P. Martinet, R. Sauvêtre, J.-F. Normant), (367) 1

Palladium(0)-mediated reactions of $\text{BrZnCH}(\text{CH}_3)\text{COO}^t\text{Bu}$ with vinyl and aryl triflates, influence of catalyst on (F. Orsini, F. Pelizzoni, L.M. Vallarino), (367) 375

Zirconium

Boron hydride insertion into Zr–C bond of dimeric (η^2 -formaldehyde)zirconocene (G. Erker, U. Hoffmann, R. Zwitter, G. Krüger), (367) C15

$(\text{C}_5\text{H}_5)_2\text{ZrCl}(\text{AlCl}_4)$, crystal and molecular structure and unusual hydrocarbon solubility properties of; relevance to Ziegler–Natta polymerization chemistry (M.V. Gaudet, M.J. Zaworotko, T.S. Cameron, A. Linden), (367) 267