

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 381 (1990)**SUBJECT INDEX****Arsenic**

- Alkyldyne-tricobalt cluster, metal exchange reactions with reactive μ_3 -CX groups of (H.T. Schacht, H. Vahrenkamp), (381) 261
 Gallium-arsenic four-membered ring compounds containing a bridging $\text{As}(\text{SiMe}_3)_2$ group, synthesis and characterization of; crystal structure of $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$ and $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Br}$ (W.K. Holley, R.L. Wells, S. Shafiezad, A.T. McPhail, C.G. Pitt), (381) 15

Beryllium

- Cyclopentadienylberyllium compounds, synthesis and characterisation of (S.J. Pratten, M.K. Cooper, M.J. Aroney), (381) 147

Boron

- Bicyclic borates containing chiral boron and nitrogen atoms, NMR and X-ray diffraction studies of (N. Farfán, T. Mancilla, D. Castillo, G. Uribe, L. Carillo, P. Joseph-Nathan, R. Contreras), (381) 1
 Molybdenum pyrazolylborato stereochemically nonrigid complexes containing a *dihapto*-thiocarbamido ligand, synthesis, structure and properties of (T. Desmond, F.J. Lalor, B. O'Sullivan, G. Ferguson), (381) C33
 2,4,6-Tri-t-butylphenyl-substituted iminoboranes, preparation, structures and reactions of (G. Elter, M. Neuhaus, A. Meller, D. Schmidt-Bäse), (381) 299

Chromium

- Alkene aminocarbene bidentate chromium complexes, alkyne insertion reactions of; formation and crystal structure of dihydroazepinetricarbonylchromium (A. Parlier, R. Yefsah, M. Rudler, H. Rudler, J.C. Daran, J. Vaissermann), (381) 191
 $\text{trans}\text{-}[(\text{CO})_5\text{M}\{\text{Ph}_2\text{PX}(\text{CH}_2)_3\text{N}=\text{CHC}_6\text{H}_4\text{-o-O}\}]_2\text{M}'$ ($\text{M} = \text{Mo}$; $\text{X} = \text{NH}$ or $\text{M} = \text{Cr}$, W ; $\text{X} = \text{CH}_2$; $\text{M}' = \text{Ni}$, Cu , Zn) complexes, synthesis and characterization of; X-ray crystal structure of $\text{trans}\text{-}[(\text{CO})_5\text{Mo}\{\text{P}(\text{OCH}_2\text{CMe}_2\text{CH}_2\text{O})\text{NH}(\text{CH}_2)_2\text{N}=\text{CHC}_6\text{H}_4\text{-o-O}\}]_2\text{Cu}$, (G.M. Gray, N. Takada, M. Jan, H. Zhang, J.L. Atwood), (381) 53
 Group 6 metal carbonyl complexes containing 2-diphenylphosphinopyridine and 3,6-bis(diphenylphosphino)pyridazine, synthesis of; molecular structure of $\text{Cr}(\text{CO})_5(3\text{-diphenylphosphino-6-diphenylphospholypyridazine})$, (Z.-Z. Zhang, H.-K. Wang, Y.-J. Shen, H.-G. Wang, R.-J. Wang), (381) 45
 Pentamethylcyclopentadienylchromium tricarbonyl dimer, isolation of; X-ray crystal structure of two polymorphs (L.Y. Goh, T.W. Hambley, D.J. Daresbourg, J. Reibenspies), (381) 349
 Trifluoromethylisocyanide complexes of chromium, molybdenum and tungsten, synthesis of hexakis(trifluoromethylisocyanide) chromium (D. Lentz), (381) 205

Cobalt

- Alkyldyne-tricobalt cluster, metal exchange reactions with reactive μ_3 -CX groups of (H.T. Schacht, H. Vahrenkamp), (381) 261
 $(\eta^4\text{-2-Bromo- and -iodo-5-trimethylsilylcyclopentadienone})(\eta^5\text{-cyclopentadienyl})\text{cobalt}$ and their 3,4-tri-, tetra-, and -penta-methylene derivatives, synthesis of, (P.C. Auderset, E.R.F. Gesing), (381) 139
 Cobaltadisiloxane cyclic compounds; crystal structures of a pyridinolithium [bis(cyclosiloxy)cobalt]cobalt chloride and a bis(tetramethylethylenediaminolithium)bis(cyclosiloxy)cobaltate (M.B. Hursthouse, M.A. Mazid, M. Mottevalli, M. Sangane, A.C. Sullivan), (381) C43

- Cobalt mediated alkene/diene coupling; documentation of scope, limitations, and regioselectivity (K.S. Feldman, K.C. Grega), (381) 251
- Cyclopentadienyldimethylsilane, bidentate silicon containing ligand, formation of and reactions with (E. Colomer, R.J.P. Corriu, R. Pleixats), (381) C1
- $\text{Fe}(\text{CO})_3[\text{Fe}(\text{CO})_3\text{-CH-CPh-S}]$, addition and substitution reactions of (T. Fässler, G. Huttner), (381) 391
- Heterotetranuclear anionic clusters, $[\text{MCo}_3(\text{CO})_{12-2n}(\text{C}_2\text{Ph}_2)_n]^-$ ($\text{M} = \text{Fe, Ru}$; $n = 0, 1$), redox behaviour of (R. Jund, J. Rimmelin, M. Gross), (381) 239
- Metallocenylcyclopentadienylides as precursors for heteropolymetallocenes, preparation and reactions of ferrocenyl- and cobaltoceniumyl-cyclopentadienylides of, (C. Rieker, G. Ingram, P. Jaitner, H. Schottenberger, K.E. Schwarzhans), (381) 127

Copper

- trans*-[(CO)₅M{Ph₂PX(CH₂)₃N=CHC₆H₄-*o*-O}]₂M' ($\text{M} = \text{Mo}$; X = NH or $\text{M} = \text{Cr, W}$; X = CH₂; M' = Ni, Cu, Zn) complexes, synthesis and characterization of; X-ray crystal structure of *trans*-[(CO)₅Mo{P(OCH₂CMe₂CH₂O)NH(CH₂)₂N=CHC₆H₄-*o*-O}]₂Cu, (G.M. Gray, N. Takada, M. Jan, H. Zhang, J.L. Atwood), (381) 53

Gallium

- Gallium-arsenic four-membered ring compounds containing a bridging As(SiMe₃)₂ group, synthesis and characterization of; crystal structure of Ph₂GaAs(SiMe₃)₂Ga(Ph)₂Cl and Ph₂GaAs(SiMe₃)₂-Ga(Ph)₂Br (W.K. Holley, R.L. Wells, S. Shafiezad, A.T. McPhail, C.G. Pitt), (381) 15

Germanium

- Perchloroorganic silicon and germanium compounds, cyclization reaction of; synthesis and crystal structure of perchloro(2,2'-biphenylene)diphenyl-silane and -germane (L. Fajari, L. Juliá, J. Riera, E. Molins, C. Miravittles), (381) 321

Gold

- [2-((Dimethylamino)methyl)phenyl](2-nitrophenyl)gold(III) complexes, synthesis of (J. Vicente, M.-D. Bermúdez, M.-T. Chicote, M.-J. Sánchez-Santano), (381) 285
- Gold(I)-chiral ferrocenylphosphine neutral complexes, enantioselective synthesis of crystal structure of the complex $\{[(\eta^5-\text{C}_5\text{H}_4\text{PPh}_2)(\eta^5-\text{C}_5\text{H}_3(\text{PPh}_2)\text{CH}(\text{Me})\text{N}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}_2]\text{Fe}\}_2(\text{AuCl})_3\cdot\text{Et}_2\text{O}$ (A. Togni, S.D. Pastor, G. Rihs), (381) C21

Iridium

- Cyclopentadienyldimethylsilane, bidentate silicon containing ligand, formation of and reactions with (E. Colomer, R.J.P. Corriu, R. Pleixats), (381) C1
- Phenylacetylido iridium(I) and rhodium(I) complexes, synthesis and reactions of (M.J. Fernández, M.A. Esteruelas, M. Covarrubias, L.A. Oro), (381) 275

Iron

- Alkene- and alkyne-pentamethylcyclopentadienylcarbonyliron complexes synthesis via the aquo derivative $[\text{Fe}(\eta\text{-C}_5\text{Me}_5)(\text{CO})_2(\text{OH}_2)]^+\text{BF}_4^-$ of; crystal structure of the ruthenium analogue (A. Tahiri, V. Guerchais, L. Toupet, C. Lapinte), (381) C47
- μ_3 -Carbynylcarbonyltriiron clusters, reactivity of and addition of alkynes under formation of η^6 -ferracyclopentadienone ligands, (D. Buchholz, G. Huttner, W. Imhof, L. Zsolnai, D. Günauer), (381) 79
- Cyclopentadienyldimethylsilane, bidentate silicon containing ligand, formation of and reactions with (E. Colomer, R.J.P. Corriu, R. Pleixats), (381) C1
- $\text{Fe}(\text{CO})_4(\eta^1\text{-dppf})$ (dppf = (Ph₂PC₅H₄)₂Fe), synthesis and conversion into $\text{Fe}_2(\text{CO})_8(\mu\text{-dppf})$ and $(\text{CO})_4\text{Fe}(\mu\text{-dppf})\text{Mo}(\text{CO})_5$ of, (T.S.A. Hor, L.-T. Phang), (381) 121
- $\text{Fe}_3(\text{CO})_{10}(\mu_3\text{-PR})$, rearrangement of carbonyl groups to carbene, carbyne and alkyne groups in (D. Buchholz, G. Huttner, L. Zsolnai) (381) 97
- $\text{Fe}(\text{CO})_3[\text{Fe}(\text{CO})_3\text{-CH-CPh-S}]$, addition and substitution reactions of (T. Fässler, G. Huttner), (381) 391

- Gold(I)-chiral ferrocenylphosphine neutral complexes, enantioselective synthesis of crystal structure of the complex $\{(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)(\eta^5\text{-C}_5\text{H}_3(\text{PPh}_2)\text{CH}(\text{Me})\text{N}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}_2)\text{Fe}\}_2(\text{AuCl})_3\} \cdot \text{Et}_2\text{O}$ (A. Togni, S.D. Pastor, G. Rihs), (381) C21
- Heterotetrานuclear anionic clusters, $[\text{MCo}_3(\text{CO})_{12-2n}(\text{C}_2\text{Ph}_2)_n]^-$ ($\text{M} = \text{Fe, Ru}; n = 0, 1$), redox behaviour of (R. Jund, J. Rimmelin, M. Gross), (381) 239
- $[\text{HFe}(\text{CO})_4]^-$ as a reagent for the synthesis of tin/iron clusters, partial crystal structure of $(\text{NEt}_4)_2[\text{SnCl}_2(\text{Fe}(\text{CO})_4)_2] \cdot \text{SnCl}_4$ (M. Ferrer, O. Rossell, M. Seco, X. Solans, M. Gómez), (381) 183
- Iron mediated diene activation; crystal structure of $[(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}(\text{methylhexa-2,4-dienoate})\text{CO}][\text{BF}_4]$ (P. McArdle, D. MacHale, M. Mahon, D. Cunningham), (381) C13
- Metalla heterocyclobutadiene complexes, $\text{Fe}(\text{CO})_3[\text{Fe}(\text{CO})_3=\text{CH}=\text{CpH}=\text{S}]$, reductive activation of (T. Fässler, G. Huttner, D. Günauer, S. Fiedler, B. Eber), (381) 409

Lithium

- Lithium dimeric derivative of tetraphenyldisiloxanediolate dianion, synthesis and structure of (M.B. Hursthouse, M.A. Hossain, M. Mottevalli, M. Sangane, A.C. Sullivan), (381) 293

Manganese

- $(\text{C}_5\text{H}_4\text{Me})(\text{CO})_2\text{Mn}[\text{CHCH}(\text{PEt}_3)]$, carbene-like ylid complexes, molecular structure of (R.D. Rogers, H.G. Alt, H.E. Maisel), (381) 233
- Cyclopentadienyl dimethylsilane, bidentate silicon containing ligand, formation of and reactions with (E. Colomer, R.J.P. Corriu, R. Pleixats), (381) C1
- Cyclopentadienyltricarbonyl-manganese and -rhenium chalcogen derivatives of $[(\text{CO})_3\text{MC}_5\text{H}_4]_2\text{E}_n$ ($\text{M} = \text{Mn, Re}; \text{E} = \text{S, Se, Te}; n = 2, 1$) M. Herberhold, M. Biersack), (381) 379
- Diterpenoids cyclomanganation, functionalization of carbon-14 (R.C. Cambie, M.R. Metzler, P.S. Rutledge, P.D. Woodgate), (381) C26

Metallocenes

- Bis(η -cyclopentadienyl)titanium(IV) bis(substituted benzoato) complexes; synthesis and spectra of (Y. Dang, H.J. Geise, R. Domisse, E. Esmans, H.O. Desseyn), (381) 333
- (η^4 -2-Bromo- and -iodo-5-trimethylsilylcyclopentadienone)(η^5 -cyclopentadienyl)cobalt and their 3,4-tri-, -tetra-, and -penta-methylene derivatives, synthesis of, (P.C. Auderset, E.R.F. Gesing), (381) 139
- Gold(I)-chiral ferrocenylphosphine neutral complexes, enantioselective synthesis of crystal structure of the complex $\{(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)(\eta^5\text{-C}_5\text{H}_3(\text{PPh}_2)\text{CH}(\text{Me})\text{N}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}_2)\text{Fe}\}_2(\text{AuCl})_3\} \cdot \text{Et}_2\text{O}$ (A. Togni, S.D. Pastor, G. Rihs), (381) C21
- Heterotetrานuclear anionic clusters, $[\text{MCo}_3(\text{CO})_{12-2n}(\text{C}_2\text{Ph}_2)_n]^-$ ($\text{M} = \text{Fe, Ru}; n = 0, 1$), redox behaviour of (R. Jund, J. Rimmelin, M. Gross), (381) 239
- Isopropyltetramethylcyclopentadienyltitanium derivatives, synthesis, X-ray structures and polymerization of (D.T. Mallin, M.D. Rausch, E.A. Mintz, A.L. Rheingold), (381) 35
- Metallocenylcyclopentadienylides as precursors for heteropolymetallocenes, preparation and reactions of ferrocenyl- and cobaltoceneumyl-cyclopentadienylides of, (C. Rieker, G. Ingram, P. Jaitner, H. Schottenberger, K.E. Schwarzhans), (381) 127

Molybdenum

- Alkyldiene-tricobalt cluster, metal exchange reactions with reactive $\mu_3\text{-CX}$ groups of (H.T. Schacht, H. Vahrenkamp), (381) 261
- trans*- $[(\text{CO})_5\text{M}(\text{Ph}_2\text{PX}(\text{CH}_2)_3\text{N}=\text{CHC}_6\text{H}_4\text{-}o\text{-O})]_2\text{M}'$ ($\text{M} = \text{Mo}; \text{X} = \text{NH}$ or $\text{M} = \text{Cr, W}; \text{X} = \text{CH}_2$; $\text{M}' = \text{Ni, Cu, Zn}$) complexes, synthesis and characterization of; X-ray crystal structure of *trans*- $[(\text{CO})_5\text{Mo}(\text{P}(\text{OCH}_2\text{CMe}_2\text{CH}_2\text{O})\text{NH}(\text{CH}_2)_2\text{N}=\text{CHC}_6\text{H}_4\text{-}o\text{-O})]_2\text{Cu}$, (G.M. Gray, N. Takada, M. Jan, H. Zhang, J.L. Atwood), (381) 53
- (η -Cyclopentadienyl)bis(arenediazo)(triphenylphosphine)molybdenum(0) cations, synthesis, structural characterisation and chemical behaviour of; synthesis of complexes containing $[\text{Mo}(\text{N}_2\text{C}_6\text{H}_4\text{R}^1)(\text{N}_2\text{C}_6\text{H}_4\text{R}^2)]^{2+}$ (G. Ferguson, B.L. Ruhl, M. Parvez, F.J. Lalor, M.E. Deane), (381) 357
- $\text{Fe}(\text{CO})_4(\eta^1\text{-dppf})$ (dppf = $(\text{Ph}_2\text{PC}_5\text{H}_4)_2\text{Fe}$), synthesis and conversion into $\text{Fe}_2(\text{CO})_8(\mu\text{-dppf})$ and $(\text{CO})_4\text{Fe}(\mu\text{-dppf})\text{Mo}(\text{CO})_5$ of, (T.S.A. Hor, L.-T. Phang), (381) 121

- Group 6 metal carbonyl complexes containing 2-diphenylphosphinopyridine and 3,6-bis(diphenylphosphino)pyridazine, synthesis of; molecular structure of $\text{Cr}(\text{CO})_5(3\text{-diphenylphosphino-6-diphenylphospholypyridazine})$, (Z.-Z. Zhang, H.-K. Wang, Y.-J. Shen, H.-G. Wang, R.-J. Wang), (381) 45
- Molybdenum(0) and tungsten(0) complexes containing $cis\text{-}[\text{M}(\text{NO})(\text{N}_2\text{Ar})]^{2+}$ structural unit, synthesis, chemistry and X-ray crystallographic characterisation of (M.E. Deane, F.J. Lalor, G. Ferguson, B.L. Ruhl, M. Parvez), (381) 213
- Molybdenum pyrazolylborato stereochemically nonrigid complexes containing a *dihapto*-thiocarbox-amido ligand, synthesis, structure and properties of (T. Desmond, F.J. Lalor, B. O'Sullivan, G. Ferguson), (381) C33
- Trifluoromethylisocyanide complexes of chromium, molybdenum and tungsten, synthesis of hexakis(trifluoromethylisocyanide chromium) (D. Lentz), (381) 205

Nickel

- Alkylidyne-tricobalt cluster, metal exchange reactions with reactive $\mu_3\text{-CX}$ groups of (H.T. Schacht, H. Vahrenkamp), (381) 261
- trans*- $[(\text{CO})_5\text{M}\{\text{Ph}_2\text{PX}(\text{CH}_2)_3\text{N}=\text{CHC}_6\text{H}_4\text{-o-O}\}]_2\text{M}'$ ($\text{M} = \text{Mo}$; $\text{X} = \text{NH}$ or $\text{M} = \text{Cr}, \text{W}$; $\text{X} = \text{CH}_2$; $\text{M}' = \text{Ni}, \text{Cu}, \text{Zn}$) complexes, synthesis and characterization of; X-ray crystal structure of *trans*- $[(\text{CO})_5\text{Mo}(\text{P}(\text{OCH}_2\text{CMe}_2\text{CH}_2\text{O})\text{NH}(\text{CH}_2)_2\text{N}=\text{CHC}_6\text{H}_4\text{-o-O})]_2\text{Cu}$, (G.M. Gray, N. Takada, M. Jan, H. Zhang, J.L. Atwood), (381) 53
- Nickel σ -sydnonyl complexes, stabilized by tertiary phosphines (L.N. Morozova, L.S. Isaeva, P.V. Petrovskii, D.N. Kravtsov, She Fan Min, V.N. Kalinin), (381) 281

Niobium

- Transition metal polysulphide complexes, chemistry of; synthesis and crystal structures of $\text{Cp}'_3\text{Nb}_3\text{S}_{12}$ and $\text{Cp}'_3\text{Nb}_3\text{S}_{10}\text{O}$ ($\text{Cp}' = t\text{-BuC}_5\text{H}_4$) (H. Brunner, W. Meier, J. Wachter, B. Nuber, M.L. Ziegler), (381) C7

Palladium

- Allylpalladium(II) cationic complexes, mechanism of phenylation by tetraphenylborate anion of. (B. Crociani, F. Di Bianca, L. Canovese, P. Uguagliati), (381) C17
- Diterpenoids cyclomanganation, functionalization of carbon-14 (R.C. Cambie, M.R. Metzler, P.S. Rutledge, P.D. Woodgate), (381) C26

Phosphorus

- Allylpalladium(II) cationic complexes, mechanism of phenylation by tetraphenylborate anion of. (B. Crociani, F. Di Bianca, L. Canovese, P. Uguagliati), (381) C17
- μ_3 -Carbynylcarbonyltriiron clusters, reactivity of and addition of alkynes under formation of η^6 -ferra-cyclopentadienone ligands, (D. Buchholz, G. Huttner, W. Imhof, L. Zsolnai, D. Günauer), (381) 79
- $(\text{C}_5\text{H}_4\text{Me})(\text{CO})_2\text{Mn}[\text{CHCH}(\text{PEt}_3)]$, carbene-like ylid complexes, molecular structure of (R.D. Rogers, H.G. Alt, H.E. Maisel), (381) 233
- Cyclopentadienyltricarbonyl-manganese and -rhenium chalcogen derivatives of $[(\text{CO})_3\text{MC}_5\text{H}_4]_2\text{E}_n$ ($\text{M} = \text{Mn, Re}; \text{E} = \text{S, Se, Te}; n = 2, 1$) M. Herberhold, M. Biersack), (381) 379
- $\text{Fe}(\text{CO})_4(\eta\text{-dppf})$ (dppf = $(\text{Ph}_2\text{PC}_5\text{H}_4)_2\text{Fe}$), synthesis and conversion into $\text{Fe}_2(\text{CO})_8(\mu\text{-dppf})$ and $(\text{CO})_4\text{Fe}(\mu\text{-dppf})\text{Mo}(\text{CO})_5$ of, (T.S.A. Hor, L.-T. Phang), (381) 121
- $\text{Fe}_3(\text{CO})_{10}(\mu_3\text{-PR})$, rearrangement of carbonyl groups to carbene, carbyne and alkyne groups in (D. Buchholz, G. Huttner, L. Zsolnai) (381) 97
- $\text{Fe}(\text{CO})_3[\text{Fe}(\text{CO})_3\text{-CH-CPh-S}]$, addition and substitution reactions of (T. Fässler, G. Huttner), (381) 391
- Gold(I)-chiral ferrocenylphosphine neutral complexes, enantioselective synthesis of crystal structure of the complex $[(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)\times(\eta^5\text{-C}_5\text{H}_3(\text{PPh}_2)\text{CH}(\text{Me})\text{N}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}_2)\text{Fe}]_2(\text{AuCl})_3 \cdot \text{Et}_2\text{O}$ (A. Togni, S.D. Pastor, G. Rihs), (381) C21
- Lithiated di-*t*-butylfluorosilylmesitylphosphane, rearrangement of (M. Andrianarison, D. Stalke, U. Klingenbriel), (381) C38
- Metalla heterocyclobutadiene complexes, $\text{Fe}(\text{CO})_3[\text{Fe}(\text{CO})_3\text{-CH-CPh-S}]$, reductive activation of (T. Fässler, G. Huttner, D. Günauer, S. Fiedler, B. Eber), (381) 409

- Nickel σ -sydnonyl complexes, stabilized by tertiary phosphines (L.N. Morozova, L.S. Isaeva, P.V. Petrovskii, D.N. Kravtsov, She Fan Min, V.N. Kalinin), (381) 281
 Triruthenium dodecacarbonyl, reactions with diphenylphosphine of (L.M. Bullock, J.S. Field, R.J. Haines, E. Minshall, M.H. Moore, F. Mulla, D.N. Smit, L.M. Steer), (381) 429

Platinum

- $\text{Fe}(\text{CO})_3[\text{Fe}(\text{CO})_3\cdots\text{CH}\cdots\text{CPh}\cdots\text{S}]$, addition and substitution reactions of (T. Fässler, G. Huttner), (381) 391

Rhenium

- Cyclopentadienyltricarbonyl-manganese and -rhenium chalcogen derivatives of $[(\text{CO})_3\text{MC}_5\text{H}_4]_2\text{E}_n$ ($\text{M} = \text{Mn, Re; E = S, Se, Te; } n = 2, 1$) M. Herberhold, M. Biersack), (381) 379

Rhodium

- Phenylacetylide iridium(I) and rhodium(I) complexes, synthesis and reactions of (M.J. Fernández, M.A. Esteruelas, M. Covarrubias, L.A. Oro), (381) 275

Ruthenium

- Alkene- and alkyne-pentamethylcyclopentadienyldicarbonyliron complexes synthesis via the aquo derivative $[\text{Fe}(\eta\text{-C}_5\text{Me}_5)(\text{CO})_2(\text{OH}_2)]^+\text{BF}_4^-$ of; crystal structure of the ruthenium analogue (A. Tahiri, V. Guerchais, L. Toupet, C. Lapinte), (381) C47

- Cyclopentadienyldimethylsilane, bidentate silicon containing ligand, formation of and reactions with (E. Colomer, R.J.P. Corriu, R. Pleixats), (381) C1

- Heterotetrานuclear anionic clusters, $[\text{MCo}_3(\text{CO})_{12-2n}(\text{C}_2\text{Ph}_2)_n]^-$ ($\text{M} = \text{Fe, Ru; } n = 0, 1$), redox behaviour of (R. Jund, J. Rimmelin, M. Gross), (381) 239

- Neutral transition metal polyhydrides, reactivity in catalytic transfer hydrogenation of ketones of, (Y. Lin, Y. Zhou), (381) 135

- Triruthenium dodecacarbonyl, reactions with diphenylphosphine of (L.M. Bullock, J.S. Field, R.J. Haines, E. Minshall, M.H. Moore, F. Mulla, D.N. Smit, L.M. Steer), (381) 429

Selenium

- Cyclopentadienyltricarbonyl-manganese and -rhenium chalcogen derivatives of $[(\text{CO})_3\text{MC}_5\text{H}_4]_2\text{E}_n$ ($\text{M} = \text{Mn, Re; E = S, Se, Te; } n = 2, 1$) M. Herberhold, M. Biersack), (381) 379

Silicon

- Alkylidyne-tricobalt cluster, metal exchange reactions with reactive $\mu_3\text{-CX}$ groups of (H.T. Schacht, H. Vahrenkamp), (381) 261

- Cobaltadisiloxane cyclic compounds; crystal structures of a pyridinolithium [bis(cyclosiloxy)cobalt]cobalt chloride and a bis(tetramethylethylenediaminolithium)bis(cyclosiloxy)cobaltate (M.B. Hursthouse, M.A. Mazid, M. Mottevalli, M. Sangane, A.C. Sullivan), (381) C43

- Cyclopentadienyldimethylsilane, bidentate silicon containing ligand, formation of and reactions with (E. Colomer, R.J.P. Corriu, R. Pleixats), (381) C1

- Fluoromethyltrimethylsilanes, synthesis of (V. Broicher, D. Geffken), (381) 315

- Gallium-arsenic four-membered ring compounds containing a bridging $\text{As}(\text{SiMe}_3)_2$ group, synthesis and characterization of; crystal structure of $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Cl}$ and $\text{Ph}_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{Ph})_2\text{Br}$ (W.K. Holley, R.L. Wells, S. Shafieezad, A.T. McPhail, C.G. Pitt), (381) 15

- Lithiated di-t-butylfluorosilylmesitylphosphane, rearrangement of (M. Andrianarison, D. Stalke, U. Klingebiel), (381) C38

- Lithium dimeric derivative of tetraphenyldisiloxanediolate dianion, synthesis and structure of (M.B. Hursthouse, M.A. Hossain, M. Mottevalli, M. Sangane, A.C. Sullivan), (381) 293

- Perchloroorganic silicon and germanium compounds, cyclization reaction of; synthesis and crystal structure of perchloro(2,2'-biphenylene)diphenyl-silane and -germane (L. Fajari, L. Juliá, J. Riera, E. Molins, C. Miravittles), (381) 321

- Poly[(disilanylene)diethynylene] with highly conducting properties, synthesis of (M. Ishikawa, Y. Hasegawa, A. Kunai, T. Yamanaka), (381) C57
- Sterically crowded aryloxide silicon compounds, preparation, characterization and X-ray structures of (M.D. Healy, A.R. Barron), (381) 165
- Trichlorosilane, γ -ray induced reduction of acid chlorides with (K. Oka, R. Nakao, Y. Abe, T. Dohmaru), (381) 155
- 2,4,6-Tri-t-butylphenyl-substituted iminoboranes, preparation, structures and reactions of (G. Elter, M. Neuhaus, A. Meller, D. Schmidt-Bäse), (381) 299

Sulphur

- Molybdenum pyrazolylborato stereochemically nonrigid complexes containing a *dihapto*-thiocarbonoamido ligand, synthesis, structure and properties of (T. Desmond, F.J. Lalor, B. O'Sullivan, G. Ferguson), (381) C33
- Transition metal polysulphide complexes, chemistry of; synthesis and crystal structures of $Cp'Nb_3S_{12}$ and $Cp'Nb_3S_{10}O$ ($Cp' = t\text{-BuC}_5H_4$) (H. Brunner, W. Meier, J. Wachter, B. Nuber, M.L. Ziegler), (381) C7

Tellurium

- Bis(aryl telluro)methanes, synthesis and characterization of (L. Torres C.), (381) 69
- Cyclopentadienyltricarbonyl-manganese and -rhenium chalcogen derivatives of $[(CO)_3MC_5H_4]_2E_n$ ($M = Mn, Re; E = S, Se, Te; n = 2, 1$) M. Herberhold, M. Biersack), (381) 379
- Metallocenylcyclopentadienylides as precursors for heteropolymetallocenes, preparation and reactions of ferrocenyl- and cobaltoceniumyl-cyclopentadienylides of, (C. Rieker, G. Ingram, P. Jaitner, H. Schottenberger, K.E. Schwarzhans), (381) 127

Thallium

- Metallocenylcyclopentadienylides as precursors for heteropolymetallocenes, preparation and reactions of ferrocenyl- and cobaltoceniumyl-cyclopentadienylides of, (C. Rieker, G. Ingram, P. Jaitner, H. Schottenberger, K.E. Schwarzhans), (381) 127

Tin

- Di-t-butylchlorostannane, reactions with sterically congested aryllithium compounds of (M. Weidenbruch, K. Schäfers, K. Peters, H.G. von Schnering), (381) 173
- 1,1,6,6,11,11-Hexamethyl-1,6,11-tristannacyclopentadecane, synthesis and complexing ability of (K. Jurkschat, A. Rühlemann, A. Tzschach), (381) C53
- $[HFe(CO)_4]^-$ as a reagent for the synthesis of tin/iron clusters, partial crystal structure of $(NEt_4)_2[SnCl_2\{Fe(CO)_4\}_2] \cdot SnCl_4$ (M. Ferrer, O. Rossell, M. Seco, X. Solans, M. Gómez), (381) 183

Titanium

- Bis(η -cyclopentadienyl)titanium(IV) bis(substituted benzoato) complexes; synthesis and spectra of (Y. Dang, H.J. Geise, R. Dommisse, E. Esmans, H.O. Desseyn), (381) 333
- $(C_5H_5)_2TiX$ ($X = Cl, Br$) applied in dehalogenation of organic halides (Q. Yanlong, L. Guisheng, Y.-Z. Huang), (381) 29
- Isopropyltetramethylcyclopentadienyltitanium derivatives, synthesis, X-ray structures and polymerization of (D.T. Mallin, M.D. Rausch, E.A. Mintz, A.L. Rheingold), (381) 35

Tungsten

- Alkylidene-tricobalt cluster, metal exchange reactions with reactive $\mu_3\text{-CX}$ groups of (H.T. Schacht, H. Vahrenkamp), (381) 261
- trans*- $[(CO)_5M\{Ph_2PX(CH_2)_3N=CHC_6H_4-o-O\}]_2M'$ ($M = Mo; X = NH$ or $M = Cr, W; X = CH_2$; $M' = Ni, Cu, Zn$) complexes, synthesis and characterization of; X-ray crystal structure of *trans*- $[(CO)_5Mo\{P(OCH_2CMe_2CH_2O)NH(CH_2)_2N=CHC_6H_4-o-O\}]_2Cu$, (G.M. Gray, N. Takada, M. Jan, H. Zhang, J.L. Atwood), (381) 53

- Group 6 metal carbonyl complexes containing 2-diphenylphosphinopyridine and 3,6-bis(diphenylphosphino)pyridazine, synthesis of; molecular structure of $\text{Cr}(\text{CO})_5(3\text{-diphenylphosphino-6-diphenylphospholypyridazine})$, (Z.-Z. Zhang, H.-K. Wang, Y.-J. Shen, H.-G. Wang, R.-J. Wang), (381) 45
- Molybdenum(0) and tungsten(0) complexes containing *cis*- $[\text{M}(\text{NO})(\text{N}_2\text{Ar})]^{2+}$ structural unit, synthesis, chemistry and X-ray crystallographic characterisation of (M.E. Deane, F.J. Lalor, G. Ferguson, B.L. Ruhl, M. Parvez), (381) 213
- Trifluoromethylisocyanide complexes of chromium, molybdenum and tungsten, synthesis of hexakis(trifluoromethylisocyanide chromium) (D. Lentz), (381) 205

Zinc

- trans*- $[(\text{CO})_5\text{M}\{\text{Ph}_2\text{PX}(\text{CH}_2)_3\text{N}=\text{CHC}_6\text{H}_4\text{-}o\text{-O}\}]_2\text{M}'$ ($\text{M} = \text{Mo}$; $\text{X} = \text{NH}$ or $\text{M} = \text{Cr}$, W ; $\text{X} = \text{CH}_2$; $\text{M}' = \text{Ni}$, Cu , Zn) complexes, synthesis and characterization of; X-ray crystal structure of *trans*- $[(\text{CO})_5\text{Mo}\{\text{POCH}_2\text{CMe}_2\text{CH}_2\text{O}\}\text{NH}(\text{CH}_2)_2\text{N}=\text{CHC}_6\text{H}_4\text{-}o\text{-O}\}]_2\text{Cu}$, (G.M. Gray, N. Takada, M. Jan, H. Zhang, J.L. Atwood), (381) 53

Book reviews

- Bioinorganic chemistry, structure and bonding, Volume 70, contributions by P. Aisen, B.C. Antanaitis, S.J. Berners-Price, K. Doi, H. Köpf-Maier, and P.J. Sadler (P.A. Chaloner), (381) C62
- Carcinogenic and mutagenic metal compounds 2, Current topics in environmental and toxicological chemistry, Volume 11, edited by E. Merian, R.W. Frei, W. Härdi, and Ch. Schlatter (P.A. Chaloner), (381) C63
- Gmelin handbook of inorganic chemistry, Si — silicon. Supplement Volume B4. Silicon–nitrogen compounds. (C. Eaborn), (381) C31
- Inorganic syntheses, Volume 25, editor-in-chief H.R. Allcock (P.A. Chaloner), (381) C61
- Modern synthetic methods, edited by R. Scheffold (P.A. Chaloner), (381) C60