

**JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 401 (1991)-425 (1992)****SUBJECT INDEX****Actinides**

Lanthanides and actinides; Annual Survey covering the years 1984-1986 (R.D. Rogers, L.M. Rogers), (416) 201

**Aluminium**

- Metalloporphyrins with metal-metal  $\sigma$ -bonds; synthesis, spectroscopic characterization, and electrochemistry of (P)MRe(CO)<sub>5</sub> where P is the dianion of octaethylporphyrin (OEP) or tetraphenylporphyrin (TPP) and M = Al, Ga, In or Tl (R. Guillard, A. Zrineh, A. Tabard, L. Courthaudon, B. Han, M. Ferhat, K.M. Kadish), (401) 227
- Ethylaluminium halides, <sup>27</sup>Al NMR spectroscopy of (Z. Černý, J. Macháček, J. Fusek, S. Heřmánek, O. Kříž, B. Čásenský), (402) 139
- Chlorinated trimethylsilyl  $\alpha$ -allenyl alcohols, regioselective synthesis via an organoaluminium compound (D. Mesnard, L. Miginiac), (403) 299
- Silazane derivatives of digallane(4) and diindane(4) with Ga-Ga and In-In bonds, respectively (M. Veith, F. Goffing, S. Becker, V. Huch), (406) 105
- Zirconocene dichloride/methylalumoxane systems, catalysts for olefin polymerization; IR and NMR studies of (L.A. Nekhaeva, G.N. Bondarenko, S.V. Rykov, A.I. Nekhaev, B.A. Krentsel, V.P. Mar'in, L.I. Vyshinskaya, I.M. Khrapova, A.V. Polonskii, N.N. Korneev), (406) 139
- Dinitrosylmolybdenum alkylidene complexes containing n-alkyl alkoxy ligands; synthesis, spectroscopic characterisation and metathesis activity of (A. Keller), (407) 237
- Silyl-substituted phosphinomethanide ligands; preparation of dimeric Al heterocycles; crystal structure of (Me<sub>2</sub>Al[CPMe<sub>2</sub>]SiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub> (H.H. Karsch, K. Zellner, J. Lachmann, G. Müller), (409) 109
- Alkylaluminiumphenoxides, three- or four-coordinated aluminium in; distinction by use of <sup>27</sup>Al NMR spectroscopy (R. Benn, E. Janssen, H. Lehmkuhl, A. Rufińska, K. Angermund, P. Betz, R. Goddard, C. Krüger), (411) 37
- Methyl- and ethylrhenium oxides, synthesis and structural aspects (W.A. Herrmann, C.C. Romao, P. Kiprof, J. Behm, M.R. Cook, M. Taillefer), (413) 11
- Platinum-indium and cobalt-gallium complexes, synthesis of; crystal structure of [Cy<sub>2</sub>P(CH<sub>2</sub>CH<sub>2</sub>)PCy<sub>2</sub>]-Pt(Ind<sub>2</sub>)<sub>2</sub>(R) (Cy = cyclo-C<sub>6</sub>H<sub>11</sub>) (R.A. Fischer, J. Behm), (413) C10
- Alumohydride group, change of bonding mode in biscyclopentadienylhydrido REM complexes from heterometallic to homometallic; crystal and molecular structures of [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Yb(μ<sub>3</sub>-H)]<sub>2</sub>[(μ<sub>2</sub>-H)AlH<sub>2</sub>·N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>]<sub>2</sub>·C<sub>6</sub>H<sub>6</sub>, [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Lu(μ<sub>2</sub>-H)]<sub>2</sub>[(μ<sub>2</sub>-H)<sub>2</sub>AlH·N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>]<sub>2</sub>·C<sub>6</sub>H<sub>6</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Lu]<sub>2</sub>[(μ<sub>2</sub>-H)<sub>2</sub>Lu(μ<sub>3</sub>-H)] (S.Ya. Knjazhanskij, B.M. Bulychev, O.K. Kireeva, V.K. Belsky, G.L. Soloveichik), (414) 11
- $\mu$ -Methylene bis[chlorobis(trimethylsilyl)methylalane] {(Me<sub>3</sub>Si)<sub>2</sub>CH(Cl)Al-CH<sub>2</sub>-Al(Cl)CH(SiMe<sub>3</sub>)<sub>2</sub>} having a heteroadamantane structure (W. Uhl, M. Layh), (415) 181
- Homogeneous Cp<sub>2</sub>ZrCl<sub>2</sub>/methylaluminoxane Ziegler-Natta catalysts, reversible and irreversible deactivation of propene polymerization (D. Fischer, R. Mühlaupt), (417) C7
- Group IIIB organometallic peroxides, synthesis and reactivity of (Yu.A. Alexandrov, N.V. Chikinova), (418) 1
- Coordinatively oversaturated metallocene derivatives, crystal and molecular structure of [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Sm(μ<sub>3</sub>-H)]<sub>2</sub>[(μ<sub>2</sub>-H)<sub>2</sub>AlH·N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>]<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>tBu)<sub>2</sub>Sm]<sub>2</sub>-(μ<sub>2</sub>-H)μ-[(μ<sub>3</sub>-H)<sub>2</sub>Al(μ<sub>2</sub>-H)<sub>2</sub>·Me<sub>2</sub>NC<sub>2</sub>H<sub>4</sub>NMe<sub>2</sub>]} complexes (V.K. Belsky, Y.K. Gun'ko, B.M. Bulychev and G.L. Soloveichik), (419) 299
- Homo- and heteronuclear hydrido lutetiecene complexes with bulky 1,3-di(tert-butyl)cyclopentadienyl (C<sub>5</sub>H<sub>3</sub>tBu<sub>2</sub>) ligands, molecular structure of [(tBu<sub>2</sub>C<sub>5</sub>H<sub>3</sub>)LuH]<sub>4</sub>[AlH<sub>4</sub>·Et<sub>2</sub>O]<sub>2</sub>[AlH<sub>4</sub>]<sub>2</sub> (S.Y. Knjazhanski, E.B. Lobkovsky, B.M. Bulychev, V.K. Belsky and G.L. Soloveichik), (419) 311

Crystal and molecular structures of the octanuclear aluminohydride samarium complex ( $\eta^5\text{-C}_5\text{H}_3\text{Bu}_2\text{Sm}[(\mu_2\text{-H})_2(\mu_3\text{-H})_2\text{Al}(\text{Me}_2\text{NC}_2\text{H}_4\text{NMe}_2)]_2$ ) [ $(\eta^5\text{-C}_5\text{H}_3\text{Bu}_2\text{SmH})_2[(\mu_2\text{-H})_3\text{Al}(\mu_2\text{-H})\text{Al}(\mu_2\text{-H})_3](\mu_3\text{-H})_2\text{Sm}(\eta^5\text{-C}_5\text{H}_3\text{Bu}_2)_2$ ] (V.K. Belsky, Y.K. Gun'ko, B.M. Bulychev and G.L. Soloveichik), (420) 43

Synthèse en une étape par voie organométallique d'amines primaires  $\beta$ -acétyléniques  $\alpha$ -ramifiées (L. Leboutet, G. Courtois and L. Miginiac), (420) 155

*Ab initio* molecular orbital study of dialuminum-ethylene complexes (S.P. So), (420) 293

Some new organotin(IV), aluminium(III)- $\mu$ -oxoisopropoxides and their benzoylacetone derivatives (P.N. Kapoor, A.K. Bhagi, R.N. Kapoor and H.K. Sharma), (420) 321

The reaction of trimethylaluminium with 1,2-dihydroxymethylbenzene (S. Pasynkiewicz and W. Ziemkowska), (423) 1

## Antimony

[ $\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{ER}_3)$ ]BF<sub>4</sub> complexes (E = N, P, As, Sb and Bi), synthesis and characterisation of; crystal structure of [ $\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{Bi}(\text{C}_6\text{H}_5)_3)$ ]BF<sub>4</sub> (H. Schumann, L. Eguren), (403) 183

Antimony; Annual Survey covering the year 1989 (L.D. Freedman, G.O. Doak), (404) 49

Ethyl-, propyl- and butyl-antimony, electrosynthesis of (Y. Mourad, Y. Mugnier, H.J. Breunig, M. Ates), (406) 323

Colourless  $\text{C}_5\text{H}_5\text{SbCl}_2$ , intra- and intermolecular antimony-Cp  $\pi$ -interactions in (W. Frank), (406) 331

Cyclopentadienyliron arene cations, photochemical arene substitution in; synthesis of cyclopentadienyl-iron bisphosphine ligand complexes with group 14 and 15 ligands (H. Schumann, L. Eguren, J.W. Ziller), (408) 361

Allylstibonium bromide, stereoselective addition to aldehydes (Y.-Z. Huang, L.-J. Zhang, C. Chen, G.-Z. Guo), (412) 47

Group 15 organometallics, high metal coordination numbers in; crystal structures of triphenylbismuth-bis(trifluoroacetate) and triphenylantimonybis(trifluoroacetate) (G. Ferguson, B. Kaitner, C. Glidewell and S. Smith), (419) 283

## Arsenic

Cationic titanocene bishexafluoroarsenate complexes (P. Gowik, T. Klapötke), (402) 349

[ $\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{ER}_3)$ ]BF<sub>4</sub> complexes (E = N, P, As, Sb and Bi), synthesis and characterisation of; crystal structure of [ $\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{Bi}(\text{C}_6\text{H}_5)_3)$ ]BF<sub>4</sub> (H. Schumann, L. Eguren), (403) 183

Titanocene polyselenide complexes, reactions with methyldichloroarsane; synthesis and properties of ( $\text{CH}_3\text{C}_5\text{H}_4$ )<sub>2</sub>Ti( $\mu_2\text{-Se}_2$ )<sub>2</sub>AsCH<sub>3</sub> and of ( $\text{CH}_3\text{As}$ )<sub>2</sub>Se<sub>3</sub> (B. Holz, R. Steudel), (406) 133

Gallium-arsenic compounds containing a four-membered Ga-As-Ga-Cl or Ga-As-Ga-As ring, synthesis and characterisation of; crystal structures of ( $\text{Me}_3\text{SiCH}_2$ )<sub>2</sub>GaAs(SiMe<sub>3</sub>)<sub>2</sub>Ga(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>Cl and [ $(\text{Me}_3\text{SiCH}_2)_2\text{GaAs}(\text{SiMe}_3)_2$ ]<sub>2</sub> (R.L. Wells, J.W. Pasterczyk, A.T. McPhail, J.D. Johansen, A. Alvanipour), (407) 17

Cyclopentadienyliron arene cations, photochemical arene substitution in; synthesis of cyclopentadienyl-iron bisphosphine ligand complexes with group 14 and 15 ligands (H. Schumann, L. Eguren, J.W. Ziller), (408) 361

(+)-*cis*-[Dicarbonyl- $\mu$ -chloro- $\mu$ -[5 $\beta$ -methyl-2 $\alpha$ -(1-methylethyl)cyclohexanethiolato]bis[tris(1,1-dimethylethyl)arsine]dirhodium, synthesis, structure determination and catalytic activity of (H. Schumann, B. Gorella, M. Eisen, J. Blum), (412) 251

Acetylstyrene- and benzoylstyrene-tricarbonylchromium, and 1-acetyl- and 1-benzoyl-2-ferrocenylethylene, preparation of (J. Huang, Y.-Z. Huang), (414) 49

( $\mu$ -tert-Butylthio)( $\mu$ -methylarsino)hexacarbonyldiiron double cluster complex, synthesis and molecular structure of (L.-C. Song, Q.-M. Hu), (414) 219

Dicarbollyl complexes of phosphor and arsenic, X-ray diffraction study on ClAs[C<sub>2</sub>Me<sub>2</sub>B<sub>9</sub>H<sub>9</sub>] (P. Jutzi, D. Wegener, M. Hursthouse), (418) 277

## Bismuth

Sterically crowded aryl bismuth compounds; synthesis and characterization of bis[2,4,6-tris(trifluoromethyl)phenyl]bismuth chloride and tris[2,4,6-tris(trifluoromethyl)phenyl]bismuth (K.H. Whitmire, D. Labahn, H.W. Roesky, M. Noltemeyer, G.M. Sheldrick), (402) 55

- $\text{BiCl}_3$ , reaction with sodium 2,4,6-tris(trifluoromethyl)phenoxide; C–F bond activation in (K.H. Whitmire, H.W. Roesky, S. Brooker, G.M. Sheldrick), (402) C4
- $[\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{ER}_3)]\text{BF}_4$  complexes ( $\text{E} = \text{N, P, As, Sb}$  and  $\text{Bi}$ ), synthesis and characterisation of; crystal structure of  $[\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{Bi}(\text{C}_6\text{H}_5)_3)]\text{BF}_4$  (H. Schumann, L. Eguren), (403) 183
- Bismuth; Annual Survey covering the year 1989 (G.O. Doak, L.D. Freedman), (404) 87
- Diphenyl(trifluoromethyl)bismuth and phenylbis(trifluoromethyl)bismuth, new organoperfluoroalkyl derivatives of tervalent bismuth (S. Pasenok, D. Naumann, W. Tyrra), (417) C47
- Group 15 organometallics, high metal coordination numbers in; crystal structures of triphenylbismuth-bis(trifluoroacetate) and triphenylantimonybis(trifluoroacetate) (G. Ferguson, B. Kaitner, C. Glidewell and S. Smith), (419) 283

## Boron

- Borylation of dicarbanions; syntheses of new five- and eight-membered boron–carbon rings (G.E. Herberich, U. Eigendorf, C. Ganter), (402) C17
- Boron; boranes in organic synthesis; Annual Survey covering the year 1988 (G.W. Kabalka, L.H.M. Guindi), (404) 1
- Boron–nitrogen–carbon heterocycles, synthesis of; crystal structure of dimethyl-2,4-*o*-tolyl-3-methyl-8-dibora-2,4-diazaro-1,3-naphthalene (A. Thozet, S. Allaoud, T. Zaïr, A. Karim, B. Frange), (406) 269
- Bis(1,3-di-*tert*-butylcyclopentadienyl)cerium chloride and borohydride, crystal and molecular structures of; a bridging tetradeinate  $\text{BH}_4^-$  group with two  $\mu_3$ -hydrogens:  $\mu:\eta^4:[(\mu_3\text{-H})_2\text{B}(\mu_2\text{-H})_2]$  (E.B. Lobkovsky, Yu.K. Gun'ko, B.M. Bulychev, V.K. Belsky, G.L. Soloveichik, M.Yu. Antipin), (406) 343
- (Ar $\text{BMe}_3$ ) $\text{Li}\cdot\text{D}$  complexes (Ar =  $\text{C}_6\text{H}_4\text{CH}(\text{X})\text{NMe}_2$ ); structure, dynamic molecular processes and the nature of the carbon–lithium bond in the light of multinuclear ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^7\text{Li}$ ,  $^{11}\text{B}$ ) NMR measurements at variable temperature (E. Kalbarczyk-Bidelska, S. Pasynkiewicz), (407) 143
- Dialkylaminobis(trifluoromethyl)boranes, [2 + 2] cycloaddition reactions with isocyanates and isothiocyanates; crystal structures of  $(\text{CF}_3)_2\text{BNMe}_2\text{CON}^+\text{Bu}$ ,  $(\text{CF}_3)_2\text{BNMe}_2\text{CSN}^+\text{Bu}$ ,  $(\text{CF}_3)_2\text{BNPhC}(\text{NMe}_2)\text{S}$  and  $(\text{CF}_3)_2\text{BNMeC}(\text{NET}_2)\text{OCONMe}$  (A. Ansorge, D.J. Brauer, H. Bürger, F. Dörrenbach, T. Hagen, G. Pawelke, W. Weuter), (407) 283
- Borido cluster  $\text{Fe}_4(\text{CO})_{12}\text{BAu}_2[\text{AsPh}_3]_2$ , synthesis and molecular structure of; investigation of the electrochemistry of  $\text{Fe}_4(\text{CO})_{12}\text{BAu}_2\text{L}_2$ , L =  $\text{AsPh}_3$  or  $\text{PPh}_3$  (C.E. Housecroft, M.S. Shongwe, A.L. Rheingold, P. Zanello), (408) 7
- Allylic organopotassium and organoboron compounds in the hydroxymethylation and formylation of cycloalkenes (M. Zajlewicz), (409) 103
- Halogenophenyl- and phenoxyhydrododecaborates,  $[(\text{XC}_6\text{H}_4)_n\text{B}_{12}\text{H}_{12-n}]^{2-}$  (X = Br, I; n = 1–3) and  $[(\text{C}_6\text{H}_5\text{O})\text{B}_{12}\text{H}_{11}]^{2-}$ ; synthesis and  $^{11}\text{B}$ ,  $^{13}\text{C}$  and  $^1\text{H}$  NMR spectra of (W. Preetz, R. von Bismarck), (411) 25
- 1-Azaadamantane, synthesis from 1-boraadamantane by a new method (Yu. N. Bubnov, M.E. Gursky, D.G. Pershin), (412) 1
- 1-Boraadamantane derivatives with functional groups in the side chain (Yu.N. Bubnov, T.V. Potapova, M.E. Gursky), (412) 311
- $[\text{NEt}_4]\text{WFe}_2(\mu_3\text{-}\eta^3\text{-CH=C=C('Bu)}\text{C}(\text{O})\text{C}_2\text{B}_9\text{H}_8\text{Me}_2)(\text{CO})_8$ , a tungstendiiiron cluster derived from  $[\text{Fe}_2(\text{CO})_9]$  and  $[\text{NEt}_4]\text{W}(\equiv\text{CC=C('Bu)}\text{C}(\text{O}))_2(\eta^5\text{-C}_2\text{B}_9\text{H}_9\text{Me}_2)$  (N. Carr, S.J. Dossett, F.G.A. Stone), (413) 223
- Diorganoindium fluorides; crystal structures of  $[{}^i\text{Pr}_2\text{In}(\text{THF})_2][\text{BF}_4]$  and  $(\text{MesBO})_3$  (B. Neumüller, F. Gahlmann), (414) 271
- Pentaphenylboracyclotrisiloxane,  $\text{BSi}_2\text{O}_3\text{Ph}_5$ , highly strained heterocyclosiloxane; synthesis and crystal structure of (D.A. Foucher, A.J. Lough, I. Manners), (414) C1
- (Ar $\text{BMe}_3$ ) $\text{Li}\cdot\text{OEt}_2\text{OEt}_2$  complexes, studies on reactions with the electron donors  $\text{OEt}_2$  and THF by means of  $^1\text{H}$ ,  $^7\text{Li}$ ,  $^{11}\text{B}$ , and  $^{13}\text{C}$  NMR spectroscopy (E. Kalbarczyk-Bidelska, S. Pasynkiewicz), (417) 1
- Group IIIB organometallic peroxides, synthesis and reactivity of (Yu.A. Alexandrov, N.V. Chikinova), (418) 1
- Dihalogenophenylhydrododecacarbonates  $[(m,m\text{-X}_2\text{C}_6\text{H}_3)_n\text{B}_{12}\text{H}_{12-n}]^{2-}$  and  $[(p,m\text{-X}_2\text{C}_6\text{H}_3)_n\text{B}_{12}\text{H}_{12-n}]^{2-}$  (X = Cl, Br; n = 1–3); synthesis,  $^{11}\text{B}$ ,  $^{13}\text{C}$  and  $^1\text{H}$  NMR spectra of (R. von Bismarck, W. Preetz), (418) 147

- Trichlorophenylhydododecarborates  $[(m,p,m\text{-Cl}_3\text{C}_6\text{H}_2)_n\text{B}_{12}\text{H}_{12-n}]^{2-}$  ( $n = 1, 2$ ); synthesis,  $^{11}\text{B}$ ,  $^{13}\text{C}$ , and  $^1\text{H}$  NMR spectra of; influence of phenyl substituents on the chemical shift of the  $^{11}\text{B}$  nuclei (R. von Bismarck, W. Preetz), (418) 157
- Dicarbollyl complexes of phosphor and arsenic, X-ray diffraction study on  $\text{ClAs}[\text{C}_2\text{Me}_2\text{B}_9\text{H}_9]$  (P. Jutzi, D. Wegener, M. Hursthouse), (418) 277
- Heterodinuclear group 6 metallacarboranes; synthesis and characterization of [*closo*-3,3,3-(CO)<sub>3</sub>-3-SnPh<sub>3</sub>-3,1,2-MC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>-</sup> (M = Cr, Mo, W) (J. Kim, Y. Do, Y.S. Sohn, C.B. Knobler, M.F. Hawthorne), (418) C1
- Carbonylation of R<sub>2</sub>Bi in the presence of NaCo(CO)<sub>4</sub> and Na<sub>2</sub>Fe(CO)<sub>4</sub>: A simple synthesis of dialkyl ketones (A. Devasagayaraj, M.L.N. Rao and M. Periasamy), (421) 147
- Determination of the activation parameters of permanent allylic rearrangement in allyl- and methallylboranes and in 1,1-bis(dipropylborylmethyl)ethylene by dynamic NMR spectroscopy (Yu.N. Bubnov, M.E. Gurskii, I.D. Gridnev, A.V. Ignatenko, Yu.A. Ustyuk and V.I. Mstislavsky), (424) 127
- Unsolvated lanthanidocene hydrides and borohydrides. X-Ray crystal structure of  $[(\eta^5\text{-C}_5\text{H}_3\text{'Bu}_2)_2\text{Ln}(\mu\text{-H})_2]$  (Ln = Ce, Sm) and  $[(\eta^5\text{-C}_5\text{H}_3\text{'Bu}_2)_2\text{Sm}(\mu\text{-BH}_4)]_2$  (Y.K. Gun'ko, B.M. Bulychev, G.L. Soloveichik and V.K. Belsky), (424) 289
- Cationic titanium alkyls as alkene polymerisation catalysts: solvent and anion dependence (M. Bochmann and A.J. Jaggar), (424) C5
- Synthesis and reactivity of hydrido, halogeno, and  $\sigma$ -organyl ruthenatetraboranes: Crystal structure of [RuH(B<sub>3</sub>H<sub>8</sub>)(CO)(PPh<sub>3</sub>)<sub>2</sub>] (I.D. Burns, A.F. Hill, A.R. Thompsett, N.W. Alcock and K.S. Claire), (425) C8

## Cadmium

- Trifluoromethyl gallium, indium, and thallium compounds, syntheses of; mechanism of polar trifluoromethyl group transfer (D. Naumann, W. Strauß, W. Tyrra), (407) 1
- Oxidative alkylation of  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiR}$  (R = Cl, Me, Et, CH=CH<sub>2</sub>, Ph, OMe, N=C(H)<sup>1</sup>Bu) to  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti(Me)R}$  by group 12 organometallic compounds MMe<sub>2</sub> (G.A. Luijstra and J.H. Teuben), (420) 337
- The adduct formed between dimethylcadmium and 1,4-dioxane, 1,4-dioxanedimethylcadmium(II): a crystallographic and spectroscopic study (M.J. Almond, M.P. Beer, M.G.B. Drew and D.A. Rice), (421) 129

## Caesium

- Allyl- and benzyl-type organoalkali intermediates, preparation via organotin compounds (O. Desponds, M. Schlosser), (409) 93
- Allyl alkali metals, comparative theoretical study of (N.J.R. van Eikema Hommes, M. Bühl, P.v.R. Schleyer, Y.-D. Wu), (409) 307
- Alkali metal derivatives of sulfinimidamides, preparation and crystal structures (F. Pauer, D. Stalke), (418) 127

## Calcium

- Bis[bis(trimethylsilyl)methyl]tin(II), examination of the reaction with calcium; molecular and crystal structure of benzyl-tris[bis(trimethylsilyl)methyl]stannane (M. Westerhausen, T. Hildenbrand), (411) 1

## Cerium

- Bis(1,3-di-tert-butylcyclopentadienyl)cerium chloride and borohydride, crystal and molecular structures of; a bridging tetradentate BH<sub>4</sub>-group with two  $\mu_3$ -hydrogens:  $\mu:\eta^4\text{-}[(\mu_3\text{-H})_2\text{B}(\mu_2\text{-H}_2)]$  (E.B. Lobkovsky, Yu.K. Gun'ko, B.M. Bulychev, V.K. Belsky, G.L. Soloveichik, M.Yu. Antipin), (406) 343
- Cationic cerium compounds  $[\text{Cp}_2^*\text{Ce(L)}_2]\text{[BPh}_4]$  (L = tetrahydrofuran or tetrahydrothiophene), synthesis of; crystal structure of the tetrahydrothiophene derivative (H.J. Heeres, A. Meetsma, J.H. Teuben), (414) 351
- Unsolvated lanthanidocene hydrides and borohydrides. X-Ray crystal structure of  $[(\eta^5\text{-C}_5\text{H}_3\text{'Bu}_2)_2\text{Ln}(\mu\text{-H})_2]$  (Ln = Ce, Sm) and  $[(\eta^5\text{-C}_5\text{H}_3\text{'Bu}_2)_2\text{Sm}(\mu\text{-BH}_4)]_2$  (Y.K. Gun'ko, B.M. Bulychev, G.L. Soloveichik and V.K. Belsky), (424) 289

## Chromium

- Metal carbonyls, rate of O-atom transfer to; effect of substituents on amine *N*-oxides on (J.-K. Shen, Y.-C. Gao, Q.-Z. Shi, F. Basolo), (401) 295
- $\alpha$ -Aryl amino acids, enantioselective synthesis of; aromatic nucleophilic substitution of fluorobenzene tricarbonylchromium by chiral enolates (M. Chaari, A. Jenhi, J.-P. Lavergne, Ph. Viallefond), (401) C10
- $\beta$ - And  $\delta$ -aryl amino esters; synthesis by reaction of methyl arylacetate tricarbonylchromium complexes with halogeno  $\alpha$ -amino esters (A. Jenhi, J.-P. Lavergne, Ph. Viallefond), (401) C14
- Alkynylalkoxycarbene metal (Cr, W) complexes, conjugate addition of alcohols to (F. Camps, A. Llebaria, J.M. Moretó, S. Ricart, J.M. Viñas, J. Ros, R. Yáñez), (401) C17
- Copper, redox behaviour in new heteropentanuclear complexes prepared from functionalised phenanthroline tweezer ligands; influence of arenechromium tricarbonyl (C. Diaz, J.-P. Gisselbrecht, M. Gross, J. Suffert, R. Ziessel), (401) C54
- Aryllithiumtricarbonylchromium, aromatic nucleophilic substitution of methoxy or chloro groups by (F. Rose-Munch, O. Bellot, L. Mignon, A. Semra, F. Robert, Y. Jeannin), (402) 1
- 2,4-Di(*t*-butyl)pentadienyl anion and its Ti(II), Cr(II), and Zn(II) complexes; synthesis of (R.D. Ernst, J.W. Freeman, P.N. Swepston, D.R. Wilson), (402) 17
- ( $\eta^4$ -Norbornadiene)tetracarbonylchromium and metal VIb hexacarbonyls, chemical shift anisotropy of carbon-13 nuclei in carbonyl groups of (A. Gryff-Keller, H. Krawczyk, P. Szczeciński), (402) 77
- [CpCr(CO)<sub>2</sub>L]<sub>2</sub> complexes [L = CO, P(OMe)<sub>3</sub>], NMR determination of metal–metal bond energies in (L.Y. Goh, Y.Y. Lim), (402) 209
- o*-Fluorobenzaldehyde-chromium-tricarbonyl, a chiral complex for highly diastereoselective nucleophilic additions (A. Solladié-Cavallo, M. Bencheqroun), (403) 159
- [( $\eta^5$ -(Dimethylphosphino)cyclopentadienyl][ $\eta^7$ -(dimethylphosphino)cycloheptatrienyl]titanium, heterobimetallic compounds derived from (M. Ogasa, M.D. Rausch, R.D. Rogers), (403) 279
- Benzyl tricarbonyl chromium allyl and crotyl ethers, 2,3-Wittig rearrangement of (M. Mahmoudi, L. Pelinski, L. Maciejewski, J. Brocard), (405) 93
- Chromium–iron–sulfide Cp<sub>3</sub>Cr<sub>3</sub>( $\mu_3$ -S)<sub>4</sub>FeSPh and [Cp<sub>3</sub>Cr<sub>3</sub>( $\mu_3$ -S)<sub>4</sub>Fe]<sub>2</sub>Fe(CO)<sub>3</sub>S·C<sub>6</sub>H<sub>6</sub> clusters with tetrahedral metal frameworks, syntheses and molecular structures of (S.E. Nefedov, A.A. Pasynskii, I.L. Eremenko, E.E. Stomakhina, A.I. Yanovsky, Yu.T. Struchkov), (405) 287
- ( $\eta^6$ -Arene)tricarbonylchromium complexes, preparation of; catalysts for complexation of arenes with Cr(CO)<sub>6</sub> (M. Hudeček, Š. Toma), (406) 147
- Allyl[N,N-bis(pyrazol-1-ylmethyl)aminomethane]dicarbonylchromium(II)hexafluorophosphate, crystal structure of; symmetric orientation of the  $\eta^3$ -allyl group with respect to the pyrazole-derived tridentate ligand (K.-B. Shiu, C.-J. Chang, Y. Wang, M.-C. Cheng), (406) 363
- (Arene)chromium complexes, regioselectivity of nucleophilic additions to; conformational effects of Cr(CO)<sub>3</sub> moieties and the nature of carbanions (M. Uemura, T. Minami, Y. Shinoda, H. Nishimura, M. Shiro, Y. Hayashi), (406) 371
- [Cp<sub>2</sub>Cr<sub>2</sub>( $\mu$ -SCMe<sub>3</sub>) $(\mu_3$ -S)<sub>2</sub>]<sub>2</sub>Fe<sup>+</sup>FeI<sub>4</sub><sup>-</sup> heterometallospirane cluster and [Cp<sub>2</sub>Cr<sub>2</sub>( $\mu$ -SCMe<sub>3</sub>) $(\mu_3$ -S)<sub>2</sub>Fe]<sub>2</sub>( $\mu$ -Se)<sub>2</sub> cluster with “kayak-paddle” framework; synthesis, magnetic properties and structures of (A.A. Pasynskii, I.L. Eremenko, E.E. Stomakhina, S.E. Nefedov, O.G. Ellert, A.I. Yanovsky, Yu.T. Struchkov), (406) 383
- Chiral arene-chromium-tricarbonyl complexes; 2-step synthesis of halostachin analogues (A. Solladié-Cavallo, M. Bencheqroun), (406) C15
- Heterometallic  $n,\pi$ -complexes of transition metals based on mesitylacetonitrile (V.S. Kaganovich, M.I. Rybinskaya), (407) 215
- Chromium and tungsten thiocarbene complexes, preparation of; insertion and annulation reactions of (K.H. Dötz, V. Leue), (407) 337
- Unidentate and heterobimetallic 1,1'-bis(diphenylphosphino)ferrocene complexes, X-ray photoelectron spectroscopic differentiation of chemically distinct phosphorus environments in (H.S.O. Chan, T.S.A. Hor, L.-T. Phang, K.L. Tan), (407) 353
- Arene tricarbonylchromium moieties, diastereoselective nucleophilic additions to imines attached to (P. Bloem, D.M. David, L.A.P. Kane-Maguire, S.G. Pyne, B.W. Skelton, A.H. White), (407) C19
- M<sup>+</sup>[M'M''(CO)<sub>9</sub>L]<sup>-</sup> (M<sup>+</sup> = Na<sup>+</sup>, PPN<sup>+</sup>; M' = Cr, W; M'' = Mn, Re; L = CO, PR<sub>3</sub>), kinetic studies of the cleavage of (Y.K. Park, S.J. Kim, J.H. Kim, I.S. Han, C.H. Lee, H.S. Choi), (408) 193

- $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{M}(\text{HgX})]$  ( $\text{M} = \text{Cr, Mo, W; X} = \text{Cl, Br, I, N}_3, \text{SCN}$ ) complexes, interaction with  $\text{M}'(\text{CO})_3(\text{bipy})$  ( $\text{M}' = \text{Mo, W}$ ) fragments; trimetallic compounds containing  $\text{M}-\text{Hg}-\text{M}'-\text{X}$  arrays (J. Granijo, M.E. Vargas), (408) 357
- $(\eta^2\text{-TCNE})\text{M}(\text{CO})_5$  ( $\text{M} = \text{Cr; W; TCNE} = \text{tetracyanoethene}$ ), resonance Raman study of; evidence for an olefin-complex  $\rightarrow$  metallacyclopropane transition upon low-energy metal-to-ligand charge-transfer excitation (D.J. Stufkens, T.L. Snoeck, W. Kaim, T. Roth, B. Olbrich-Deussner), (409) 189
- Carbonylchromium, carbonylmolybdenum and carbonyltungsten complexes of the asymmetrically substituted phosphino sulfur diimides  ${}^1\text{Bu}_2\text{P}(\text{NSN})\text{PPPh}_2$  and  ${}^1\text{Bu}_2\text{P}(\text{NSN})\text{P}({}^1\text{Bu})\text{Ph}$ ;  $^1\text{H}, ^{13}\text{C}$  and  $^{31}\text{P}$  NMR spectroscopic characterisation of (M. Herberhold, S.M. Frank, B. Wrackmeyer), (410) 159
- Naphthaleneytterbium, reactions with bis(cyclopentadienyl)complexes of cobalt, nickel, chromium and vanadium; crystal structure of the triple-decker compound  $\text{CpVC}_{10}\text{H}_8\text{VCp}$  (M.N. Bochkarev, I.L. Fedushkin, H. Schumann, J. Loebel), (410) 321
- $\text{CH}_3\text{C}[\text{CH}_2\text{NC}-\text{M}(\text{CO})_x]_3$  complexes ( $\text{M} = \text{Cr, W, } x = 5; \text{M} = \text{Fe, } x = 4$ ), synthesis of; crystal structure of  $\text{CH}_3\text{C}[\text{CH}_2\text{NC}-\text{W}(\text{CO})_5]_3$  (F.E. Hahn, M. Tamm), (410) C9
- $\text{Cr}(\text{CO})_5\text{CNCCl}_3$ , reactions with primary amines; synthesis of aliphatic, aromatic, functional, and chiral isocyanides (W.P. Fehlhammer, S. Ahn, G. Beck), (411) 181
- Bi- and trinuclear cyclopentadienyl clusters of chromium with nitrene bridges (I.L. Eremenko, A.A. Pasynskii, E.A. Vas'utinskaya, A.S. Katugin, S.E. Nefedov, O.G. Ellert, V.M. Novotortsev, A.F. Shestakov, A.I. Yanovsky, Yu.T. Struchkov), (411) 193
- $(\text{CO})_{5-n}(\text{R}'_3\text{P})_n\text{M}(\text{H})\text{SiR}_3$  ( $n = 2, 3$ ), synthesis of phosphane substituted hydridosilyl complexes of chromium, molybdenum and tungsten; synthesis of  $(\text{CO})_3(\text{L-L})\text{M}(\text{H})\text{Cl}$  by conversion of  $(\text{CO})_3(\text{L-L})\text{M}(\text{NCR}')$  with  $\text{HSiCIR}_2$  ( $\text{L-L} = \text{tmida, bipy, dppe, (PPh}_3)_2$ ) (H. Piana, U. Schubert), (411) 303
- Highly substituted tricarbonyl(cyclobutabenzeno)chromium(0) derivative; synthesis, structure, and solid state  $^{13}\text{C}$  NMR spectroscopy of (H.G. Wey, P. Betz, I. Topalović, H. Butenschön), (411) 369
- Indenyl anions, mechanism of migration of the  $\text{Cr}(\text{CO})_3$  unit from the six- to the five-membered ring (A. Cecon, A. Gambaro, F. Gottardi, S. Santi), (412) 85
- $(\eta^5\text{-Pentadienyl})\text{Cr}^{II}$  complexes, preparation and reactions of; crystal structure of  $(\eta^5\text{-2,4-Me}_2\text{C}_5\text{H}_5)\text{X}(\eta^3\text{-C}_3\text{H}_5)\text{CrPMe}_3$  (P.W. Jolly, C. Krüger, U. Zakrzewski), (412) 371
- $(\eta^5\text{-Cyclopentadienyl})\text{dicarbonylnitrosylchromium}$  (cynichrodene), functionally substituted derivatives; formation and reactions of (M.D. Rausch, Y.-P. Wang), (413) 111
- ortho*- And *meta*-substituted tricarbonylchromium benzaldehydes, enantioselective reduction by bakers' yeast (S. Top, G. Jaouen, C. Baldoli, P. Del Buttero, S. Maiorana), (413) 125
- 1,2:9,10-dibenzo[2.2]paracyclophane-1,9-dienes, regioselective complexation of (O. Reiser, A. de Meijere), (413) 137
- Chromium aminocarbene complexes in organic synthesis (M.A. Schwint, J.R. Miller, L.S. Hegedus), (413) 143
- $\text{Cr}(\text{CO})_6$ , complexation with arenes; catalyst effects (M. Hudeček, V. Gajda, Š. Toma), (413) 155
- $\text{Cr}(\text{CO})_6$ , acid catalysed complexation with arenes (P. Hrnčiar, Š. Toma), (413) 161
- $(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4)\text{CH}_2(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)$ , Friedel-Crafts cynichrodenylation of; crystal structures of  $(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4)\text{CH}_2(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)\text{C}(\text{O})(\eta^5\text{-C}_5\text{H}_4)$   $\text{Cr}(\text{CO})_2(\text{NO})$  and  $(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4)\text{CH}_2(\eta^5\text{-}\{2\text{-C}(\text{O})\text{-}\}(\eta^5\text{-C}_5\text{H}_4)\text{Cr}(\text{CO})_2(\text{NO}))\text{-C}_5\text{H}_3\text{Fe}(\eta^5\text{-C}_5\text{H}_5)$  (Y.-P. Wang, J.-M. Hwu, S.-L. Wang, Y.-J. Wu), (414) 33
- Acetylstyrene- and benzoylstyrene-tricarbonylchromium, and 1-acetyl- and 1-benzoyl-2-ferrocenylethylene, preparation of (J. Huang, Y.-Z. Huang), (414) 49
- Triangular cluster  $\text{Cp}_2\text{Cr}_2(\mu\text{-SCMe}_3)(\mu_3\text{-S})_2\text{Ir}(\text{CO})(\text{PPh}_3)$  and octanuclear cluster  $\text{Cp}_2\text{Cr}_2(\mu_3\text{-S}_2)(\mu_4\text{-S})\text{Rh}_2[\text{Cp}_2\text{Cr}_2(\mu\text{-SCMe}_3)(\mu_3\text{-S})_2]_2$  with unusual triangular heterometalloornament framework; synthesis, molecular structure and magnetic properties of (A.A. Pasynskii, I.L. Eremenko, V.R. Zalmanovitch, V.V. Kavarin, B. Orazsakhatov, S.E. Nefedov, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky, Yu.T. Struchkov), (414) 55
- Complexed triphosphanes, preparation and characterisation of; crystal structure of  $\text{HP}[\text{PPh}_2\text{Cr}(\text{CO})_5]_2$  (M. Scheer, S. Gremler, E. Herrmann, P.G. Jones), (414) 337
- Haloarenetricarbonylchromium, complexes obtained after treatment with anions of  $\alpha$ -imino esters or nitriles; synthesis and conformational study of (F. Rose-Munch, K. Aniss, E. Rose, J. Vaisserman), (415) 223
- Metal complexed indoles, chemistry and synthetic utility of (U.S. Gill, R.M. Moriarty, Y.Y. Ku, I.R. Butler), (417) 313

- Tetracarbonyl-aminocarbene complexes of chromium, synthesis and aminolysis to give coordinated iminonitrides (K.H. Dötz, A. Rau), (418) 219
- Aminocarbene complexes of chromium and molybdenum derived from primary amines, preparation, structure and reactivity of (B. Denise, P. Dubost, A. Parlier, M. Rudler, H. Rudler, J.C. Daran, J. Vaissermann, F. Delgado, A.R. Arevalo, R.A. Toscano, C. Albarez), (418) 377
- Heterodinuclear group 6 metallacarboranes; synthesis and characterization of [*clos*-3,3,3-(CO)<sub>3</sub>-3-SnPh<sub>3</sub>-3,1,2-MC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>-</sup> (M = Cr, Mo, W) (J. Kim, Y. Do, Y.S. Sohn, C.B. Knobler, M.F. Hawthorne), (418) C1
- ( $\eta^5$ -Cyclopentadienyl)tricarbonylmethylchromium, photochemistry in gas matrices and polymer films at 12–298 K; infrared spectroscopic evidence for reversible carbon monoxide ejection and  $\alpha$ -H elimination from the methyl group (R.H. Hooker, K.A. Mahmoud, A.J. Rest and H.G. Alt), (419) 101
- Group VI metal carbonyls, ( $\eta^6$ -C<sub>6</sub>H<sub>6</sub>)Cr(CO)<sub>3</sub>, ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Mn(CO)<sub>3</sub>, ( $\eta^5$ -CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Mn(CO)<sub>3</sub> and ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Re(CO)<sub>3</sub>, photolysis in Nujol at 77 K (T.E. Bitterwolf, K.A. Lott, A.J. Rest and J. Mascetti), (419) 113
- [ $\eta^5$ -(C<sub>5</sub>H<sub>4</sub>COOCH<sub>3</sub>)]Cr(NO)<sub>2</sub>L (L = Cl, I, CH<sub>3</sub>), syntheses and spectra of; crystal structures of [ $\eta^5$ -(C<sub>5</sub>H<sub>4</sub>COOCH<sub>3</sub>)]Cr(NO)<sub>2</sub>L (L = Cl, I) (Y. P. Wang, Y.-H. Yang, S.-L. Wang and F.-L. Liao), (419) 325
- Tricarbonyl(benzene)chromium and ferrocene containing unsaturated amides and nitriles, preparation of (J. Huang, H.-X. Jiang and Y.-Z. Huang), (419) 337
- Enaminoketene and arenetricarbonyl nitrogen ylid chromium complexes, successive formation during intramolecular alkyne insertions into aminocarbene complexes (E. Chelain, A. Parlier, H. Rudler, J.C. Daran and J. Vaissermann), (419) C5
- Déprotonation des allylbénzènes chrometricarbonyle en milieu basique. Influence du système basique et du groupement chrometricarbonyle sur la régiosélectivité de l'addition des dérivés carbonylés et sur la nature des produits isolés (M.-C. Séchéchal-Tocquer, D. Séchéchal, J.-Y. Le Bihan, D. Gentric and B. Caro), (420) 185
- Heteronucleare Komplexverbindungen mit Metall–Metall-Bindungen. VI. Kleine heteronucleare Cluster mit Kupfer und HCr(CO)<sub>5</sub>-Fragmenten. Die Strukturen von [(N<sub>2</sub>)Cu(μ<sub>3</sub>-H)Cr<sub>2</sub>(CO)<sub>10</sub>]<sub>2</sub> (N<sub>2</sub> = 2 NH<sub>3</sub>, bpy, 1,2-Diaminobenzol), [{(py)Cu(μ<sub>3</sub>-H)Cr(CO)<sub>5</sub>}<sub>2</sub>] und [Cu(PPh<sub>3</sub>)<sub>3</sub>][Cr<sub>2</sub>(μ-H)(CO)<sub>10</sub>] sowie die Struktur von [Zn(NH<sub>3</sub>)<sub>4</sub>][Cr<sub>2</sub>(μ-H)(CO)<sub>10</sub>]<sub>2</sub> (P. Klüfers and U. Wilhelm), (421) 39
- Kohlenwasserstoffverbrückte Komplexe. XXIII. Heterobimetallische Komplexe mit Ferrocenyl- und (Ph<sub>3</sub>P)(OC)(Cp)Fe(CO)CH<sub>2</sub>-Gruppen; Darstellung und Struktur von ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe[ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)Re(CO)<sub>5</sub>], (PPh<sub>3</sub>)(OC)(Cp)FeC(O)CH<sub>2</sub>( $\eta^6$ -C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub> und [ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)CH<sub>2</sub>( $\eta^6$ -C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub>] (M = Cr, Mo) (J. Breimair, M. Wieser, B. Wagner, K. Polborn and W. Beck), (421) 55
- Carbon-13 spin-lattice relaxation in organometallic complexes (H. Schumann), (421) C7
- Application of scalar <sup>13</sup>C, <sup>19</sup>F spin–spin couplings between carbonyl carbons and aromatic fluorine to investigation of conformation of tricarbonylchromium complexes of fluorobenzenes (P. Szczeciński), (423) 23
- Electrochemical generation of 17-electron cation radicals from arene–M(CO)<sub>3</sub> (M = Cr, Mo, W) and thiophene–Cr(CO)<sub>3</sub> complexes in MeCN: Conventional cyclic voltammetric studies with digital simulation and microelectrode voltammetry in the absence of supporting electrolyte (F. Rourke, R. Gash and J.A. Crayston), (423) 223
- Scalar <sup>13</sup>C–<sup>19</sup>F spin–spin couplings between carbonyl carbons and aromatic fluorines in tricarbonylchromium complexes of biphenyl derivatives (P. Szczeciński and K. Wiśniewski), (423) C13
- Metal-side chain interaction in tricarbonylchromium complexes of phenylacetylenes studied by <sup>13</sup>C NMR and IR spectroscopy (J. Szewczyk and A. Gryff-Keller), (424) 41
- Organometallic chemistry of diphosphazanes. IV. Reactions of the cyclodiphosphazane, *cis*-[<sup>1</sup>BuNP(OPh)]<sub>2</sub> with M(CO)<sub>6</sub> (M = Cr, Mo, W), rhodium(I), palladium(II) and platinum(II) derivatives (M.S. Balakrishna and S.S. Krishnamurthy), (424) 243
- Synthesis, characterizations, and structures of group IV metal–chromium complexes bridged by the benzyloxide group (H.-M. Gau, C.-T. Chen and C.-C. Schei), (424) 307
- Structures of (*N,N*-dicyclohexyl)benzenesulfenamide and its pentacarbonylchromium(0) complex via the sulfur atom (M.L. Rodríguez, I. Brito, C. Diaz, G. González and V. Manríquez), (425) 49

## Cobalt

- Transition metal complexes as electrochemical markers for steroids (D. Osella, E. Stein, G. Jaouen, P. Zanello), (401) 37
- ( $\eta^6$ -Phenyl)Cr(CO)<sub>3</sub>- and ( $\eta^6$ -phenyl)Co<sub>4</sub>(CO)<sub>9</sub>-substituted metallocenylcarbinols of the iron subgroup; synthesis, structure and hydrogen bonding in (E.S. Shubina, L.M. Epstein, Yu.L. Slovokhotov, A.V. Mironov, Yu.T. Struchkov, V.S. Kaganovich, A.Z. Kreindlin, M.I. Rybinskaya), (401) 155
- $\eta^1$ - And  $\eta^3$ -cobalt(I) allyl complexes, reactions with bis(phosphine) ligands; crystal structure of {( $\eta^3$ -C<sub>3</sub>H<sub>5</sub>)Co(CO)<sub>2</sub>}<sub>2</sub>( $\mu$ -dppe-PP) (C. Loubser, H.M. Roos, S. Lotz), (402) 393
- Cobalt, ruthenium, and rhodium complexes as catalysts in the reaction of terminal olefins with a hydrosilane and with carbon monoxide (Y. Seki, K. Kawamoto, N. Chatani, A. Hidaka, N. Sonoda, K. Ohe, Y. Kawasaki, S. Murai), (403) 73
- Bis[tris(*m*-(sodium sulfonato)phenyl)phosphine] hexacarbonyl dicobalt, Co<sub>2</sub>(CO)<sub>6</sub>(P(*m*-C<sub>6</sub>H<sub>4</sub>SO<sub>3</sub>Na)<sub>3</sub>)<sub>2</sub>, in a supported aqueous phase for the hydroformylation of 1-hexene (I. Guo, B.E. Hanson, I. Tóth, M.E. Davis), (403) 221
- C-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325
- Co<sub>2</sub>(CO)<sub>8</sub> and NaCo(CO)<sub>4</sub> reagents, synthesis by the reduction of CoCl<sub>2</sub> or CoBr<sub>2</sub> under carbon monoxide at atmospheric pressure (A. Devasagayaraj, S.A. Rao, M. Periasamy), (403) 387
- Stable titanium-cobalt complex with non-bridged Ti-Co bond, synthesis and characterisation of (D. Selent, R. Beckhaus, T. Bartik), (405) C15
- Bisalkynyltitanocene complexes as bidentate chelating ligands (H. Lang, L. Zsolnai), (406) C5
- Co<sub>4</sub>(CO)<sub>12</sub>, reactions with polydentate ligands; fragmentation kinetic studies of substituted derivatives (C. Moreno, Ma.J. Macazaga, S. Delgado), (407) 125
- Cobalt-catalysed conversion of methyl formate into acetic acid (G. Jenner, E.M. Nahmed), (407) 135
- Heterometallic *n*, $\pi$ -complexes of transition metals based on mesitylacetonitrile (V.S. Kaganovich, M.I. Rybinskaya), (407) 215
- Co-C bond lengths in vitamin B<sub>12</sub> model compounds, evidence of steric influences on (S. Geremia, M. Mari, L. Randaccio, E. Zangrando), (408) 95
- Iron, cobalt and rhodium complexes of the optically active diolefins (+)-nopadiene and its derivatives; crystal structure of C<sub>5</sub>Me<sub>5</sub>Rh(nopadiene) (A. Salzer, H. Schmalle, R. Stauber, S. Streiff), (408) 403
- [{Co<sub>2</sub>(CO)<sub>6</sub>PhC<sub>2</sub>}<sub>2</sub>] acetylene-linked dimeric cluster, synthesis and structural characterisation of (B.F.G. Johnson, J. Lewis, P.R. Raithby, D.A. Wilkinson), (408) C9
- Homogeneous metal complex catalysts in the synthesis of cyclobutane and cyclopentane compounds (U.M. Dzhemilev, R.I. Khusnutdinov, G.A. Tolstikov), (409) 15
- Naphthaleneytterbium, reactions with bis(cyclopentadienyl)complexes of cobalt, nickel, chromium and vanadium; crystal structure of the triple-decker compound CpVC<sub>10</sub>H<sub>8</sub>VCp (M.N. Bochkarev, I.L. Fedushkin, H. Schumann, J. Loebel), (410) 321
- Octacarbonyldicobalt, reductive dehalogenation of chloro(alkynyl)phosphines by (H. Lang, M. Leise, L. Zsolnai), (410) 379
- Bi- and trinuclear cyclopentadienyl clusters of chromium with nitrene bridges (I.L. Eremenko, A.A. Pasynskii, E.A. Vas'utinskaya, A.S. Katugin, S.E. Nefedov, O.G. Ellert, V.M. Novotortsev, A.F. Shestakov, A.I. Yanovsky, Yu.T. Struchkov), (411) 193
- Actinide complexes with tripod ligands, synthesis and structure of [CpCo{P(O)(OEt)<sub>2</sub>}<sub>3</sub>]UCl<sub>3</sub>(THF) (M. Wedler, J.W. Gilje, M. Noltemeyer, F.T. Edelmann), (411) 271
- (OC)<sub>2</sub>LCo( $\eta^2$ -PR<sub>2</sub><sup>1</sup>-PR<sub>2</sub><sup>2</sup>); synthesis, properties and structures of this type of ( $\eta^2$ -phosphanediyi-phosphinato)cobalt complexes (E. Lindner, T. Funk, W. Hiller, R. Fawzi), (411) 491
- Aldol and retroaldol reactions, catalysis by cobalt compounds (Y. Tencer, M. Michman, I. Goldenfeld), (412) 203
- $\sigma$ -Bonded alkynylcobalt(III) complexes of 1,3-bis(diacetylmonoximeimino)propane, synthesis and crystal structure of (B. Giese, M. Zehnder, M. Neuburger, F. Trach), (412) 415
- Thiophene complexes of cobalt (R. Albrecht, E. Weiss), (413) 355
- Four-membered cobalta heterocycles from C<sub>5</sub>H<sub>5</sub>Co(CNR)(PMe<sub>3</sub>) and isocyanates or isothiocyanates, synthesis and reactivity of; unusual fragmentation of the chiral isocyanide ligand (*S*)-CNCH(Me)Ph (H. Werner, B. Strecker), (413) 379
- Acetylenes and diazoketones in organic synthesis (F. Serratosa), (413) 445
- Pauson-Khand reaction, application to the synthesis of pentalenic acid (E.G. Rowley, N.E. Schore), (413) C5

- Platinum–indium and cobalt–gallium complexes, synthesis of; crystal structure of  $[\text{Cy}_2\text{P}(\text{CH}_2\text{CH}_2)\text{PCy}_2]\text{Pt}(\text{InR}_2)(\text{R})$  ( $\text{Cy}$  = cyclo- $\text{C}_6\text{H}_{11}$ ) (R.A. Fischer, J. Behm), (413) C10
- Organometallic cobaloximes containing an equatorial diphenylglyoximato(-1) ligand; comparison between their properties and those of other  $\text{B}_{12}$  model compounds; crystal structure of *trans*- $[\text{Co}(\text{dpgH})_2(\text{CH}_3\text{Xpyridine})]$  (C. López, S. Alvarez, M. Font-Bardía, X. Solans), (414) 245
- ( $\eta^5\text{-C}_5\text{H}_4\text{CH}_3$ )<sub>4</sub> $\text{Co}_2\text{Fe}_2(\mu_3\text{-NO})_2$  and trinuclear methylcyclopentadienyl hetero-metal clusters with  $\mu_3\text{-NH}$  and  $\mu_3\text{-NO}$  ligands, synthesis and structures of (J. Müller, I. Sonn, T. Akhnoukh), (414) 381
- Enantioselective Co-catalysed homo Diels–Alder reaction of norbornadiene with acetylenes (H. Brunner, F. Prester), (414) 401
- Substituted  $\eta^4$ -cyclooctadiene-1,5- $\eta^5$ -cyclopentadienyl-cobalt complexes, preparation by reaction of cobaltocenes with alkene cobaltates(1–) (H.-U. Jürgens, J. Alm), (414) C61
- Metal carbonyl dimers, new mechanism for substitution reactions of (B.F.G. Johnson), (415) 109
- 20-Electron halfsandwich compounds, sterical stabilization of; crystal structure of  $[(\text{C}_5\text{H}_5)\text{Co}(\text{P}(\text{OMe})_2\text{O})_3\text{Ni}(\text{C}_5\text{Ph}_5)]$  (W. Klaui, M. Huhn, R. Herbst-Irmer), (415) 133
- Carbomethoxycyclopentadienylcarbonylcobalt(I), carbonyl substitution reactions with bidentate phosphines; kinetic and mechanistic study (C. Moreno, Ma.J. Macazaga, S. Delgado), (415) 271
- Cobalt(O) and cobalt(I) complexes, reactivity with diynes towards  $\text{C}\equiv\text{C}$ ,  $\text{C}\equiv\text{N}$ ,  $\text{C}\equiv\text{C}$  bonds; X-ray structure of a cyclopentadienylcobaltacyclopentadiene complex (Z. Zhou, L.P. Battaglia, G.P. Chiusoli, M. Costa, M. Nardelli, C. Pelizzi, G. Predieri), (417) 51
- Cobalt-catalyzed hydroformylation, activation of molecular hydrogen (P. Pino, A. Major, F. Spindler, R. Tannenbaum, G. Bor, I.T. Horváth), (417) 65
- Hydroformylation of olefins in the presence of dicobalt octacarbonyl; some considerations (F. Piacenti, M. Bianchi, P. Frediani, G. Menchi, U. Matteoli), (417) 77
- Hydrido tetracarbonyl cobalt kinetics of the ring-opening carbonylation of ethyloxirane with (J. Kreisz, F. Ungváry, A. Sisak, L. Markó), (417) 89
- $\alpha$ -Amidocarboxylic acids, amidocarbonylation as a route to (J.J. Lin, J.F. Knifton), (417) 99
- The use of  $^2\text{H}$  NMR in the elucidation of the catalytic pathway of the hydroformylation reaction (G. Uccello-Barretta, R. Lazzaroni, R. Settambolo, P. Salvadori), (417) 111
- Dihydrogen activation by  $\text{Co}_2\text{Rh}_2(\text{CO})_{12}$  disproportionation and a second order cluster fragmentation, kinetics of (M. Garland, P. Pino), (417) 121
- $\text{M}_4(\text{CO})_{11}\text{L}$  ( $\text{M}$  = Co, Ir;  $\text{L}$  =  $\text{PPh}_3$ ,  $\text{P}^n\text{Bu}_3$ ,  $\text{P}(\text{OMe})_3$ ,  $\text{P}(\text{OEt})_3$ ); kinetics and mechanisms of CO substitution with L in the presence of  $\text{Me}_3\text{NO}$  (J.-Q. Wang, J.-K. Shen, Y.-C. Gao, Q.-Z. Shi, F. Basolo), (417) 131
- Enantioselective hydrogenation catalysts, some general regularities in the action of (E.I. Klabunovskii), (417) 181
- Alkynyl substituted chlorosilanes as cluster units (H. Lang, U. Lay, L. Zsolnai), (417) 377
- Cobalt-catalysed dimerization of styrenes under syngas (M. Fontaine, A.J. Hubert, A.F. Noels, A. Demonceau, P. Teyssié), (417) C28
- (Dimethoxycarbonyl)tricarbonyl cobaltates, reductive elimination of dimethylcarbonate from; isolation and crystal structures of  $\text{Cs}[\text{Co}(\text{COOCH}_3)_2(\text{CO})_3]$  and  $\text{K}[\text{dibenzo-18-crown-6}][\text{Co}(\text{COOCH}_3)_2(\text{CO})_3]$  (G. Fachinetti, T. Funaioli, D. Masi, C. Mealli), (417) C32
- Bis(fulvalene)diiron and bis(fulvalene)iron–cobalt monohexafluorophosphates, crystallographic comparison of the cation (P. Brüggeller, P. Jaitner, H. Schottenberger, K.E. Schwarzhans), (417) C53
- Pyridinyloxazoline cobalt(I) complexes as catalysts, enantioselective hydrosilyierung of acetophenone (H. Brunner, K. Amberger), (417) C63
- Phenylethynyl chlorosilanes, complex-bonded; reactivity of (U. Lay, H. Lang), (418) 79
- $\text{PhCCo}_3(\text{CO})_9$ , bis(dimethylphosphino)ethane substitution in; synthesis and X-ray structure of the phosphine-bridged cluster  $\text{PhCCo}_3(\text{CO})_7(\text{dmpe})$  (M.-J. Don, M.G. Richmond, W.H. Watson, M. Krawiec, R.P. Kashyap), (418) 231
- $[\text{RuCo}_3\text{H}(\text{CO})_{12}]$ , disordered crystal structure of (J. Pursiainen, P. Hirva and T.A. Pakkanen), (419) 193
- Alkylcobalt carbonyls, ligand-assisted carbonylation–decarbonylation reactions of (J. Somlyai-Haász, F. Haász, V. Galamb, A. Benedetti, C. Zucchi, G. Pályi, T. Krümmling, B. Happ and T. Bartik), (419) 205
- Some chiral alkyne-cobalt complexes (J.A. Dunn and P.L. Pauson), (419) 383
- Cluster chemistry. LXXIII. Preparation and X-ray structure of the hexanuclear cobalt-ruthenium cluster,  $\text{CoRu}_5(\mu_4\text{-PPh})\mu_4\text{-C}_2\text{Ph}(\mu\text{-PPh}_2)(\text{CO})_{12}(\eta\text{-C}_5\text{H}_5)$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (420) 95

- Reactivity of Main-Group-transition-metal bonds. IX. The kinetics of iodination of compounds containing two or more tin-transition-metal bonds (J.R. Chipperfield, S. Clark, D.E. Webster and H. Yusof), (421) 205**
- Alkoxy- and aryloxycarbonylcobalt carbonyls. IV. Preparation and structure of derivatives with tertiary pnictogen ligands (T. Bartik, T. Krümmeling, C. Krüger, L. Markó, R. Boese, G. Schmid, P. Vivarelli and G. Pályi), (421) 323**
- Reactivity of phosphinoalkynes toward  $[HFe_3(CO)_6(C=CH_2)]^-$  (E. Louattani, J. Suades and R. Mathieu), (421) 335**
- Carbon-13 spin-lattice relaxation in organometallic complexes (H. Schumann), (421) C7**
- Substitution reactions of dinuclear alkenyl-bridged iron-cobalt complexes with phosphines. Crystal structures of  $[(CO)_3Fe-Co(CO)_2(PMe_2Ph)(\mu-C(CO_2Me)=C(CO_2Me)H)$  and  $[(CO)_2(PMe_2Ph)Fe-Co(CO)_2(PMe_2Ph)(\mu-C(CO_2Me)=C(CO_2Me)H)$  (I. Moldes, J. Ros, R. Mathieu, X. Solans and M. Font-Bardía), (423) 65**
- Formation of metallacycles from terminal diynes with geminal methyl groups  $\alpha$  to the triple bonds. Synthesis and crystal structure of  $Co_2(CO)_5[(HC\equiv CCMe_2)_2NMe]$ , an intermediate in cobalt-catalyzed organic syntheses (G. Predieri, A. Tiripicchio, M.T. Camellini, M. Costa and E. Sappa), (423) 129**
- Reactions of dithioleenes with diazo compounds: cycloaddition and dissociation of alkylidene groups (M. Sakurada, M. Kajitani, K. Dohki, T. Akiyama and A. Sugimori), (423) 141**
- Novel class of transition metal coordination compounds with macrocyclic organosiloxanolate ligands; their synthesis and crystal structure (V.A. Igonin, O.I. Shchegolikhina, S.V. Lindeman, M.M. Levitsky, Yu.T. Struchkov and A.A. Zhdanov), (423) 351**
- $\mu$ -[3-4- $\eta$ -(1-Alken-3-yne)]hexacarbonylcobalt complexes: radical cyclocondensation mediated by manganese(III) acetate (G.G. Melikyan, O. Vostrowsky, W. Bauer and H.J. Bestmann), (423) C24**
- Propargyl complexes of ruthenium (C.E. Shuchart, R.R. Willis and A. Wojcicki), (424) 185**
- Heteronucleare Komplexverbindungen mit Metall-Metall-Bindungen. VII. Umsetzungen von  $(NH_3)_3CuCo(CO)_4$  mit Methylendiaminen. Die Strukturen von  $[(NH_3)\eta^2,\mu_2-Me_3triaz]Cu_2(\mu-Co(CO)_4)\{Co(CO)_4\}$  ( $Me_3triaz = 1,3,5$ -Trimethyl-1,3,5-triaza-cyclohexan),  $[(HN(CH_2NMe_2)_2)_2CuCo(CO)_4]$ ,  $[(^nPrN(CH_2NMe_2)_2)_2CuCo(CO)_4]$ ,  $[(Me_2NH)CuCo(CO)_4]_2$  und  $[(\mu-N : N' - hmt)Cu_2(Co(CO)_4)_2]_{n/n'}THF$  ( $hmt =$  Hexamethylentetramin) (R. Fuchs und P. Klüfers), (424) 353**
- Evidence of steric influences on the Co-C bond lengths in vitamin  $B_{12}$  model compounds. II. Crystal structures of two adamantlycobaloximes with the neutral ligands  $H_2O$  and  $PPh_2Et$  (S. Géremia, L. Randaccio, E. Zangrando and L. Antolini), (425) 131**
- $[(C_5Me_4R)Co(As_2)]$ -Molekülbausteine (O.J. Scherer, K. Pfeiffer, G. Heckmann and G. Wolmershäuser), (425) 141**
- Cobalt isocyanide complexes (I. Beaumont and A.H. Wright), (425) C11**

## Copper

- Copper, redox behaviour in new heteropentanuclear complexes prepared from functionalised phenanthroline tweezer ligands; influence of arenechromium tricarbonyl (C. Diaz, J.-P. Gisselbrecht, M. Gross, J. Suffert, R. Ziessel), (401) C54**
- Phenylcoppermagnesium and methylcoppermagnesium reagents,  $^1H$  and  $^{13}C$  NMR spectra of (S.K. Nahar, H.P. Nur, M.T. Rahman, M. Nilsson, T. Olsson), (408) 261**
- Group IB metal heteronuclear cluster compounds in the solid state, high-resolution  $^{31}P$ - $\{^1H\}$  nuclear magnetic resonance studies of (S.S.D. Brown, I.D. Salter, D.J. Smith, N.J. Clayden, C.M. Dobson), (408) 439**
- Homogeneous metal complex catalysts in the synthesis of cyclobutane and cyclopentane compounds (U.M. Dzhemilev, R.I. Khusnutdinov, G.A. Tolstikov), (409) 15**
- Bis(alkynyl) titanocene complexes of  $FeCl_2$  and  $CuCl$  (H. Lang, M. Herres, L. Zsolnai, W. Imhof), (409) C7**
- 1,5,9,13-Tetraselenacyclohexadecane selenium coronand; syntheses,  $^{77}Se$  CP-MAS solid state NMR spectra and crystal structures of adducts with copper(I) trifluoromethanesulfonate and mercury(II) cyanide (R.J. Batchelor, F.W.B. Einstein, I.D. Gay, J.-H. Gu, B.M. Pinto), (411) 147**
- Binuclear transition metal complexes of  $Cu^I$ ,  $Mo^0$ , and  $Re^I$  with 2,2'-bipyrimidine (P.S. Braterman, J.-I. Song, S. Kohlmann, C. Vogler, W. Kaim), (411) 207**

- Three-membered metallacycle  $\text{Cp}_3\text{Rh}_3(\mu\text{-CO})_3$ , cluster and benzenoid system in reactions with 12-electron particles  $\text{PPh}_3\text{M}^+$  ( $\text{M} = \text{Cu, Ag, Au}$ ) and  $\text{CpM}^+$  ( $\text{M} = \text{Fe, Ru}$ ) (M.I. Rybinskaya, A.R. Kudinov, D.V. Muratov, N.P. Gambaryan, I.V. Stankevich, A.L. Chistyakov), (413) 419
- Enantioselective hydrogenation catalysts, some general regularities in the action of (E.I. Klabunovskii), (417) 181
- Alkyocuprates, reaction with 1-bromo-1,2-dienes as a regio- and stereo-selective route to acetylenic or allenic compounds (C. Polizzi, C. Consoloni, L. Lardicci, A.M. Caporusso), (417) 289
- Palladium-catalyzed reactions of 2-alkynyl caronates with terminal acetylenes, synthesis of 1,2-dien-4-ynes (T. Mandai, H. Murayama, T. Nakata, H. Yamaoki, M. Ogawa, M. Kawada, J. Tsuji), (417) 305
- Copper(I) halogenides, complexes with 3,3,6,6-tetramethyl-1-thia-4-cyclo-heptines; synthesis and structure of (F. Olbrich, G. Schmidt, U. Behrens, E. Weiss), (418) 421
- A comparative study of 1,7-octadiene and 1,7-octadiyne complexes of copper(I) chloride: preparation and molecular structures (M. Håkansson, K. Wettström and S. Jagner), (421) 347
- Conjugate addition of organocuprates to  $\gamma$ -methyl- $\delta$ -oxy- $\alpha,\beta$ -enones. Influence of the alkoxy substituent on the diastereoselection (G. Courtemanche, A. Alexakis, J. Vaissermann and J.-F. Normant), (423) 281
- Novel class of transition metal coordination compounds with macrocyclic organosiloxanolate ligands; their synthesis and crystal structure (V.A. Igonin, O.I. Shchegolikhina, S.V. Lindeman, M.M. Levitsky, Yu.T. Struchkov and A.A. Zhdanov), (423) 351
- Heteronucleare Komplexverbindungen mit Metall–Metall-Bindungen. VII. Umsetzungen von  $(\text{NH}_3)_x\text{CuCo}(\text{CO})_4$  mit Methylen diaminen. Die Strukturen von  $[(\text{NH}_3)(\eta^2,\mu_2\text{-Me}_3\text{triaz})\text{Cu}_2\{\mu\text{-Co}(\text{CO})_4\}\{\text{Co}(\text{CO})_4\}]$  ( $\text{Me}_3\text{triaz} = 1,3,5\text{-Trimethyl-1,3,5-triazacyclohexan}$ ),  $[(\text{HN}(\text{CH}_2\text{NMe}_2)_2)\text{-CuCo}(\text{CO})_4]$ ,  $[{}^n\text{PrN}(\text{CH}_2\text{NMe}_2)_2]\text{CuCo}(\text{CO})_4$ ,  $[(\text{Me}_2\text{NH})\text{CuCo}(\text{CO})_4]_2$  und  $[(\mu\text{-N} : N'\text{-hmt})\text{Cu}_2\{\text{Co}(\text{CO})_4\}_2]_{n/2}\cdot\text{THF}$  (hmt = Hexamethylenetetramin) (R. Fuchs und P. Klüfers), (424) 353
- Ligand-stabilized copper(I) hexafluoroacetylacetone complexes: NMR spectroscopy and the nature of the copper–alkene bond (T.H. Baum, C.E. Larson and G. May), (425) 189
- The reaction of arylcoppermagnesium and other organometallic reagents with phthalic anhydride (M.T. Rahman and S.K. Nahar), (425) 201

## Dysprosium

- Organo rare earth hydrides, synthesis and reactivity of (C. Ye, C. Qian, X. Yang), (407) 329

## Erbium

- Organo rare earth hydrides, synthesis and reactivity of (C. Ye, C. Qian, X. Yang), (407) 329
- Triindenyllanthanide complexes  $(\eta^5\text{-C}_9\text{H}_7)_3\text{Ln}\cdot\text{OC}_4\text{H}_8$  ( $\text{Ln} = \text{Nd, Gd, Er}$ ), syntheses and crystal structures of (J. Xia, Z. Jin, G. Lin, W. Chen), (408) 173
- Organolanthanide hydrides,  $[\text{O}(\text{CH}_2\text{CH}_2\text{C}_5\text{H}_4)_2\text{LnH}]_2$  ( $\text{Ln} = \text{Gd, Er, Yb, Lu, Y}$ ); synthesis and reactivity towards alkenes, alkynes and organic halides (Z. Xie, C. Qian, Y. Huang), (412) 61

## Gadolinium

- Tricyclopentadienyllanthanides/sodium hydride, reaction with organic halides (C. Qian, D. Zhu, Y. Gu), (401) 23
- Bis(t-butylcyclopentadienyl)lanthanoid chloride complexes, syntheses of; crystal structures of bis(t-butylcyclopentadienyl)chlorobistetrahydrofuran)praseodymium and bis(t-butylcyclopentadienyl)-chlorotetrahydrofuranterbium (Q. Shen, M. Qi, J. Guan, Y. Lin), (406) 353
- Triindenyllanthanide complexes  $(\eta^5\text{-C}_9\text{H}_7)_3\text{Ln}\cdot\text{OC}_4\text{H}_8$  ( $\text{Ln} = \text{Nd, Gd, Er}$ ), syntheses and crystal structures of (J. Xia, Z. Jin, G. Lin, W. Chen), (408) 173
- Organolanthanide hydrides,  $[\text{O}(\text{CH}_2\text{CH}_2\text{C}_5\text{H}_4)_2\text{LnH}]_2$  ( $\text{Ln} = \text{Gd, Er, Yb, Lu, Y}$ ); synthesis and reactivity towards alkenes, alkynes and organic halides (Z. Xie, C. Qian, Y. Huang), (412) 61
- Methylbis(t-butyl-cyclopentadienyl)neodymium and -gadolinium, preparation and molecular structure of (Q. Shen, Y. Cheng and Y. Lin), (419) 293

## Gallium

- Metalloporphyrins with metal–metal  $\sigma$ -bonds; synthesis, spectroscopic characterization, and electrochemistry of (P)MRe(CO)<sub>5</sub> where P is the dianion of octaethylporphyrin (OEP) or tetraphenylporphyrin (TPP) and M = Al, Ga, In or Tl (R. Guilard, A. Zrineh, A. Tabard, L. Courthaudon, B. Han, M. Ferhat, K.M. Kadish), (401) 227

- Trifluoromethyl gallium, indium, and thallium compounds, syntheses of; mechanism of polar trifluoromethyl group transfer (D. Naumann, W. Strauß, W. Tyrra), (407) 1
- Gallium-arsenic compounds containing a four-membered Ga–As–Ga–Cl or Ga–As–Ga–As ring, synthesis and characterisation of; crystal structures of  $(\text{Me}_3\text{SiCH}_2)_2\text{GaAs}(\text{SiMe}_3)_2\text{Ga}(\text{CH}_2\text{SiMe}_3)_2\text{Cl}$  and  $[(\text{Me}_3\text{SiCH}_2)_2\text{GaAs}(\text{SiMe}_3)_2]_2$  (R.L. Wells, J.W. Pasterczyk, A.T. McPhail, J.D. Johansen, A. Alvanipour), (407) 17
- Platinum–indium and cobalt–gallium complexes, synthesis of; crystal structure of  $[\text{Cy}_2\text{P}(\text{CH}_2\text{CH}_2)\text{PCy}_2]\text{Pt}(\text{InR}_2)(\text{R})$  ( $\text{Cy} = \text{cyclo-C}_6\text{H}_{11}$ ) (R.A. Fischer, J. Behm), (413) C10
- Group IIIB organometallic peroxides, synthesis and reactivity of (Yu.A. Alexandrov, N.V. Chikinova), (418) 1
- Mono- and di-methyl gallium tetraaza macrocycles  $[\text{MeGa}(\text{C}_{22}\text{H}_{22}\text{N}_4)]$  and  $[\text{Me}_2\text{Ga}(\text{C}_{22}\text{H}_{23}\text{N}_4)]$ , synthesis of; molecular structure of the dimethyl derivative (N.W. Alcock, N.C. Blacker, M.G.H. Wallbridge and J. Barker), (419) C23
- Structural characterization of a dialkylgallium cation: X-ray crystal structure of  $[\text{Me}_2\text{Ga}(\text{'BuNH}_2)_2]\text{Br}$  (D.A. Atwood, R.A. Jones, A.H. Cowley, S.G. Bott and J.L. Atwood), (425) C1

## Germanium

- Group IVB organometallic radicals, regioselective addition to substituted benzof[2,1,3]thiadiazoles; an ESR investigation (A. Alberti, M. Benaglia, D.F. Dal Monte, M. Lucarini, G.F. Pedulli), (401) 13
- 3-Thia-7,7-dimethyl-7-germabicyclo[3.3.0]oct-1(5)-ene and its selenium analogue, synthesis and reactivity of (P. Mazerolles, C. Laurent), (402) 35
- Dimethylgermathione,  $\text{Me}_2\text{Ge=S}$ ; study by vacuum pyrolysis-MS and low-temperature matrix IR spectroscopy (V.N. Khabashesku, S.E. Boganov, P.S. Zuev, O.M. Nefedov, J. Tamás, A. Gömöry, I. Besenyei), (402) 161
- 1-Halo- and 1-organoxygermatrane, synthesis of (M. Nasim, L.I. Livantsova, G.S. Zaitseva, J. Lorberth), (403) 85
- Difluorenyl- and tert-butylfluorenyl(fluorenylidene)germenes; synthesis, stabilisation and reactivity of (G. Anselme, J. Escudié, C. Couret, J. Satgé), (403) 93
- 2,3-Bis(bromomethyl)-1,4-dibromo-2-butene, reduction in the presence of organosilicon and organogeranium dihalides (P. Mazerolles, C. Laurent), (406) 119
- Bis(dimethylgermyl)alkaneruthenium tetracarbonyls, reactions with unsaturated compounds (J. Barrau, N. Ben Hamida, J. Satgé), (406) 123
- spiro*-Germanium compounds, synthesis by 1,4-cycloaddition of cyclic diaminogermyle to 1,3-butadienes and hetero-1,3-butadienes (G. Billeb, K. Bootz, W.P. Neumann, G. Steinhoff), (406) 303
- Bis(trimethylsilyl)- and bis(trimethylgermyl)carbodiimide, electron diffraction study of the gas-phase structure of (A. Hammel, H.V. Volden, A. Haaland, J. Weidlein, R. Reischmann), (408) 35
- (O–Ge)-chelate 1-(dimethylchlorogermylmethyl)pyrrolidone-2, 1-(dimethylchlorogermylmethyl)piperidone-2, (N–Ge)-chelate  $O$ -(dimethylchlorogermylmethyl)- $\delta$ -valeroactim and 2-(chlorodimethylgermylmethylthio)pyrrolidine-1, crystal and molecular structures of; structural correlations between derivatives of four- and five-coordinated germanium (A.O. Mozzchukhin, A.A. Macharashvili, V.E. Shklover, Yu.T. Struchkov, A.G. Shipov, V.N. Sergeev, S.A. Artamkin, S.V. Pestunovich, Yu.I. Baukov), (408) 305
- Group 14 metals, NQR studies of intramolecular complexes in organic compounds of (V.P. Feshin, G.A. Polygalova), (409) 1
- Triphenylsilanol, molecular and crystal structure of (H. Puff, K. Braun, H. Reuter), (409) 119
- 3-Organochlorogermylamino 2-methylthiophoates, synthesis of (M. Rivière-Baudet, P. Rivière, A. Castel, A. Morère, C. Abdennhader), (409) 131
- (Ferrocenylacyl)germanes, chemical and structural investigations of (H.K. Sharma, F. Cervantes-Lee, K.H. Pannell), (409) 321
- Group 14 triflates, synthesis of (W. Uhlig), (409) 377
- Methylethylnylgermanes;  $^{73}\text{Ge}$ ,  $^{13}\text{C}$  and  $^1\text{H}$  NMR spectroscopy of (E. Liepiņš, M.V. Petrova, E.T. Bogoradovsky, V.S. Zavgorodny), (410) 287
- $\text{PhMe}_2\text{E-E'Me}_3$  ( $\text{E}, \text{E}' = \text{Si}$  and  $\text{Ge}$ ), aryl-substituted catenates of group 4B elements; photochemical reactions, formation of a radical pair (K. Mochida, H. Kikkawa, Y. Nakadaira), (412) 9
- Trifluoromethylgermanes, properties of; adduct formation and substitution reactions of  $(\text{CF}_3)_n\text{GeX}_{4-n}$  ( $n = 2-4$ ) with  $\text{NH}_3$  (R. Eujen, F.E. Laufs), (415) 47

- Dimethylgermylene, photo-generated from dodecamethylcyclohexagermane; reactivity, absorption and emission (K. Mochida, N. Kanno, R. Kato, M. Kotani, S. Yamauchi, M. Wakasa, H. Hayashi), (415) 191
- Germaporphene, stable complex; chemical properties (H. Ranaivonjatovo, J. Escudié, C. Couret, J. Satgé), (415) 327
- Some silicon, germanium and tin acyloxyderivatives;  $^{13}\text{C}$ ,  $^{17}\text{O}$ ,  $^{29}\text{Si}$  NMR study of (I. Zicmane, E. Liepiņš, L.M. Ignatovich, E. Lukevics), (417) 355
- A new spirogermole: its synthesis, and some aspects of its reactivity (J.-D. Andriamizaka, C. Couret, J. Escudié, A. Laporterie, G. Manuel and M. Regitz), (419) 57
- On carbon-13 spectra of thio- and dithiocarboxylic acids triorgano group 14 metal esters (S. Kato, A. Hori, M. Mizuta, T. Katada, H. Ishihara, K. Fujeda and Y. Ikebe), (420) 13
- "Head-to-head" dimerization and dehydrodimerization of vinyl ketones catalyzed by modified rhodium(I) complexes (I.P. Kovalev, Y.N. Kolmogorov, Y.A. Strelenko, A.V. Ignatenko, M.G. Vinogradov and G.I. Nikishin), (420) 125
- Zum gezielten Aufbau oligomerer Strukturen der Elemente der 14. Gruppe mittels ihrer Triflatderivate (W. Uhlig), (421) 189
- Pyrrolyl compounds of main group elements. Synthesis of group 14 5-metallated pyrrole-2-carbaldehydes (F. Denat, H. Gaspard-Iloughmane and J. Dubac), (423) 173
- Synthesis, reactivity and crystal structure of trimesitylgermylamine,  $\text{Mes}_3\text{GeNH}_2$  (M. Rivière-Baudet, A. Morère, J.F. Britten and M. Onyszchuk), (423) C5
- Zur Stereochemie der mikrobiellen Reduktion von *rac*-Acetyl(t-butyl)methylphenylsilan mit *Trigonopsis variabilis* (DSM 70714) und *Corynebacterium dioxydans* (ATCC 21766): Aufklärung der absoluten Konfiguration der Biotransformationsprodukte (*SiR,CR*)- und (*SiS,CR*)-t-Butyl(1-hydroxyethyl)-methylphenylsilan (R. Tacke, F. Wuttke und H. Henke), (424) 273

## Gold

- [ $\text{Os}_6\text{AuC}(\text{CO})_{20}(\mu\text{-OMe})$ ]; isolation, characterisation, and crystal and molecular structure of (C.M. Hay, B.F.G. Johnson, J. Lewis, N.D. Prior, P.R. Raithby, W.T. Wong), (401) C20
- Borido cluster  $\text{Fe}_4(\text{CO})_{12}\text{BHAu}_2(\text{AsPh}_3)_2$ , synthesis and molecular structure of; investigation of the electrochemistry of  $\text{Fe}_4(\text{CO})_{12}\text{BHAu}_2\text{L}_2$ ,  $\text{L} = \text{AsPh}_3$  or  $\text{PPh}_3$  (C.E. Housecroft, M.S. Shongwe, A.L. Rheingold, P. Zanello), (408) 7
- Reactions of some Fe–Ir clusters; crystal structures of  $\text{Fe}_2\text{Ir}(\mu\text{-H})(\mu_3\text{-CCHPh})(\text{CO})_8(\text{PPh}_3)_2$  and  $\text{Au}_2\text{Fe}_2\text{Ir}(\mu_4\text{-C}_2\text{Ph})(\text{CO})_7(\text{PPh}_3)_3$  (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tiekkink), (408) 77
- $\text{R}_3\text{PAu}^{\text{I}}$  C-, S-, and N-derivatives, gold-197 Mössbauer investigation of (F. Bonati, A. Burini, B.R. Pietroni, E. Torregiani, S. Calogero, F.E. Wagner), (408) 125
- Symmetric C-imidazolylgold(I), reactions leading to  $\text{Au}^{\text{I}}$  carbene complexes or mixed valence or  $\text{Au}^{\text{III}}$  imidazolyl derivatives; crystal structure of [1-benzyl-3-(carboethoxy)imidazolin-2-yliden]chlorogold(I) (F. Bonati, A. Burini, B.R. Pietroni, B. Bovio), (408) 271
- Group IB metal heteronuclear cluster compounds in the solid state, high-resolution  $^{31}\text{P}\text{-}\{^1\text{H}\}$  nuclear magnetic resonance studies of (S.S.D. Brown, I.D. Salter, D.J. Smith, N.J. Clayden, C.M. Dobson), (408) 439
- [ $\{\text{Co}_2(\text{CO})_6\text{PhC}_2\}_2$ ] acetylene-linked dimeric cluster, synthesis and structural characterisation of (B.F.G. Johnson, J. Lewis, P.R. Raithby, D.A. Wilkinson), (408) C9
- Iridium-193 and gold-197 Mössbauer investigations of iridium(I), iridium(III) and iridium(I)/gold(I) complexes (A.L. Bandini, G. Banditelli, F. Bonati, S. Calogero, F.E. Wagner), (410) 241
- Three-membered metallacycle  $\text{Cp}_3\text{Rh}_3(\mu\text{-CO})_3$ , cluster and benzenoid system in reactions with 12-electron particles  $\text{PPh}_3\text{M}^+$  ( $\text{M} = \text{Cu, Ag, Au}$ ) and  $\text{CpM}^+$  ( $\text{M} = \text{Fe, Ru}$ ) (M.I. Rybinskaya, A.R. Kudinov, D.V. Muratov, N.P. Gambaryan, I.V. Stankevich, A.L. Chistyakov), (413) 419
- [ $\text{TiAu}(\text{C}_5\text{H}_4\text{PPh}_2)_2\right]_2$  tetrametallic complex, synthesis and structure of (G.K. Anderson, N.P. Rath), (414) 129
- $\mu\text{-}(1,2\text{-Diisocyanobenzol})\text{bis}(\text{chlorogold})$ , addition of amines to; participation of neighbouring isocyano functions and  $[\text{Au}]C$  transfer (W.P. Fehlhammer, W. Finck), (414) 261
- Gold(I) complexes of 1,1'-bis(diphenylphosphino)ferrocene, studies on (A. Houlton, R.M.G. Roberts, J. Silver, R.V. Parish), (418) 269

## Hafnium

- Zirconium and hafnium hexakis(trimethylsilyl)metallocene dichlorides, hindered cyclopentadienyl rotation in; evidence for complete rotation of the tris(trimethylsilyl)cyclopentadienyl ligands (C.H. Winter, D.A. Dobbs, X.-X. Zhou), (403) 145
- Bis(cyclopentadienyl)hafnium(IV) dichloride, heterocarboxylates of (P. Lukose, S.C. Bhatia, A.K. Narula), (403) 153
- Heterobimetallic Zr,Hf( $\mu$ -formaldehyde) complex from  $[(\text{CH}_2\text{O})\text{ZrCp}_2]$  dimer and hafnocene dichloride (G. Erker, M. Mena, M. Bendix), (410) C5
- Bis(trimethylsilylcyclopentadienyl) complexes of hafnium, synthesis of (J.B. Hoke, E.W. Stern), (412) 77
- Hafnium and titanium catalysts, comparative behaviour in copolymerization of  $\alpha$ -olefins (A. Altomare, R. Solaro, F. Ciardelli, L. Barazzoni, F. Menconi, S. Malquori, F. Masi), (417) 29
- Group 4 1,1'-bis(trimethylsilyl)metallocenes, binuclear benzyl-1,2,4,5-tetrathiolato complexes of; synthesis and structure of  $[(\eta^5\text{-Me}_3\text{SiC}_5\text{H}_4)_2\text{TiS}_2]_2\text{TiS}_2\text{C}_6\text{H}_2$  and  $[(\eta^5\text{-Me}_3\text{SiC}_5\text{H}_4)_2\text{HfS}_2]_2\text{C}_6\text{H}_2$  (H. Balz, H. Köpf, J. Pickardt), (417) 397
- Molekulares Design von Übergangsmetallkomplexverbindungen zur Aktivierung kleiner Moleküle. II. Zur Carbonylierung von Bis( $\pi$ -pentamethylcyclopentadienyl)metall-1,3-Dien-Komplexen  $\text{Cp}^*_2\text{M}$ -(1,3-Dien) ( $\text{M} = \text{Zr}, \text{Hf}$ )—Bildung einkerniger Metall-Endiolate (R. Beckhaus, D. Wilbrandt, S. Flatau und W.-H. Böhmer), (423) 211

## Indium

- Metalloporphyrins with metal–metal  $\sigma$ -bonds; synthesis, spectroscopic characterization, and electrochemistry of (P)MRe(CO)<sub>5</sub> where P is the dianion of octaethylporphyrin (OEP) or tetraphenylporphyrin (TPP) and M = Al, Ga, In or Tl (R. Guillard, A. Zrinch, A. Tabard, L. Courthaudon, B. Han, M. Ferhat, K.M. Kadish), (401) 227
- Trialkylindium compounds, reaction with (cyclopentadienyl)(tricarbonyl)metal radicals (D.L. Thorn), (405) 161
- Trifluoromethyl gallium, indium, and thallium compounds, syntheses of; mechanism of polar trifluoromethyl group transfer (D. Naumann, W. Strauß, W. Tyrra), (407) 1
- In–Fe heteronuclear carbonyl compounds, synthesis and chemical characterisation of; crystal structures of  $[\text{NMe}_3\text{CH}_2\text{Ph}]_3[\text{In}(\text{Fe}(\text{CO}))_3]$  and  $[\text{NEt}_4]_2[\text{In}_2\text{Br}_4(\mu\text{-Fe}(\text{CO}))_2]$  (V.G. Albano, M. Cané, M.C. Iapalucci, G. Longoni, M. Monari), (407) C9
- Dimethylindium- and bis(trimethylsilylmethyl)indium-dimesylamide, synthesis of; crystal structure of  $[(\text{Me}_3\text{SiCH}_2)_2\text{In}(\text{SO}_2\text{Me})_2]_2$  (A. Blaschette, A. Michalides, P.G. Jones), (411) 57
- Platinum–indium and cobalt–gallium complexes, synthesis of; crystal structure of  $[\text{Cy}_2\text{P}(\text{CH}_2\text{CH}_2)\text{PCy}_2]\text{Pt}(\text{InR}_2)(\text{R})$  ( $\text{Cy} = \text{cyclo-C}_6\text{H}_{11}$ ) (R.A. Fischer, J. Behm), (413) C10
- Diorganindium fluorides; crystal structures of  $[{}^1\text{Pr}_2\text{In}(\text{THF})_2]\text{[BF}_4]$  and  $(\text{MesBO})_3$  (B. Neumüller, F. Gahlmann), (414) 271
- Organoindium derivatives, preparation and characterisation of; crystal structure of dimethyl(salicyl-aldehyde)indium (N.W. Alcock, I.A. Degnan, S.M. Roe, M.G.H. Wallbridge), (414) 285
- Quinones, allylation by allylic reagents (S. Araki, N. Katsumura, Y. Butsugan), (415) 7
- Group IIIIB organometallic peroxides, synthesis and reactivity of (Yu.A. Alexandrov, N.V. Chikinova), (418) 1
- (Tert-butyl) cyclopentadienylindium(I),  $\text{In}(\text{C}_5\text{H}_4\text{CMe}_3)$ ; synthesis, characterization and X-ray structural study (O.T. Beachley, Jr., J.F. Lees, R.D. Rogers), (418) 165
- Dialkyindium derivatives of unsaturated nitrogen heterocycles—preparation, spectroscopic properties and the use in MOVPE (K. Locke, J. Weidlein, F. Scholz, N. Bouanah, N. Brianese, P. Zanella and Y. Gao), (420) 1
- Die Molekül- und Kristallstruktur von  $(\text{CH}_3)_2\text{InNC}_4\text{H}_4$  (H.-D. Hausen, K. Locke und J. Weidlein), (423) C1
- Diethylindium(III) derivatives: synthesis and characterization of diethylindium(III) dialkylamides and diethylindium(III) pyrrolide. X-Ray crystal structures of diethylindium(III) pyrrolide and diethylindium(III) bromide (M. Porchia, F. Benetollo, N. Brianese, G. Rossetto, P. Zanella and G. Bombieri), (424) 1

## Iridium

- Heteronuclear and heterovalent diphenylphosphido-bridged complexes of rhodium, iridium and ruthenium; crystal structure of  $[(\text{C}_5\text{Me}_5)\text{ClRh}(\mu\text{-Cl})(\mu\text{-PPh}_2)\text{Rh}(\text{COD})]\cdot\frac{1}{2}\text{Et}_2\text{O}$  ( $\text{COD} = 1,5$ -cyclooctadiene) (M. Esteban, A. Pequerul, D. Carmona, F.J. Lahoz, A. Martin, L.A. Oro), (402) 421

- Rhodium(I) and iridium(I) complexes of  $\text{Ph}_2\text{P}(\text{S})\text{CH}_2\text{P}(\text{S})\text{Ph}_2$ , bis(diphenylphosphino)methane disulfide (M.S. Abbassioun, P.A. Chaloner, C. Claver, P.B. Hitchcock, A.M. Masdeu, A. Ruiz, T. Saballs), (403) 229
- Coordinated water complexes of rhodium and iridium, formation and characterisation of; crystal structure of  $\text{trans}\text{-}[\text{Rh}(\text{PPh}_3)_2(\text{CO})(\text{H}_2\text{O})]^+\text{CF}_3\text{SO}_3^-$  (P.J. Stang, L. Song, Y.-H. Huang, A.M. Arif), (405) 403
- Vinylidene complexes, nucleophilic behavior of (F. Delbecq), (406) 171
- Iron–iridium clusters; X-ray structures of  $\text{Fe}_2\text{Ir}(\mu_3\text{-C}_2\text{Ph})(\text{CO})_8(\text{PPh}_3)$ ,  $\text{FeIr}_2(\mu_3\text{-}\eta^2\text{-PhC}_2\text{C}_2\text{Ph})(\text{CO})_7$  ( $\text{PPh}_3$ )<sub>2</sub> and  $\text{Fe}_2\text{Ir}(\mu_3\text{-PhC}_2\text{PEt}_3)(\text{CO})_7(\text{PEt}_3)$  (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tiekkink), (407) 391
- Reactions of some Fe–Ir clusters; crystal structures of  $\text{Fe}_2\text{Ir}(\mu\text{-H})(\mu_3\text{-CCHPh})(\text{CO})_8(\text{PPh}_3)$  and  $\text{Au}_2\text{Fe}_2\text{Ir}(\mu_4\text{-C}_2\text{Ph})(\text{CO})_7(\text{PPh}_3)_3$  (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tiekkink), (408) 77
- Coordinatively unsaturated  $\sigma$ -aryl iridium(III) complexes containing iridium–mercury bonds (W.R. Roper, G.C. Saunders), (409) C19
- Chiral chloro(dimethylsulfoxide)( $\eta^5$ -pentamethylcyclopentadienyl)thyminatoiridium(III), preparation and structure of (R. Krämer, K. Polborn, W. Beck), (410) 111
- Iridium-193 and gold-197 Mössbauer investigations of iridium(I), iridium(III) and iridium(I)/gold(I) complexes (A.L. Bandini, G. Banditelli, F. Bonati, S. Calogero, F.E. Wagner), (410) 241
- Tris(diphenylthiophosphinoyl)methane and tris(diphenylthiophosphinoyl)methanide complexes of rhodium and iridium, synthesis and structural characterisation of; crystal structures of  $[\text{Rh}(\text{C}_8\text{H}_{12})(\eta^2\text{-C}(\text{P}(\text{S})\text{Ph}_2)_3\text{S}, \text{S})]$  and  $[\text{Ir}(\text{CO})_2(\eta^2\text{-C}(\text{P}(\text{S})\text{Ph}_2)_3\text{S}, \text{S})]$  (J. Browning, K.R. Dixon, R.W. Hiltz, N.J. Meanwell, F. Wang), (410) 389
- Highly reduced metalcarbonyl  $[\text{Ir}(\text{CO})_3]^{13}$ , reaction with the ethylenepentacarbonylrhenium cation; formation of the trinuclear hydride  $(\text{OC})_5\text{Re}-\text{Ir}(\text{CO})_3(\text{H})-\text{Re}(\text{CO})_5$  and the trimetalated carbonate  $\{\mu_3\text{-CO}_3(\text{Re}(\text{CO})_5)_3\}^+$  (J. Breimair, C. Robl, W. Beck), (411) 395
- $[\text{M}_6(\text{CO})_{15}]^{2-}$  anionic clusters ( $\text{M} = \text{Co}$  and  $\text{Ir}$ ), electrochemistry of; preparation and chemical characterisation of  $[\text{Ir}_6(\text{CO})_{14}(\text{NCO})]^-$  and of the radical anion  $[\text{Ir}_6(\text{CO})_{15}]^-$ ; proposed mechanism of formation of the hexanuclear substituted cluster  $[\text{Ir}_6(\text{CO})_{15}\text{Cl}]^-$  (A. Cinquantini, P. Zanello, R. Della Pergola, L. Garlaschelli, S. Martinengo), (412) 215
- Pentamethylcyclopentadienyliridium complexes, ring-methyl activation in; synthesis of  $[(\text{C}_5\text{Me}_4\text{CH}_2\text{E})\text{IrL}_n]$  ( $\text{E} = \text{SiMe}_3$ ,  $\text{PhCH}_2$ ,  $\text{CH}_2=\text{CHCH}_2$ ,  $\text{PPh}_2$ ,  $\text{Pt}(\text{PEt}_3)_2\text{Cl}$ ); synthesis and crystal structure of  $[(1,3\text{-C}_5\text{Me}_3(\text{CH}_2\text{SiMe}_3)_2)\text{Ir}(\text{CO})\text{Me}(\text{Cl})]$  (J.A. Miguel-Garcia, H. Adams, N.A. Bailey, P.M. Maitlis), (413) 427
- Triangular cluster  $\text{Cp}_2\text{Cr}_2(\mu\text{-SCMe}_3)(\mu_3\text{-S})_2\text{Ir}(\text{CO})(\text{PPh}_3)$  and octanuclear cluster  $\text{Cp}_2\text{Cr}_2(\mu_3\text{-S})_2(\mu_4\text{-S})\text{Rh}_2[\text{Cp}_2\text{Cr}_2(\mu\text{-SCMe}_3)(\mu_3\text{-S})_2]_2$  with unusual triangular heterometalloornament framework; synthesis, molecular structure and magnetic properties of (A.A. Pasynskii, I.L. Eremenko, V.R. Zalmanovich, V.V. Kavarin, B. Orazsakhatov, S.E. Nefedov, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky, Yu.T. Struchkov), (414) 55
- Rhodium(I) and iridium(I) complexes with 1,2-bis(dicyclohexylphosphino)ethane ligands; their reactions with carbon monoxide (A. Del Zotto, L. Costella, A. Mezzetti, P. Rigo), (414) 109
- $\text{M}_4(\text{CO})_{11}\text{L}$  ( $\text{M} = \text{Co}$ ,  $\text{Ir}$ ;  $\text{L} = \text{PPh}_3$ ,  $\text{P}^n\text{Bu}_3$ ,  $\text{P}(\text{OMe})_3$ ,  $\text{P}(\text{OEt})_3$ ); kinetics and mechanisms of CO substitution with  $\text{L}$  in the presence of  $\text{Me}_3\text{NO}$  (J.-Q. Wang, J.-K. Shen, Y.-C. Gao, Q.-Z. Shi, F. Basolo), (417) 131
- Iridium hydrogenation catalysts, hybrid P–N phosphines as participative ligands (E. Farnetti, G. Nardin, M. Graziani), (417) 163
- $\beta$ -diketonato-1,5-cyclooctadieneiridium(I) complexes, oxidative addition of  $\text{Hg}(\text{CN})_2$  to; mechanism of (G.J.J. Steyn, S.S. Basson, J.G. Leipoldt, G.J. van Zyl), (418) 113
- $\beta$ -Diketonato-1,5-cyclo-octadieneiridium(I) complexes, kinetics of the substitution reactions with derivatives of 1,10-phenanthroline and 2,2'-dipyridyl (J.G. Leipoldt, S.S. Basson, G.J. van Zyl, G.J.J. Steyn), (418) 241
- Protonation of cationic hydride and alkyl complexes of Ir(III) (T.J. Johnson, B.E. Hauger, E.B. Lobkovsky and K.G. Caulton), (424) 371

## Iron

Half-sandwich and tripledecker *nido* and *clos* compounds, a simple MO model to explain the redox behaviour of clusters (B. Eber, D. Buchholz, G. Huttner, Th. Fässler, W. Imhof, M. Fritz, J.C. Jochims, J.C. Daran, Y. Jeannin), (401) 49

- Reaction of norbornadiene with sulfur and dodecacarbonyltriiron; crystal structure of one of the isomeric products (A.I. Nekhaev, S.D. Alekseeva, B.I. Kolobkov, G.G. Aleksandrov, M.T. Toshev, H.B. Dustov, N.A. Parpiev), (401) 65
- $\text{Fe}_3(\eta^4\text{C}_6\text{H}_8)(\mu\text{-CO})_2(\text{CO})_6(\mu_3\text{-S})$  trinuclear cluster, synthesis and structure of (A.I. Nekhaev, S.D. Alekseeva, B.I. Kolobkov, G.G. Aleksandrov, M.T. Toshev, H.B. Dustov), (401) 75
- Ferrocene, monosubstituted derivatives of; ferrocene-containing penicillins and cephalosporins (D. Scutaru, L. Tătaru, I. Mazilu, E. Diaconu, T. Lixandru, Cr. Simionescu), (401) 81
- Ferrocene, heterodisubstituted derivatives of; ferrocene-containing penicillins and cephalosporins (D. Scutaru, I. Mazilu, M. Văjă, L. Tătaru, A. Vlase, T. Lixandru, Cr. Simionescu), (401) 87
- Cyclic and acyclic (diene)Fe(CO)<sub>2</sub>L complexes (L = phosphine, phosphite, isonitrile), solution structure and fluxional behaviour of (J.A.S. Howell, P. McArdle, D. Cunningham, Z. Goldschmidt, H. Gottlieb, G. Strul), (401) 91
- Chiral ( $\eta^6$ -arene)( $\eta^5$ -cyclopentadienyl)iron(II) complexes, synthesis of; their characterisation by <sup>13</sup>C NMR spectroscopy (K. Bambridge, R.M.G. Roberts), (401) 125
- Permethylated  $\alpha$ -metallocenylcarbinols (M = Os, Ru, Fe), hydrogen bonds in and conformations of (E.S. Shubina, L.M. Epstein, T.V. Timofeeva, Yu. T. Struchkov, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 133
- Permethylated  $\alpha$ -metallocenylcarbinols in the solid state (M = Fe, Ru, Os), IR spectroscopic study of hydrogen bonds in (E.S. Shubina, L.M. Epstein, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 145
- ( $\eta^6$ -Phenyl)Cr(CO)<sub>3</sub>- and ( $\eta^6$ -phenyl)Co<sub>4</sub>(CO)<sub>9</sub>-substituted metallocenylcarbinols of the iron subgroup; synthesis, structure and hydrogen bonding in (E.S. Shubina, L.M. Epstein, Yu.L. Slovokhotov, A.V. Mironov, Yu.T. Struchkov, V.S. Kaganovich, A.Z. Kreindlin, M.I. Rybinskaya), (401) 155
- (CO)<sub>6</sub>Fe<sub>2</sub>( $\mu_3$ -Se)( $\mu_3$ -Te)Pt(PPh<sub>3</sub>)<sub>2</sub> and Fe<sub>3</sub>(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub>( $\mu_3$ -Se)( $\mu_3$ -Te), synthesis and characterisation of (P. Mathur, D. Chakrabarty, Md.M. Hossain), (401) 167
- Organotin hexacyanoferrates, synthesis and characterisation of (A. Bonardi, C. Carini, C. Pelizzi, G. Pelizzi, G. Predieri, P. Tarasconi, M.A. Zoroddu, K.C. Molloy), (401) 283
- Fe<sub>3</sub>(CO)<sub>9</sub>( $\mu_3$ -Te)<sub>2</sub>, addition to methylene groups to (P. Mathur, V.D. Reddy, R. Bohra), (401) 339
- Pt(II) complexes of tetrathia[n](1,1')ferrocenophanes, preparation and properties of (M. Sato, H. Asano, S. Akabori), (401) 363
- ( $\eta^3$ -Allyl)Fe(CO)<sub>4</sub><sup>+</sup>, reactions with allyl nucleophiles; approaches to selective isoprenologation (Z. Li, K.M. Nicholas), (402) 105
- ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe(CO)<sub>2</sub>Si(Ph)(SiMe<sub>3</sub>)<sub>2</sub> bulky iron-substituted trisilane, X-ray structural study of (R. West, E.K. Pham), (402) 215
- Carbene-substituted sulphur-bridged carbonyl iron clusters, synthesis of (B. Eber, G. Huttner, L. Zsolnai, W. Imhof), (402) 221
- Thiolato bridged-iron carbonyl complexes, synthesis of (P.M. Treichel, R.A. Crane, R. Matthews, K.R. Bonnin, D. Powell), (402) 233
- 1,2,3,4,5-Pentaethylcyclopentadiene, synthesis of (D. Stein, H. Sitzmann), (402) 249
- Cyclopentadienyliron(II) arene cations, photochemistry of; analysis of the reaction with epoxides by means of time-resolved laser spectroscopy (K.M. Park, G.B. Schuster), (402) 355
- trans-[FeH(P≡CBu')(dppe)<sub>2</sub>]BF<sub>4</sub>  $\eta^1$ -phosphaalkyne complex, synthesis of; its conversion into a  $\eta^1$ -fluorophosphaalkene complex; crystal structure of trans-[FeH(PF=CIIBu'<sup>1</sup>(dppe)<sub>2</sub>]FeCl<sub>2</sub>F<sub>2</sub>] (P.B. Hitchcock, M.A.N.D.A. Lemos, M.F. Meidine, J.F. Nixon, A.J.L. Pombeiro), (402) C23
- (RR,SS)-[( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe(CO)(PPh<sub>3</sub>)COCH<sub>2</sub>]CH<sub>2</sub>, self-recognition by the iron chiral auxiliary [( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe(CO)(PPh<sub>3</sub>)] in the formation of (G.J. Bodwell, S.G. Davies, S.C. Preston), (402) C56
- Tricarbonyliron complexes as intermediates to isoquinuclidines (G.R. Stephenson, R.D. Thomas, F. Cassidy), (402) C59
- Cyclopentadienyliron complexes, monosubstituted nitrogen donors as ligands in; synthesis, reactivity, ligand properties, and crystal structure of [C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub>(C<sub>5</sub>H<sub>5</sub>N)]SbF<sub>6</sub> (H. Schumann, M. Speis, W.P. Bosman, J.M.M. Smits, P.T. Beurskens), (403) 165
- [C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub>(ER<sub>3</sub>)]BF<sub>4</sub> complexes (E = N, P, As, Sb and Bi), synthesis and characterisation of; crystal structure of [C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub>(Bi(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>)]BF<sub>4</sub> (H. Schumann, L. Eguren), (403) 183
- Substituted ferrocenes and ruthenocenes, dynamic NMR studies of ring rotation in (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šík), (403) 195
- [( $\eta^5$ -Dimethylphosphino)cyclopentadienyl][ $\eta^7$ -(dimethylphosphino)cycloheptatrienyl]titanium, heterobimetallic compounds derived from (M. Ogasa, M.D. Rausch, R.D. Rogers), (403) 279

- C*-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325
- HFe<sub>3</sub>(CO)<sub>9</sub>(C≡CSiMe<sub>3</sub>), synthesis and crystal structure of (J.J. Schneider, M. Nolte, C. Krüger), (403) C4
- Organoiron chemistry; Annual Survey for the year 1989 (R.C. Kerber), (404) 107
- Titanium, iron and thallium vinylcyclopentadienyl derivatives, synthesis and structure of (M. Ogasa, D.T. Mallin, D.W. Macomber, M.D. Rausch, R.D. Rogers, A.N. Rollins), (405) 41
- 1,*n*-Dicarbonyl compounds, one-step synthesis from carboxylic acid derivatives and di-Grignard reagents in the presence of transition metal catalysts (F. Babudri, A. D'Ettore, V. Fiandanese, G. Marchese, F. Naso), (405) 53
- Clusters with Re<sub>2</sub>Fe and Re<sub>2</sub>Ni<sub>2</sub> cores, phenylacetylene transformations in; synthesis and molecular structure of Re<sub>2</sub>Fe(CO)<sub>6</sub>(μ<sub>3</sub>,η<sup>4</sup>-CH=CHC<sub>6</sub>H<sub>4</sub>) and Cp<sub>2</sub>Ni<sub>2</sub>Re<sub>2</sub>(CO)<sub>6</sub>(μ<sub>4</sub>-PhC≡CCH=CHPh) (A.D. Shaposhnikova, G.L. Kamalov, R.A. Stadnichenko, A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, Yu.T. Struchkov, A.I. Yanovsky), (405) 111
- (Pentadienyl)iron[1,2-bis(diethylphosphino)ethane] electron-rich complexes, synthesis and characterisation of (J.R. Bleeke, R.J. Wittenbrink), (405) 121
- Ferrocenethioles, reactions with norbornadiene and norbornene (M. Herberhold, O. Nuyken, T. Pöhlmann), (405) 217
- Ferrocenecarboxaldoxime, crystal structure of the low-melting form of (G. Ferguson, W. Bell, C. Glidewell), (405) 229
- [CpM(CO)<sub>2</sub>R] long-chain alkyl complexes (Cp = η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>; M=Fe or Ru; R=n-C<sub>6</sub>H<sub>13</sub> to n-C<sub>12</sub>H<sub>25</sub>), synthesis, characterisation and properties of (A. Emeran, M.A. Gafoor, J.K.I. Goslett, Y.-H. Liao, L. Pimble, J.R. Moss), (405) 237
- Chromium–iron–sulfide Cp<sub>3</sub>Cr<sub>3</sub>(μ<sub>3</sub>-S)<sub>4</sub>FeSPh and [Cp<sub>3</sub>Cr<sub>3</sub>(μ<sub>3</sub>-S)<sub>4</sub>Fe]<sub>2</sub>Fe(CO)<sub>3</sub>S·C<sub>6</sub>H<sub>6</sub> clusters with tetrahedral metal frameworks, syntheses and molecular structures of (S.E. Nefedov, A.A. Pasynskii, I.L. Eremenko, E.E. Stomakhina, A.I. Yanovsky, Yu.T. Struchkov), (405) 287
- σ-Alkenyl cyclopentadienyl iron(II) complexes [Fe(C≡CR)L<sub>2</sub>(η-C<sub>5</sub>H<sub>5</sub>)] [L = CO; L<sub>2</sub> = bis(di-phenylphosphino)methane (dppm)], synthesis and characterisation of; crystal structure of [Fe(C≡CC<sub>6</sub>H<sub>5</sub>)(dppm)(η-C<sub>5</sub>H<sub>5</sub>)] (M.P. Gamasa, J. Gimeno, E. Lastra, M. Lanfranchi, A. Tiripicchio), (405) 333
- Iron sandwich complexes and related species, carbon-13 spin-lattice relaxation in (R.M.G. Roberts, J.F. Warmsley), (405) 347
- Ferrocene complexes with cyclodextrins in DMSO, carbon-13 spin-lattice relaxation studies of (R.M.G. Roberts, J.F. Warmsley), (405) 357
- Metallocene-containing cryptands, NMR studies on metal complexes of (C.D. Hall, N.W. Sharpe), (405) 365
- Platinum(II) chloride complexes of 1,1'-bis(methylthio)ferrocene, 1,1'-bis(methylseleno)ferrocene and 1,1'-bis(methylthio)ruthenocene, synthesis and stereodynamics of (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šík), (405) 375
- Vinylidene complexes, nucleophilic behavior of (F. Delbecq), (406) 171
- Ferrocene-containing rifamycins, synthesis and characterisation of (D. Scutaru, I. Mazilu, L. Tătaru, M. Vătă, T. Lixandru), (406) 183
- [Cp<sub>2</sub>Cr<sub>2</sub>(μ-SCMe<sub>3</sub>)(μ<sub>3</sub>-S)<sub>2</sub>]<sub>2</sub>Fe<sup>+</sup>I<sub>4</sub><sup>-</sup> heterometallospirane cluster and [Cp<sub>2</sub>Cr<sub>2</sub>(μ-SCMe<sub>3</sub>)(μ<sub>3</sub>-S)<sub>2</sub>Fe]<sub>2</sub>(μ-Se)<sub>2</sub> cluster with "kayak-paddle" framework; synthesis, magnetic properties and structures of (A.A. Pasynskii, I.L. Eremenko, E.E. Stomakhina, S.E. Nefedov, O.G. Ellert, A.I. Yanovsky, Yu.T. Struchkov), (406) 383
- 1,2,4,5-Tetrakis(diphenylphosphino)benzene complexes; synthesis and X-ray crystal structure of [(CO)<sub>3</sub>Fe]<sub>2</sub>(μ-η<sup>2</sup>-η<sup>2</sup>-(Ph<sub>2</sub>P)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>(PPh<sub>2</sub>)<sub>2</sub>]] (G. Hogarth), (406) 391
- Simple iron alkyls, reactions with a thiosulfinate ester capable of serving as a disulfur monoxide source (K.R. Powell, W.J. Elias, M.E. Welker), (407) 81
- Di-iron centre-bridging ethenyl ligand, generation via carbon–phosphorus bond cleavage (G. Hogarth), (407) 91
- Heterometallic *n,π*-complexes of transition metals based on mesitylacetonitrile (V.S. Kaganovich, M.I. Rybinskaya), (407) 215
- Cyclopalladated *N,N*-dimethylferrocenecarbothioamide, derivatives of (M. Nonoyama, K. Hamamura), (407) 271

- Unidentate and heterobimetallic 1,1'-bis(diphenylphosphino)ferrocene complexes, X-ray photoelectron spectroscopic differentiation of chemically distinct phosphorus environments in (H.S.O. Chan, T.S.A. Hor, L.-T. Phang, K.L. Tan), (407) 353
- [Rh(dppf)( $\eta^6$ -C<sub>6</sub>H<sub>5</sub>)B(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>] (dppf = 1,1'-bis(diphenylphosphino)ferrocene), synthesis and crystal structure of (B. Longato, G. Pilloni, R. Graziani, U. Casellato), (407) 369
- $\mu_3$ -Phosphinidene-bridged iron and ruthenium clusters, selective hydrogenation of diphenylacetylene and isomerisation of *cis*-stilbene in the presence of (M. Castiglioni, R. Giordano, E. Sappa), (407) 377
- Iron–iridium clusters; X-ray structures of Fe<sub>2</sub>Ir( $\mu_3$ -C<sub>2</sub>Ph)(CO)<sub>8</sub>(PPh<sub>3</sub>), FeIr<sub>2</sub>( $\mu_3$ - $\eta^2$ -PhC<sub>2</sub>C<sub>2</sub>Ph)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub> and Fe<sub>2</sub>Ir( $\mu_3$ -PhC<sub>2</sub>PEt<sub>3</sub>)(CO)<sub>7</sub>(PEt<sub>3</sub>) (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tieckink), (407) 391
- In–Fe heteronuclear carbonyl compounds, synthesis and chemical characterisation of; crystal structures of [NMe<sub>3</sub>CH<sub>2</sub>Ph]<sub>2</sub>[In{Fe(CO)<sub>4</sub>}<sub>3</sub>] and [NEt<sub>4</sub>]<sub>2</sub>[In<sub>2</sub>Br<sub>4</sub>{ $\mu$ -Fe(CO)<sub>4</sub>}<sub>2</sub>] (V.G. Albano, M. Cané, M.C. Iapalucci, G. Longoni, M. Monari), (407) C9
- Borido cluster Fe<sub>4</sub>(CO)<sub>12</sub>BHAu<sub>2</sub>[AsPh<sub>3</sub>]<sub>2</sub>, synthesis and molecular structure of; investigation of the electrochemistry of Fe<sub>4</sub>(CO)<sub>12</sub>BHAu<sub>2</sub>L<sub>2</sub>, L = AsPh<sub>3</sub> or PPh<sub>3</sub> (C.E. Housecroft, M.S. Shongwe, A.L. Rheingold, P. Zanello), (408) 7
- Reactions of some Fe–Ir clusters; crystal structures of Fe<sub>2</sub>Ir( $\mu$ -H)( $\mu_3$ -CCHPh)(CO)<sub>8</sub>(PPh<sub>3</sub>) and Au<sub>2</sub>Fe<sub>2</sub>Ir( $\mu_4$ -C<sub>2</sub>Ph)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>3</sub> (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tieckink), (408) 77
- Dicarbonylbis(phosphorus-donor)iron complexes, trimerisation of methyl propiolate on (U. Grossmann, H.-U. Hund, H.W. Bosch, H. Schmalke, H. Berke), (408) 203
- Diferrocenylketones, new synthesis of (D.C. O'Connor Salazar, D.O. Cowan), (408) 219
- Diferrocenylsulfide, new synthesis and study of (D.C. O'Connor Salazar, D.O. Cowan), (408) 227
- Iron(II), tungsten(II) and molybdenum(II) ( $\eta^2$ -iodocyclopentadienyl)metallo methyl derivatives, methyl–iodine exchange in the presence of palladium (C. Lo Sterzo) (408) 253
- Silyl-bridged di- to tetranuclear ferrocenes; synthesis, cyclic voltammetry and oxidation of (H. Atzkern, J. Hiermeier, F.H. Köhler, A. Steck), (408) 281
- Cyclopentadienyliron arene cations, photochemical arene substitution in; synthesis of cyclopentadienyl-iron bisphosphine ligand complexes with group 14 and 15 ligands (H. Schumann, L. Eguren, J.W. Ziller), (408) 361
- Heterodichalcogene ferrocenes with open or bridging chains, synthesis by addition of Fe(C<sub>5</sub>H<sub>4</sub>ELi)<sub>2</sub>·2THF (E = S, Se) to propynoates (R. Broussier, M. El Mjidi, B. Gautheron), (408) 381
- Iron, cobalt and rhodium complexes of the optically active diolefin (+)-nopadiene and its derivatives; crystal structure of C<sub>5</sub>Me<sub>5</sub>Rh(nopadiene) (A. Salzer, H. Schmalke, R. Stauber, S. Streiff), (408) 403
- [( $\mu$ -CH<sub>3</sub>Te)Fe<sub>2</sub>(CO)<sub>6</sub>]<sub>2</sub>[ $\mu$ -Te(CH<sub>2</sub>)Te- $\mu$ ] methylene-bridged double butterfly shaped complex, synthesis and structural characterisation of (P. Mathur, V.D. Reddy, K. Das, U.C. Sinha), (409) 255
- (Ferrocenylacyl)germanes, chemical and structural investigations of (H.K. Sharma, F. Cervantes-Lee, K.H. Pannell), (409) 321
- Iron and molybdenum cyclometallated diazene complexes (II. Kisch, D. Garn), (409) 347
- Iron and ruthenium cationic ( $\eta^6$ -chloroarene) complexes, reactions with some O-silyl and C-silyl compounds (R.C. Cambie, S.A. Coulson, L.G. Mackay, S.J. Janssen, P.S. Rutledge, P.D. Woodgate), (409) 385
- Bis(alkinyl) titanocene complexes of FeCl<sub>2</sub> and CuCl (H. Lang, M. Herres, L. Zsolnai, W. Imhof), (409) C7
- Rigid rod, iron-metal containing polyyne polymers, synthesis of (B.F.G. Johnson, A.K. Kakkar, M.S. Khan, J. Lewis), (409) C12
- Fe<sub>2</sub>Ru( $\mu_3$ -Se)( $\mu_3$ -Te)(CO)<sub>9</sub> and ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)CoFe<sub>2</sub>( $\mu_3$ -Se)( $\mu_3$ -Te)(CO)<sub>6</sub>, synthesis and characterisation of (D. Chakrabarty, Md.M. Hossain, R.K. Kumar, P. Mathur), (410) 143
- Ferrocenyl-CO-CH=CH-aryl-type chalcone-analogous ferrocenes *ortho*-substituted in the aromatic ring, spectroscopic investigation of (Á.G. Nagy, P. Sohár, J. Márton), (410) 357
- Cyclopentadienyliron bis(phosphine) complexes with dimethylchalcogenide ligands; synthesis, reactivity and crystal structure of [C<sub>5</sub>H<sub>5</sub>Fe(P(OCH<sub>3</sub>)<sub>3</sub>)<sub>2</sub>(S(CH<sub>3</sub>)<sub>2</sub>)]PF<sub>6</sub> (H. Schumann, J.W. Ziller), (410) 365
- CH<sub>3</sub>C[CH<sub>2</sub>NC-M(CO)<sub>x</sub>]<sub>3</sub> complexes (M = Cr, W, x = 5; M = Fe, x = 4), synthesis of; crystal structure of CH<sub>3</sub>C[CH<sub>2</sub>NC-W(CO)<sub>5</sub>]<sub>3</sub> (F.E. Hahn, M. Tamm), (410) C9
- Mixed-valence 1',1''-dibenzylferrocenium cations, micromodulation of electron transfer rates in (T.-Y. Dong, C.-C. Schei, M.-Y. Hwang, T.-Y. Lee), (410) C39

- cis*-Fe(CO)<sub>4</sub>(SiCl<sub>3-n</sub>Me<sub>n</sub>)<sub>2</sub> (*n* = 1–3), conversion with phosphines; competition between CO substitution, SiR<sub>3</sub> elimination and formation of SiR<sub>2</sub> bridged complexes (U. Schubert, M. Knorr, C. Straßer), (411) 75
- 1,3-Ditellura[3]ferrocenophanes, Fe(C<sub>5</sub>H<sub>4</sub>Te)<sub>2</sub>E (E = S, Se, Te or CH<sub>2</sub>) (M. Herberhold, P. Leitner), (411) 233
- Te<sub>2</sub>S, Te<sub>2</sub>Se and Te<sub>3</sub> bridged compounds, energies of; CNDO/2 investigation of the mechanism of the bridge reversal process (E.W. Abel, K.G. Orrell, A.G. Osborne, V. Šík, W. Guoxiong), (411) 239
- 1,1'-Di-tert-butyl-3,3'-bis(2,2-dimethylpropionyl)ferrocene, diastereoisomeric forms of (W. Bell, C. Glidewell), (411) 251
- [{(Me<sub>3</sub>Sn<sup>IV</sup>)<sub>4</sub>Fe<sup>II</sup>(CN)<sub>6</sub>·4H<sub>2</sub>O]<sub>x</sub>, an organotin Berlin blue analogue with coordinative O–H ··· N hydrogen bonds (U. Behrens, A.K. Brimah, R.D. Fischer), (411) 325
- [Fe<sub>2</sub>(CN)(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>3</sub>]<sup>-</sup>, synthesis, structure and dynamics of (W.P. Fehlhammer, A. Schröder, F. Schoder, J. Fuchs, A. Völkl, B. Boyadjiev, S. Schrökamp), (411) 405
- Dicyclohexylammonium hydridoundecacarbonyltriferrate; X-ray diffraction study and Mössbauer effect spectral parameters of (R.B. King, G.S. Chorghade, N.K. Bhattacharyya, E.M. Holt, G.J. Long), (411) 419
- Fe<sub>3</sub>(CO)<sub>11</sub><sup>2-</sup> and Fe<sub>4</sub>(CO)<sub>13</sub><sup>2-</sup>, transformation to give trinuclear clusters (W. Deck, A.K. Powell, H. Vahrenkamp), (411) 431
- Bis[dicarbonyl(pentamethylcyclopentadienyl)iron], electron transfer reactions with 2,3-dichloro-5,6-disubstituted benzoquinone (F. Pradella, M. Scoponi, S. Sostero), (412) 137
- Decaethylferrocene, reaction with 2,3,5,6-tetrafluoro-7,7',8,8'-tetracyano-*p*-chinodimethane gives a charge-transfer complex (D. Stein, H. Sitzmann, R. Boese, E. Dormann, H. Winter), (412) 143
- Ferrocene-1,1'-dicarbaldehyde, improved synthesis of (G.G.A. Balavoine, G. Doisneau, T. Fillebeek-Khan), (412) 381
- Ditertiary phosphine and disphosphazane ligand-bridged derivatives of di-iron and diruthenium nonacarbonyl, electrochemical behaviour of (J.S. Field, A.M.A. Francis, R.J. Haines, S.F. Woollam), (412) 383
- [(Cp(OC)<sub>2</sub>Fe(olefin)]<sup>+</sup> and (1,5-hexadiene)PtCl<sub>2</sub>, addition of α-amino acids to coordinated olefins of (I. Zahn, K. Polborn, W. Beck), (412) 397
- (C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Fe(CO)(SiHPh<sub>2</sub>)<sub>2</sub>H, a catalytically active intermediate in the hydrosilylation of acetophenone by diphenylsilane; synthesis of (H. Brunner, K. Fisch), (412) C11
- "Flyover-bridged" complex [Fe<sub>2</sub>(CO)<sub>5</sub>(P-dppm){C<sub>2</sub>Et<sub>2</sub>(CO)C<sub>2</sub>Et<sub>2</sub>}], unexpected formation by reaction of a "ferrole" derivative with Ph<sub>2</sub>PCH<sub>2</sub>PPh<sub>2</sub> (dppm) and Me<sub>3</sub>NO; crystal structure of (R. Giordano, E. Sappa, D. Cauzzi, G. Predieri, A. Tiripicchio, M. Tiripicchio Camellini), (412) C14
- CpFe(CO)LX complexes (Cp = η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>; L = CO, PPh<sub>3</sub>; X = Br, I), photochemical halide exchange in; tandem photosubstitution of X<sup>-</sup> and CO (J. Zakrzewski), (412) C23
- Oligonuclear ferrocene derivatives with chalcogenide bridges (M. Herberhold, H.-D. Brendel, O. Nuyken, T. Pöhlmann), (413) 65
- [NEt<sub>4</sub>]<sub>2</sub>WFe<sub>2</sub>{μ<sub>3</sub>-η<sup>3</sup>,η<sup>5</sup>-CH=C(<sup>1</sup>Bu)C(O)C<sub>2</sub>B<sub>9</sub>H<sub>9</sub>Me<sub>2</sub>}(CO)<sub>8</sub>], a tungstendioiron cluster derived from [Fe<sub>2</sub>(CO)<sub>9</sub>] and [NEt<sub>4</sub>]<sub>2</sub>W(≡CC≡C<sup>1</sup>Bu)(CO)<sub>2</sub>(η<sup>5</sup>-C<sub>2</sub>B<sub>9</sub>H<sub>9</sub>Me<sub>2</sub>) (N. Carr, S.J. Dossett, F.G.A. Stone), (413) 223
- Deuterated sandwich- and piano-stool organo-iron complexes and polyalkylbenzenes, syntheses of (J.-R. Hamon, P. Hamon, S. Sinbandhit, P. Guenot, D. Astruc), (413) 243
- Ferrocenyl(alkoxy)methylium cations, generation and isolation of salts of; their intermediacy in acid-promoted acetal hydrolysis (T.P.E. Kenny, A.C. Knipe, W.E. Watts), (413) 257
- 4-Alkoxy-4'-cyanobiphenyl esters of ferrocene dicarboxylic acid, preparation and thermal properties of (J. Bhatt, B.M. Fung, K.M. Nicholas), (413) 263
- (η<sup>4</sup>-Diene)Fe(CO)<sub>2</sub>L complexes, evidence for correlated rotation and diastereoisomer discrimination in; crystal and solution structures of (cyclohexadiene)Fe(CO)<sub>2</sub>P(*o*-tolyl)<sub>3</sub> and (cycloheptadiene)Fe(CO)<sub>2</sub>(+)-PPh<sub>2</sub>(neomenthyl) (J.A.S. Howell, M.G. Palin, M.-C. Tirvengadum, D. Cunningham, P. McArdle, Z. Goldschmidt, H.E. Gottlieb), (413) 269
- Stacked face-to-face ferrocene polymer, segment of; crystal and molecular structure of (B.M. Foxman, D.A. Gronbeck, M. Rosenblum), (413) 287
- Chiral ferrocenylbisphosphines containing an aminoalkyl side chain, preparation of (T. Hayashi, A. Yamazaki), (413) 295

- Three-membered metallacycle  $\text{Cp}_3\text{Rh}_3(\mu\text{-CO})_3$ , cluster and benzenoid system in reactions with 12-electron particles  $\text{PPh}_3\text{M}^+$  ( $\text{M} = \text{Cu, Ag, Au}$ ) and  $\text{CpM}^+$  ( $\text{M} = \text{Fe, Ru}$ ) (M.I. Rybinskaya, A.R. Kudinov, D.V. Muratov, N.P. Gambaryan, I.V. Stankevich, A.L. Chistyakov), (413) 419
- Acetylenes and diazoketones in organic synthesis (F. Serratosa), (413) 445
- $(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4)\text{CH}_2(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)$ , Friedel-Crafts cynichrodenylation of; crystal structures of  $(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4)\text{CH}_2(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)\text{C}(\text{O})(\eta^5\text{-C}_5\text{H}_4)\text{Cr}(\text{CO})_2(\text{NO})$  and  $(\text{CO})_2(\text{NO})\text{Cr}(\eta^5\text{-C}_5\text{H}_4)\text{CH}_2(\eta^5\text{-C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)\text{C}(\text{O})(\eta^5\text{-C}_5\text{H}_4)\text{Cr}(\text{CO})_2(\text{NO})$  (Y.-P. Wang, J.-M. Hwu, S.-L. Wang, Y.-J. Wu), (414) 33
- Acetylstyrene- and benzoylstyrene-tricarbonylchromium, and 1-acetyl- and 1-benzoyl-2-ferrocenylethylene, preparation of (J. Huang, Y.-Z. Huang), (414) 49
- $[\text{Fe}_4(\text{CO})_{10}(\text{PR})_2(\mu_2\text{-COEt})]$ , functionalisation of the CO group of an anionic radical cluster (T. Feng, P. Lau, W. Imhof, G. Huttner), (414) 89
- $[(\eta\text{-C}_5\text{R}_5)\text{M}^+]$  fragments ( $\text{M} = \text{Fe, R} = \text{H}; \text{M} = \text{Ru, R} = \text{H, Me}$ ) in synthesis of cationic clusters; crystal structure of  $[\text{RuRh}_3(\eta\text{-C}_5\text{Me}_5)(\eta\text{-C}_5\text{H}_5)_3(\mu_3\text{-CO})_3]\text{PF}_6$  (A.R. Kudinov, D.V. Muratov, M.I. Rybinskaya, P.V. Petrovskii, A.V. Mironov, T.V. Timofeeva, Yu.L. Slovokhotov, Yu.T. Struchkov), (414) 97
- Anionic alkenyl-bridged diiron complexes, new routes to; their reactions with propargyl chloride (R. Yáñez, J. Ros, R. Mathieu), (414) 209
- $(\mu\text{-tert-Butylthio})(\mu\text{-methylarsino})\text{hexacarbonyldiiron}$  double cluster complex, synthesis and molecular structure of (L.-C. Song, Q.-M. Hu), (414) 219
- Interannular bridged ferrocenophanium salts, physical properties of; structural, NMR, EPR, and Mössbauer studies (T.-Y. Dong, H.-M. Lin, M.-Y. Hwang, T.-Y. Lee, L.-H. Tseng, S.-M. Peng, G.-H. Lee), (414) 227
- Complexed triphosphanes, preparation and characterisation of; crystal structure of  $\text{HP}[\text{PPh}_2\text{Cr}(\text{CO})_5]_2$  (M. Scheer, S. Gremler, E. Herrmann, P.G. Jones), (414) 337
- closo*-Clusters, carbene-ligand assisted formation of; test of a model to clarify the redox activity of *nido*- and *closo*-compounds (B. Eber, G. Huttner, D. Günauer, W. Imhof, L. Zsolnai), (414) 361
- Chiral iron secondary methoxycarbene  $[\text{Fe}(\text{C}_5\text{Me}_5)(\text{CO})(\text{PMcPh}_2)(\text{CHOMe})]\text{SO}_3\text{CF}_3$ , synthesis and characterisation of (M.-J. Tudoret, V. Guerchais, C. Lapinte), (414) 373
- $(\eta^5\text{-C}_5\text{H}_4\text{CH}_3)_4\text{Co}_2\text{Fe}_2(\mu_3\text{-NO})_2$  and trinuclear methylcyclopentadienyl hetero-metal clusters with  $\mu_3\text{-NH}$  and  $\mu_3\text{-NO}$  ligands, synthesis and structures of (J. Müller, I. Sonn, T. Akhnoukh), (414) 381
- Ferrocene; Annual Survey covering the year 1989 (B.W. Rockett, G. Marr), (416) 327
- Tetracarbonyl(1,1',3,3'-tetra-tert-butyl-5,5'-pentafulvalene)diiron, reactions at the iron–iron bond (P. Jutzi, J. Schnittger, J. Dahlhaus, D. Gestmann, H.-C. Leue), (415) 117
- Carbonyl metallate ions, coordinatively unsaturated and electron deficient, chemistry of; gas phase reactions of  $[\text{Fe}(\text{CO})_2]^-$  with hydrocarbons (I.K. Gregor), (415) 257
- $[\text{Fe}_2(\mu\text{-CCN)SMe}_2(-\text{CO})(\text{CO})_2(\text{Cp})_2]\text{SO}_3\text{CF}_3$  ( $\text{Cp} = \eta\text{-C}_5\text{H}_5$ ), reactions with alcohols (L. Busetto, M.C. Cassani, V. Zanotti, V.G. Albano, D. Braga), (415) 395
- Iron(0)-phthalocyanine and bis(dimethylglyoximate)-triphenylphosphinerhodate(I), conversion of 1-bromoalkines (D. Steinborn, U. Sedlak, M. Dargatz), (415) 407
- Di- and tri-phosphollide ions and their ferrocene analogues; synthesis of (R. Bartsch, J.F. Nixon), (415) C15
- $(\eta^3\text{-C}_3\text{H}_5)\text{Fe}(\text{CO})_3\text{X}$  ( $\text{X} = \text{Cl, Br, I}$ ) complexes, electrochemical reduction of; evidence of formation of surface heterometallic compounds (D. Osella, M. Ravera, S.V. Kukharenko, V.V. Strelets, C.E. Housecroft), (417) 421
- Iron and ruthenium complexes with 1,1-difluoroethene (W. Keim, W. Fischer), (417) C20
- Bis(fulvalene)diiron and bis(fulvalene)iron–cobalt monohexafluorophosphates, crystallographic comparison of the cation (P. Brüggeller, P. Jaitner, H. Schottenberger, K.E. Schwarzhan), (417) C53
- Iron sandwich compounds, studies on the anti-tumour activity of (A. Houlton, R.M.G. Roberts, J. Silver), (418) 107
- Gold(I) complexes of 1,1'-bis(diphenylphosphino)ferrocene, studies on (A. Houlton, R.M.G. Roberts, J. Silver, R.V. Parish), (418) 269
- Unsaturated mixed metal clusters  $\text{Fe}_2\text{Ru}_2(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})(\mu_4\text{-Te})$  and  $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})(\mu_4\text{-Te})$ , construction of (P. Mathur, D. Chakrabarty, Md. M. Hossain), (418) 415
- $[\eta^5\text{-Indenyl}\text{Fe}(\text{CO})_3]\text{BF}_4$ , evidence for  $\eta^3$ -indenyl intermediates (H. Ahmed, D.A. Brown, N.J. Fitzpatrick, W.K. Glass), (418) C14

- Opening of cluster frameworks, increase in skeletal electrons via amino substituents (B. Eber, G. Huttner, C. Emmerich, J.C. Daran, B. Heim and Y. Jeannin), (419) 43
- Tricarbonyliron complex of tricyclo[5.3.1.0<sup>4,9</sup>]undeca-2,5-diene, a strongly homo-conjugated nonparallel 1,4-diene; preparation, structure, and reaction of (N. Sakai, K. Mashima, H. Takaya, R. Yamaguchi and S. Kozima), (419) 181
- Alkyl-Fp [Fp = (C<sub>5</sub>H<sub>5</sub>)Fe(CO)<sub>2</sub>] complexes, preparation via organometallic reagents; reactions of RMgX and RLi with Fp-I and Fp-THF<sup>+</sup>BF<sub>4</sub><sup>-</sup> (H.-j. Li and M.M. Turnbull), (419) 245
- Tricarbonyl(benzene)chromium and ferrocene containing unsaturated amides and nitriles, preparation of (J. Huang, H.-X. Jiang and Y.-Z. Huang), (419) 337
- Tricarbonyl(η<sup>1</sup>,η<sup>2</sup>-but-3-en-1-yl)iron(0) anion complexes, bromination and alkylation of (M.-C.P. Yeh and C.-C. Hwu), (419) 341
- Cyclopentadienyliron-mediated introduction of functionalized alkyl or alkynyl groups to arenes in an addition-demetallation sequence (R.G. Sutherland, C. Zhang and A. Piórko), (419) 357
- Tricarbonyliron complexes, regioselective access to; controlled preparation and reactions of trifluoromethyl-substituted complexes (G.R. Stephenson, P.W. Howard and S.C. Taylor), (419) C14
- [(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)Fe(CO)<sub>3</sub>]BF<sub>4</sub>, direct observation of ion pair formation in dichloromethane solution and the crystal structure of (P. McArdle, D. MacHale, D. Cunningham and A.R. Manning), (419) C18
- Heteropolymetallic complexes of 1,1'-bis(diphenylphosphino)ferrocene (dppf). VII. Redox behaviour of dppf (G. Pilloni, B. Longato and B. Corain), (420) 57
- Ultraviolet vapour-phase absorption spectra of *d*<sup>6</sup> metallocenes (S.Y. Ketkov and G.A. Domrachev), (420) 67
- Mixed chalcogen carbonyl compounds. IV. Reactivity of Fe<sub>2</sub>(CO)<sub>6</sub>(μ-STe) and Fe<sub>3</sub>(CO)<sub>9</sub>(μ<sub>3</sub>-S)(μ<sub>3</sub>-Te) towards coordinatively unsaturated species (P. Mathur, D. Chakrabarty, M.M. Hossain and R.S. Rashid), (420) 79
- Syntheses of dibenzo[*b,e*][1,4]dioxin derivatives via iron complexes, and further functionalizations via directed metallation (R.C. Cambie, S.J. Janssen, P.S. Rutledge and P.D. Woodgate), (420) 387
- Kohlenwasserstoffverbrückte Komplexe. XXIII. Heterobimetallische Komplexe mit Ferrocenyl- und (Ph<sub>3</sub>P)(OC)(Cp)FeC(O)CH<sub>2</sub>-Gruppen; Darstellung und Struktur von (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)Fe[η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>C(O)Re(CO)<sub>5</sub>], (PPh<sub>3</sub>)(OC)(Cp)FeC(O)CH<sub>2</sub>(η<sup>6</sup>-C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub> und [η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>C(O)CH<sub>2</sub>(η<sup>6</sup>-C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub>] (M = Cr, Mo) (J. Breimair, M. Wieser, B. Wagner, K. Polborn und W. Beck), (421) 55
- Redox potential and substituent effects at ferrocene derivatives. Estimates of Hammett σ<sub>p</sub> and Taft polar σ<sup>\*</sup> substituent constants (M.E.N.P.R.A. Silva, A.J.L. Pombeiro, J.J.R.F. da Silva, R. Herrmann, N. Deus, T.J. Castilho and M.F.C.G. Silva), (421) 75
- Reactivity of Main-Group-transition-metal bonds. IX. The kinetics of iodination of compounds containing two or more tin-transition-metal bonds (J.R. Chipperfield, S. Clark, D.E. Webster and H. Yusof), (421) 205
- Preparation and spectroscopic properties of a series of new bimetallic phosphine-bridged seven-coordinate complexes of the type [M<sub>2</sub>I<sub>4</sub>(CO)<sub>6</sub>L<sub>2</sub>(μ-L̄L̄)] (M = Mo or W; L = PPh<sub>3</sub>, AsPh<sub>3</sub> or SbPh<sub>3</sub>; L̄L̄ = Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub> (*n* = 1, 2 or 4); L̄L̄ = [Fe(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>]) (P.K. Baker, M. van Kampen and D. ap Kendrick), (421) 241
- Dialkylaminophosphorus metal carbonyls. X. Phosphorus-bridging carbonyl expulsion in reactions of aldehydes and ketones with (iPr<sub>2</sub>NP)<sub>2</sub>COFe<sub>2</sub>(CO)<sub>6</sub> (R.B. King, N.K. Bhattacharyya and E.M. Holt), (421) 247
- Synthesis of tetracarbonyl(η<sup>3</sup>-2-hydroxypropenyl)iron complexes (M. Frey and T.A. Jenny), (421) 257
- Bis-ferrocene derivatives of bridged calix[4]arenes: synthesis, X-ray crystal structure and electrochemical properties (P.D. Beer, A.D. Keefe, V. Böhmer, H. Goldmann, W. Vogt, S. Lecocq and M. Perrin), (421) 265
- Charge Transfer-Komplexe des Octaethylferrocens (D. Stein, H. Sitzmann und R. Boese), (421) 275
- Heterometallic analogues of [(M(CO)<sub>2</sub>(η-C<sub>5</sub>H<sub>5</sub>))<sub>2</sub>] (M = Fe, Ru, Os). The synthesis of the tetracarbonylcyclopentadienylcycloheptatrienyl complexes [M(CO)<sub>2</sub>(η-C<sub>5</sub>H<sub>5</sub>)M'(CO)<sub>2</sub>(η-C<sub>7</sub>H<sub>7</sub>)] (M = Fe, Ru; M' = Mo, W). Crystal structures of [Ru(CO)<sub>2</sub>(η-C<sub>5</sub>H<sub>5</sub>)Mo(CO)<sub>2</sub>(η-C<sub>7</sub>H<sub>7</sub>)] (Ru-Mo) and [Mo(CO)<sub>3</sub>(μ-η<sup>6</sup>, η<sup>1</sup>-C<sub>7</sub>H<sub>7</sub>)Ru(CO)<sub>2</sub>(η-C<sub>5</sub>Me<sub>5</sub>)] (R.L. Beddoes, E.S. Davies, M. Helliwell and M.W. Whitley), (421) 285
- Synthesis and nonlinear optical properties of new bimetallic iron/palladium complexes (G. Doisneau, G. Balavoine, T. Fillebeen-Khan, J.-C. Clinet, J. Delaire, I. Ledoux, R. Loucif and G. Puccetti), (421) 299

- Reactivity of phosphinoalkynes toward  $[HFe_3(CO)_9(C=CH_2)]^-$  (E. Louattani, J. Suades and R. Mathieu), (421) 335
- Carbon-13 spin-lattice relaxation in organometallic complexes (H. Schumann), (421) C7
- Organoiron chemistry; Annual Survey for the year 1990 (R.C. Kerber), (422) 209
- Substitution reactions of dinuclear alkenyl-bridged iron-cobalt complexes with phosphines. Crystal structures of  $[(CO)_3Fe-CO(PMe_2Ph)(\mu-CO_2Me)\equiv C(CO_2Me)H]$  and  $[(CO)_2(PMe_2Ph)Fe-CO(CO_2Ph)(\mu-CO_2Me)\equiv C(CO_2Me)H]$  (I. Moldes, J. Ros, R. Mathieu, X. Solans and M. Font-Bardía), (423) 65
- Cluster chemistry. LXXVII. Carbide clusters from  $Ru_5(\mu_5-C_2PPh_2)-(\mu-PPh_2)(CO)_{13}$  and  $Fe_2(CO)_9$ : X-ray structures of  $MRu_4(\mu_5-C(\mu-dppm)(CO)_{13}\cdot 0.5CH_2Cl_2$  ( $M = Fe$  and  $Ru$ ) and  $FeRu_5(\mu_6-C(\mu-PPh_2)_2(CO)_{13}(PMe_2Ph)$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (423) 105
- Rapid  $[SR]^+$  exchange between metal-thiolate and metal-organic disulphide complexes (P.M. Treichel and E.K. Rublein), (423) 391
- Dodecacarbonyl triiron, an efficient catalyst for photochemical isomerization of unsaturated alcohols, ethers and ester to their corresponding carbonyl compounds, enol ethers and esters (N. Iranpoor and E. Mottaghinejad), (423) 399
- Syntheses and properties of isocyanide complexes of iron,  $trans-[FeH(CNR)(Ph_2PCH_2CH_2PPh_2)_2]A$  ( $A = BF_4$  or  $PF_6^-$ ) (M.B. Baptista, M.A.N.D.A. Lemos, J.J.R. Fraústo da Silva and A.J.L. Pombeiro), (424) 49
- Synthesis and crystal structures of the isoelectronic monocyclopentadienyliron complexes  $[\eta^5-C_5H_5]Fe(P(OMe)_3)(CO)I$  and  $[(\eta^5-C_5H_5)Fe(P(OMe)_3)(NO)I]BF_4^-$  (J.E. King and S.J. Simpson), (424) 57
- Use of  $[Me_2SSMe]BF_4$  to prepare thiolate complexes of iron and ruthenium (P.M. Treichel and E.K. Rublein), (424) 71
- Reactions of acyl(carbonyl)ruthenium(II) and acyl(carbonyl)iron(II) complexes with amines. Preparation of novel propionyl(carbamoyl)ruthenium(II) complexes (F. Ozawa, S. Chikaoka, X.-Z. Cao and A. Yamamoto), (424) 173
- Flash vacuum pyrolysis study of iron cyclopentadienyl complexes and the crystal structure of *O*-acetyl-*N*-(1-ferrocenylpropylidene)hydroxylamine (J.C. Barnes, W. Bell, C. Glidewell and R.A. Howie), (424) 343
- Synthesis and some reactions of ferrocenylacetylenes (G. Doisneau, G. Balavoine and T. Fillebeen-Khan), (425) 113
- Enantioselektive Katalysen. LXXVIII. Der optisch aktive Phosphido-verbrückte Enantioselektive Katalysen Eisenkomplex  $[(\eta^5-C_5H_5)_2Fe_2(\mu-H)(\mu-PMen_2)(CO)_2]$  — Synthese, Isomerisierung, Katalysen (H. Brunner and M. Rötzer), (425) 119
- Enantioselective reduction of benzoyl derivatives of ferrocene and ruthenocene; optically active carbinols and carbenium ions (M.N. Nefedova, I.A. Mamedyarova, P.P. Petrovski and V.I. Sokolov), (425) 125

## Lanthanides

- Lanthanides and actinides; Annual Survey covering the years 1984–1986 (R.D. Rogers, L.M. Rogers), (416) 201

## Lanthanum

- Tricyclopentadienyllanthanides/sodium hydride, reaction with organic halides (C. Qian, D. Zhu, Y. Gu), (401) 23
- Tris( $\eta^5$ -cyclopentadienyl)neodymium(III), neutral base adducts of; interpretation of their optical magnetochemical, ESR and NMR properties (H. Reddmann, H. Schultze, H.-D. Amberger, G.V. Shalimoff, N.M. Edelstein), (411) 331
- $(MeCp)_3La$  ( $MeCp = CH_3C_5H_4$ ), a tetrameric complex of the type  $[(MeCp)_3La]_4$ , synthesis and molecular structure of (Z. Xie, F.E. Hahn, C. Qian), (414) C12

## Lead

- ortho*-Palladated benzylamines and acetanilides, comparative study of the mechanism of alkenylation of (A.D. Ryabov, I.K. Sakodinskaya, A.K. Yatsimirsky), (406) 309
- Group 14 triflates, synthesis of (W. Uhlig), (409) 377

- Diphenyllead(IV) dichloride, molecular electronic structure of; comparison with diphenyltin(IV) dichloride by the SCF-MS molecular-orbital method (E.M. Berksoy, M.A. Whitehead), (410) 293
- Paramagnetic organolead complexes; EPR investigation of their formation and structures (H.B. Stegmann, M. Schenkl, K. Scheffler), (414) 145
- p*-Tol<sub>6</sub>Sn<sub>2</sub>, *p*-Tol<sub>6</sub>PbSn, and *p*-Tol<sub>6</sub>Pb<sub>2</sub>, hexa-*p*-tolylethane derivatives; synthesis, homoeotype structures, NMR couplings, and valence vibrations (C. Schneider, M. Dräger), (415) 349
- On carbon-13 spectra of thio- and dithiocarboxylic acids triorgano group 14 metal esters (S. Kato, A. Hori, M. Mizuta, T. Katada, H. Ishihara, K. Fujieda and Y. Ikebe), (420) 13

## Lithium

- Elementary lithium, reaction with diphenylmethylenecyclobutane; reductive cleavage of a cyclobutane  $\sigma$ -bond and rearrangement with 1,7-proton shift of the 1,4-dilithio compound that is formed primarily (A. Maercker, K.-D. Klein), (401) C1
- Aryllithiumtricarbonylchromium, aromatic nucleophilic substitution of methoxy or chloro groups by (F. Rose-Munch, O. Bellot, L. Mignon, A. Semra, F. Robert, Y. Jeannin), (402) 1
- Lithiation of 3-bromo-3-hexene; synthesis of 1,2,3,4,5-pentaethylcyclopentadiene (D. Stein, H. Sitzmann), (402) C1
- N,N*-Dimethyl-3,5-bis(trifluoromethyl)aniline, regioselective metalation of (N. Kuhn, A. Kuhn, M. Schulten), (402) C41
- [ $(\eta^5\text{-Dimethylphosphino})\text{cyclopentadienyl}][\eta^7\text{-}(dimethylphosphino)\text{cycloheptatrienyl}]$ titanium, heterobimetallic compounds derived from (M. Ogasa, M.D. Rausch, R.D. Rogers), (403) 279
- Titanium, iron and thallium vinylcyclopentadienyl derivatives, synthesis and structure of (M. Ogasa, D.T. Mallin, D.W. Macomber, M.D. Rausch, R.D. Rogers, A.N. Rollins), (405) 41
- Benzyl tricarbonyl chromium allyl and crotyl ethers, 2,3-Wittig rearrangement of (M. Mahmoudi, L. Pelinski, L. Maciejewski, J. Brocard), (405) 93
- Phenyllithium, metallation of substituted pyridine derivatives by (M. Mallet), (406) 49
- (ArBMe<sub>3</sub>)Li·D complexes (Ar = C<sub>6</sub>H<sub>4</sub>CH(X)NMe<sub>2</sub>-2); structure, dynamic molecular processes and the nature of the carbon–lithium bond in the light of multinuclear (<sup>1</sup>H, <sup>13</sup>C, <sup>7</sup>Li, <sup>11</sup>B) NMR measurements at variable temperature (E. Kalbarczyk-Bidelska, S. Pasynkiewicz), (407) 143
- [Li(Et<sub>2</sub>O)<sub>2</sub>·2,4,6-Ph<sub>3</sub>C<sub>6</sub>H<sub>2</sub>] monomeric,  $\sigma$ -bonded aryllithium etherate complex, isolation and X-ray crystal structure of (M.M. Olmstead, P.P. Power), (408) 1
- Butyllithium-induced dimerisation of pent-3-en-1-yne and related additions (P.A.A. Klusener, H. Hommes, J.C. Hanekamp, A.C.H.T.M. van der Kerk, L. Brandsma), (409) 67
- 1,( $\omega$  - 1)-Dienes, solvent controlled unilateral or bilateral metalation of (E. Moret, O. Desponds, M. Schlosser), (409) 83
- Allyl- and benzyl-type organoalkali intermediates, preparation via organotin compounds (O. Desponds, M. Schlosser), (409) 93
- Allyl alkali metals, comparative theoretical study of (N.J.R. van Eikema Hommes, M. Bühl, P.v.R. Schleyer, Y.-D. Wu), (409) 307
- Group 14 triflates, synthesis of (W. Uhlig), (409) 377
- Organolithium and -magnesium compounds, conjugated addition to  $\alpha$ -ethylenic mesitylketones (F. Barbot, D. N'Goma, Ph. Miginiac), (410) 277
- 1,6-Dilithio-3-hexyne, preparation by cleavage of 1,1'-bicyclopropylidene with lithium powder; isolation of the intermediate (1,3-dilithiopropylidene)cyclopropane (A. Maercker, K.-D. Klein), (410) C35
- cis-Bis(tetrahydrothiophene)bis( $\eta^2$ -biphenyl-2,2'-diyl)platinum(IV) complexes, synthesis of (H.-A. Brune, H. Roth, G. Schmidberg), (412) 237
- sym*-Disubstituted pyrazines, lithiation of (A. Turck, D. Trohay, L. Mojovic, N. Plé, G. Quéguiner), (412) 301
- Oligonuclear ferrocene derivatives with chalcogene bridges (M. Herberhold, H.-D. Brendel, O. Nuyken, T. Pöhlmann), (413) 65
- Lithium-coordinated phosphinomethanides, influence of Co-ligands upon; crystal structure of ((THF)Li[C(SiMe<sub>3</sub>)<sub>2</sub>(PMe<sub>2</sub>)])<sub>2</sub> (H.H. Karsch, K. Zellner, S. Gamper, G. Müller), (414) C39
- Alkylarenes, optional *ortho* or alpha hydroxymethylation (Y. Guggisberg, F. Faigl, M. Schlosser), (415) 1
- 3,3'-Di-tert-butyltitanocene dichloride, tetramethylethanobridged; stereoselectivity in the synthesis of (M.S. Erickson, F.R. Fronczek, M.L. McLaughlin), (415) 75

- (ArBMe<sub>3</sub>)Li·OEt<sub>2</sub>OEt<sub>2</sub> complexes, studies on reactions with the electron donors OEt<sub>2</sub> and THF by means of <sup>1</sup>H, <sup>7</sup>Li, <sup>11</sup>B, and <sup>13</sup>C NMR spectroscopy (E. Kalbarczyk-Bidelska, S. Pasynkiewicz), (417) 1
- Alkali metal derivatives of sulfinimidamides, preparation and crystal structures (F. Pauer, D. Stalke), (418) 127
- (Alkylthio)fluorobenzenes, regioselective metalation of (S. Cabiddu, C. Fattuoni, C. Floris, G. Gelli and S. Melis), (419) 1
- The structure of 2-(diphenylphosphino)phenyllithium: the significance of P-Li bonding (S. Harder, L. Brandsma, J.A. Kanters, A. Duisenberg and J.H. van Lenthe), (420) 143
- The reaction of arylcoppermagnesium and other organometallic reagents with phthalic anhydride (M.T. Rahman and S.K. Nahar), (425) 201

## Lutetium

- Tris(2,4-dimethylpentadienyl)lutetium, synthesis and crystal structure of (H. Schumann, A. Dietrich), (401) C33
- Organo rare earth hydrides, synthesis and reactivity of (C. Ye, C. Qian, X. Yang), (407) 329
- Organolanthanide hydrides, [O(CH<sub>2</sub>CH<sub>2</sub>C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>LnH]<sub>2</sub> (Ln = Gd, Er, Yb, Lu, Y); synthesis and reactivity towards alkenes, alkynes and organic halides (Z. Xie, C. Qian, Y. Huang), (412) 61
- Alumohydride group, change of bonding mode in biscyclopentadienylhydrido REM complexes from heterometallic to homometallic; crystal and molecular structures of [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Yb(μ<sub>3</sub>-H)]<sub>2</sub>[(μ<sub>2</sub>-H)AlH<sub>2</sub>·N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>]<sub>2</sub>·C<sub>6</sub>H<sub>6</sub>, [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Lu(μ<sub>2</sub>-H)]<sub>2</sub>[(μ<sub>2</sub>-H)<sub>2</sub>AlH·N(C<sub>2</sub>H<sub>5</sub>)<sub>3</sub>]<sub>2</sub>·C<sub>6</sub>H<sub>6</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Lu]<sub>3</sub>(μ<sub>2</sub>-H)<sub>2</sub>(μ<sub>3</sub>-H) (S.Ya. Knjazhanskij, B.M. Bulychev, O.K. Kireeva, V.K. Belsky, G.L. Soloveichik), (414) 11
- Homo- and heteronuclear hydrido lutetocene complexes with bulky 1,3-di(tert-butyl)cyclopentadienyl (C<sub>5</sub>H<sub>3</sub><sup>t</sup>Bu<sub>2</sub>) ligands, molecular structure of [(<sup>t</sup>Bu<sub>2</sub>C<sub>5</sub>H<sub>3</sub>)LuH]<sub>4</sub>[AlH<sub>4</sub>·Et<sub>2</sub>O]<sub>2</sub>[AlH<sub>4</sub>]<sub>2</sub> (S.Y. Knjazhanski, E.B. Lobkovsky, B.M. Bulychev, V.K. Belsky and G.L. Soloveichik), (419) 311

## Magnesium

- Titanocene-catalysed hydromagnesation of silylacetylenes, reversibility of (M.A. Djadchenko, K.K. Pivnitsky, J. Spanig, H. Schick), (401) 1
- Arylmagnesium compounds, intramolecularly coordinated; effects on the Schlenk equilibrium (P.R. Markies, R.M. Altink, A. Villena, O.S. Akkerman, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek), (402) 289
- 1,*n*-Dicarbonyl compounds, one-step synthesis from carboxylic acid derivatives and di-Grignard reagents in the presence of transition metal catalysts (F. Babudri, A. D'Ettore, V. Fiandanese, G. Marchese, F. Naso), (405) 53
- Non-Grignard organomagnesium reagents obtained from 1,3-dienes, synthesis and transformations of (U.M. Dzhemilev, A.G. Ibragimov, G.A. Tolstikov), (406) 1
- Organomagnesium reagents in the regioselective synthesis of 2-(1,1-dialkylprop-2-enyl)-4,5-dihydro-thiazoles and oxazoles, precursors of α,α-disubstituted β-ethylenic carboxylic acids (J. Laduranty, F. Barbot, L. Miginiac), (406) 57
- Ru<sup>II</sup>-alkyl-butadiene complexes, formation from dihalogeno-Ru<sup>IV</sup>-allyl precursors by the reaction with BrMg(CH<sub>2</sub>)<sub>4</sub>MgBr; decomposition pathway for ruthenium(IV)cyclopentane intermediates (H. Nagashima, Y. Michino, K. Ara, T. Fukahori, K. Itoh), (406) 189
- Pentamethylcyclopentadienyl Grignard reagent, [Cp<sup>\*</sup>Mg(thf)μ-Cl]<sub>2</sub>, crystal and molecular structure of (R.E. Cramer, P.N. Richmann, J.W. Gilje), (408) 131
- Phenylcoppermagnesium and methylcoppermagnesium reagents, <sup>1</sup>H and <sup>13</sup>C NMR spectra of (S.K. Nahar, H.P. Nur, M.T. Rahman, M. Nilsson, T. Olsson), (408) 261
- Organolithium and -magnesium compounds, conjugated addition to α-ethylenic mesitylketones (F. Barbot, D. N'Goma, Ph. Miginiac), (410) 277
- Dimeric [bis(*ortho*-anisyl)magnesium·THF]<sub>2</sub>, crystal structure of (P.R. Markies, G. Schat, A. Villena, O.S. Akkerman, F. Bickelhaupt, W.J.J. Smeets, A.L. Spek), (411) 291
- Magnesium bromide and iodide in diethyl ether and tetrahydrofuran; extended X-ray absorption fine structure (EXAFS) and large angle X-ray scattering (LAXS) study (A. Wellmar, I. Persson), (415) 143

- Organomagnesium iodides in diethyl ether; large angle scattering (LAXS) study (A. Wellmar, I. Persson), (415) 155
- Organomagnesium bromides in diethyl ether; extended X-ray absorption fine structure (EXAFS) and large angle X-ray scattering (LAXS) study (A. Wellmar, A. Hallberg, I. Persson), (415) 167
- Nickel complexes containing (*R*, *R*)-1,2-cyclopentanediyldibis(diphenylphosphine) and related ligands, enantioselective cross-coupling of vinyl-, aryl and allyl-electrophiles catalyzed by (G. Consiglio, A. Indolese), (417) C36
- Synthèse en une étape par voie organométallique d'amines primaires  $\beta$ -acétyléniques  $\alpha$ -ramifiées (L. Leboutet, G. Courtois and L. Miginiac), (420) 155
- Heterocyclen als Liganden. XII. Synthese und Struktur von  $(C_{28}H_{40}N)MgC_2H_5 \cdot 2C_4H_8O$ —Stabilisierung eines monomeren Organylmagnesiumamids durch sterische Überfrachtung (N. Kuhn, M. Schulten, R. Boese und D. Bläser), (421) 1
- The reaction of arylcoppermagnesium and other organometallic reagents with phthalic anhydride (M.T. Rahman and S.K. Nahar), (425) 201

## Manganese

- $M_2(CO)_{10}$  ( $M = Mn, Re$ ), mechanisms of the photochemical reactions with *ortho*- and *para*-quinones; a combined EPR and FTIR study (T. van der Graaf, D.J. Stukens, J. Víčová, A. Vlček, Jr.), (401) 305
- Linked metal acetylide complexes for nonlinear optics, synthesis of (S.J. Davies, B.F.G. Johnson, J. Lewis, M.S. Khan), (401) C43
- $\eta^5-C_5H_4CH_2CH(CH_3)N(CH_3)(C_3H_5)Mn(CO)_2$ , synthesis of a novel bidentate complex of (T.-F. Wang, T.-Y. Lee, Y.-S. Wen, L.-K. Liu), (403) 353
- Manganese cyclopentadienyldicarbonyl complexes of 1,2,3-selenadiazoles and thiadiazoles; structural comparison of free and complexed 1,2,3-benzothiadiazole and 4-phenyl-1,2,3-thiadiazole (A.J. Mayr, B. Carrasco-Flores, F. Cervantes-Lee, K.H. Pannell, L. Párkányi, K. RaghuVeer), (405) 309
- Tricarbonyl- $\eta^5$ -2,4-pentadienylmanganese, [4+5]-cycloaddition of conjugated dienes to (C.G. Kreiter, K. Lehr), (406) 159
- Vinylidene complexes, nucleophilic behavior of (F. Delbecq), (406) 171
- Heterometallic  $n,\pi$ -complexes of transition metals based on mesitylacetonitrile (V.S. Kaganovich, M.I. Rybinskaya), (407) 215
- Hydroxycarbene complex  $[(DPPP)(CO)_3Mn=C(OH)CH_3]^+[CF_3SO_3]^-$ , {*facial*-[1,3-bis(diphenylphosphino)propane(methylhydroxycarbene)tricarbonylmanganese(I)]} triflate; X-ray structure of (P.L. Motz, D.M. Ho, M. Orchin), (407) 259
- Tetrahydrofurfurylcyclopentadiene and its thallium, manganese, and rhodium complexes, synthesis of; low temperature matrix photolysis of the manganese complex (T.E. Bitterwolf, K.A. Lott, A.J. Rest), (408) 137
- $M^+[M'M''(CO)_6L]^-$  ( $M^+ = Na^+$ , PPN $^+$ ;  $M' = Cr, W$ ;  $M'' = Mn, Re$ ;  $L = CO, PR_3$ ), kinetic studies of the cleavage of (Y.K. Park, S.J. Kim, J.H. Kim, I.S. Han, C.H. Lee, H.S. Choi), (408) 193
- Triorganostannyldiazo acetates, reactions with manganese pentacarbonyl bromide (H. Kandler, H.W. Bosch, V. Shklover, H. Berke), (409) 233
- Metal carbonyl anions,  $I^+$ -abstraction versus  $I^-$ -displacement in their reactions with diiodoacetylene; X-ray structure of  $[(OC)_5MnC\equiv CMn(CO)_5]$  (J.A. Davies, M. El-Ghanam, A.A. Pinkerton, D.A. Smith), (409) 367
- Carbonyl- $\eta^4$ -dien- $\eta^5$ -2,4-dimethyl-2,4-pentadienyl-manganese complexes, photochemical reactions with olefins (C.G. Kreiter, K. Lehr, R. Exner), (411) 225
- Complexes with two  $-SMn(CO)_2Cp$  centres, separated radicals or S-S bonds (H. Braunwarth, P. Lau, G. Huttner, M. Minelli, D. Günauer, L. Zsolnai, I. Jibril, K. Evertz), (411) 383
- $Fe_3(CO)_{11}^{2-}$  and  $Fe_4(CO)_{13}^{2-}$ , transformation to give trinuclear clusters (W. Deck, A.K. Powell, H. Vahrenkamp), (411) 431
- $(CO)_n(L)Mn^I$  complexes ( $L = S_2CPCy_3$  or diphenylphosphine), ring opening and closing in complexes containing the  $Mn(\mu-L)AuC_6F_5$  entity (D. Miguel, V. Riera, J.A. Miguel), (412) 127
- $\eta^6$ -Arenetricarbonylmanganese cations, nucleophilic addition to; role of intermediates in reactions with phosphines (D.A. Brown, W.K. Glass, K.M. Kreddan), (413) 233
- ( $\eta$ -Methylcyclopentadienyl)manganese tricarbonyl, reactions with primary amines (S.C. Srivastava, A.K. Shrimal, A. Srivastava), (414) 65

- $[(\text{Cp}'(\text{CO})_2\text{Mn})_2\text{SnHal}]^-$ , trigonal-planar coordinated  $\text{Sn}^0$  in; "inidene" type tin compounds (F. Ettel, G. Huttner, L. Zsolnai, C. Emmerich), (414) 71
- 1,3-Propanediylbis(pentacarbonylmanganese), preparation via the bistriflate method (E. Lindner, M. Pabel), (414) C19
- Manganese, technetium, and rhenium; Annual Survey covering the year 1987 (P.M. Treichel), (416) 291
- Metal carbonyl dimers, new mechanism for substitution reactions of (B.F.G. Johnson), (415) 109
- $\eta^6$ -Arenetricarbonylmanganese cations, reactions with (D.A. Brown, W.K. Glass, K.M. Kreddan, D. Cunningham, P.A. McArdle, T. Higgins), (418) 91
- $[(2,4,6\text{-}^1\text{Bu}_3\text{C}_6\text{H}_2\text{O})(^6\text{C}_5\text{Me}_5)]\text{P}=\text{Mn}(\text{CO})_4$ , a stable  $\lambda^4$ -phosphanediyi complex containing a multiple phosphor-manganese bond (H. Lang, M. Leise, C. Emmerich), (418) C9
- Group VI metal carbonyls,  $(\eta^6\text{C}_6\text{H}_6)\text{Cr}(\text{CO})_3$ ,  $(\eta^5\text{C}_5\text{H}_5)\text{Mn}(\text{CO})_3$ ,  $(\eta^5\text{CH}_3\text{C}_5\text{H}_4)\text{Mn}(\text{CO})_3$ , and  $(\eta^5\text{C}_5\text{H}_5)\text{Re}(\text{CO})_3$ , photolysis in Nujol at 77 K (T.E. Bitterwolf, K.A. Lott, A.J. Rest and J. Mascetti), (419) 113
- Unsaturated dihydrides  $[\text{Mn}_2(\mu\text{-H})_2(\text{CO})_6(\mu\text{-L}_2)]$  ( $\text{L}_2 = (\text{EtO})_2\text{POP}(\text{OEt})_2$ ,  $\text{Ph}_2\text{PCH}_2\text{PPh}_2$ ), reactions with phosphines containing P-H bonds (R. Carreño, V. Riera and M.A. Ruiz), (419) 163
- Silylphosphinites and their complexation to manganese, crystal structure of *cis*- $\text{MnBr}(\text{CO})_4\text{P}(4\text{-CH}_3\text{C}_6\text{H}_4)_2\text{OSiMe}_3$  (K.M. Cooke, T.P. Kee, A.L. Langton and M. Thornton-Pett), (419) 171
- A study of  $\eta^3\text{-allyl}(\eta^5\text{-cyclopentadienyl})$ -dicarbonylmanganese salts (G.R. Knox, P.L. Pauson and J. Rooney), (420) 379
- Novel heterobimetallic complexes with  $\text{S}_2\text{CPR}_3(\kappa\text{-S},\text{S}'\text{)(}\kappa\text{-S,C},\text{S}')$  bridges. X-Ray structure of  $[\text{MnMo}(\text{CO})_6(\mu\text{-Br})(\mu\text{-S}_2\text{CP}^1\text{Pr}_3)]$  (D. Miguel, J.A. Pérez-Martínez, V. Riera and S. García-Granda), (420) C12
- Reactivity of Main-Group-transition-metal bonds. IX. The kinetics of iodination of compounds containing two or more tin-transition-metal bonds (J.R. Chipperfield, S. Clark, D.E. Webster and H. Yusof), (421) 205
- Carbon-13 spin-lattice relaxation in organometallic complexes (H. Schumann), (421) C7
- Unexpected formation of hydrido-hydroxy derivatives as direct products in the reactions of the unsaturated dihydrides  $[\text{Mn}_2(\mu\text{-H})_2(\text{CO})_6(\mu\text{-L-L})]$  ( $\text{L-L} = \text{Ph}_2\text{PCH}_2\text{PPh}_2$ ,  $(\text{EtO})_2\text{POP}(\text{OEt})_2$ ) with carbon dioxide (F.J. García Alonso, M. García Sanz and V. Riera), (421) C12
- Novel class of transition metal coordination compounds with macrocyclic organosiloxanolate ligands; their synthesis and crystal structure (V.A. Igonin, O.I. Shchegolikhina, S.V. Lindeman, M.M. Levitsky, Yu.T. Struchkov and A.A. Zhdanov), (423) 351
- Übergangsmetall-Heteroallen-Komplexe. XXIV. Synthese und Struktur von Dicarbonylcyclopenta-dienyl( $\eta^2(\text{C},\text{C})$ -Ketenimin)mangan (W. Ziegler und U. Behrens), (423) C16
- Synthesis and characterization of  $\text{M}[\text{Mn}(\text{CO})_4\text{DF}]$  and  $\text{M}[\text{Mn}(\text{CO})_4\text{DM}]$  ( $\text{M} = \text{Li}$ ,  $\text{Na}$ ,  $(\text{PPh}_3)_2\text{N}$ ; DF = dimethyl fumarate, DM = dimethyl maleate) and the role of these complexes in the isomerization of free DM to DF (L.L. Padolik, F. Ungváry and A. Wojciecki), (424) 319
- Organomanganese complexes of podocarpic acid derivatives (R.C. Cambie, M.R. Metzler, C.E.F. Rickard, P.S. Rutledge and P.D. Woodgate), (425) 59
- Temperature and pressure dependence of the addition of 4-cyanopyridine to tricarbonyl(1-5- $\eta$ -diaryl)iron(II) complexes (diaryl =  $\text{C}_6\text{H}_5$  or  $\text{C}_7\text{H}_9$ ) (T.I. Odiaka and R. van Eldik), (425) 89

## Mercury

- Mercuration of n-butyrin, n-valeric and isovaleric acid; crystal structure of  $\text{C}_2\text{H}_5\text{C}(\text{HgCl})_2\text{COOH}\cdot\text{DMSO}$  (B. Korpar-Čolig, Z. Popović, M. Sikirica, D. Grdenić), (405) 59
- Methyl- and phenylmercury(II) diphenyldithiophosphinates, synthesis and structural characterisation of; crystal structure of  $\text{MeHgS}_2\text{PPh}_2$  (J. Zukerman-Schpector, E.M. Vázquez-López, A. Sánchez, J.S. Casas, J. Sordo), (405) 67
- Alkymercury(II) compound-derived ions and radicals, molecular and electronic structures of; an SCF-MO study (C. Glidewell), (405) 141
- $[(\text{HOMe}_2\text{Si})(\text{Me}_3\text{Si})_2\text{Cl}_2\text{Zn}$ ,  $[(\text{MeOMe}_2\text{Si})(\text{Me}_3\text{Si})_2\text{Cl}_2\text{Zn}$ , and  $[(\text{MeOMe}_2\text{Si})(\text{Me}_3\text{Si})_2\text{Cl}_2]\text{Hg}$ , crystal structures of (F.I. Aigbirhio, S.S. Al-Juaid, C. Eaborn, A. Habtemariam, P.B. Hitchcock, J.D. Smith), (405) 149
- $[(\eta^5\text{C}_5\text{H}_5)(\text{CO})_3\text{M(HgX)}]$  ( $\text{M} = \text{Cr}$ ,  $\text{Mo}$ ,  $\text{W}$ ;  $\text{X} = \text{Cl}$ ,  $\text{Br}$ ,  $\text{I}$ ,  $\text{N}_3$ ,  $\text{SCN}$ ) complexes, interaction with  $\text{M}'(\text{CO})_3\text{bipy}$  ( $\text{M}' = \text{Mo}$ ,  $\text{W}$ ) fragments; trimetallic compounds containing  $\text{M}-\text{Hg}-\text{M}'-\text{X}$  arrays (J. Granizo, M.E. Vargas), (408) 357

- $[(\text{Co}_2(\text{CO})_6\text{PhC}_2)_2]$  acetylene-linked dimeric cluster, synthesis and structural characterisation of (B.F.G. Johnson, J. Lewis, P.R. Raithby, D.A. Wilkinson), (408) C9
- Coordinatively unsaturated  $\sigma$ -aryl iridium(III) complexes containing iridium–mercury bonds (W.R. Roper, G.C. Saunders), (409) C19
- trans*-NiI<sub>2</sub>(RNC)<sub>2</sub> complex, unusual reductive behaviour at mercury and platinum electrodes (K. Ehara, K. Kumagaya, Y. Yamamoto, K. Takahashi, H. Yamazaki), (410) C49
- Monomercurated acetic acid; crystal structure of NO<sub>3</sub>HgCH<sub>2</sub>COOH (D. Grdenić, B. Korpar-Čolig, D. Matković-Čalogović, M. Sikirica, Z. Popović), (411) 19
- 1,5,9,13-Tetraselenacyclohexadecane selenium coronand; syntheses, <sup>77</sup>Se CP-MAS solid state NMR spectra and crystal structures of adducts with copper(I) trifluoromethanesulfonate and mercury(II) cyanide (R.J. Batchelor, F.W.B. Einstein, I.D. Gay, J.-H. Gu, B.M. Pinto), (411) 147
- Organic calomels, recent developments in studies of (O.A. Reutov, K.P. Butin), (413) 1
- M–Hg–M (M = Cr, Mo, W) structural unit, unexpected metal–metal bond formation in; synthesis and characterisation of  $[\eta^5\text{-RC}_5\text{H}_4(\text{CO})_3\text{M}]_2\text{Hg}$  complexes (R = H, M = W; R = CH<sub>3</sub>CO, M = Cr, Mo, W) (L.-C. Song, H. Yang, Q. Dong, Q.-M. Hu), (414) 137
- Mercury(II) and thallium(III) organometallic derivatives of 2-mercaptopbenzoxazole (M.V. Castaño, H. Calvo, A. Sánchez, J.S. Casas, J. Sordo, Y.P. Mascarenhas, C. de O.P. Santos), (417) 327
- Trimeric perfluoro-*o*-phenylenemercury, unusual complex with the bromide anion having a polydecker sandwich structure, (V.B. Shur, I.A. Tikhanova, A.I. Yanovsky, Yu.T. Struchkov, P.V. Petrovskii, S.Yu. Panov, G.G. Furin, M.E. Vol'pin), (418) C29
- Oxidative alkylation of  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiR}$  (R = Cl, Me, Et, CH=CH<sub>2</sub>, Ph, OMe, N=C(H)<sup>t</sup>Bu) to  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti(Me)R}$  by group 12 organometallic compounds MMe<sub>2</sub> (G.A. Luinstra and J.H. Teuben), (420) 337
- Mercury–ruthenium mixed-metal carbonyl clusters containing 2-amido-6-methylpyridine (ampy) as a  $\mu_3\text{-}\eta^2$ -ligand. Crystal structures of  $[\text{Ru}_6(\mu_4\text{-Hg})(\mu_3\text{-ampy})_2(\text{CO})_{18}]\cdot 2\text{C}_4\text{H}_8\text{O}$  and  $[\text{Ru}_3(\mu\text{-HgBr})(\mu_3\text{-ampy})(\text{CO})_9]$  (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, C. Bois and Y. Jeannin), (420) 431
- Raman spectroscopic study of the complexation of the methylmercury(II) cation, CH<sub>3</sub>Hg<sup>+</sup>, by ligands containing oxygen and sulphur (C.A. Barradell, H.G.M. Edwards and D.N. Smith), (421) 137
- Investigations on the electronic influence of organyl ligands. IX. *trans*-Influence in mercury organyl compounds. A pseudopotential *ab initio* study (J. Reinhold, N. Steinfeldt, M. Schüler and D. Steinborn), (425) 1

## Metallocenes

- Titanocene-catalysed hydromagnesation of silylacetylenes, reversibility of (M.A. Djadchenko, K.K. Pivnitsky, J. Spanig, H. Schick), (401) 1
- Ferrocene, monosubstituted derivatives of; ferrocene-containing penicillins and cephalosporins (D. Scutaru, L. Tătaru, I. Mazilu, E. Diaconu, T. Lixandru, Cr. Simionescu), (401) 81
- Ferrocene, heterodisubstituted derivatives of; ferrocene-containing penicillins and cephalosporins (D. Scutaru, I. Mazilu, M. Văjă, L. Tătaru, A. Vlase, T. Lixandru, Cr. Simionescu), (401) 87
- Permethylated  $\alpha$ -metallocenylcarbinols (M = Os, Ru, Fe), hydrogen bonds in and conformations of (E.S. Shubina, L.M. Epstein, T.V. Timofeeva, Yu. T. Struchkov, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 133
- Permethylated  $\alpha$ -metallocenylcarbinols in the solid state (M = Fe, Ru, Os), IR spectroscopic study of hydrogen bonds in (E.S. Shubina, L.M. Epstein, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 145
- ( $\eta^6$ -Phenyl)Cr(CO)<sub>3</sub>- and ( $\eta^6$ -phenyl)Co<sub>4</sub>(CO)<sub>9</sub>-substituted metallocenylcarbinols of the iron subgroup; synthesis, structure and hydrogen bonding in (E.S. Shubina, L.M. Epstein, Yu.L. Slovokhotov, A.V. Mironov, Yu.T. Struchkov, V.S. Kaganovich, A.Z. Kreindlin, M.I. Rybinskaya), (401) 155
- Pt(II) complexes of tetrathia[n](1,1')ferrocenophanes, preparation and properties of (M. Sato, H. Asano, S. Akabori), (401) 363
- ansa*-Titanocene and zirconocene complexes with asymmetrically substituted carbon bridges (G.S. Herrmann, H.G. Alt, M.D. Rausch), (401) C5
- Dinuclear ( $\eta^2$ -formaldehyde)zirconocene dimer t-butylisothiocyanate complexes, structure of (G. Erker, M. Mena, C. Krüger, R. Noe), (402) 67
- Cationic titanocene bishexafluoroarsenate complexes (P. Gowik, T. Klapötke), (402) 349

- Zirconium and hafnium hexakis(trimethylsilyl)metallocene dichlorides, hindered cyclopentadienyl rotation in; evidence for complete rotation of the tris(trimethylsilyl)cyclopentadienyl ligands (C.H. Winter, D.A. Dobbs, X.-X. Zhou), (403) 145
- Substituted ferrocenes and ruthenocenes, dynamic NMR studies of ring rotation in (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šík), (403) 195
- Titanium, iron and thallium vinylcyclopentadienyl derivatives, synthesis and structure of (M. Ogasa, D.T. Mallin, D.W. Macomber, M.D. Rausch, R.D. Rogers, A.N. Rollins), (405) 41
- Bis( $\eta^5$ -cyclopentadienyl)methyl(alkoxy)- and bis( $\eta^5$ -cyclopentadienyl)methyl(acyloxy)titanium(IV) complexes, synthesis and reactions of (R. Schobert), (405) 201
- Ferrocenethioles, reactions with norbornadiene and norbornene (M. Herberhold, O. Nuyken, T. Pöhlmann), (405) 217
- Ferrocenecarboxaldoxime, crystal structure of the low-melting form of (G. Ferguson, W. Bell, C. Glidewell), (405) 229
- Iron sandwich complexes and related species, carbon-13 spin-lattice relaxation in (R.M.G. Roberts, J.F. Warmsley), (405) 347
- Ferrocene complexes with cyclodextrins in DMSO, carbon-13 spin-lattice relaxation studies of (R.M.G. Roberts, J.F. Warmsley), (405) 357
- Metallocene-containing cryptands, NMR studies on metal complexes of (C.D. Hall, N.W. Sharpe), (405) 365
- Platinum(II) chloride complexes of 1,1'-bis(methylthio)ferrocene, 1,1'-bis(methylseleno)ferrocene and 1,1'-bis(methylthio)ruthenocene, synthesis and stereodynamics of (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šík), (405) 375
- Titanocene polyselenide complexes, reactions with methyldichloroarsane; synthesis and properties of  $(CH_3C_5H_4)_2Ti(\mu_2-Se_2)_2AsCH_3$  and of  $(CH_3As)_2Se_3$  (B. Holz, R. Steudel), (406) 133
- Zirconocene dichloride/methylalumoxane systems, catalysts for olefin polymerization; IR and NMR studies of (L.A. Nekhaeva, G.N. Bondarenko, S.V. Rykov, A.I. Nekhaev, B.A. Krentsel, V.P. Mar'in, L.I. Vyshinskaya, I.M. Krapova, A.V. Polonskii, N.N. Korneev), (406) 139
- Ferrocene-containing rifamycins, synthesis and characterisation of (D. Scutaru, I. Mazilu, L. Tătaru, M. Văjă, T. Lixandru), (406) 183
- Bisalkynyltitanocene complexes as bidentate chelating ligands (H. Lang, L. Zsolnai), (406) C5
- 1,1'-(1,4,10,13-Tetraoxa-7,16-diazacyclooctadecane-7,16-diyldicarbonyl)ruthenocene, crystal structure of (C.D. Hall, A.W. Parkins, S.C. Nyburg, N.W. Sharpe), (407) 107
- Etheno-bridged *ansa*-titanocene derivatives, syntheses by the McMurry reaction (P. Burger, H.H. Brintzinger), (407) 207
- Cyclopalladated *N,N*-dimethylferrocenecarbothioamide, derivatives of (M. Nonoyama, K. Hamamura), (407) 271
- Unidentate and heterobimetallic 1,1'-bis(diphenylphosphino)ferrocene complexes, X-ray photoelectron spectroscopic differentiation of chemically distinct phosphorus environments in (H.S.O. Chan, T.S.A. Hor, L.-T. Phang, K.L. Tan), (407) 353
- $[Rh(dppf)(\eta^6-C_6H_5)_2B(C_6H_5)_3]$  (dppf = 1,1'-bis(diphenylphosphino)ferrocene), synthesis and crystal structure of (B. Longato, G. Pilloni, R. Graziani, U. Casellato), (407) 369
- Diferrocenylketones, new synthesis of (D.C. O'Connor Salazar, D.O. Cowan), (408) 219
- Diferrocenylsulfide, new synthesis and study of (D.C. O'Connor Salazar, D.O. Cowan), (408) 227
- Silyl-bridged di- to tetranuclear ferrocenes; synthesis, cyclic voltammetry and oxidation of (H. Atzkern, J. Hiermeier, F.H. Köhler, A. Steck), (408) 281
- $Cp_2VCl(AsF_6)$  and other vanadocene(IV) hexafluoropnictogenate species, synthesis and EPR spectroscopic characterisation of; HF-catalysed oxidation to cationic vanadocenium(V) complexes; structure of  $[Cp_2VCl_2]^+[AsF_6]^-$  (F.H. Görlitz, P.K. Gowik, T.M. Klapötke, D. Wang, R. Meier, J. v. Welzen), (408) 343
- Heterodichalcogene ferrocenes with open or bridging chains, synthesis by addition of  $Fe(C_5H_4ELi)_2 \cdot 2THF$  ( $E = S, Se$ ) to propynoates (R. Broussier, M. El Mjidi, B. Gautheron), (408) 381
- ( $\eta^2$ -Phenyltrimethylsilylacetylene)zirconocene generated *in situ*, reactions of (G. Erker, R. Zwettler), (409) 179
- (Ferrocenylacyl)germanes, chemical and structural investigations of (H.K. Sharma, F. Cervantes-Lee, K.H. Pannell), (409) 321
- Bis(alkinyl) titanocene complexes of  $FeCl_2$  and  $CuCl$  (H. Lang, M. Herres, L. Zsolnai, W. Imhof), (409) C7

- $\text{Cp}_2^*\text{Sm}(\text{THF})_2$ , reactions with 1,4-diazadienes and cyclooctatetraene (A. Recknagel, M. Noltemeyer, F.T. Edelmann), (410) 53
- ( $\mu$ -Alkylideneamido)titanocene complex  $[(\text{C}_5\text{H}_5)_2\text{TiCl}]_2[\mu\text{-}\{\text{N}=\text{C}(\text{CH}_2\text{C}_6\text{H}_5)\text{C}(\text{CH}_2\text{C}_6\text{H}_5)=\text{N}\}]$ , preparation and crystal structure of (F. Rehbaum, K.-H. Thiele, S.I. Trojanov), (410) 327
- Disilanes, dehydrogenative polymerisation catalysed by titanocene and zirconocene complexes; novel route to oligo- and poly-silanes (E. Henge, M. Weinberger, Ch. Jammegg), (410) C1
- Heterobimetallic  $\text{Zr},\text{H}(\mu\text{-formaldehyde})$  complex from  $[(\text{CH}_2\text{O})\text{ZrCp}_2]$  dimer and hafnocene dichloride (G. Erker, M. Mena, M. Bendix), (410) C5
- Mixed-valence 1',1''-dibenzylferrocenium cations, micromodulation of electron transfer rates in (T.-Y. Dong, C.-C. Schei, M.-Y. Hwang, T.-Y. Lee), (410) C39
- Bis( $\eta^5$ -pentamethylcyclopentadienyl)zirconium(IV) dialkyls, autoxidations of (J.M. Atkinson, P.B. Brindley), (411) 131
- Bis( $\eta^5$ -cyclopentadienyl)molybdenum(IV) dialkyls, autoxidations of (J.M. Atkinson, P.B. Brindley), (411) 139
- Tris(cyclopentadienyl)technetium(III),  $(\eta^5\text{-C}_5\text{H}_5)_2\text{Tc}(\eta^1\text{-C}_5\text{H}_5)$ , X-ray diffraction study of; dipole moment and charge distribution of  $(\text{C}_5\text{H}_5)_3\text{M}$  ( $\text{M} = \text{V}, \text{Tc}, \text{Re}$ ) (C. Apostolidis, B. Kanellakopulos, R. Maier, J. Rebizant, M.L. Ziegler), (411) 171
- 1,3-Ditellura[3]ferrocenophanes,  $\text{Fe}(\text{C}_5\text{H}_4\text{Te})_2\text{E}$  ( $\text{E} = \text{S}, \text{Se}, \text{Te}$  or  $\text{CH}_2$ ) (M. Herberhold, P. Leitner), (411) 233
- 1,1'-Di-tert-butyl-3,3'-bis(2,2-dimethylpropionyl)ferrocene, diastereoisomeric forms of (W. Bell, C. Glidewell), (411) 251
- Bis(trimethylsilylcyclopentadienyl) complexes of hafnium, synthesis of (J.B. Hoke, E.W. Stern), (412) 77
- Decaethylferrocene, reaction with 2,3,5,6-tetrafluoro-7,7',8,8'-tetracyano-*p*-chinodimethane gives a charge-transfer complex (D. Stein, H. Sitzmann, R. Boese, E. Dormann, H. Winter), (412) 143
- Ferrocene-1,1'-dicarbaldehyde, improved synthesis of (G.G.A. Balavoine, G. Doisneau, T. Fillebeeen-Khan), (412) 381
- Oligonuclear ferrocene derivatives with chalcogene bridges (M. Herberhold, H.-D. Brendel, O. Nuyken, T. Pöhlmann), (413) 65
- 4-Alkoxy-4'-cyanobiphenyl esters of ferrocene dicarboxylic acid, preparation and thermal properties of (J. Bhatt, B.M. Fung, K.M. Nicholas), (413) 263
- Stacked face-to-face ferrocene polymer, segment of; crystal and molecular structure of (B.M. Foxman, D.A. Gronbeck, M. Rosenblum), (413) 287
- Chiral ferrocenylbisphosphines containing an aminoalkyl side chain, preparation of (T. Hayashi, A. Yamazaki), (413) 295
- Acetylstyrene- and benzoylstyrene-tricarbonylchromium, and 1-acetyl- and 1-benzoyl-2-ferrocenylethylene, preparation of (J. Huang, Y.-Z. Huang), (414) 49
- Bis( $\eta^5$ -cyclopentadienyl)dihydridotungsten, insertion reactions with activated olefins and with acetylenes; preparation of tungstenocene alkyl hydrides, olefin and acetylene complexes (G.E. Herberich, W. Barlage, K. Linn), (414) 193
- Interannular bridged ferrocenophanium salts, physical properties of; structural, NMR, EPR, and Mössbauer studies (T.-Y. Dong, H.-M. Lin, M.-Y. Hwang, T.-Y. Lee, L.-H. Tseng, S.-M. Peng, G.-H. Lee), (414) 227
- Substituted  $\eta^4$ -cyclooctadiene-1,5- $\eta^5$ -cyclopentadienyl-cobalt complexes, preparation by reaction of cobaltocenes with alkene cobaltates(1-) (H.-U. Jürgens, J. Alm), (414) C61
- Ferrocene; Annual Survey covering the year 1989 (B.W. Rockett, G. Marr), (416) 327
- 3,3'-Di-tert-butyltitanocene dichloride, tetramethylethanobridged; stereoselectivity in the synthesis of (M.S. Erickson, F.R. Fronczek, M.L. McLaughlin), (415) 75
- $[\text{Fe}_2(\mu\text{-C}(\text{CN})\text{SMc}_2)(-\text{CO})(\text{CO})_2(\text{Cp})_2]\text{SO}_3\text{CF}_3$  ( $\text{Cp} = \eta\text{-C}_5\text{H}_5$ ), reactions with alcohols (L. Busetto, M.C. Cassani, V. Zanotti, V.G. Albano, D. Braga), (415) 395
- Aryl-substituted titanocene and zirconocene complexes with interannular tetramethylethano- or dimethylsilano-bridges; crystal structures of representative examples (P. Burger, K. Hortmann, J. Diebold, H.-H. Brintzinger), (417) 9
- Metal complexed indoles, chemistry and synthetic utility of (U.S. Gill, R.M. Moriarty, Y.Y. Ku, I.R. Butler), (417) 313
- Permethylscandocene complex, investigations of the magnitude of steric and  $\alpha$  deuterium kinetic isotope effects in a carbon–carbon bond-forming reaction of (W.D. Cotter, J.E. Bercaw), (417) C1

- Zirconocene complexes, olefin hydrosilation catalyzed by (M.R. Kesti, M. Abdulrahman, R.M. Waymouth), (417) C12
- Iron sandwich compounds, studies on the anti-tumour activity of (A. Houlton, R.M.G. Roberts, J. Silver), (418) 107
- $\text{Cp}_2\text{Ti}(\mu_2\mu_1\text{-SO}_4)\text{Ti}(\text{H}_2\text{O})\text{Cp}_2$ , synthesis and structure of (U. Thewalt, P. Schinnerling), (418) 191
- Homo- and heteronuclear hydrido lutetium complexes with bulky 1,3-di(tert-butyl)cyclopentadienyl ( $\text{C}_5\text{H}_3^1\text{Bu}_2$ ) ligands, molecular structure of [ $(^1\text{Bu}_2\text{C}_5\text{H}_3)\text{LuH}_4[\text{AlH}_4 \cdot \text{Et}_2\text{O}]_2[\text{AlH}_4]_2$ ] (S.Y. Kniazhanski, E.B. Lobkovsky, B.M. Bulychev, V.K. Belsky and G.L. Soloveichik), (419) 311
- Tricarbonyl(benzene)chromium and ferrocene containing unsaturated amides and nitriles, preparation of (J. Huang, H.-X. Jiang and Y.-Z. Huang), (419) 337
- Heteropolymetallic complexes of 1,1'-bis(diphenylphosphino)ferrocene (dppf). VII. Redox behaviour of dppf (G. Pilloni, B. Longato and B. Corain), (420) 57
- Ultraviolet vapour-phase absorption spectra of  $d^6$  metallocenes (S.Y. Ketkov and G.A. Domrachev), (420) 67
- Acetylen-Komplexe von Di( $\eta^5$ -cyclopentadienyl)molybdän: Synthesen und Protonierung. Struktur von  $[\text{Cp}_2\text{MoH}(\eta^2\text{-C}_2\text{SiMe}_3)_2]\text{PF}_6$  (G.E. Herberich, U. Englert and W. Faßbender), (420) 303
- Kohlenwasserstoffverbrückte Komplexe. XXIII. Heterobimetallische Komplexe mit Ferrocenyl- und ( $\text{Ph}_3\text{P}(\text{OC})(\text{Cp})\text{FeC(O)CH}_2$ -Gruppen; Darstellung und Struktur von ( $\eta^5\text{-C}_5\text{H}_5$ ) $\text{Fe}[\eta^5\text{-C}_5\text{H}_4\text{C(O)CH}_2]$  ( $\text{Re}(\text{CO})_5$ ), ( $\text{PPh}_3(\text{OC})(\text{Cp})\text{FeC(O)CH}_2(\eta^6\text{-C}_7\text{H}_7)$ ) $\text{M}(\text{CO})_3$  und  $[\eta^5\text{-C}_5\text{H}_4\text{C(O)CH}_3]\text{Fe}[\eta^5\text{-C}_5\text{H}_4\text{C(O)CH}_2(\eta^6\text{-C}_7\text{H}_7)]\text{M}(\text{CO})_3$  ( $\text{M} = \text{Cr, Mo}$ ) (J. Breimair, M. Wieser, B. Wagner, K. Polborn und W. Beck), (421) 55
- Redox potential and substituent effects at ferrocene derivatives. Estimates of Hammett  $\sigma_p$  and Taft polar  $\sigma^*$  substituent constants (M.E.N.P.R.A. Silva, A.J.L. Pombeiro, J.J.R.F. da Silva, R. Herrmann, N. Deus, T.J. Castilho and M.F.C.G. Silva), (421) 75
- Preparation of metallacyclic titanocene hydrocarbyl complexes and their use in propene polymerization (G. Erker, U. Korek, R. Petrenz and A.L. Rheingold), (421) 215
- Charge Transfer-Komplexe des Octaethylferrocens (D. Stein, H. Sitzmann and R. Boese), (421) 275
- Unsolvated lanthanidocene hydrides and borohydrides. X-Ray crystal structure of  $[(\eta^5\text{-C}_5\text{H}_3^1\text{Bu}_2)_2\text{Ln}(\mu\text{-H})_2]$  ( $\text{Ln} = \text{Ce, Sm}$ ) and  $[(\eta^5\text{-C}_5\text{H}_3^1\text{Bu}_2)_2\text{Sm}(\mu\text{-BH}_4)]_2$  (Y.K. Gun'ko, B.M. Bulychev, G.L. Soloveichik and V.K. Belsky), (424) 289
- Frontier occupied orbitals in methyl-substituted fulvene and dimethylenecyclopentenyltitanium complexes by UV-photoelectron spectroscopy and EHT calculations (T. Vondrák, K. Mach, V. Varga and A. Terpstra), (425) 27
- Enantioselective reduction of benzoyl derivatives of ferrocene and ruthenocene; optically active carbinols and carbonium ions (M.N. Nefedova, I.A. Mamedyarova, P.P. Petrovski and V.I. Sokolov), (425) 125
- Bis( $\eta^5$ -cyclopentadienyl)tungsten chemistry: hydration reactions of functional alkyl hydrides and olefin complexes (G.E. Herberich and K. Linn), (425) C4

## Molybdenum

- $\text{Mo}(\text{CO})_6$ , photochemical reactions with *o*-semiquinones; EPR studies of the paramagnetic complexes formed (M. Hanaya, M. Iwaizumi), (401) 31
- Transition metal complexes as electrochemical markers for steroids (D. Osella, E. Stein, G. Jaouen, P. Zanello), (401) 37
- Metal carbonyls, rate of O-atom transfer to; effect of substituents on amine *N* oxides on (J.-K. Shen, Y.-C. Gao, Q.-Z. Shi, F. Basolo), (401) 295
- ( $\eta^4$ -Norbornadiene)tetracarbonylchromium and metal VIb hexacarbonyls, chemical shift anisotropy of carbon-13 nuclei in carbonyl groups of (A. Gryff-Keller, H. Krawczyk, P. Szczeciński), (402) 77
- $[\text{Mo}(\text{C}\equiv\text{CPh})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)(\eta\text{-C}_7\text{H}_7)]$  and  $[\text{Mo}(\text{C}\equiv\text{CPh})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)(\eta\text{-C}_7\text{H}_7)][\text{BF}_4^-]$  ( $\text{C}_7\text{H}_7$  = cycloheptatrienyl), X-ray crystal structures of; an investigation of metal-alkynyl bonding in (R.L. Beddoes, C. Bitcon, M.W. Whiteley), (402) 85
- Molybdenum(0) and tungsten(0) ( $\eta^4$ -1-oxa-1,3-diene) complexes; compounds with *R*-(+)-pulegone ligands (T. Schmidt, C. Krüger, P. Betz), (402) 97
- 1,2,3,4,5-Pentaethylcyclopentadiene, synthesis of (D. Stein, H. Sitzmann), (402) 249
- Molybdenum and tungsten cycloheptadienyl complexes, interconversion of  $\eta^5$  and  $\eta^3$  hapticities in (J.R. Hinchliffe, M.W. Whiteley), (402) C50

- $[(\eta^5\text{-}(\text{dimethylphosphino)cyclopentadienyl})\eta^7\text{-}(\text{dimethylphosphino)cycloheptatrienyl}]$ titanium, heterobimetallic compounds derived from (M. Ogasa, M.D. Rausch, R.D. Rogers), (403) 279
- Bis(pentamethylcyclopentadienyl)dioxodimolybdenum(VI) oxide,  $[\text{Cp}^*\text{MoO}_2]_2\text{O}$ , comparison of the solid-state structures of four crystallographically independent molecules of (A.L. Rheingold, J.R. Harper), (403) 335
- Dimolybdenum oxo-imido complexes; crystal structures of  $[(\text{MeC}_5\text{H}_4)_2\text{Mo}_2\text{O}_2(\mu\text{-O})(\mu\text{-NPh})]$  and  $[(\text{MeC}_5\text{H}_4)_2\text{MoO}(\mu\text{-NPh})_2]$  (J. Fletcher, G. Hogarth, D.A. Tocher), (403) 345
- Molybdenum(0) and tungsten(0) tricarbonylphosphine complexes with chelating uninegative (X,Y)-donor ligands (X, Y = S, N, O) (A.E. Sánchez-Peláez, M.F. Perpiñán), (405) 101
- Trialkylindium compounds, reaction with (cyclopentadienyl)(tricarbonyl)metal radicals (D.L. Thorn), (405) 161
- Dimolybdenum oxo-imido complexes; reactivity of  $[(\text{MeC}_5\text{H}_4)_2\text{Mo}_2\text{O}_2(\mu\text{-O})(\mu\text{-NPh})]$  and crystal structure of  $[(\text{MeC}_5\text{H}_4)_2\text{Mo}_2\text{O}(\text{S})(\mu\text{-O})(\mu\text{-NPh})]$  (J. Fletcher, G. Hogarth, D.A. Tocher), (405) 207
- Carbonyl carbene complex of molybdenum, preparation of naphthoquinones and indanones by anellation of (K.H. Dötz, H. Larbig), (405) C38
- Dimolybdenum complexes containing oxo, imido, and sulphido moieties, electrochemistry of; a measure of their relative  $\pi$ -donor abilities (G. Hogarth, P.C. Konidaris, G.C. Saunders), (406) 153
- Heterometallic  $n,\pi$ -complexes of transition metals based on mesitylacetonitrile (V.S. Kaganovich, M.I. Rybinskaya), (407) 215
- Molybdenum  $\pi$ -allyldicarbonyl complexes containing pyrazole-derived bidentate ligands; synthesis, structure and behavior in solution of (K.-B. Shiu, C.-J. Chang, S.-L. Wang, F.-L. Liao), (407) 225
- Dinitrosymolybdenum alkylidene complexes containing n-alkyl alkoxy ligands; synthesis, spectroscopic characterisation and metathesis activity of (A. Keller), (407) 237
- $[\text{CpM}(\text{CO})_3\text{PPh}_2]$  ( $\text{Cp} = \eta\text{-C}_5\text{H}_5$ , M = Mo, W) metallophosphines, reactivity towards alkynes; X-ray crystal structure of  $[\text{CpMo}(\text{CO})_2\{\text{Ph}_2\text{PCH}=\text{C}(\text{Ph})\text{CO}\}]$  (H. Adams, N.A. Bailey, A.N. Day, M.J. Morris, M.M. Harrison), (407) 247
- Unidentate and heterobimetallic 1,1'-bis(diphenylphosphino)ferrocene complexes, X-ray photoelectron spectroscopic differentiation of chemically distinct phosphorus environments in (H.S.O. Chan, T.S.A. Hor, L.-T. Phang, K.L. Tan), (407) 353
- Diisocyanide complexes  $\text{trans-}[\text{Mo}(\text{CNR})_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$  and derived aminocarbyne compounds, molecular orbital study of the bonding and reactivity of (E.G. Bakalbassis, C.A. Tsipis, A.J.L. Pombeiro), (408) 181
- Iron(II), tungsten(II) and molybdenum(II) ( $\eta^5$ -iodocyclopentadienyl)metallo methyl derivatives, methyl–iodine exchange in the presence of palladium (C. Lo Sterzo), (408) 253
- $[(\eta^5\text{-C}_5\text{H}_5)\text{CO}_3\text{M}(\text{HgX})]$  (M = Cr, Mo, W; X = Cl, Br, I, N<sub>3</sub>, SCN) complexes, interaction with  $\text{M}'(\text{CO})_3(\text{bipy})$  ( $\text{M}' = \text{Mo, W}$ ) fragments; trimetallic compounds containing  $\text{M}-\text{Hg}-\text{M}'-\text{X}$  arrays (J. Granero, M.E. Vargas), (408) 357
- ( $\eta^5\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_{13}\text{H}_9\text{M}(\text{CO})_3\text{Me}$  complexes (M = Mo, W), photoinduced reactions with acetylenes (H.G. Alt, J.S. Han, H.E. Maisel), (409) 197
- Seven-coordinate molybdenum(II) and tungsten(II) organophosphine carbonyl iodide derivatives, NMR study of (P.K. Baker, M.A. Beckett, L.M. Severs), (409) 213
- Molybdenum and tungsten monoalkyne complexes, syntheses and NMR studies of (J.L. Davidson, F. Sence), (409) 219
- Stereochemically non-rigid molybdenum-allyl complexes containing the bis(3,5-dimethylpyrazolyl)-methane ligand, synthesis of (V.S. Joshi, A. Sarkar, P.R. Rajamohanan), (409) 341
- Iron and molybdenum cyclometallated diazene complexes (H. Kisch, D. Garn), (409) 347
- $[(\text{OC})_3\text{M}(\mu\text{-Cl})(\mu\text{-CO})(\mu\text{-dppm})\text{Rh}(\text{NBD})]$  (M = Mo, W) tribridged derivatives, formation by ring opening of  $[(\text{OC})_4\text{M}(\text{dppm-PP})]$ ; crystal structure of  $[(\text{OC})_3\text{Mo}(\mu\text{-Cl})(\mu\text{-CO})(\mu\text{-dppm})\text{Rh}(\text{NBD})]$  (M. Cano, J.V. Heras, P. Ovejero, E. Pinilla, A. Monge), (410) 101
- Carbonylchromium, carbonylmolybdenum and carbonyltungsten complexes of the asymmetrically substituted phosphino sulfur diimides  ${}^1\text{Bu}_2\text{P}(\text{NSN})\text{PPh}_2$  and  ${}^1\text{Bu}_2\text{P}(\text{NSN})\text{P}({}^1\text{Bu})\text{Ph}$ ;  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{31}\text{P}$  NMR spectroscopic characterisation of (M. Herberhold, S.M. Frank, B. Wrackmeyer), (410) 159
- High-valent molybdenum complexes, reaction with carbon monoxide (R. Laï, S. Le Bot, F. Djafri), (410) 335
- Tetra-tert-butyl-pentafulvalene as ligand in dinuclear molybdenum complexes; *cis*-configuration without a metal–metal bond (P. Jutzi, J. Schnittger, B. Neumann, H.-G. Stammler), (410) C13

- $\text{CpMH}=\text{C}(\text{NHEt})\text{Ph})(\text{CO})_2$  ( $\text{M} = \text{Mo, W}$ ) system, hydride to carbene migration in; crystal structure of  $\text{CpMo}(\eta^2\text{-CH}(\text{Ph})\text{NHEt})(\text{CO})_2$  (H. Adams, N.A. Bailey, M.J. Winter, S. Woodward), (410) C21
- $\text{Bis}(\eta^5\text{-cyclopentadienyl})$ molybdenum(IV) dialkyls, autoxidations of (J.M. Atkinson, P.B. Brindley), (411) 139
- Binuclear transition metal complexes of  $\text{Cu}^1$ ,  $\text{Mo}^0$ , and  $\text{Re}^1$  with 2,2'-bipyrimidine (P.S. Braterman, J.-I. Song, S. Kohlmann, C. Vogler, W. Kaim), (411) 207
- Molybdenum(II) and tungsten(II) tricyclohexylphosphinecarbonylsulphide seven-coordinate complexes (P.K. Baker, D. ap Kendrick), (411) 215
- $(\text{CO})_{5-n}(\text{R}'_3\text{P})_n\text{M}(\text{H})\text{SiR}_3$  ( $n = 2, 3$ ), synthesis of phosphane substituted hydridosilyl complexes of chromium, molybdenum and tungsten; synthesis of  $(\text{CO})_3(\text{L-L})\text{M}(\text{H})\text{Cl}$  by conversion of  $(\text{CO})_3(\text{L-L})\text{M}(\text{NCR}')$  with  $\text{HSiCIR}_2$  ( $\text{L-L} = \text{tmida, bipy, dppe, (PPh}_3)_2$ ) (H. Piana, U. Schubert), (411) 303
- $N$ -Allyl-trichloroacetamide, enantioselective epoxydation with optically active Mo catalysts (H. Brunner, H. Zintl), (411) 375
- Highly reduced metalcarbonyl  $[\text{Ir}(\text{CO})_3]^{3-}$ , reaction with the ethylenepentacarbonylrhenium cation; formation of the trinuclear hydride  $(\text{OC})_5\text{Re}-\text{Ir}(\text{CO})_3(\text{H})-\text{Re}(\text{CO})_5$  and the trismetalated carbonate  $\{\mu_3\text{-CO}_3\text{[Re(CO)}_5\text{]}_3\}^+$  (J. Breimair, C. Robl, W. Beck), (411) 395
- Binuclear oxo complexes of molybdenum and tungsten with chalcogenes (O, S, Se, Te) as bridging ligands; synthesis and characterisation of (M. Gorzellik, H. Bock, L. Gang, B. Nuber, M.L. Ziegler), (412) 95
- $\text{Cp}_2\text{MM}'(\text{Cp})(\text{CO})_3$  ( $\text{M} = \text{Nb, Ta}; \text{M}' = \text{Mo, W}$ ), heteronuclear complexes with bridging and semi-bridging carbonyl groups; synthesis and dynamic behaviour (D. Perrey, J.C. Leblanc, C. Moïse, J. Martin-Gil), (412) 363
- $(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\text{Mo}\equiv\text{CNEt}_2$ , low-valent diethylaminocarbyne complex of molybdenum with a pentamethylcyclopentadienyl ligand; synthesis and crystal structure of (A.C. Filippou, W. Grünleitner, E.O. Fischer, W. Imhof, G. Huttner), (413) 165
- $\eta^2\text{-Iminoacyl}$ ,  $\eta^3\text{-1-azaallyl}$  and  $\eta^2\text{-1,4-diazabutadien-2-yl}$  complexes of molybdenum, selective formation from ethyl isocyanide complexes (A.C. Filippou, W. Grünleitner, C. Völkl, P. Kiprof), (413) 181
- $[\text{M}(\eta^3\text{-C}(\text{CF}_3)\text{C}(\text{CF}_3)\text{SR})(\text{CF}_3\equiv\text{CCF}_3)(\eta^5\text{-C}_5\text{H}_5)]$ , molybdenum and tungsten  $\eta^2\text{-C,C-vinyl}$  complexes; reactions with alkynes leading to novel metallacycles (L. Carlton, N.M. Agh-Atabay, J.L. Davidson), (413) 205
- Molybdenum(II) carbonyl complexes, chelate-assisted synthesis of; crystal structure of  $[\text{Mo}(\eta^2\text{-dppm})(\eta^1\text{-dppm})(\text{CO})_2\text{Br}_2]$  (dppm = bis(diphenylphosphino)methane) (K.-B. Shiu, K.-H. Yih, S.-L. Wang, F.-L. Liao), (414) 165
- Alkylidene dinitrosyl molybdenum complexes; synthesis and characterization of ethylidene complexes of the type  $[(\text{Et}_3\text{Sn}\cdot\text{AlCl}_3)_2\text{Mo}(\text{NO})_2(\text{=CHMe})]$  (A. Keller), (415) 97
- $[\text{W}_2(\text{CO})_{10}]^{2-}$  and  $[\text{Mo}_2(\text{CO})_8(\mu\text{-H})(\mu\text{-L})]^-$  ( $\text{L} = \text{bidentate phosphines}$ ) anions, their use in the synthesis of mercury-transition metal clusters (R. Reina, O. Rossell, M. Seco, A. Perales), (415) 101
- $[\text{Mo}(\text{CO})_2(\text{L})^{\text{thu}}\text{S}_4]'$ , complexes with sterical tetradentate thioether thiolate ligands  ${}^{\text{thu}}\text{S}_4^{2-} = 1,2\text{-bis}(2\text{-mercapto-3,5-di-t-butylphenylthio})\text{ethan}(2-)$  ( $\text{L} = \text{CO, PMe}_3, \text{PPh}_3$ ) (D. Sellmann, F. Grasser, M. Moll), (415) 367
- $[\text{MeCpMo}(\text{CO})_3]_2$ , photochemical reactions with quinones; EPR studies of (M. Hanaya, M. Iwaizumi), (417) 407
- Aminocarbene complexes of chromium and molybdenum derived from primary amines, preparation, structure and reactivity of (B. Denise, P. Dubost, A. Parlier, M. Rudler, H. Rudler, J.C. Daran, J. Vaissermann, F. Delgado, A.R. Arevalo, R.A. Toscano, C. Albarez), (418) 377
- Tungsten and molybdenum alkylidyne complexes  $[\text{M}(\text{=CR})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ , reactions with the bidentate secondary phosphine  $\text{C}_6\text{H}_4(\text{PH}_2)_2\text{-1,2}$ ; crystal structure of the phospha-allyl complex  $[\text{W}\{\sigma, \eta^3\text{-C}_6\text{H}_4(\text{PH}_2)\text{XPC(OH)CH}(\text{C}_6\text{H}_4\text{Me-4})\}\text{-1,2}(\text{CO})(\eta\text{-C}_5\text{H}_5)]$  (J.C. Jeffery, D.W.I. Sams, K.D.V. Weerasuria), (418) 395
- Heterodinuclear group 6 metallacboranes; synthesis and characterization of [*cis*-3,3,3-(CO)<sub>3</sub>-3-SnPh<sub>3</sub>-3,1,2-MC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>] ( $\text{M} = \text{Cr, Mo, W}$ ) (J. Kim, Y. Do, Y.S. Sohn, C.B. Knobler, M.F. Hawthorne), (418) C1
- $(\mu\text{-}\eta^2\text{,}\eta^3\text{-propargyl})$ -bis(cyclopentadienyl)tetracarbonyldimolybdenum tetrafluoroborate and  $(\mu\text{-}\eta^2\text{,}\eta^3\text{-1,1-dimethylpropargyl})$ -bis(cyclopentadienyl)tetracarbonyldimolybdenum tetrafluoroborate, crystal and molecular structure of (I.V. Barinov, O.A. Reutov, A.V. Polyakov, A.I. Yanovsky, Yu.T. Struchkov, V.I. Sokolov), (418) C24

- CpMH{=C(OEt)Ph}(CO)<sub>2</sub> (M = Mo, W) system, hydride to carbene migration and subsequent rearrangement of the consequent products to  $\eta^3$ -benzyl complexes; crystal structure of the  $\eta^3$ -benzyl CpW(CO)<sub>2</sub>{ $\eta^3$ -C<sub>6</sub>H<sub>5</sub>CH(OEt)} (H. Adams, N.A. Bailey, M.J. Winter, S. Woodward), (418) C39
- Group VI metal carbonyls, ( $\eta^6$ -C<sub>6</sub>H<sub>6</sub>)Cr(CO)<sub>3</sub>, ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Mn(CO)<sub>3</sub>, ( $\eta^5$ -CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Mn(CO)<sub>3</sub> and ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Re(CO)<sub>3</sub>, photolysis in Nujol at 77 K (T.E. Bitterwolf, K.A. Lott, A.J. Rest and J. Mascetti), (419) 113
- Cyclopentadienyltrichloromolybdenum(IV), synthesis, characterization, and reactivity of; X-ray structure of CpMoCl<sub>3</sub>[P(OCH<sub>2</sub>)<sub>3</sub>CEt]<sub>2</sub> (Cp =  $\eta^5$ -C<sub>5</sub>H<sub>5</sub>) (R. Poli and M.A. Kelland), (419) 127
- Alkyne, cyclobutadiene and cyclopentadienone complexes of molybdenum and tungsten (J.L. Davidson), (419) 137
- Organometallic compounds, oxidation with polychalcogenides; preparation and structure of [(Me<sub>5</sub>C<sub>5</sub>)Mo(S<sub>4</sub>)S<sub>2</sub>CO)]<sup>-</sup> containing a dithiocarbonate ligand (S.L. Gales, W.T. Pennington and J.W. Kolis), (419) C10
- Organotransition-metal complexes of multidentate ligands. XVI. On the nature of the sigma-donicity of the saturated nitrogen ligands; chelate-assisted weakening of the  $\alpha$ -N-H bond: synthesis, and spectral and structural study of [Mo(N-N)(CO)<sub>4</sub>] (N-N = saturated nitrogen bidentate ligands) (K.-B. Shiu, S.-L. Wang and F.-L. Liao), (420) 207
- Acetylen-Komplexe von Di( $\eta^5$ -cyclopentadienyl)molybdän: Synthesen und Protonierung. Struktur von [Cp<sub>2</sub>MoH( $\eta^2$ -C<sub>2</sub>(SiMe<sub>3</sub>)<sub>2</sub>)]PF<sub>6</sub> (G.E. Herberich, U. Englert and W. Faßbender), (420) 303
- Organotransition-metal complexes of multidentate ligands. XVII. Synthesis and characterization of some  $\eta^3$ -allylcarbonyl complexes of molybdenum containing dithiocarbamates as the uni-negative sulphur-bidentate ligands: crystal structure of [Mo( $\eta^2$ -S<sub>2</sub>CNEt<sub>2</sub>)CO]- ( $\eta^3$ -allyl){ $\eta^2$ -bis(diphenylphosphino)methane}] (K.-B. Shiu, K.-H. Yih, S.-L. Wang and F.-L. Liao), (420) 359
- An unexpected molybdenum(0) complex with "MoP<sub>6</sub>" coordination: crystal structure of [Mo(P(CH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>3</sub>)<sub>2</sub>]<sup>.</sup>C<sub>5</sub>H<sub>10</sub> (M. García-Basallote, P. Valerga, M.C. Puerta-Vizcaíno, A. Romero, A. Vegas and M. Martínez-Ripoll), (420) 371
- Kohlenwasserstoffverbrückte Komplexe. XXIII. Heterobimetallische Komplexe mit Ferrocenyl- und (Ph<sub>3</sub>P)(OC)(Cp)FeC(O)CH<sub>2</sub>-Gruppen; Darstellung und Struktur von ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe[ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)Re(CO)<sub>5</sub>], (PPPh<sub>3</sub>)(OC)(Cp)FeC(O)CH<sub>2</sub>( $\eta^6$ -C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub> und [ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)CH<sub>2</sub>( $\eta^6$ -C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub>] (M = Cr, Mo) (J. Breimair, M. Wieser, B. Wagner, K. Polborn und W. Beck), (421) 55
- <sup>1</sup>H and <sup>13</sup>C NMR study of low-energy intramolecular transformations of coordinated propargyl cations [( $\mu$ - $\eta^2$ , $\eta^3$ -HCCCR<sub>1</sub>R<sub>2</sub>)Cp<sub>2</sub>Mo<sub>2</sub>(CO)<sub>4</sub>]<sup>+</sup> BF<sub>4</sub><sup>-</sup> (M.V. Galakhov, V.I. Bakhmutov, I.V. Barinov and O.A. Reutov), (421) 65
- Reactivity of Main-Group-transition-metal bonds. IX. The kinetics of iodination of compounds containing two or more tin-transition-metal bonds (J.R. Chipperfield, S. Clark, D.E. Webster and H. Yusoff), (421) 205
- Synthesis and electrochemistry of dimolybdenum crown ether imido complexes (M.L.H. Green, G. Hogarth and G.C. Saunders), (421) 233
- Preparation and spectroscopic properties of a series of new bimetallic phosphine-bridged seven-coordinate complexes of the type [M<sub>2</sub>I<sub>4</sub>(CO)<sub>6</sub>L<sub>2</sub>( $\mu$ -L-L)] {M = Mo or W; L = PPh<sub>3</sub>, AsPh<sub>3</sub> or SbPh<sub>3</sub>; L-L = Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub> (n = 1, 2 or 4); L-L = [Fe( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>]} (P.K. Baker, M. van Kampen and D. ap Kendrick), (421) 241
- Half-sandwich complex with intramolecular amino group coordination: synthesis of molybdenum iodide complexes (T.-F. Wang, T.-Y. Lee, J.-W. Chou and C.-W. Ong), (423) 31
- Preparation and properties of molybdenum and tungsten dinitrogen complexes. XXXIV. Nucleophilic addition reactions of hydrazido(2-) complexes with diphenylketene: characterization of new acylhydrazido(2-) complexes of molybdenum and tungsten (T. Aoshima, Y. Mizobe, M. Hidai and J. Tsuchiya), (423) 39
- Ti<sup>IV</sup> and Mo<sup>V</sup> complexes with diesters. The crystal structure of [ $\sigma$ -C<sub>6</sub>H<sub>4</sub>(CO<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>Ph)<sub>2</sub>Cl<sub>4</sub>Ti], [C<sub>2</sub>O<sub>4</sub>(CH<sub>2</sub>CH<sub>2</sub>Ph)<sub>2</sub>Cl<sub>4</sub>Ti] and [ $\sigma$ -C<sub>6</sub>H<sub>4</sub>(CO<sub>2</sub>Et)<sub>2</sub>Cl<sub>3</sub>MoO]<sup>.</sup>C<sub>6</sub>H<sub>6</sub> (P. Sobota, S. Szafert, J. Utko and T. Lis), (423) 195
- Electrochemical generation of 17-electron cation radicals from arene-M(CO)<sub>3</sub> (M = Cr, Mo, W) and thiophene-Cr(CO)<sub>3</sub> complexes in MeCN: Conventional cyclic voltammetric studies with digital simulation and microelectrode voltammetry in the absence of supporting electrolyte (F. Rourke, R. Gash and J.A. Crayston), (423) 223

Synthesis and reactivity of molybdenocene isocyanide complexes; crystal structure of  $(\eta^5\text{-C}_5\text{H}_5)_2\text{MoCN}^+\text{Bu}$  (A.M. Martins, M.J. Calhorda, C.C. Romão, C. Völkl, P. Kiprof and A.C. Filippou), (423) 367

Photochemistry of  $(\text{Me}_3\text{P})_4\text{Mo}(\eta^2\text{-CO}_2)_2$ : deoxygenation of coordinated carbon dioxide and phosphine oxidation (W. Ziegler and K.M. Nicholas), (423) C35

Organometallic chemistry of diphosphazanes. IV. Reactions of the cyclodiphosphazane, *cis*- $[(\text{BuNP(OPh)})_2\text{M}(\text{CO})_6]$  ( $\text{M} = \text{Cr, Mo, W}$ ), rhodium(I), palladium(II) and platinum(II) derivatives (M.S. Balakrishna and S.S. Krishnamurthy), (424) 243

## Neodymium

Bis(t-butylcyclopentadienyl)lanthanoid chloride complexes, syntheses of; crystal structures of bis(t-butylcyclopentadienyl)chloro(bistetrahydrofuran)praseodymium and bis(t-butylcyclopentadienyl)-chlorotetrahydrofuran ytterbium (Q. Shen, M. Qi, J. Guan, Y. Lin), (406) 353

Triindenyl lanthanide complexes  $(\eta^5\text{-C}_9\text{H}_7)_3\text{Ln}\cdot\text{OC}_4\text{H}_8$  ( $\text{Ln} = \text{Nd, Gd, Er}$ ), syntheses and crystal structures of (J. Xia, Z. Jin, G. Lin, W. Chen), (408) 173

Bis(cyclopentadienyl)(2,6-diphenylphenoxy)bis(tetrahydrofuran)neodymium(III), preparation and crystal structure of; an unexpected example of formal nine coordination (G.B. Deacon, S. Nickel, E.R.T. Tiekkink), (409) C1

Tris( $\eta^5$ -cyclopentadienyl)neodymium(III), neutral base adducts of; interpretation of their optical magnetic, ESR and NMR properties (H. Reddmann, H. Schultze, H.-D. Amberger, G.V. Shalimoff, N.M. Edelstein), (411) 331

Neodymium(III) complexes with bulky cyclopentadienyl and benzamidinate ligands, a comparison (A. Recknagel, F. Knösel, H. Gornitzka, M. Noltemeyer, F.T. Edelmann, U. Behrens), (417) 363

Methylbis(t-butyl-cyclopentadienyl)neodymium and -gadolinium, preparation and molecular structure of (Q. Shen, Y. Cheng and Y. Lin), (419) 293

## Nickel

$[(\text{PhSiO})_6(\mu_4\text{-O})_2(\mu_3\text{-O})_4[\text{Ni}_8(\mu_3\text{-O})_2](\mu_3\text{-O})_4(\mu_4\text{-O})_2]\text{[PhSiO}]_6 \cdot 14\eta\text{-BuOH} \cdot 10\text{H}_2\text{O} \cdot 2\text{Me}_2\text{CO}$  sandwich-type nickel oxide-siloxanolate complex; synthesis and crystal structure of (M.M. Levitsky, O.I. Schegolikhina, A.A. Zhdanov, V.A. Igonin, Yu.E. Ovchinnikov, V.E. Shklover, Yu.T. Struchkov), (401) 199

Nickel-catalysed electrochemical Reformatsky reaction with methyl chlorodifluoroacetate (S. Mcharek, S. Sibille, J.-Y. Nédélec, J. Péridon), (401) 211

Chiral metal complexes of 1-dimethylphosphino-3-dimethylamines, homogeneous asymmetric catalysis by means of (C. Döbler, A. Kinting), (401) C23

$[(\eta^5\text{-Dimethylphosphino)cyclopentadienyl}][\eta^7\text{-}(dimethylphosphino)cycloheptatrienyl]titanium$ , heterobimetallic compounds derived from (M. Ogasa, M.D. Rausch, R.D. Rogers), (403) 279

$\text{Ni}^0$ -induced and -catalysed preparation of unsaturated adipic acid anilides (H. Hoberg, D. Bärhausen), (403) 401

Nickel(0)-ligand complex, formation of C-C bonds between carbon dioxide and cycloalkenes on (H. Hoberg, A. Ballesteros, A. Sigan), (403) C19

1,*n*-Dicarbonyl compounds, one-step synthesis from carboxylic acid derivatives and di-Grignard reagents in the presence of transition metal catalysts (F. Babudri, A. D'Ettole, V. Fiandanese, G. Marchese, F. Naso), (405) 53

Clusters with  $\text{Re}_2\text{Fe}$  and  $\text{Re}_2\text{Ni}_2$  cores, phenylacetylene transformations in; synthesis and molecular structure of  $\text{Re}_2\text{Fe}(\text{CO})_9(\mu_3\text{-}, \eta^4\text{-CH=CH=CH}_2\text{C}_6\text{H}_4)$  and  $\text{Cp}_2\text{Ni}_2\text{Re}_2(\text{CO})_6(\mu_4\text{-PhC}\equiv\text{CCH=CHPh})$  (A.D. Shaposhnikova, G.L. Kamalov, R.A. Stadnichenko, A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, Yu.T. Struchkov, A.I. Yanovsky), (405) 111

Nickela compounds with a five-membered ring, ring contraction of; use of the resulting four-membered ring compounds in preparative chemistry (H. Hoberg, A. Ballesteros, A. Sigan, C. Jégat, D. Bärhausen, A. Milchereit), (407) C23

Infrared spectra of triple-decker sandwich complexes  $(\text{Ni}_2\text{Cp}_3)\text{A}$  ( $\text{A} = \text{BF}_4^-$ ,  $\text{PF}_6^-$ ) (I.A. Garbuzova, O.G. Garkusha, B.V. Lokshin, A.R. Kudinov, M.I. Rybinskaya), (408) 247

(2.6- $i\text{-Pr}_2\text{Ph}$ -dad) $\text{Ni}(\text{C}_2\text{F}_4)$ , synthesis and reactivity of (W. Schröder, W. Bonrath, K.R. Pörschke), (408) C25

Homogeneous metal complex catalysts in the synthesis of cyclobutane and cyclopentane compounds (U.M. Dzhemilev, R.I. Khusnutdinov, G.A. Tolstikov), (409) 15

- 2-3- $\eta^2$ -(Butyne-1,4-diol)bis(triisopropylphosphane)nickel(0), ( $^i\text{Pr}_3\text{P}$ )<sub>2</sub>Ni(HOCH<sub>2</sub>C≡CCH<sub>2</sub>OH), crystal and molecular structure of; some peculiarities of the complexation of 2-butyne-1,4-diol with nickel(0) complexes (U. Rosenthal, G. Oehme, H. Görls, V.V. Burlakov, A.V. Polyakov, A.I. Yanovsky, Yu.T. Struchkov), (409) 299
- Ni<sup>0</sup> catalysed reactions between 1,3-butadiene and phenylisocyanate (H. Hoberg, D. Bärhausen, R. Mynott, G. Schroth), (410) 117
- Naphthaleneytterbium, reactions with bis(cyclopentadienyl)complexes of cobalt, nickel, chromium and vanadium; crystal structure of the triple-decker compound CpVC<sub>10</sub>H<sub>8</sub>VCp (M.N. Bochkarev, I.L. Fedushkin, H. Schumann, J. Loebel), (410) 321
- $\eta^3,\eta^2,\eta^2$ -Dodeca-2(E),6(E),10(Z)-trien-1-ylnickel(II) complexes [Ni(C<sub>12</sub>H<sub>19</sub>)]X (X = BF<sub>4</sub>, B(O<sub>2</sub>C<sub>6</sub>H<sub>4</sub>)<sub>2</sub>, CF<sub>3</sub>SO<sub>3</sub>, AlBr<sub>4</sub>) as catalysts for 1,4-polymerisation of butadiene; anion influence on catalytic structure-reactivity relationships (R. Taube, J.-P. Gehrke, P. Böhme, K. Scherzer), (410) 403
- trans*-NiI<sub>2</sub>(RNC)<sub>2</sub> complex, unusual reductive behaviour at mercury and platinum electrodes (K. Ehara, K. Kumagaya, Y. Yamamoto, K. Takahashi, H. Yamazaki), (410) C49
- Fe<sub>3</sub>(CO)<sub>11</sub><sup>2-</sup> and Fe<sub>4</sub>(CO)<sub>13</sub><sup>2-</sup>, transformation to give trinuclear clusters (W. Deck, A.K. Powell, H. Vahrenkamp), (411) 431
- Binuclear nickel complex [(depe)NiBr]<sub>2</sub>( $\mu$ -Z,Z-PhC=CH-CH=CPh), facile cleavage of C-C bond (C.J. Lawrie, H.E. Dankosh, B.K. Carpenter), (411) C7
- Ni<sup>0</sup> induced synthesis of cyclic C<sub>8</sub> carbon acids from cyclooctenes and carbon dioxide (H. Hoberg, A. Ballesteros), (411) C11
- Ni<sup>0</sup> induced CC-coupling between phenylisocyanate and five-membered cyclic alkenes, catalytic formation of  $\beta,\gamma$ -unsaturated carboxylic acid anilides (H. Hoberg, M. Nohlen), (412) 225
- Acetylenes and diazoketones in organic synthesis (F. Serratosa), (413) 445
- Zerovalent nickel complex mediated carbon-carbon coupling reactions of haloaromatic compounds; preparation of cyclic oligomers of thiophene and benzene and stable anthrylnickel(II) complexes (Z. Zhou, T. Yamamoto), (414) 119
- [Ni( $\eta$ -CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)XX(PPh<sub>3</sub>)<sub>2</sub>] and functionally substituted monocyclopentadienyl derivatives with other phosphines, preparation of; crystal structure of [Ni( $\eta$ -CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)X(PPh<sub>3</sub>)<sub>2</sub>] (L. Ballester, S. Perez, A. Gutierrez, M.F. Perpiñan, E. Gutierrez-Puebla, A. Monge, C. Ruiz), (414) 411
- 20-Electron halfsandwich compounds, sterical stabilization of; crystal structure of [(C<sub>5</sub>H<sub>5</sub>)Co{P(OMe)<sub>2</sub>O<sub>3</sub>Ni(C<sub>5</sub>Ph<sub>5</sub>)}] (W. Klaui, M. Huhn, R. Herbst-Irmer), (415) 133
- Para- and diamagnetic nickel complexes, tetra-tert-butyl-dihydropentafulvene as educt for (P. Jutzi, J. Schnittger, W. Wieland, B. Neumann, H.-G. Stammler), (415) 425
- P-substituted olefins, addition of optically active amines to (H. Brunner, S. Limmer), (417) 173
- Enantioselective hydrogenation catalysts, some general regularities in the action of (E.I. Klabunovskii), (417) 181
- Nickel complexes containing (R, R)-1,2-cyclopentanediylibis(diphenylphosphine) and related ligands, enantioselective cross-coupling of vinyl-, aryl and allyl-electrophiles catalyzed by (G. Consiglio, A. Indolese), (417) C36
- Oxidative carbonylation of aliphatic amines catalysed by nickel-complexes (P. Giannoccaro, C.F. Nobile, P. Mastrolilli and N. Ravasio), (419) 251
- Novel class of transition metal coordination compounds with macrocyclic organosiloxanolate ligands; their synthesis and crystal structure (V.A. Igonin, O.I. Shchegolikhina, S.V. Lindeman, M.M. Levitsky, Yu.T. Struchkov and A.A. Zhdanov), (423) 351
- New C-C bond formation through the nickel-catalysed electrochemical coupling of 1,3-enynes and carbon dioxide (S. Dérien, J.-C. Clinet, E. Duñach and J. Périchon), (424) 213

## Niobium

- Bis-silylated cyclopentadienyl niobium trihydride, electrochemical behaviour of; ESR spectroscopic characterisation of the dimeric hydride niobium(IV) complex [Nb( $\eta^5$ -Me<sub>3</sub>SiC<sub>5</sub>H<sub>4</sub>)<sub>2</sub>H]<sub>2</sub>( $\mu$ -H)<sub>2</sub> (L. Roullier, D. Lucas, Y. Mugnier, A. Antónolo, M. Fajardo, A. Otero), (412) 353
- Cp<sub>2</sub>MM'(Cp)(CO)<sub>3</sub> (M = Nb, Ta; M' = Mo, W), heteronuclear complexes with bridging and semi-bridging carbonyl groups; synthesis and dynamic behaviour (D. Perrey, J.C. Leblanc, C. Moïse, J. Martin-Gil), (412) 363
- $\eta^6$ -Arene derivatives of niobium(II), niobium(I), and niobium(0); synthesis and reactivity of (F. Calderazzo, G. Pampalini, L. Rocchi, J. Strähle, K. Wurst), (413) 91

Bis-silylated cyclopentadienylniobium(V) cationic complexes, synthesis and electrosynthesis of; their reactions and structural characterisation (A. Antíñolo, M. Fajardo, A. Otero, Y. Mugnier, H. Nabouï, H. Mourad), (414) 155

Synthesis and characterization of the paramagnetic niobium(III) complex  $[\text{Nb}(\eta\text{-C}_5\text{Me}_5)\text{Cl}_2(\text{PMe}_3)_2]$  and its reaction with carbon monoxide (U. Siemeling and V.C. Gibson), (424) 159

## Osmium

Permethylated  $\alpha$ -metallocenylcarbinols ( $M = \text{Os, Ru, Fe}$ ), hydrogen bonds in and conformations of (E.S. Shubina, L.M. Epstein, T.V. Timofeeva, Yu. T. Struchkov, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 133

Permethylated  $\alpha$ -metallocenylcarbinols in the solid state ( $M = \text{Fe, Ru, Os}$ ), IR spectroscopic study of hydrogen bonds in (E.S. Shubina, L.M. Epstein, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 145

( $\eta^6$ -Phenyl)Cr(CO)<sub>3</sub>- and ( $\eta^6$ -phenyl)Co<sub>4</sub>(CO)<sub>9</sub>-substituted metallocenylcarbinols of the iron subgroup; synthesis, structure and hydrogen bonding in (E.S. Shubina, L.M. Epstein, Yu.L. Slovokhotov, A.V. Mironov, Yu.T. Struchkov, V.S. Kaganovich, A.Z. Kreindlin, M.I. Rybinskaya), (401) 155

Ester and amide functionalized alkylidene triosmium clusters; rearrangement to acyl-coordinated alkylidene complexes (D.A. Strickland, J.R. Shapley), (401) 187

OSNC<sub>6</sub>H<sub>4</sub>Me-4, pseudo-olefinic coordination to zerovalent osmium; crystal structure of  $[\text{OsCl}(\text{NO})(\text{PPh}_3)_2(\text{OSNC}_6\text{H}_4\text{Me}-4)]$  (A.F. Hill, G.R. Clark, C.E.F. Rickard, W.R. Roper, M. Herberhold), (401) 357

[Os<sub>6</sub>Au(CO)<sub>20</sub>( $\mu$ -OMe)]; isolation, characterisation, and crystal and molecular structure of (C.M. Hay, B.F.G. Johnson, J. Lewis, N.D. Prior, P.R. Raithby, W.T. Wong), (401) C20

Osmium heptanuclear clusters; crystal structure of  $[\text{Os}_7(\text{CO})_{20}\text{P}(\text{OMe})_3]$  (B.F.G. Johnson, J. Lewis, J. Luniss, D. Braga, F. Grepioni), (401) C46

Osmium carbonyl anion  $[\text{HOs}_3(\text{CO})_{10}\cdot\text{O}_2\text{C}\cdot\text{Os}_6(\text{CO})_{20}]^-$ , synthesis and X-ray structure of (B.F.G. Johnson, J. Lewis, P.R. Raithby, W.T. Wong), (401) C50

Osmium(II) complexes; reactions of coordinated phosphorus trichloride, phenylphosphonous dichloride, 1-phenyl-3,4-dimethylphosphole and mesitylphosphine in (D.S. Bohle, G.R. Clark, C.E.F. Rickard, W.R. Roper, W.B. Shepard), (402) 375

Osmium clusters with 1,2-dipolar ligands; synthesis and crystal structure of  $\text{HOs}_3(\text{CO})_9(\mu_3\text{-}\eta^2\text{-PhNCHCPPh}_3)$  (M. Langenbahn, H. Stoeckli-Evans, G. Süss-Fink), (402) C12

Os<sub>6</sub>(CO)<sub>18</sub>, reaction with Me<sub>3</sub>NO; isolation and NMR study of the new hexaosmium cluster anion  $[\text{HOs}_6(\text{CO})_{17}]^-$  (B.F.G. Johnson, J. Lewis, M.-A. Pearsall, L.G. Scott), (402) C27

Os<sub>3</sub>(CO)<sub>10</sub>[ $(\text{CF}_3)_2\text{P}=\text{N=PPPh}_3$ ]<sub>2</sub> novel delocalised diphosphazene, preparation and X-ray structure of (H.G. Ang, Y.M. Cai, W.L. Kwik, E.C. Morrison, D.A. Tocher), (403) 383

Os<sub>3</sub>(CO)<sub>9</sub>[1,1,1-tris(isocyanomethyl)ethane], synthesis and crystal and molecular structure of (E.C. Constable, B.F.G. Johnson, F.K. Khan, J. Lewis, P.R. Raithby, P. Mikulcik), (403) C15

Annual Survey of ruthenium and osmium for the year 1988 (M.G. Richmond), (404) 213

Tetraosmium cluster, displacement of benzene by an alkyne in; synthesis and structural characterisation of  $[\text{H}_2\text{Os}_4(\text{CO})_9(\mu_3\text{-}\eta^2\text{-Ph}_2\text{C}_2)\ (\eta^2\text{-Ph}_2\text{C}_2)]$  (H. Chen, B.F.G. Johnson, J. Lewis, D. Braga, F. Grepioni, P. Sabatino), (405) C22

Hg(O<sub>2</sub>CCF<sub>3</sub>)<sub>2</sub>, reaction with [Os<sub>6</sub>(CO)<sub>18</sub>]; synthesis and crystal structures of  $[\text{HOs}_5\text{Hg}_2(\text{CO})_{15}(\text{O}_2\text{CCF}_3)_3]$  and *cis*-[Os<sub>4</sub>(CO)<sub>12</sub>(O<sub>2</sub>CCF<sub>3</sub>)<sub>2</sub>] (M.P. Diebold, B.F.G. Johnson, J. Lewis, V.P. Saharan, M. McPartlin, H.R. Powell), (405) C25

Triosmium and triruthenium clusters, reactions with the isomers methoxyallene and methyl propargyl ether; crystal structure of the methoxyallene cluster  $[\text{Os}_3(\text{CO})_{11}(\mu,\eta^1,\eta^3\text{-CH}_2\text{CCHOME})]$  (A.J. Arce, Y. De Sanctis, A.J. Deeming, K.I. Hardcastle, R. Lee), (406) 209

Triosmium clusters of the type Os<sub>3</sub>( $\mu$ -H)<sub>2</sub>(CO)<sub>9</sub>Ar (Ar = C<sub>6</sub>H<sub>4</sub>, C<sub>9</sub>H<sub>6</sub>, C<sub>4</sub>H<sub>3</sub>N, or C<sub>4</sub>H<sub>2</sub>NMe), reactions with phosphines and phosphites; X-ray crystal structure of Os<sub>3</sub>( $\mu$ -H)<sub>2</sub>(CO)<sub>7</sub>[P(OMe)<sub>3</sub>]<sub>2</sub> (C<sub>4</sub>H<sub>3</sub>N) (H. Chen, B.F.G. Johnson, J. Lewis, P.R. Raithby), (406) 219

Triosmium clusters with  $\mu_3$ -AsR group as a building block for hexaosmium-arsenic cluster synthesis; synthesis and characterisation of [(CO)<sub>11</sub>Os<sub>3</sub>As(Os<sub>3</sub>(CO)<sub>9</sub>H<sub>3</sub>)] and [(CO)<sub>9</sub>H<sub>3</sub>Os<sub>3</sub>As(Os<sub>3</sub>(CO)<sub>9</sub>H<sub>3</sub>)]; crystal structure of [(CO)<sub>11</sub>Os<sub>3</sub>As(Os<sub>3</sub>(CO)<sub>9</sub>H<sub>3</sub>)] (K. Guldner, B.F.G. Johnson, J. Lewis, A.D. Massey, S. Bott), (408) C13

- Phenylphosphole, oxidative addition of; X-ray crystal structures of two ring-opened products  $[\text{Os}_3(\mu_3-\text{PhPC}_4\text{H}_4\text{XCO})_x]$  where  $x = 8$  or  $9$  (A.J. Arce, Y. De Sanctis, J. Manzur, A.J. Deeming, N.I. Powell), (408) C18
- $\text{Os}_3(\text{CO})_{10}(\mu_2-\eta^3-\text{C}_3\text{H}_5)\text{X}(\text{AuPEt}_3)$ , a triosmium cluster exhibiting an allyl ligand with a novel mode of coordination; synthesis, molecular and electronic structure of (C.E. Housecroft, B.F.G. Johnson, J. Lewis, J.A. Luniss, S.M. Owen, P.R. Raithby), (409) 271
- Mesitylene osmium complexes with hydrido, carboxylato, carbonyl and vinylidene ligands (H. Werner, S. Stahl, W. Kohlmann), (409) 285
- $\text{Os}_3(\text{CO})_{10}(\mu\text{-Br}\text{CH}=\text{CHPh})$  triosmium complex, reactions with benzyl isonitrile and triphenylphosphine; crystal structure and solution fluxionality of the ethenyl derivatives (S.-H. Chuang, Y. Chi, F.-L. Liao, S.-L. Wang, S.-M. Peng, G.-H. Lee, J.-C. Wu, K.-M. Horng), (410) 85
- Triosmium and triruthenium carbon radicals, reactivity toward fluoro- and hexafluorobenzene (N. Hadj-Bagheri, D.S. Strickland, S.R. Wilson, J.R. Shapley), (410) 231
- $\text{L}_4\text{Ru}^0$  complex cores, inter- and intramolecular cleavage of  $sp^2$ - and  $sp^3\text{-C-H}$  bonds with  $\text{P}(\text{o-C}_6\text{H}_4\text{PMe}_2)_3$ ,  $\text{N}(\text{CH}_2\text{CH}_2\text{PMe}_2)_3$ ,  $\text{MeP}(\text{CH}_2\text{CH}_2\text{CH}_2\text{PMe}_2)_2/\text{PR}_3$  and  $\text{MeSi}(\text{CH}_2\text{PMe}_2)_3/\text{PR}_3$  ( $\text{R} = \text{Me}, \text{OMe}$ ) as supporting ligands (L. Dahlenburg, S. Kerstan, D. Werner), (411) 457
- $[\text{C}_6\text{H}_6\text{Os}(\text{PR}_3)]$  as structural element; synthesis and reactions with alkinyl, vinylidene, and vinylosmium(II) complexes (W. Knaup, H. Werner), (411) 471
- $[\text{Os}_3(\mu\text{-H})(\mu\text{-X})(\text{CO})_{10}]$  clusters ( $\text{X} = \text{furyl, thienyl, pyrrolyl, or indolyl}$ ), comparison of *monohapto* versus *dihapto* and *exo* versus *endo* coordination (A.J. Arce, J. Manzur, M. Marquez, Y. De Sanctis), (412) 177
- $[\text{HOs}_3(\text{CO})_{11}]^-$ , reactions with disubstituted acetylenes; synthesis and structural characterisation of the pentanuclear species  $[\text{H}_2\text{Os}_3(\text{CO})_{13}(\text{Ph}_2\text{C}_2\text{XPhC}_2\text{H})(\text{C}_6\text{H}_4)]$  (B.F.G. Johnson, J. Lewis, J. Luniss, D. Braga, F. Grepioni), (412) 195
- Bis(pentafluorophenyl)dioxobis(pyridine)osmium(VI),  $(\text{C}_6\text{F}_5)_2\text{OsO}_2\cdot(\text{py})_2$ , synthesis, structure and properties of (W.A. Herrmann, S.J. Eder, P. Kiprof), (412) 407
- Organic osmium oxide, pyridine complexes of; synthesis, spectra and crystal structure of  $\text{R}_2\text{OsO}_2\cdot(\text{L})$  and  $\text{R}_2\text{OsO}_2\cdot(\text{L})_2$  (W.A. Herrmann, S.J. Eder, P. Kiprof), (413) 27
- Tertiary-phosphine, arsine and stibine complexes of hexaosmium clusters; preparation and aspects of the chemistry of (B.F.G. Johnson, J. Lewis, M.-A. Pearsall, L.G. Scott), (413) 337
- Trisulphur- and triselenium-bridged [3]metallocenophanes, comparison of the energy barriers to bridge reversal in (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne and V. Šík), (419) 375
- Ultraviolet vapour-phase absorption spectra of  $d^6$  metallocenes (S.Y. Ketkov and G.A. Domrachev), (420) 67
- Cyclopentadienyl-ruthenium and -osmium chemistry. XXXVI. Oligomerisation of phenylacetylidyne residues on ruthenium. Crystal structures of  $\{\text{Ru}(\text{PPh}_3)(\eta\text{-C}_5\text{H}_5)\}_2(\mu\text{-C}_8\text{Ph}_4)$  and  $\{\text{Ru}(\text{PPh}_3)(\eta\text{-C}_5\text{H}_5)\}_2(\mu\text{-C}_{10}\text{Ph}_4(\text{C}_6\text{H}_4))$  (M.I. Bruce, G.A. Koutsantonis, M.J. Liddell and E.R.T. Tiekkink), (420) 253
- Cyclopentadienyl-ruthenium and -osmium chemistry. XXXVII. Oligomerisation of  $\text{C}_2(\text{CO}_2\text{Me})_2$  at a cyclopentadienyl-ruthenium centre: X-ray structures of  $\text{RuI}[\eta^4\text{-CH}(\text{CO}_2\text{Me})=\text{C}(\text{CO}_2\text{Me})\text{C}(\text{CO}_2\text{Me})=\text{CH}(\text{CO}_2\text{Me})](\eta\text{-C}_5\text{H}_5)$ ,  $\text{Ru}(\eta^5\text{-C}_5\text{H}_5)(\eta^5\text{-C}_6\text{H}(\text{CO}_2\text{Me})_6)$  and  $\text{Ru}(\eta^5\text{-C}_5\text{H}_5)[\eta^5\text{-C}_6\{\text{C}(\text{CO}_2\text{Me})=\text{CH}(\text{CO}_2\text{Me})\}(\text{CO}_2\text{Me})_6]$  (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tiekkink and B.K. Nicholson), (420) 271
- Preferential attack of the terminal hydrogen in  $(\mu\text{-H})\text{Os}_3(\text{CO})_{11}\text{H}$  by the  $(\text{CF}_3)_2\text{NO}$  radical and X-ray structure of  $\{(\mu\text{-H})\text{Os}_3(\text{CO})_{11}\}[\text{ON}(\text{CF}_3)_2]$  (H.G. Ang, W.L. Kwik, C.H. Koh, W.K. Leong, B.F.G. Johnson, J. Lewis and P.R. Raithby), (420) 289
- Heterometallic analogues of  $[(\text{M}(\text{CO})_2(\eta\text{-C}_5\text{H}_5))_2]$  ( $\text{M} = \text{Fe, Ru, Os}$ ). The synthesis of the tetracarbonylcyclopentadienylcycloheptatrienyl complexes  $[\text{M}(\text{CO})_2(\eta\text{-C}_5\text{H}_5)\text{M}'(\text{CO})_2(\eta\text{-C}_7\text{H}_7)]$  ( $\text{M} = \text{Fe, Ru}; \text{M}' = \text{Mo, W}$ ). Crystal structures of  $[\text{Ru}(\text{CO})_2(\eta\text{-C}_5\text{H}_5)\text{Mo}(\text{CO})_2(\eta\text{-C}_7\text{H}_7)]$  (Ru-Mo) and  $[\text{Mo}(\text{CO})_3(\mu\text{-}\eta^6\text{-}, \eta^1\text{-C}_7\text{H}_7)\text{Ru}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)]$  (R.L. Beddoes, E.S. Davies, M. Helliwell and M.W. Whittlesey), (421) 285
- Unexpected formation of hydrido-hydroxy derivatives as direct products in the reactions of the unsaturated dihydrides  $[\text{Mn}_2(\mu\text{-H})_2(\text{CO})_6(\mu\text{-L-L})]$  ( $\text{L-L} = \text{Ph}_2\text{PCH}_2\text{PPh}_2$ ,  $(\text{EtO})_2\text{POP(OEt)}_2$ ) with carbon dioxide (F.J. García Alonso, M. García Sanz and V. Riera), (421) C12
- Electrochemical behaviour of the electronically and coordinatively unsaturated cluster  $[\text{Os}_3(\mu\text{-H})(\text{CO})_8\{\text{Ph}_2\text{PCH}_2\text{P}(\text{Ph})\text{C}_6\text{H}_4\}]$  (D. Osella, M. Ravera, A.K. Smith, A.J. Mathews and P. Zanello), (423) 255

Reaktionen von  $[(\text{Mes})\text{Os}(\text{CO})\text{Cl}_2]$  mit Alkyl-Grignardreagenzien: Ein einfacher Weg zu Carbonyl(olefin)osmium(0)-Komplexen (U. Wecker und H. Werner), (424) 199

## Palladium

- 2-Substituted 8-quinolinols containing C, N, O-donor atoms and their Pd<sup>II</sup> or Pt<sup>II</sup> complexes, preparation of (A. Yoneda, G.R. Newkome, K.J. Theriot), (401) 217
- Cyclometallated derivatives of palladium(II) with 1,4-benzodiazepin-2-ones; crystal structure of (L-H)Pd(PPh<sub>3</sub>)Cl·CHCl<sub>3</sub> (L = Prazepam: 7-chloro-1-cyclopropylmethyl-1,3-dihydro-5-phenyl-2H-1,4-benzodiazepin-2-one); synthesis of isoindolo[2,1-d][1,4]benzodiazepine derivatives by reaction of [(L-H)PdCl]<sub>2</sub> species with carbon monoxide (M.A. Cinelli, S. Gladiali, G. Minghetti, S. Stoccoro, F. Demartin), (401) 371
- Cyclometallated complexes of palladium(II), synthesis of; X-ray crystal structure of di- $\mu$ -bromo-bis[N-(3,4-dimethoxybenzylidene)cyclohexylaminato-C,N]dipalladium(II) (J.M. Vila, M. Gayoso, M.T. Pereira, A. Romar, J.J. Fernández, M. Thornton-Pett), (401) 385
- Chiral metal complexes of 1-dimethylphosphino-3-dimethylamines, homogeneous asymmetric catalysis by means of (C. Döbler, A. Kinting), (401) C23
- Palladium(0) complex as catalyst in a tandem arylsulfonylation and cyclization of arylazoxy aryl sulfones with norbornene (N. Kamigata, M. Satoh, M. Yoshida), (401) C26
- Linked metal acetylide complexes for nonlinear optics, synthesis of (S.J. Davies, B.F.G. Johnson, J. Lewis, M.S. Khan), (401) C43
- Platinum or palladium 3-diphenylphosphinocamphor complexes including acetylides (S.D. Perera, B.L. Shaw), (402) 133
- Organopalladium(IV) complexes, synthesis by halogen exchange; structure of the azidopalladium(II) complex formed on reductive elimination of ethane from Pd(N<sub>3</sub>)Me<sub>2</sub>(CH<sub>2</sub>Ph)(bpy) (A.J. Canty, P.R. Traill, B.W. Skelton, A.H. White), (402) C33
- Mono- and polynuclear perhalophenyl palladium–platinum acetylide complexes, synthesis of; molecular structure of (NBu<sub>4</sub>)<sub>2</sub>[Pt<sub>2</sub>Ag<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>(C≡CPh)<sub>4</sub>]·4CH<sub>2</sub>Cl<sub>2</sub> (P. Espinet, J. Forniés, F. Martínez, M. Sotes, E. Lalinde, M.T. Moreno, A. Ruiz, A.J. Welch), (403) 253
- Five-coordinate trigonal-bipyramidal palladium(II) species, stabilisation of; crystal structure of (2,9-dimethyl-1,10-phenanthroline)methylchloropalladium(II) (V. De Felice, V.G. Albano, C. Castellari, M.E. Cucciolito, A. De Renzi), (403) 269
- C-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325
- Palladium-mediated *ortho*-alkylation of 2-aryloxazolines (J.C. Clinet, G. Balavoine), (405) C29
- Palladium-catalysed dimerisation of isoprene with carbon dioxide in the presence of organotin ethoxide and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (H. Hoberg, M. Minato), (406) C25
- Thiophene, oligomerisation in the presence of PdCl<sub>4</sub><sup>2-</sup> (D.-L. Wang, W.-S. Hwang), (406) C29
- Cyclopalladated *N,N*-dimethylferrocenecarbothioamide, derivatives of (M. Nonoyama, K. Hamamura), (407) 271
- Palladium-catalysed addition of tin–tin and tin–silicon bonds to allenes (T.N. Mitchell, U. Schneider), (407) 319
- Diazadiene-stabilised palladacyclopent-2-enes; catalytic addition of various allylic systems to dimethyl acetylenedicarboxylate (C. Munz, C. Stephan, H. tom Dieck), (407) 413
- Palladium-catalysed reactions involving oxidation state IV, changing selectivity in; new synthesis from iodo benzene and norbornene (M. Catellani, G.P. Chiusoli, C. Castagnoli), (407) C30
- Palladium(0) compounds, oxidative addition of aryl–halogen bonds of *N*-benzylidenbenzylamines to (J. Albert, J. Barro, J. Granell), (408) 115
- Iron(II), tungsten(II) and molybdenum(II) ( $\eta^5$ -iodocyclopentadienyl)metallo methyl derivatives, methyl–iodine exchange in the presence of palladium (C. Lo Sterzo), (408) 253
- [Pd(C≡N(C<sub>6</sub>F<sub>5</sub>)(NCMe))]<sub>2</sub> (C≡N = dimethylbenzylamine-*C*<sup>2</sup>,*N*; 8-quinolylmethyl-*C*,*N*), synthesis and reactivity of; crystal structure of (NBu<sub>4</sub>)<sub>2</sub>[{Pd(8-quinolylmethyl-*C*,*N*)(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>( $\mu$ -Cl)] (J. Forniés, R. Navarro, V. Sicilia, F. Martinez, A.J. Welch), (408) 425
- Homogeneous metal complex catalysts in the synthesis of cyclobutane and cyclopentane compounds (U.M. Dzhemilev, R.I. Khusnutdinov, G.A. Tolstikov), (409) 15
- Palladium-catalysed reactions of vinyl bromides with disubstituted alkynes; new synthesis of fulvenes (L.J. Silverberg, G. Wu, A.L. Rheingold, R.F. Heck), (409) 411

- Double cyclopalladation of diacetylbenzene dioximes and 1,5-bis(dimethylamino)naphthalene (I.G. Phillips, P.J. Steel), (410) 247
- Palladium(II) complexes of 3,3', 5,5'-, and 7,7'-dimethyl-2,2'-biindazole (A.C. Albéniz, J.C. Cuevas, P. Espinet, J. de Mendoza, P. Prados), (410) 257
- Binuclear palladium complexes of isocyanides with novel bridging  $\eta^3$ -indenyl ligands, synthesis and characterisation of; crystal structure of  $[Pd_2(\mu\text{-}\eta^3\text{-indenyl})_2(RNC)_2]$  ( $R = 2,6\text{-dimethylphenyl}$ ) (T. Tanase, T. Nomura, Y. Yamamoto, K. Kobayashi), (410) C25
- Di- $\mu$ -hydroxo-bis[bis(pentachlorophenyl)palladate(II)] ion, synthesis and reactivity of (G. López, J. Ruiz, G. García, J.M. Martí, G. Sánchez, J. García), (412) 435
- Acetylenes and diazoketones in organic synthesis (F. Serratosa), (413) 445
- Depalladation of organopalladium compounds; annulation of aryl rings and formation of pyrones in (M. Pfeffer, M.A. Rotteveel, A. de Cian, J. Fisher, G. le Borgne), (413) C15
- Platinum(IV), platinum(II) and palladium(II) cyclometallates of benzylthio- or benzosulphinyl-substituted azobenzenes, properties of (S. Chattopadhyay, C. Sinha, P. Basu, A. Chakravorty), (414) 421
- Platinum and palladium compound catalysed formation of acetals from alcohols and acetylene (D. Steinborn, R. Nünthel, K. Krause), (414) C54
- Cyclopalladated *p*-azoxyanisole complexes; crystal structure of [*N*-(phenyl)-2-salicylideneaminato]-[4'-(methoxyphenyl-*NNO*-azoxy-*N*<sup>2</sup>)-4-methoxyphenyl-2-ato]palladium(II) (M. Ghedini, S. Morrone, G. De Munno, A. Crispini), (415) 281
- 3,3-Diarylallyl acetates, palladium catalyzed cyclocarbonylation of (M. Iwasaki, Y. Ishii, M. Hidai), (415) 435
- N-bonded  $\eta^1$ -1-azaallyl type complexes of platinum(II) and palladium(II), synthesis and reactivity of; crystal structure of  $[Pt(CH=C(NHCOMe)C(O)\bar{O})(PM_2Ph)_2]$  (R.D.W. Kemmitt, S. Mason, D.R. Russell), (415) C9
- Enantioselective hydrogenation catalysts, some general regularities in the action of (E.I. Klabunovskii), (417) 181
- Palladium carbene cluster; synthesis, structure and reactivity (T.A. Stromnova, I.N. Busygina, D.I. Kochubey, I.I. Moiseev), (417) 193
- Allylamine, the Wacker-type oxidation of (W. Kaufmann, L.M. Venanzi), (417) 205
- Pd<sup>II</sup> and Pd<sup>0</sup> complexes as homogeneous catalysts, reductive carbonylation of nitroaromatic compounds to isocyanates (R. Ugo, R. Psaro, M. Pizzotti, P. Nardi, C. Dossi, A. Andreetta, G. Capparella), (417) 211
- Efficient palladium catalysts for the copolymerization of carbon monoxide with olefins to produce perfectly alternating polyketones (E. Drent, J.A.M. van Broekhoven, M.J. Doyle), (417) 235
- N-alkenylamides and  $\alpha$ -alkenyllactams, amide-directed hydrocarbonylation of (I. Ojima, Z. Zhang), (417) 253
- Palladium-catalyzed reactions of 2-alkynyl caronates with terminal acetylenes, synthesis of 1,2-dien-4-yne (T. Mandai, H. Murayama, T. Nakata, H. Yamaoki, M. Ogawa, M. Kawada, J. Tsuji), (417) 305
- Pd(P<sup>t</sup>Bu<sub>3</sub>)<sub>2</sub>, reactivity towards the oxonium ion; crystal structure of *trans*-[(<sup>t</sup>Bu<sub>3</sub>P)<sub>2</sub>Pd(H)(CH<sub>3</sub>CN)]-BPh<sub>4</sub> (M. Sommovigo, M. Pasquali, P. Leoni, P. Sabatino, D. Braga), (418) 119
- $\sigma$ -Alkyl diketonato and monothio  $\beta$ -diketonato complexes of platinum(II) and palladium(II), synthesis of (H. Jin and K.J. Cavell), (419) 259
- Mixed chalcogen carbonyl compounds. IV. Reactivity of Fe<sub>2</sub>(CO)<sub>6</sub>( $\mu$ -STe) and Fe<sub>3</sub>(CO)<sub>9</sub>( $\mu_3$ -S)( $\mu_3$ -Te) towards coordinatively unsaturated species (P. Mathur, D. Chakrabarty, M.M. Hossain and R.S. Rashid), (420) 79
- Synthesis and nonlinear optical properties of new bimetallic iron/palladium complexes (G. Doisneau, G. Balavoine, T. Fillebeen-Khan, J.-C. Clinet, J. Delaire, I. Ledoux, R. Loucif and G. Puccetti), (421) 299
- A simple synthesis and crystal structure of the dinuclear diphosphido-bridged palladium(I) complex [Pd(P<sup>t</sup>Bu<sub>2</sub>H)( $\mu$ -P<sup>t</sup>Bu<sub>2</sub>)]<sub>2</sub> (P. Leoni, M. Sommovigo, M. Pasquali, P. Sabatino and D. Braga), (423) 263
- Metallation of aliphatic carbon atoms. VI. Reactions of 2-pivaloylpyridine with palladium(II) acetate resulting in cyclopalladation and nucleophilic attacks on carbonyl carbon (K. Hiraki, M. Nakashima, T. Uchiyama and Y. Fuchita), (423) 271
- Chemistry of Pd<sup>II</sup> complexes containing both BINAP (2'2'-bis(diphenylphosphino)binaphthyl) and a  $\eta^3$ -pinene or -allyl ligand. NMR studies on  $[Pd(\eta^3\text{-}C_{10}H_{15})(S(-)BINAP)](CF_3SO_3)$ . The crystal structure of  $[Pd(\eta^3\text{-}C_{10}H_{15})(4,4'\text{-dimethylbipyridine})](CF_3SO_3)$  (C.J. Ammann, P.S. Pregosin, H. Rüegger, A. Albinati, F. Lanza and R.W. Kunz), (423) 415

- Styryl coupling vs. styryl acetate formation in reactions of styryl tellurides with palladium(II) salts (S.U.H. Takahashi and K. Ohe), (423) C9
- Synthesis of monomeric, oligomeric and polymeric  $\sigma$ -acetylide complexes of platinum, palladium, nickel and rhodium (M.S. Khan, S.J. Davies, A.K. Kakkar, D. Schwartz, B. Lin, B.F.G. Johnson and J. Lewis), (424) 87
- Synthesis of aquanitratato(1-methyl-2,2'-bipyridin-3-yl-ium)-palladium(II) perchlorate hydrate (F.L. Wimmer, S. Wimmer and P. Castan), (424) 99
- Organometallic chemistry of diphosphazanes. IV. Reactions of the cyclodiphosphazane, *cis*-[ $^t$ BuNP(OPh)]<sub>2</sub> with M(CO)<sub>6</sub> (M = Cr, Mo, W), rhodium(I), palladium(II) and platinum(II) derivatives (M.S. Balakrishna and S.S. Krishnamurthy), (424) 243
- Alkene insertion reactions of nitrogen-coordinated acylpalladium(II) complexes. The crystal structure of the dicyclopentadiene insertion product [Pd(C<sub>10</sub>H<sub>12</sub>COMe)(bpy)]SO<sub>3</sub>CF<sub>3</sub> (B.A. Markies, M.H.P. Rietveld, J. Boersma, A.L. Spek and G. van Koten), (424) C12
- Carbonylation of alkyltinanium compounds with allylic derivatives (Y. Tsuji and T. Ishii), (425) 41
- Temperature and pressure dependence of the addition of 4-cyanopyridine to tricarbonyl(1-5- $\eta$ -diienyl)iron(II) complexes (diienyl = C<sub>6</sub>H<sub>7</sub> or C<sub>7</sub>H<sub>9</sub>) (T.I. Odiaka and R. van Eldik), (425) 89
- Palladacycle formation by electrophilic aromatic substitution, as monitored by <sup>1</sup>H NMR spectroscopy (M. Catellani and G.P. Chiusoli), (425) 151
- Nucleophilic attack by 2-pyridylpalladium(II) and platinum(II) complexes on the organic chlorides CICH<sub>2</sub>R (R = COMe, CN, Ph, Cl) (B. Crociani, F. Di Bianca, A. Fontana and R. Bertani), (425) 155

## Phosphorus

- Half-sandwich and tripledecker *nido* and *clos*o compounds, a simple MO model to explain the redox behaviour of clusters (B. Eber, D. Buchholz, G. Huttner, Th. Fässler, W. Imhof, M. Fritz, J.C. Jochims, J.C. Daran, Y. Jeannin), (401) 49
- Cyclopentadienyl-ruthenium t-butyl thiol complexes, synthesis of (P.M. Treichel, R.A. Crane, K.N. Haller), (401) 173
- Cyclometallated derivatives of palladium(II) with 1,4-benzodiazepin-2-ones; crystal structure of (L-H)Pd(PPh<sub>3</sub>)Cl·CHCl<sub>3</sub> (L = Prazepam: 7-chloro-1-cyclopropylmethyl-1,3-dihydro-5-phenyl-2*H*-1,4-benzodiazepin-2-one); synthesis of isoindolo[2,1-d][1,4]benzodiazepine derivatives by reaction of [(L-H)PdCl]<sub>2</sub> species with carbon monoxide (M.A. Cinelli, S. Gladiali, G. Minghetti, S. Stoccoro, F. Demartin), (401) 371
- Electron-rich tungsten diethylaminocarbyne complexes, synthesis from Cp(Br)<sub>2</sub>(CO)W≡CNEt<sub>2</sub> (A.C. Filippou, W. Grünleitner, E.O. Fischer), (401) C37
- Dibenzoplatinol system, photochemical and UV-spectroscopic studies of (H.-A. Brune, R. Hohenadel, G. Schmidtberg, U. Ziegler), (402) 179
- Carbene-substituted sulphur-bridged carbonyl iron clusters, synthesis of (B. Eber, G. Huttner, L. Zsolnai, W. Imhof), (402) 221
- Osmium(II) complexes; reactions of coordinated phosphorus trichloride, phenylphosphonous dichloride, 1-phenyl-3,4-dimethylphosphole and mesitylphosphine in (D.S. Bohle, G.R. Clark, C.E.F. Rickard, W.R. Roper, W.B. Shepard), (402) 375
- $\eta^1$ - And  $\eta^3$ -cobalt(I) allyl complexes, reactions with bis(phosphine) ligands; crystal structure of {( $\eta^3$ -C<sub>3</sub>H<sub>5</sub>)Co(CO)<sub>2</sub>}( $\mu$ -dppe-PP) (C. Loubser, H.M. Roos, S. Lotz), (402) 393
- Pyrazolate-bridged dirhodium(I) complex Rh<sub>2</sub>(CO)<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>( $\mu$ -3,5-Me<sub>2</sub>pz)<sub>2</sub>, electrochemistry of (R. Seeber, G. Minghetti, M.I. Pilo, G. Banditelli, S. Zamponi), (402) 413
- Osmium clusters with 1,2-dipolar ligands; synthesis and crystal structure of HOs<sub>3</sub>(CO)<sub>9</sub>( $\mu_3$ - $\eta^2$ -PhNCHCPPh<sub>3</sub>) (M. Langenbahn, H. Stoeckli-Evans, G. Süss-Fink), (402) C12
- trans*-[FeH(P≡CBu<sup>t</sup>)(dppe)<sub>2</sub>](BF<sub>4</sub>)  $\eta^1$ -phosphaphalkyne complex, synthesis of; its conversion into a  $\eta^1$ -fluorophosphaphalkene complex; crystal structure of *trans*-[FeH(PF=CHBu<sup>t</sup>)(dppe)<sub>2</sub>](FeCl<sub>2</sub>F<sub>2</sub>) (P.B. Hitchcock, M.A.N.D.A. Lemos, M.F. Meidine, J.F. Nixon, A.J.L. Pombeiro), (402) C23
- N,N*-Dimethyl-3,5-bis(trifluoromethyl)aniline, regioselective metalation of (N. Kuhn, A. Kuhn, M. Schulten), (402) C41
- [C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub>(ER<sub>3</sub>)]BF<sub>4</sub> complexes (E = N, P, As, Sb and Bi), synthesis and characterisation of; crystal structure of [C<sub>5</sub>H<sub>5</sub>Fe(CO)<sub>2</sub>(Bi(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>)]BF<sub>4</sub> (H. Schumann, L. Eguren), (403) 183
- Ni<sup>0</sup>-induced and -catalysed preparation of unsaturated adipic acid anilides (H. Hoberg, D. Bärhausen), (403) 401

- Nickel(0)-ligand complex, formation of C-C bonds between carbon dioxide and cycloalkenes on (H. Hoberg, A. Ballesteros, A. Sigan), (403) C19
- (Pentadienyl)iron[1,2-bis(diethylphosphino)ethane] electron-rich complexes, synthesis and characterisation of (J.R. Bleeke, R.J. Wittenbrink), (405) 121
- Phosphasilene, synthesis and reactivity of; an improved route to 1-(2,4,6-tri-tert-butylphenyl)-2-tert-butyl-2-(2,4,6-tri-isopropylphenyl)phosphasilene (Y. van den Winkel, H.M.M. Bastiaans, F. Bickelhaupt), (405) 183
- [(-)-Norphos]Ru<sup>II</sup>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)I diastereomers, synthesis and structure of (H. Nishiyama, H. Brunner, P.G. Jones), (405) 247
- Chirally modified ruthenium clusters in the catalytic hydrogenation of racemic mixtures of terpenes; kinetic resolution of the enantiomers in (T. Jenke, G. Süss-Fink), (405) 383
- (Ph<sub>3</sub>P)<sub>2</sub>Pt(H)SiR<sub>3</sub> hydridosilyl complexes, base-induced (Ph<sub>3</sub>P)<sub>2</sub>Pt release from (C. Müller, U. Schubert), (405) C1
- Ketenimine-rhodium complexes, preparation by [2+1]-cycloaddition of vinylidene-rhodium precursors; crystal and molecular structure of C<sub>5</sub>H<sub>5</sub>Rh[η<sup>2</sup>-N,C-TosN=C=CHPh](P<sup>i</sup>Pr<sub>3</sub>) (H. Werner, U. Brekau, M. Dziallas), (406) 237
- 1,2,4,5-Tetrakis(diphenylphosphino)benzene complexes; synthesis and X-ray crystal structure of [{(CO)<sub>3</sub>Fe}<sub>2</sub>{μ-η<sup>2</sup>-η<sup>2</sup>-(Ph<sub>2</sub>P)<sub>2</sub>C<sub>6</sub>H<sub>2</sub>(PPh<sub>2</sub>)<sub>2</sub>}]} (G. Hogarth), (406) 391
- cis-Arylhydridoruthenium(II) chelate complexes as catalysts for the dimerisation of 1-alkynes to linear 1-en-3-yne; molecular structure of (pp<sub>3</sub>)RuH(C<sub>6</sub>H<sub>4</sub>CF<sub>3</sub>)<sub>3</sub> (pp<sub>3</sub> = P(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>e)<sub>3</sub>) (L. Dahlenburg, K.-M. Frosin, S. Kerstan, D. Werner), (407) 115
- Iron-iridium clusters; X-ray structures of Fe<sub>2</sub>Ir(μ<sub>3</sub>-C<sub>2</sub>Ph)(CO)<sub>8</sub>(PPh<sub>3</sub>), FeIr<sub>2</sub>(μ<sub>3</sub>-η<sup>2</sup>-PhC<sub>2</sub>C<sub>2</sub>Ph)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub> and Fe<sub>2</sub>Ir(μ<sub>3</sub>-PhC<sub>2</sub>PEt<sub>3</sub>)(CO)<sub>7</sub>(PEt<sub>3</sub>) (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tiekkink), (407) 391
- Nickela compounds with a five-membered ring, ring contraction of; use of the resulting four-membered ring compounds in preparative chemistry (H. Hoberg, A. Ballesteros, A. Sigan, C. Jégat, D. Bärhausen, A. Milchereit), (407) C23
- Reactions of some Fe-Ir clusters; crystal structures of Fe<sub>2</sub>Ir(μ-H)(μ<sub>3</sub>-CCHPh)(CO)<sub>8</sub>(PPh<sub>3</sub>) and Au<sub>2</sub>Fe<sub>2</sub>Ir(μ<sub>4</sub>-C<sub>2</sub>Ph)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>3</sub> (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tiekkink), (408) 77
- Dicarbonylbis(phosphorus-donor)iron complexes, trimerisation of methyl propiolate on (U. Grössmann, H.-U. Hund, H.W. Bosch, H. Schmalle, H. Berke), (408) 203
- Cyclopentadienyliron arene cations, photochemical arene substitution in; synthesis of cyclopentadienyl-iron bisphosphine ligand complexes with group 14 and 15 ligands (H. Schumann, L. Eguren, J.W. Ziller), (408) 361
- Trimethylphosphine-substituted η<sup>2</sup>-thiocarbene complexes of tungsten, preparation and reactions of (W. Schütt, N. Ullrich, F.R. Kreißl), (408) C5
- Phenylphosphole, oxidative addition of; X-ray crystal structures of two ring-opened products [Os<sub>3</sub>(μ<sub>3</sub>-PhPC<sub>4</sub>H<sub>4</sub>)(CO)<sub>x</sub>] where x = 8 or 9 (A.J. Arce, Y. De Sanctis, J. Manzur, A.J. Deeming, N.I. Powell), (408) C18
- Silyl-substituted phosphinomethanide ligands; preparation of dimeric Al heterocycles; crystal structure of {Me<sub>2</sub>Al[C(PMe<sub>2</sub>)<sub>2</sub>(SiMe<sub>3</sub>)<sub>2</sub>]}<sub>2</sub> (H.H. Karsch, K. Zellner, J. Lachmann, G. Müller), (409) 109
- Vinylphosphines, preparation by means of free radical addition of diphenylphosphine to alkynes and allenes (T.N. Mitchell, K. Heesche), (409) 163
- Diphosphine-derived radical cations, molecular and electronic structures of; a semi-empirical PM3 study (C. Glidewell), (409) 171
- Mesitylene osmium complexes with hydrido, carboxylato, carbonyl and vinylidene ligands (H. Werner, S. Stahl, W. Kohlmann), (409) 285
- 2-3-η<sup>2</sup>-(Butyne-1,4-diol)bis(triisopropylphosphane)nickel(0), (iPr<sub>3</sub>P)<sub>2</sub>Ni(HOCH<sub>2</sub>C≡CCH<sub>2</sub>OH), crystal and molecular structure of; some peculiarities of the complexation of 2-butyne-1,4-diol with nickel(0) complexes (U. Rosenthal, G. Oehme, H. Görts, V.V. Burlakov, A.V. Polyakov, A.I. Yanovsky, Yu.T. Struchkov), (409) 299
- Ni<sup>0</sup> catalysed reactions between 1,3-butadiene and phenylisocyanate (H. Hoberg, D. Bärhausen, R. Mynott, G. Schroth), (410) 117
- Carbonylchromium, carbonylmolybdenum and carbonyltungsten complexes of the asymmetrically substituted phosphino sulfur diimides <sup>1</sup>Bu<sub>2</sub>P(NSN)PPh<sub>2</sub> and <sup>1</sup>Bu<sub>2</sub>P(NSN)P(<sup>1</sup>Bu)Ph; <sup>1</sup>H, <sup>13</sup>C and <sup>31</sup>P NMR spectroscopic characterisation of (M. Herberhold, S.M. Frank, B. Wrackmeyer), (410) 159

- 9,9-(1,2,5,6- $\eta^4$ -Cycloocta-1,5-diene)-3,6-bis(tert-butyl)dibenzoplatinol and 9,9-bis(triphenylphosphine)-3,6-bis(trifluoromethyl)dibenzoplatinol, crystal and molecular structures of (T. Debaerdemaeker, R. Hohenadel, H.-A. Brune), (410) 265
- Cyclopentadienyliron bis(phosphine) complexes with dimethylchalcogenide ligands; synthesis, reactivity and crystal structure of  $[C_5H_5Fe(P(OCH_3)_3)_2(S(CH_3)_2)]PF_6$  (H. Schumann, J.W. Ziller), (410) 365
- Octacarbonyldicobalt, reductive dehalogenation of chloro(alkynyl)phosphines by (H. Lang, M. Leise, L. Zsolnai), (410) 379
- cis*-Fe(CO)<sub>4</sub>(SiCl<sub>3-n</sub>Me<sub>n</sub>)<sub>2</sub> ( $n = 1-3$ ), conversion with phosphines; competition between CO substitution, SiR<sub>3</sub> elimination and formation of SiR<sub>2</sub> bridged complexes (U. Schubert, M. Knorr, C. Straßer), (411) 75
- (CO)<sub>5-n</sub>(R'<sub>3</sub>P)<sub>n</sub>M(H)SiR<sub>3</sub> ( $n = 2,3$ ), synthesis of phosphane substituted hydridosilyl complexes of chromium, molybdenum and tungsten; synthesis of (CO)<sub>3</sub>(L-L)M(H)Cl by conversion of (CO)<sub>3</sub>(L-L)M(NCR') with HSiClR<sub>2</sub> (L-L = tmeda, bipy, dppe, (PPh<sub>3</sub>)<sub>2</sub>) (H. Piana, U. Schubert), (411) 303
- Alkylvanadium complexes, synthesis and characterisation of; molecular structure of *trans*-[V(CO)<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>] (D. Rehder, F. Süßmilch, W. Priebisch, M. Fornalczyk), (411) 357
- L<sub>4</sub>Ru<sup>0</sup> complex cores, inter- and intramolecular cleavage of  $sp^2$ - and  $sp^3$ -C-H bonds with P(o-C<sub>6</sub>H<sub>4</sub>PM<sub>2</sub>e<sub>2</sub>)<sub>3</sub>, N(CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>e<sub>2</sub>)<sub>3</sub>, MeP(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>e<sub>2</sub>)<sub>2</sub>/PR<sub>3</sub> and MeSi(CH<sub>2</sub>PM<sub>2</sub>e<sub>2</sub>)<sub>3</sub>/PR<sub>3</sub> (R = Me, OMe) as supporting ligands (L. Dahlenburg, S. Kerstan, D. Werner), (411) 457
- [C<sub>6</sub>H<sub>6</sub>Os(PR<sub>3</sub>)] as structural element; synthesis and reactions with alkinyl, vinylidene, and vinylosmium(II) complexes (W. Knaup, H. Werner), (411) 471
- (OC)<sub>2</sub>LCO( $\eta^2$ -PR'<sub>2</sub>  $\rightleftharpoons$  PR<sup>2</sup>); synthesis, properties and structures of this type of ( $\eta^2$ -phosphanediylphosphinito)cobalt complexes (E. Lindner, T. Funk, W. Hiller, R. Fawzi), (411) 491
- Polycyclic silacyclobutanes, Rh-catalyzed synthesis from norbornadiene and diphenylsilane (H. Brunner, F. Prester), (411) C1
- Ni<sup>0</sup> induced synthesis of cyclic C<sub>8</sub> carbon acids from cyclooctenes and carbon dioxide (H. Hoberg, A. Ballesteros), (411) C11
- [C<sub>5</sub>H<sub>5</sub>]Ru(dppm)(CH<sub>2</sub>=SO<sub>2</sub>)<sup>+</sup>, a strong organometallic electrophile (W.A. Schenk, P. Urban), (411) C27
- Ni<sup>0</sup> induced CC-coupling between phenylisocyanate and five-membered cyclic alkenes, catalytic formation of  $\beta,\gamma$ -unsaturated carboxylic acid anilides (H. Hoberg, M. Nohlen), (412) 225
- P-substituted olefins, base-catalysed addition of optically active P-H complexes (H. Brunner, S. Limmer), (413) 55
- Chiral ferrocenylbisphosphines containing an aminoalkyl side chain, preparation of (T. Hayashi, A. Yamazaki), (413) 295
- Tetranuclear rhodium-triruthenium carbonyl clusters containing phosphorus(III) ligands; crystal and molecular structure of [RhRu<sub>3</sub>( $\mu$ -H)<sub>2</sub>( $\mu$ -CO)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub>( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)] (J.-L. Le Grand, W.E. Lindsell, K.J. McCullough), (413) 321
- Thiophene complexes of cobalt (R. Albrecht, E. Weiss), (413) 355
- Four-membered cobaltahterocycles from C<sub>5</sub>H<sub>5</sub>Co(CNR)(PM<sub>2</sub>e<sub>3</sub>) and isocyanates or isothiocyanates, synthesis and reactivity of; unusual fragmentation of the chiral isocyanide ligand (S)-CNCH(Me)Ph (H. Werner, B. Strecker), (413) 379
- Platinum-indium and cobalt-gallium complexes, synthesis of; crystal structure of [Cy<sub>2</sub>P(CH<sub>2</sub>CH<sub>2</sub>)PCy<sub>2</sub>]-Pt(Ind<sub>2</sub>)(R) (Cy = cyclo-C<sub>6</sub>H<sub>11</sub>) (R.A. Fischer, J. Behm), (413) C10
- [Fe<sub>4</sub>(CO)<sub>10</sub>(PR)<sub>2</sub>( $\mu_2$ -COEt)<sub>2</sub>]; functionalisation of the CO group of an anionic radical cluster (T. Feng, P. Lau, W. Imhof, G. Huttner), (414) 89
- Lithium, samarium and ytterbium diiminophosphinates; crystal structures of Li[Ph<sub>2</sub>P(NSiMe<sub>3</sub>)<sub>2</sub>](THF)<sub>2</sub> and [Ph<sub>2</sub>P(NSiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub>Sm( $\mu$ -I)<sub>2</sub>Li(THF)<sub>2</sub> (A. Recknagel, A. Steiner, M. Noltemeyer, S. Brooker, D. Stalke, F.T. Edelmann), (414) 327
- Complexed triphosphanes, preparation and characterisation of; crystal structure of HP[PPh<sub>2</sub>Cr(CO)<sub>5</sub>]<sub>2</sub> (M. Scheer, S. Gremler, E. Herrmann, P.G. Jones), (414) 337
- closو*-Clusters, carbene-ligand assisted formation of; test of a model to clarify the redox activity of *nido*- and *closو*-compounds (B. Eber, G. Huttner, D. Günauer, W. Imhof, L. Zsolnai), (414) 361
- Enantioselective Co-catalysed homo Diels-Alder reaction of norbornadiene with acetylenes (H. Brunner, F. Prester), (414) 401
- Lithium-coordinated phosphinomethanides, influence of Co-ligands upon; crystal structure of ((THF)Li[C(SiMe<sub>3</sub>)<sub>2</sub>(PM<sub>2</sub>e<sub>2</sub>)])<sub>2</sub> (H.H. Karsch, K. Zellner, S. Gamper, G. Müller), (414) C39

- [ $\text{Mo}(\text{CO})_2(\text{L})^{\text{bu}}\text{S}_4$ ]), complexes with sterical tetradeinate thioether thiolate ligands  ${}^{\text{bu}}\text{S}_4^{2-}$  = 1,2-bis(2-mercapto-3,5-di-t-butylphenylthio)ethan(2-) ( $\text{L} = \text{CO, PMe}_3, \text{PPh}_3$ ) (D. Sellmann, F. Grasser, M. Moll), (415) 367
- Para- and diamagnetic nickel complexes, tetra-tert-butyl-dihydropentafulvene as educt for (P. Jutzi, J. Schnittger, W. Wieland, B. Neumann, H.-G. Stammler), (415) 425
- Di- and tri-phosphollide ions and their ferrocene analogues; synthesis of (R. Bartsch, J.F. Nixon), (415) C15
- P*-substituted olefins, addition of optically active amines to (H. Brunner, S. Limmer), (417) 173
- Dicarbollyl complexes of phosphor and arsenic, X-ray diffraction study on  $\text{ClAs}[\text{C}_2\text{Me}_2\text{B}_9\text{H}_9]$  (P. Jutzi, D. Wegener, M. Hursthouse), (418) 277
- (dppe) $\text{Pt}(\text{Me}_2\text{SiCH}_2\text{CH}_2\text{SiMe}_2)$ , synthesis by exchange of silyl groups (U. Schubert, C. Müller), (418) C6
- [ $(2,4,6-{}^1\text{Bu}_3\text{C}_6\text{H}_2\text{OX}^c\text{C}_5\text{Me}_5)\text{P}=\text{Mn}(\text{CO})_4$ ], a stable  $\lambda^4$ -phosphanediyi complex containing a multiple phosphor-manganese bond (H. Lang, M. Leise, C. Emmerich), (418) C9
- Opening of cluster frameworks, increase in skeletal electrons via amino substituents (B. Eber, G. Huttner, C. Emmerich, J.C. Daran, B. Heim and Y. Jeannin), (419) 43
- New family of water-soluble tertiary phosphine ligands and their rhodium(I) complexes, synthesis of; olefin hydrogenation in aqueous and biphasic media (E. Renaud, R.B. Russell, S. Fortier, S.J. Brown and M.C. Baird), (419) 403
- Crystal structure and reactivity of [ $(\text{pp}_3)\text{Rh}(\text{H})(\text{SeCH}_3)_3\text{BPh}_4$ ]. An organometallic complex from metal-assisted Se-H activation (M. Di Vaira, D. Rovai, P. Stoppioni and M. Peruzzini), (420) 135
- The structure of 2-(diphenylphosphino)phenyllithium: the significance of P-Li bonding (S. Harder, L. Brandsma, J.A. Kanders, A. Duisenberg and J.H. van Lenthe), (420) 143
- $[\text{Ph}_4\text{P}] [\text{CpTi}(1,2-\text{S}_2\text{C}_6\text{H}_4)_2]$ : Synthese und Molekülstruktur eines anionischen, tetragonal-pyramidal koordinierten Benzol-1,2-dithiolato-Komplexes des Titans (H. Köpf, K. Lange and J. Pickardt), (420) 345
- Heteronucleare Komplexverbindungen mit Metall-Metall-Bindungen. VI. Kleine heteronucleare Cluster mit Kupfer und  $\text{HCr}(\text{CO})_5$ -Fragmenten. Die Strukturen von  $[(N_2)\text{Cu}(\mu_3\text{-H})\text{Cr}_2(\text{CO})_{10}]$ , ( $N_2 = 2 \text{ NH}_3$ , bpy, 1,2-Diaminobenzol),  $[(\text{py})\text{Cu}(\mu_3\text{-H})\text{Cr}(\text{CO})_5]_2$  und  $[\text{Cu}(\text{PPh}_3)_3][\text{Cr}_2(\mu\text{-H})(\text{CO})_{10}]$  sowie die Struktur von  $[\text{Zn}(\text{NH}_3)_4][\text{Cr}_2(\mu\text{-H})(\text{CO})_{10}]_2$  (P. Klüfers und U. Wilhelm), (421) 39
- Kohlenwasserstoffverbrückte Komplexe. XXIII. Heterobimetallische Komplexe mit Ferrocenyl- und  $(\text{Ph}_3\text{P})(\text{OC})(\text{Cp})\text{Fe}(\text{C})(\text{O})\text{CH}_2$ -Gruppen; Darstellung und Struktur von  $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}[\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{Re}(\text{CO})_5]$ ,  $(\text{PPh}_3)(\text{OC})(\text{Cp})\text{Fe}(\text{C})(\text{O})\text{CH}_2[\eta^6\text{-C}_7\text{H}_7]\text{M}(\text{CO})_3$  und  $[\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{CH}_3]\text{Fe}[\eta^5\text{-C}_5\text{H}_4\text{C}(\text{O})\text{CH}_2][\eta^6\text{-C}_7\text{H}_7]\text{M}(\text{CO})_3$  ( $\text{M} = \text{Cr, Mo}$ ) (J. Breimair, M. Wieser, B. Wagner, K. Polborn und W. Beck), (421) 55
- Darstellung und Struktur von  $(\text{EtO})_2\text{P}(\text{O})\text{CH}_2\text{Si}(\text{Me})_2\text{CH}_2\text{SnMe}_2\text{Cl}$ —ein Sechsringchelat mit Seskelkonformation und  $\text{P}=\text{O} \cdots \text{Sn}(\text{Cl})\text{Me}_2\text{CH}_2$ -trigonaler Bipyramide am Lewis-aciden Zinn (U. Kolb, M. Dräger, E. Fischer und K. Jurkschat), (423) 339
- Untersuchungen zum Mechanismus der reduktiven Photo-Eliminierung des Biphenyl-Systems aus *cis*-Bis(phenyl)platin(II)-Verbindungen (H.-A. Brune, E. Hupfer, G. Schmidtberg und A. Baur), (424) 225
- Organometallic chemistry of diphosphazanes. IV. Reactions of the cyclodiphosphazane, *cis*-[ $'\text{BuNP}(\text{OPh})_2$ ] with  $\text{M}(\text{CO})_6$  ( $\text{M} = \text{Cr, Mo, W}$ ), rhodium(I), palladium(II) and platinum(II) derivatives (M.S. Balakrishna and S.S. Krishnamurthy), (424) 243
- Zur Kenntnis funktionalisierter Zinn-Phosphorverbindungen: Darstellung und Reaktionen von Chlorstannylyphosphinen (D. Hänssgen, E. Stahlhut, H. Aldenhoven and A. Dörr), (425) 19
- Synthesis and reactivity of rhenium(I) isocyanide complexes. Crystal and molecular structures of  $\text{ReBr}(\text{CO})_4(\text{CNHPh})(\text{NHCHMe}_2)$  and  $\text{ReBr}(\text{CO})_3(\text{PPh}_3)(\text{CNHPh})(\text{NHCHMe}_2)$  (L.-C. Chen, M.-Y. Chen, J.-H. Chen, Y.-S. Wen and K.-L. Lu), (425) 99
- Enantioselektive Katalysen. LXXVIII. Der optisch aktive Phosphido-verbrückte Enantioselektive Katalysen Eisenkomplex  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Fe}_2(\mu\text{-H})(\mu\text{-PMen}_2)(\text{CO})_2]$  — Synthese, Isomerisierung, Katalysen (H. Brunner and M. Rötzer), (425) 119

## Platinum

- $(\text{CO})_6\text{Fe}_2(\mu_3\text{-Se})(\mu_3\text{-Te})\text{Pt}(\text{PPh}_3)_2$  and  $\text{Fe}_3(\text{CO})_7(\text{PPh}_3)_2(\mu_3\text{-Se})(\mu_3\text{-Te})$ , synthesis and characterisation of (P. Mathur, D. Chakrabarty, Md.M. Hossain), (401) 167

- 2-Substituted 8-quinolinols containing C, N, O-donor atoms and their Pd<sup>II</sup> or Pt<sup>II</sup> complexes, preparation of (A. Yoneda, G.R. Newkome, K.J. Theriot), (401) 217
- Pt(II) complexes of tetrathia[n](1,1')ferrocenophanes, preparation and properties of (M. Sato, H. Asano, S. Akabori), (401) 363
- Chiral metal complexes of 1-dimethylphosphino-3-dimethylamines, homogeneous asymmetric catalysis by means of (C. Döbler, A. Kinting), (401) C23
- Platinum-catalysed elimination-addition reactions of trisilylethenes (T. Suzuki, P.Y. Lo), (402) 27
- Platinum or palladium 3-diphenylphosphinocamphor complexes including acetylides (S.D. Perera, B.L. Shaw), (402) 133
- Dibenzoplatinol system, photochemical and UV-spectroscopic studies of (H.-A. Brune, R. Hohenadel, G. Schmidtberg, U. Ziegler), (402) 179
- Bis[tetra(n-butyl)ammonium]bis[ $\{\mu\text{-bromo}\}2,2'\text{-}\eta^2\text{-}[4,4'\text{-bis(trifluoromethyl)biphenyl}]platinate(II)$ ], synthesis and structure of (H.-A. Brune, H. Roth, T. Debaerdemaeker, H.-M. Schiebel), (402) 435
- Five-coordinate stable [Pt( $\overline{NN'}$ Xolefin)(R)X] complexes formed by oxidative addition to [Pt( $\overline{NN'}$ Xolefin)] precursors (V. De Felice, M. Funicello, A. Panunzi, F. Ruffo), (403) 243
- Mono- and polynuclear perhalophenyl palladium-platinum acetylide complexes, synthesis of; molecular structure of ( $\text{NBu}_4$ )<sub>2</sub>[Pt<sub>2</sub>Ag<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>(C≡CPh)<sub>4</sub>]·4CH<sub>2</sub>Cl<sub>2</sub> (P. Espinet, J. Forniés, F. Martínez, M. Sotes, E. Lalinde, M.T. Moreno, A. Ruiz, A.J. Welch), (403) 253
- Five-coordinate trigonal-bipyramidal palladium(II) species, stabilisation of; crystal structure of (2,9-dimethyl-1,10-phenanthroline)methylchloropalladium(II) (V. De Felice, V.G. Albano, C. Castellari, M.E. Cucciolito, A. De Renzi), (403) 269
- C-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325
- Low-valent platinum complexes, ketene formation by reactions with acid chlorides (Y. Ishii, Y. Kobayashi, M. Iwasaki, M. Hidai), (405) 133
- Phenylselenolato-bridged dinuclear platinum(II) complexes, preparation and characterisation of (V.K. Jain, S. Kannan), (405) 265
- Platinum(II) chloride complexes of 1,1'-bis(methylthio)ferrocene, 1,1'-bis(methylseleno)ferrocene and 1,1'-bis(methylthio)ruthenocene, synthesis and stereodynamics of (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šik), (405) 375
- (Ph<sub>3</sub>P)<sub>2</sub>Pt(H)SiR<sub>3</sub> hydridosilyl complexes, base-induced (Ph<sub>3</sub>P)<sub>2</sub>Pt release from (C. Müller, U. Schubert), (405) C1
- Platinum(II), binuclear methyl and phenyl derivatives of (M. Rashidi, Z. Fakhroian, R.J. Puddephatt), (406) 261
- Platinum-silver cluster compound, [Pt<sub>3</sub>Ag(O<sub>3</sub>SCF<sub>3</sub>)(μ-CO)<sub>3</sub>(PCy<sub>3</sub>)<sub>2</sub>(dppp)] (Cy = C<sub>6</sub>H<sub>11</sub>, dppp =  $\{(C_6H_5)_2P\}_2C_3H_6$ ), with a distorted tetrahedral geometry; synthesis and crystal structure of (A.D. Burrows, J.G. Jeffrey, J.C. Machell, D.M.P. Mingos), (406) 399
- Hydrosilylation of the C=C double bond employing homogeneous catalysis with transition metals (R. Skoda-Földes, L. Kollár, B. Heil), (408) 297
- 9,9-(1,2,5,6- $\eta^4$ -Cycloocta-1,5-diene)-3,6-bis(tert-butyl)dibenzoplatinol and 9,9-bis(triphenylphosphine)-3,6-bis(trifluoromethyl)dibenzoplatinol, crystal and molecular structures of (T. Debaerdemaeker, R. Hohenadel, H.-A. Brune), (410) 265
- [PtMe(pp<sub>3</sub>)]<sup>+</sup> (pp<sub>3</sub> = P(CH<sub>2</sub>ClI<sub>2</sub>PPH<sub>2</sub>)<sub>3</sub>) five-coordinate organoplatinum cation, synthesis by oxidative addition of methyl iodide to [Pt<sub>2</sub>(μ-pp<sub>3</sub>)<sub>2</sub>] (M. Peter, M. Probst, P. Peringer, E.P. Müller), (410) C29
- cis-Bis(tetrahydrothiophene)bis( $\eta^2$ -biphenyl-2,2'-diyl)platinum(IV) complexes, synthesis of (H.-A. Brune, H. Roth, G. Schmidtberg), (412) 237
- cis-Bis(tetrahydrothiophene)bis( $\eta^2$ -[4,4'-bis(trifluoromethyl)biphenyl-2,2'-diyl])platinum(IV), crystal and molecular structure of (T. Debaerdemaeker, H. Roth, H.-A. Brune), (412) 243
- [Cp(OC)<sub>2</sub>Fe(olefin)]<sup>+</sup> and (1,5-hexadiene)PtCl<sub>2</sub>, addition of α-amino acids to coordinated olefins of (I. Zahn, K. Polborn, W. Beck), (412) 397
- Platina(IV)cyclopentane complex, synthesis and characterisation of (S. Tangestaninejad, M. Rashidi, R.J. Puddephatt), (412) 445
- Five-coordinate platinum(II) hydride complexes, synthesis and NMR spectroscopy of (A. Handler, P. Peringer, E.P. Müller), (412) 451
- Platinum-indium and cobalt-gallium complexes, synthesis of; crystal structure of [Cy<sub>2</sub>P(CH<sub>2</sub>CH<sub>2</sub>)PCy<sub>2</sub>]-Pt(InR<sub>2</sub>)<sub>x</sub>(R) (Cy = cyclo-C<sub>6</sub>H<sub>11</sub>) (R.A. Fischer, J. Behm), (413) C10
- Platinum(IV), platinum(II) and palladium(II) cyclometallates of benzylthio- or benzosulphinyl-substituted azobenzenes, properties of (S. Chattopadhyay, C. Sinha, P. Basu, A. Chakravorty), (414) 421

- Platinum and palladium compound catalysed formation of acetals from alcohols and acetylene (D. Steinborn, R. Nünthel, K. Krause), (414) C54
- $\alpha$ -Ketoacyl complex with planar *s-cis* oxaryl configuration; crystal structure of *trans*-[Pt(COCOOMe)(OH<sub>2</sub>)(PPh<sub>3</sub>)<sub>2</sub>]CF<sub>3</sub>SO<sub>3</sub> (J.-T. Chen, Y.-S. Yeh, G.-H. Lee, Y. Wang), (414) C64
- N-bonded  $\eta^1$ -1-azaallyl type complexes of platinum(II) and palladium(II), synthesis and reactivity of; crystal structure of [Pt(CH=C(NHCOME)C(O)O)(PMe<sub>2</sub>Ph)<sub>2</sub>] (R.D.W. Kemmitt, S. Mason, D.R. Russell), (415) C9
- 2-Pyridyl(phosphine) complexes of platinum(0), synthesis and chemistry of (Y. Xie, B.R. James), (417) 277
- Alkene complexes of platinum(0), hydroformylation of olefins catalyzed by (C. Botteghi, S. Paganelli), (417) C41
- Facile cycloplatination of nitrogen compounds; crystal structure of the cycloplatinated Schiff's base tetralone derivative PtCl(cyclohexy)N=C(CH<sub>2</sub>)<sub>3</sub>C<sub>6</sub>H<sub>3</sub>(CO) (P.S. Pregosin, F. Wombacher, A. Albinati, F. Lianza), (418) 249
- Aryltellurolate-bridged dinuclear complexes of platinum(II), preparation and characterization of (V.K. Jain, S. Kannan), (418) 349
- Chiral derivatising agent  $\eta^2$ -ethene platinum-DIOP, crystal structure and further applications of (R. Fulwood, D. Parker, G. Ferguson and B. Kaltner), (419) 269
- Mixed chalcogen carbonyl compounds. IV. Reactivity of Fe<sub>2</sub>(CO)<sub>6</sub>( $\mu$ -STe) and Fe<sub>3</sub>(CO)<sub>9</sub>( $\mu_3$ -S)( $\mu_3$ -Te) towards coordinatively unsaturated species (P. Mathur, D. Chakrabarty, M.M. Hossain and R.S. Rashid), (420) 79
- Hydrosilylation of olefins with monosilane catalyzed by transition metal complexes (M. Itoh, K. Iwata, R. Takeuchi and M. Kobayashi), (420) C5
- New [PtCl(monophosphine)(chiral bisphosphine)]<sup>+</sup> cations (L. Kollár and G. Szalontai), (421) 341
- Synthesis of new vinylsilanes containing an asymmetric silicon via platinum catalyzed hydrosilylation of acetylene and monosubstituted alkynes (V. Gevorgyan, L. Borisova, J. Popelis, E. Lukevics, Z. Foltynowic, J. Gulinski and B. Marciniec), (424) 15
- The synthesis and reactivity of perfluorinated undecafluorobicyclo[2.2.1]-heptan-1-yl platinum(II), palladium(II) and rhenium(V) complexes and the X-ray crystal structure of *cis*-PtCl(C<sub>7</sub>F<sub>11</sub>XPPPh<sub>3</sub>)<sub>2</sub> (P.P.K. Claire, C.J. Jones, J.A. McCleverty, P.L. Coe and M.G.B. Drew), (424) 105
- Synthesis and characterization of trimethylplatinum(IV) complexes with diphenyldithiophosphinic acid. Crystal and molecular structures of [PtMe<sub>3</sub>(SSPPPh<sub>2</sub>)<sub>2</sub>] and [PtMe<sub>3</sub>(py)(SSPPPh<sub>2</sub>)] (S. Chaudhury, V.K. Jain, V.S. Jakkal and K. Venkatasubramanian), (424) 115
- Untersuchungen zum Mechanismus der reduktiven Photo-Eliminierung des Biphenyl-Systems aus *cis*-Bis(phenyl)platin(II)-Verbindungen (H.-A. Brune, E. Hupfer, G. Schmidtberg und A. Baur), (424) 225
- Organometallic chemistry of diphosphazanes. IV. Reactions of the cyclodiphosphazane, *cis*-[<sup>1</sup>BuNP(OPh)]<sub>2</sub> with M(CO)<sub>6</sub> (M = Cr, Mo, W), rhodium(I), palladium(II) and platinum(II) derivatives (M.S. Balakrishna and S.S. Krishnamurthy), (424) 243
- Structural studies of unusually disordered diorganoplatinum(IV) complexes containing the cations [PtR<sub>2</sub>(L-N,N',N'')]<sup>+</sup>, where the ligands L are facially coordinated (A.J. Canty, R.T. Honeyman, B.W. Skelton and A.H. White), (424) 381
- Nucleophilic attack by 2-pyridylpalladium(II) and platinum(II) complexes on the organic chlorides CICH<sub>2</sub>R (R = COMe, CN, Ph, Cl) (B. Crociani, F. Di Bianca, A. Fontana and R. Bertani), (425) 155
- Di-, tri-, pseudo-di- and pseudo-tetra-acetylenic polymers of platinum: synthesis, characterization and optical spectra (J. Lewis, M.S. Khan, A.K. Kakkar, B.F.G. Johnson, T.B. Marder, H.B. Fyfe, F. Wittmann, R.H. Friend and A.E. Dray), (425) 165
- Synthesis and characterization of five-coordinate platinum(II) complexes [Pt(2,9-dimethyl-1,10-phenanthroline)(SnR<sub>n</sub>X<sub>3-n</sub>)X(olefin)] (X = Cl or Br; R = Ph or Me). Molecular structure of [Pt(2,9-dimethyl-1,10-phenanthroline)(SnPh<sub>2</sub>Cl)Cl(ethylene)] (V.G. Albano, C. Castellari, V. De Felice, A. Panunzi and F. Ruffo), (425) 177

## Potassium

- 2,4-Di(t-butyl)pentadienyl anion and its Ti(II), Cr(II), and Zn(II) complexes; synthesis of (R.D. Ernst, J.W. Freeman, P.N. Swepston, D.R. Wilson), (402) 17

- Sodium and potassium pentamethylcyclopentadienyl pyridine solvates, preparation and crystal structures of (G. Rabe, H.W. Roesky, D. Stalke, F. Pauer, G.M. Sheldrick), (403) 11
- Potassium/naphthalene induced carbon–hydrogen bond activation; selective catalytic dehydrogenation of 1,4-cyclohexadiene to benzene; isomerisation of linear 1,4- to 2,4-hexadienes (G. Fochi, L. Nucci), (407) 279
- Allyl- and benzyl-type organoalkali intermediates, preparation via organotin compounds (O. Desponts, M. Schlosser), (409) 93
- Allyl alkali metals, comparative theoretical study of (N.J.R. van Eikema Hommes, M. Bühl, P.v.R. Schleyer, Y.-D. Wu), (409) 307
- Tetra-tert-butyl-pentafulvalene as ligand in dinuclear molybdenum complexes; *cis*-configuration without a metal–metal bond (P. Jutzi, J. Schnittger, B. Neumann, H.-G. Stammmer), (410) C13
- Potassium alcoholate/crown ether dissolved in hydrocarbon phase as catalyst system for the synthesis of vinyl ethers from acetylenes and alcohols (D. Steinborn, H. Mosinski, T. Rosenstock), (414) C45
- Alkylarenes, optional *ortho* or alpha hydroxymethylation (Y. Guggisberg, F. Faigl, M. Schlosser), (415) 1
- Alkali metal derivatives of sulfinimidamides, preparation and crystal structures (F. Pauer, D. Stalke), (418) 127

### Praseodymium

- Bis(t-butylcyclopentadienyl)lanthanoid chloride complexes, syntheses of; crystal structures of bis(t-butylcyclopentadienyl)chloro(bistetrahydrofuran)praseodymium and bis(t-butylcyclopentadienyl)-chlorotetrahydrofuranytterbium (Q. Shen, M. Qi, J. Guan, Y. Lin), (406) 353
- Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXX. Kristall- und Molekülstrukturen ungewöhnlich gut kristallisierender Addukte des Grundkörpers Tris( $\eta^5$ -cyclopentadienyl)-lanthanoid(III) (H. Schulz, H. Schultze, H. Reddmann, M. Link und H.-D. Amberger), (424) 139
- Synthesis and reactivity of rhenium(I) isocyanide complexes. Crystal and molecular structures of  $\text{ReBr}(\text{CO})_4\{\text{CNHPh}\}\{\text{NHCHMe}_2\}$  and  $\text{ReBr}(\text{CO})_3(\text{PPh}_3)\{\text{CNHPh}\}\{\text{NHCHMe}_2\}$  (L.-C. Chen, M.-Y. Chen, J.-H. Chen, Y.-S. Wen and K.-L. Lu), (425) 99

### Rhenium

- Metalloporphyrins with metal–metal  $\sigma$ -bonds; synthesis, spectroscopic characterization, and electrochemistry of (P)MRe(CO)<sub>5</sub> where P is the dianion of octaethylporphyrin (OEP) or tetraphenylporphyrin (TPP) and M = Al, Ga, In or Tl (R. Guilard, A. Zrineh, A. Tabard, L. Courthaudon, B. Han, M. Ferhat, K.M. Kadish), (401) 227
- $\text{M}_2(\text{CO})_{10}$  (M = Mn, Re), mechanisms of the photochemical reactions with *ortho*- and *para*-quinones; a combined EPR and FTIR study (T. van der Graaf, D.J. Stufkens, J. Víchová, A. Vlček, Jr.), (401) 305
- Re(2,4-C<sub>7</sub>H<sub>11</sub>)(H)<sub>2</sub>(P(C<sub>6</sub>H<sub>5</sub>)<sub>3</sub>)<sub>2</sub>; synthesis, characterization, and structural study of (T.E. Waldman, A.L. Rheingold, R.D. Ernst), (401) 331
- [Re(NCC<sub>6</sub>H<sub>4</sub>Me-4)<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>]ReF<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>[BF<sub>4</sub>]<sub>2</sub> complex double salt, synthesis and crystal structure of (M.F.C.G. Silva, A.J.L. Pombeiro, A. Hills, D.L. Hughes, R.L. Richards), (403) C1
- Clusters with Re<sub>2</sub>Fe and Re<sub>2</sub>Ni<sub>2</sub> cores, phenylacetylene transformations in; synthesis and molecular structure of Re<sub>2</sub>Fe(CO)<sub>9</sub>( $\mu_3$ , $\eta^4$ -CH=CHC<sub>6</sub>H<sub>4</sub>) and Cp<sub>2</sub>Ni<sub>2</sub>Re<sub>2</sub>(CO)<sub>9</sub>( $\mu_4$ -PhC≡CCH=CHPh) (A.D. Shaposhnikova, G.L. Kamalov, R.A. Stadnichenko, A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, Yu.T. Struchkov, A.I. Yanovsky), (405) 111
- PdO catalysed reaction between Re(CO)<sub>5</sub>X (X = Cl, Br, I) and L (L = group 15 donor ligand); synthesis of Re(CO)<sub>4</sub>LX (A.E. Leins, N.J. Coville), (407) 359
- LTc(CO)<sub>3</sub> compounds (L = C<sub>5</sub>Me<sub>5</sub>, C<sub>5</sub>Me<sub>4</sub>Et, C<sub>9</sub>H<sub>7</sub>), preparation and characterisation of; crystal structures of ( $\eta^5$ -C<sub>5</sub>Me<sub>5</sub>)M(CO)<sub>3</sub> (M = Tc, Re), ( $\eta^5$ -C<sub>5</sub>Me<sub>4</sub>Et)Tc(CO)<sub>3</sub> and ( $\eta^5$ -C<sub>9</sub>H<sub>7</sub>)Tc(CO)<sub>3</sub> (K. Raptis, E. Dornberger, B. Kanellakopulos, B. Nuber, M.L. Ziegler), (408) 61
- M<sup>+</sup>[M'M"(CO)<sub>9</sub>L]<sup>-</sup> (M<sup>+</sup> = Na<sup>+</sup>, PPN<sup>+</sup>; M' = Cr, W; M" = Mn, Re; L = CO, PR<sub>3</sub>), kinetic studies of the cleavage of (Y.K. Park, S.J. Kim, J.H. Kim, I.S. Han, C.H. Lee, H.S. Choi), (408) 193
- Bis(cyclopentadienyl)rhenium(III) chloride, crystal structure, properties and reactivity of; preparation and characterisation of Cp<sub>3</sub>Re and Cp<sub>2</sub>ReH (C. Apostolidis, B. Kanellakopulos, R. Maier, J. Rebizant, M.L. Ziegler), (409) 243

- Metal carbonyl anions, I<sup>+</sup>-abstraction versus I<sup>-</sup>-displacement in their reactions with diiodoacetylene; X-ray structure of [(OC)<sub>5</sub>MnC≡CMn(CO)<sub>5</sub>] (J.A. Davies, M. El-Ghanam, A.A. Pinkerton, D.A. Smith), (409) 367
- Iron and ruthenium cationic ( $\eta^6$ -chloroarene) complexes, reactions with some O-silyl and C-silyl compounds (R.C. Cambie, S.A. Coulson, L.G. Mackay, S.J. Janssen, P.S. Rutledge, P.D. Woodgate), (409) 385
- Rhenium cyanamide-isocyanide complexes *trans*-[ReL(CNR)(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub>]<sup>+</sup> (L = NCNH<sub>2</sub> or NCNH<sup>-</sup>), synthesis and electrochemical behaviour of (M.F.N.N. Carvalho, A.J.L. Pombeiro), (410) 347
- Tris(cyclopentadienyl)technetium(III), ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Tc( $\eta^1$ -C<sub>5</sub>H<sub>5</sub>), X-ray diffraction study of; dipole moment and charge distribution of (C<sub>5</sub>H<sub>5</sub>)<sub>3</sub>M (M = V, Tc, Re) (C. Apostolidis, B. Kanellakopoulos, R. Maier, J. Rebantz, M.L. Ziegler), (411) 171
- Binuclear transition metal complexes of Cu<sup>1</sup>, Mo<sup>0</sup>, and Re<sup>1</sup> with 2,2'-bipyrimidine (P.S. Braterman, J.-I. Song, S. Kohlmann, C. Vogler, W. Kaim), (411) 207
- Highly reduced metalcarbonyl [Ir(CO)<sub>3</sub>]<sup>3-</sup>, reaction with the ethylenepentacarbonylrhenium cation; formation of the trinuclear hydride (OC)<sub>5</sub>Re-Ir(CO)<sub>3</sub>(H)-Re(CO)<sub>5</sub> and the trismetalated carbonate { $\mu_3$ -CO<sub>3</sub>[Re(CO)<sub>5</sub>]<sub>3</sub>}<sup>+</sup> (J. Breimair, C. Robl, W. Beck), (411) 395
- Fe<sub>3</sub>(CO)<sub>11</sub><sup>2-</sup> and Fe<sub>4</sub>(CO)<sub>13</sub><sup>2-</sup>, transformation to give trinuclear clusters (W. Deck, A.K. Powell, H. Vahrenkamp), (411) 431
- Hydro-carbonyl rhenium cluster, transformation of a coordinated carbon monoxide into a oxyethyl bridging group; synthesis and crystal structure of the anion [Re<sub>3</sub>( $\mu$ -H)<sub>3</sub>(CO)<sub>9</sub>( $\mu_3$ - $\eta^2$ -C(H(Me)O))<sup>2-</sup> (T. Beringhelli, G. D'Alfonso, M. Freni, G. Ciani, M. Moret, A. Sironi), (412) C4
- [( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Re(NO)(PPh<sub>3</sub>)(O=CRR')<sup>+</sup> X<sup>-</sup>,  $\pi/\sigma$  equilibria in rhenium ketone complexes; a marked electronic effect upon the binding model (D.P. Klein, D.M. Dalton, N. Quirós Méndez, A.M. Arif, J.A. Gladysz), (412) C7
- Methyl- and ethylrhenium oxides, synthesis and structural aspects (W.A. Herrmann, C.C. Romao, P. Kiprof, J. Behm, M.R. Cook, M. Taillefer), (413) 11
- eq, eq*-Re<sub>2</sub>(CO)<sub>8</sub>(MeCN)<sub>2</sub>, reactions with phenylacetylene and  $\alpha$ -ethynylestradiol; new synthesis of acetylide complexes (S. Top, M. Gunn, G. Jaouen, J. Vaissermann, J.-C. Daran, J.R. Thornback), (414) C22
- Metal carbonyl dimers, new mechanism for substitution reactions of (B.F.G. Johnson), (415) 109
- Manganese, technetium, and rhenium; Annual Survey covering the year 1987 (P.M. Treichel), (416) 291
- Group VI metal carbonyls, ( $\eta^6$ -C<sub>6</sub>H<sub>6</sub>)Cr(CO)<sub>3</sub>, ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Mn(CO)<sub>3</sub>, ( $\eta^5$ -CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Mn(CO)<sub>3</sub> and ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Re(CO)<sub>3</sub>, photolysis in Nujol at 77 K (T.E. Bitterwolf, K.A. Lott, A.J. Rest and J. Mascetti), (419) 113
- Kohlenwasserstoffverbrückte Komplexe. XXIII. Heterobimetallische Komplexe mit Ferrocenyl- und (Ph<sub>3</sub>P)(OC)(Cp)Fe(CO)CH<sub>2</sub>-Gruppen; Darstellung und Struktur von ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Fe[ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)Re(CO)<sub>5</sub>], (PPh<sub>3</sub>)(OC)(Cp)FeC(O)CH<sub>2</sub>( $\eta^6$ -C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub> und [ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)CH<sub>3</sub>]Fe[ $\eta^5$ -C<sub>5</sub>H<sub>4</sub>C(O)CH<sub>2</sub>( $\eta^6$ -C<sub>7</sub>H<sub>7</sub>)M(CO)<sub>3</sub>] (M = Cr, Mo) (J. Breimair, M. Wieser, B. Wagner, K. Polborn und W. Beck), (421) 55
- Site selectivity studies on homonuclear metal dimer complexes: substitution reactions of Re<sub>2</sub>(CO)<sub>9</sub>L (W.L. Ingham and N.J. Coville), (423) 51
- Photochemische Reaktion von Pentacarbonyl(1-ethoxy-2-methyl-2-propenyliden)wolfram und Dekacarbonyldirhenium (C.G. Kreiter, S. Schufft und G. Heckmann), (424) 163
- Relaxation mechanisms, bonding, and molecular reorientation in CpRe(CO)<sub>3</sub> (P. Yuan, M.G. Richmond and M. Schwartz), (424) 331
- Synthesis and reactivity of rhenium(I) isocyanide complexes. Crystal and molecular structures of ReBr(CO)<sub>4</sub>{C(NHPh)(NHCHMe<sub>2</sub>)} and ReBr(CO)<sub>3</sub>(PPh<sub>3</sub>){C(NHPh)(NHCHMe<sub>2</sub>)} (L.-C. Chen, M.-Y. Chen, J.-H. Chen, Y.-S. Wen and K.-L. Lu), (425) 99

## Rhodium

- Chiral metal complexes of 1-dimethylphosphino-3-dimethylamines, homogeneous asymmetric catalysis by means of (C. Döbler, A. Kinting), (401) C23
- Acetylacetato-1,5-cyclooctadienerhodium(I), kinetics of the substitution of acetylacetone by derivatives of 1,10-phenanthroline and 2,2'-dipyridyl (J.G. Leipoldt, G.J. Lamprecht, E.C. Steynberg), (402) 259

- Diene- and carbonyl-rhodium(I) complexes containing bridging adeninate ligands, synthesis and structural characterisation of (W.S. Sheldrick, B. Günther), (402) 265
- Pyrazolate-bridged dirhodium(I) complex  $\text{Rh}_2(\text{CO})_2(\text{PPh}_3)_2(\mu\text{-}3\text{-Me}_2\text{pz})_2$ , electrochemistry of (R. Seeber, G. Minghetti, M.I. Pilo, G. Banditelli, S. Zamponi), (402) 413
- Heteronuclear and heterovalent diphenylphosphido-bridged complexes of rhodium, iridium and ruthenium; crystal structure of  $[(\text{C}_5\text{Me}_5)_2\text{Cl}\text{Rh}(\mu\text{-Cl})(\mu\text{-PPH}_2)\text{Rh}(\text{COD})]\cdot\frac{1}{2}\text{Et}_2\text{O}$  ( $\text{COD} = 1,5\text{-cyclooctadiene}$ ) (M. Esteban, A. Pequerul, D. Carmona, F.J. Lahoz, A. Martin, L.A. Oro), (402) 421
- Cobalt, ruthenium, and rhodium complexes as catalysts in the reaction of terminal olefins with a hydrosilane and with carbon monoxide (Y. Seki, K. Kawamoto, N. Chatani, A. Hidaka, N. Sonoda, K. Ohe, Y. Kawasaki, S. Murai), (403) 73
- Rhodium(I) and iridium(I) complexes of  $\text{Ph}_2\text{P}(\text{S})\text{CH}_2\text{P}(\text{S})\text{Ph}_2$ , bis(diphenylphosphino)methane disulfide (M.S. Abbassioun, P.A. Chaloner, C. Claver, P.B. Hitchcock, A.M. Masdeu, A. Ruiz, T. Saballs), (403) 229
- C*-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325
- Cationic dinuclear rhodium(I) complexes as catalysts in the hydroformylation of alkenes; effect of counter ions (J.C. Bayón, P. Esteban, J. Real, C. Claver, A. Polo, A. Ruiz, S. Castillón), (403) 393
- Chirally modified ruthenium clusters in the catalytic hydrogenation of racemic mixtures of terpenes; kinetic resolution of the enantiomers in (T. Jenke, G. Süss-Fink), (405) 383
- Bis( $\mu$ -methylenepentamethylcyclopentadienyl)dirhodium di-, tri- and tetra-nuclear cationic complexes, synthesis and reactions of; crystal structure of tetrานuclear  $[(\text{C}_5\text{Me}_5)_2\text{Rh}-\mu\text{-CH}_2)_2]_2(\mu\text{-CN})_2](\text{PF}_6)_2$  (J. Martinez, H. Adams, N.A. Bailey, P.M. Maitlis), (405) 393
- Coordinated water complexes of rhodium and iridium, formation and characterisation of; crystal structure of *trans*- $[\text{Rh}(\text{PPh}_3)_2(\text{CO})(\text{H}_2\text{O})]^+\text{CF}_3\text{SO}_3^-$  (P.J. Stang, L. Song, Y.-H. Huang, A.M. Arif), (405) 403
- Vinylidene complexes, nucleophilic behavior of (F. Delbecq), (406) 171
- Ketenimine-rhodium complexes, preparation by [2+1]-cycloaddition of vinylidene-rhodium precursors; crystal and molecular structure of  $\text{C}_5\text{H}_5\text{Rh}[\eta^2\text{-N,C-TosN=C=CHPh}](\text{P}^{\text{i}}\text{Pr}_3)$  (H. Werner, U. Brekau, M. Dziallas), (406) 237
- Dirhodium(II) compound with bridging orthometallated dimethylphenylphosphine ligands, synthesis and characterisation of (E.C. Morrison, D.A. Tocher), (408) 105
- Tetrahydrofurlylcyclopentadiene and its thallium, manganese, and rhodium complexes, synthesis of; low temperature matrix photolysis of the manganese complex (T.E. Bitterwolf, K.A. Lott, A.J. Rest), (408) 137
- $[(\eta^5\text{-C}_5\text{Me}_5)\text{RhCl}(\mu\text{-Cl})_2]$ , reactivity towards some potentially bidentate ligands (G. García, G. Sánchez, I. Romero, I. Solano, M.D. Santana, G. López), (408) 241
- Hydrosilylation of the C=C double bond employing homogeneous catalysis with transition metals (R. Skoda-Földes, L. Kollár, B. Heil), (408) 297
- Rhodium trichloride, cyclometallation with 1-phenylpyrazole; X-ray crystal structure of the resulting product (P.J. Steel), (408) 395
- Iron, cobalt and rhodium complexes of the optically active diolefin (+)-nopadiene and its derivatives; crystal structure of  $\text{C}_5\text{Me}_5\text{Rh}(\text{nopadiene})$  (A. Salzer, H. Schmalke, R. Stauber, S. Streiff), (408) 403
- $[(\text{OC})_3\text{M}(\mu\text{-Cl})(\mu\text{-CO})(\mu\text{-dppm})\text{Rh}(\text{NBD})]$  ( $\text{M} = \text{Mo, W}$ ) tribridged derivatives, formation by ring opening of  $[(\text{OC})_4\text{M}(\text{dppm-PP})]$ ; crystal structure of  $[(\text{OC})_3\text{Mo}(\mu\text{-Cl})(\mu\text{-CO})(\mu\text{-dppm})\text{Rh}(\text{NBD})]$  (M. Cano, J.V. Heras, P. Ovejero, E. Pinilla, A. Monge), (410) 101
- Tris(diphenylthiophosphinoyl)methane and tris(diphenylthiophosphinoyl)methanide complexes of rhodium and iridium, synthesis and structural characterisation of; crystal structures of  $[\text{Rh}(\text{C}_8\text{H}_{12})(\eta^2\text{-C}(\text{P}(\text{S})\text{Ph}_2)_3\text{S},\text{S})]$  and  $[\text{Ir}(\text{CO})_2(\eta^2\text{-C}(\text{P}(\text{S})\text{Ph}_2)_3\text{S},\text{S})]$  (J. Browning, K.R. Dixon, R.W. Hiltz, N.J. Meanwell, F. Wang), (410) 389
- Dirhodium(II) aqua-cation, reaction with carbon monoxide; new pathway for  $\text{Rh}_4(\text{CO})_{12}$  synthesis (M. Moszner, J.J. Ziolkowski), (411) 281
- Triazenido-bridged  $[\text{Rh}_2]^{4+}$  unit, superoxide and semi-quinone derivatives of (N.G. Connelly, A.C. Loynes), (411) 285
- Polycyclic silacyclobutanes, Rh-catalyzed synthesis from norbornadiene and diphenylsilane (H. Brunner, F. Prester), (411) C1

- (+)-*cis*-[Dicarbonyl- $\mu$ -chloro- $\mu$ -[5 $\beta$ -methyl-2 $\alpha$ -(1-methylethyl)cyclohexanethiolato]]bis[tris(1,1-dimethylethyl)arsine]dirhodium, synthesis, structure determination and catalytic activity of (H. Schumann, B. Gorella, M. Eisen, J. Blum), (412) 251
- Rhodium(I) complexes with the polydentate ligand 3,5-bis(4-methylpyrazol-1-yl)-4-methylpyrazole (R.M. Claramunt, C. Lopez, D. Sanz, J. Elguero, D. Carmona, M. Esteban, L.A. Oro, M. Begtrup), (412) 259
- Chiral cyclopentadienyl hydrogenation catalysts; crystal structures of [Rh(C<sub>5</sub>R<sub>4</sub>R<sup>\*</sup>)Cl<sub>2</sub>]<sub>2</sub> (R = H, R<sup>\*</sup> = neomenthyl; R = Me, R<sup>\*</sup> = menthyl) (P.A. Schofield, H. Adams, N.A. Bailey, E. Cesarotti, C. White), (412) 273
- Rhodium carbonyl catalyst modified with phosphines, cross-hydrocarbylation of 1-alkyne and ethylene (P. Hong, T. Mise, H. Yamazaki), (412) 291
- $\mu$ -Propylidene(2)-bis[carbonyl( $\eta^5$ -pentamethylcyclopentadienyl)rhodium(Rh–Rh), ( $\mu$ -C(CH<sub>3</sub>)<sub>2</sub>)<sub>2</sub>{ $\eta^5$ -C<sub>5</sub>(CH<sub>3</sub>)<sub>5</sub>}Rh(CO)<sub>2</sub>], neutron diffraction study at 20 K (R.C. Stevens, J.S. Ricci, T.F. Koetzle, W.A. Herrmann), (412) 425
- P*-substituted olefins, base-catalysed addition of optically active P–H complexes (H. Brunner, S. Limmer), (413) 55
- Tetranuclear rhodium-triruthenium carbonyl clusters containing phosphorus(III) ligands; crystal and molecular structure of [RhRu<sub>3</sub>( $\mu$ -H)<sub>2</sub>( $\mu$ -CO)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub>( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)] (J.-L. Le Grand, W.E. Lindsell, K.J. McCullough), (413) 321
- Unsaturated amines, rhodium-catalysed reactions with hydrogen and carbon monoxide; new synthesis of 2-piperidinones (D. Anastasiou, W.R. Jackson), (413) 399
- Phosphaalkene complexes, catalytic hydrogenation of the phosphorus–carbon double bond (R. de Vaumas, A. Marinetti, F. Mathey), (413) 411
- Three-membered metallacycle Cp<sub>3</sub>Rh<sub>3</sub>( $\mu$ -CO)<sub>3</sub>, cluster and benzenoid system in reactions with 12-electron particles PPh<sub>3</sub>M<sup>+</sup> (M = Cu, Ag, Au) and CpM<sup>+</sup> (M = Fe, Ru) (M.I. Rybinskaya, A.R. Kudinov, D.V. Muratov, N.P. Gambaryan, I.V. Stankevich, A.L. Chistyakov), (413) 419
- Triangular cluster Cp<sub>2</sub>Cr<sub>2</sub>( $\mu$ -SCMe<sub>3</sub>)<sub>2</sub>( $\mu$ <sub>3</sub>-S)Ir(CO)(PPh<sub>3</sub>) and octanuclear cluster Cp<sub>2</sub>Cr<sub>2</sub>( $\mu$ <sub>3</sub>-S)<sub>2</sub>( $\mu$ <sub>4</sub>-S)Rh<sub>2</sub>[Cp<sub>2</sub>Cr<sub>2</sub>( $\mu$ -SCMe<sub>3</sub>)<sub>2</sub>( $\mu$ <sub>3</sub>-S)]<sub>2</sub> with unusual triangular heterometalloornament framework; synthesis, molecular structure and magnetic properties of (A.A. Pasynskii, I.L. Eremenko, V.R. Zalmanovitch, V.V. Kavarin, B. Orazsakhatov, S.E. Nefedov, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky, Yu.T. Struchkov), (414) 55
- Rhodium(I) and iridium(I) complexes with 1,2-bis(dicyclohexylphosphino)ethane ligands; their reactions with carbon monoxide (A. Del Zotto, L. Costella, A. Mezzetti, P. Rigo), (414) 109
- Iron(0)-phthalocyanine and bis(dimethylglyoximate)-triphenylphosphinerhodate(I), conversion of 1-bromoalkynes (D. Steinborn, U. Sedlak, M. Dargatz), (415) 407
- trans*-Chloroethanebis(piperidine)rhodium(I), *trans*-[RhCl(C<sub>2</sub>H<sub>4</sub>)(C<sub>5</sub>H<sub>10</sub>NH)<sub>2</sub>], selective catalyst for the hydroamination of ethene; molecular and crystal structure of (D. Selent, D. Scharfenberg-Pfeiffer, G. Reck, R. Taube), (415) 417
- Rhodium pentafluorophenylthiolato complexes, steric effects in oxidative addition and reductive elimination reactions of (L. Carlton), (415) C19
- $\alpha$ -Amidocarboxylic acids, amidocarbonylation as a route to (J.J. Lin, J.F. Knifton), (417) 99
- The use of <sup>2</sup>H NMR in the elucidation of the catalytic pathway of the hydroformylation reaction (G. Uccello-Barretta, R. Lazzaroni, R. Settambolo, P. Salvadori), (417) 111
- Dihydrogen activation by Co<sub>2</sub>Rh<sub>2</sub>(CO)<sub>12</sub> disproportionation and a second order cluster fragmentation, kinetics of (M. Garland, P. Pino), (417) 121
- <sup>n</sup>Bu<sub>4</sub>N[Rh(CO)<sub>2</sub>(OAc)<sub>2</sub>]<sub>n</sub>, synthesis, properties, and crystal structure of; <sup>n</sup>Bu<sub>4</sub>N[Rh(CO)<sub>2</sub>(X)<sub>2</sub>]<sub>n</sub>, exchange of acetate, chloride, and iodide in (A. Fulford, N.A. Bailey, H. Adams, P.M. Maitlis), (417) 139
- Ylide rhodium(I) and rhodium(III) complexes, synthesis, structure, and reactivity of (H. Werner, O. Schippe, J. Wolf, M. Schulz), (417) 149
- N-alkenylamides and  $\alpha$ -alkenyllactams, amide-directed hydrocarbylation of (I. Ojima, Z. Zhang), (417) 253
- Dimethylsulfide derivatives of [Rh<sub>6</sub>(CO)<sub>16</sub>], crystal structures of [Rh<sub>6</sub>(CO)<sub>15</sub>(SMe<sub>2</sub>)<sub>1</sub>] and [Rh<sub>6</sub>(CO)<sub>12</sub>(SMe<sub>2</sub>)<sub>4</sub>] (S. Rossi, K. Kallinen, J. Pursiainen, T.T. Pakkanen and T.A. Pakkanen), (419) 219
- Isomeric bidiazine complexes [(C<sub>5</sub>Me<sub>5</sub>)ClRh(bdz)]<sup>+</sup> and (C<sub>5</sub>Me<sub>5</sub>)Rh(bdz), spectroscopic and electrochemical properties of; their relevance to the catalysis of the 2 H<sup>+</sup> → H<sub>2</sub> reaction by 2,2'-bipyridine analogues (M. Ladwig and W. Kaim), (419) 233

- RhCl(CO)(PR<sub>3</sub>)<sub>2</sub>, substitution of CO by picolines and amines in; synthesis and crystal structure of *cis*-RhCl(3-pic){P(OPh)<sub>3</sub>}<sub>2</sub> (A.M. Trzeciak, J.J. Ziolkowski and T. Lis), (419) 391
- Rh<sup>1</sup> square planar derivative, fast olefin site interchange of cycloocta-1,5-diene in (E. Rotondo, G. Battaglia, C.G. Arena and F. Faraonc), (419) 399
- New family of water-soluble tertiary phosphine ligands and their rhodium(I) complexes, synthesis of; olefin hydrogenation in aqueous and biphasic media (E. Renaud, R.B. Russell, S. Fortier, S.J. Brown and M.C. Baird), (419) 403
- The role of alcohol in the catalytic reductive carbonylation of nitrobenzenes to carbamates in the presence of Rh(CO)<sub>4</sub><sup>-</sup> or Ru<sub>3</sub>(CO)<sub>12</sub> (C.-H. Liu and C.-H. Cheng), (420) 119
- "Head-to-head" dimerization and dehydromerization of vinyl ketones catalyzed by modified rhodium(I) complexes (I.P. Kovalev, Y.N. Kolmogorov, Y.A. Strelenko, A.V. Ignatenko, M.G. Vinogradov and G.I. Nikishin), (420) 125
- Crystal structure and reactivity of [(pp<sub>3</sub>)Rh(H)(SeCH<sub>3</sub>)]BPh<sub>4</sub>. An organometallic complex from metal-assisted Se-H activation (M. Di Vaira, D. Rovai, P. Stopponi and M. Peruzzini), (420) 135
- A mechanistic study of rhodium tri(*o*-t-butylphenyl)phosphite complexes as hydroformylation catalysts (T. Jongsma, G. Challa and P.W.N.M. van Leeuwen), (421) 121
- Direct binding of  $\eta^5$ -pentamethylcyclopentadienyl- rhodium(III) to nicotinamide cofactors: a step towards NAD<sup>+</sup>/NADH recycling (A.D. Ryabov, D.L. Menglet and M.D. Levi), (421) C16
- The coordination of functionalised diphosphines to rhodium. Reactions with [(RhCl(CO))<sub>2</sub>] to give binuclear complexes. Crystal structure of *cis*-[(RhCl(CO))<sub>2</sub>( $\mu$ -(C<sub>7</sub>H<sub>4</sub>NS)<sub>2</sub>PCH<sub>2</sub>P(C<sub>7</sub>H<sub>4</sub>S)<sub>2</sub>)<sub>2</sub>] $\cdot$ 2C<sub>4</sub>H<sub>8</sub>O (M.F.M. Al-Dulayymi, D.L. Hughes and R.L. Richards), (424) 79

## Rubidium

- Allyl alkali metals, comparative theoretical study of (N.J.R. van Eikema Hommes, M. Bühl, P.v.R. Schleyer, Y.-D. Wu), (409) 307
- Alkali metal derivatives of sulfinimidamides, preparation and crystal structures (F. Pauer, D. Stalke), (418) 127

## Ruthenium

- Permethylated  $\alpha$ -metallocenylcarbinols (M = Os, Ru, Fe), hydrogen bonds in and conformations of (E.S. Shubina, L.M. Epstein, T.V. Timofeeva, Yu. T. Struchkov, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 133
- Permethylated  $\alpha$ -metallocenylcarbinols in the solid state (M = Fe, Ru, Os), IR spectroscopic study of hydrogen bonds in (E.S. Shubina, L.M. Epstein, A.Z. Kreindlin, S.S. Fadeeva, M.I. Rybinskaya), (401) 145
- ( $\eta^6$ -Phenyl)Cr(CO)<sub>3</sub> and ( $\eta^6$ -phenyl)Co<sub>4</sub>(CO)<sub>9</sub>-substituted metallocenylcarbinols of the iron subgroup; synthesis, structure and hydrogen bonding in (E.S. Shubina, L.M. Epstein, Yu.L. Slovokhotov, A.V. Mironov, Yu.T. Struchkov, V.S. Kaganovich, A.Z. Kreindlin, M.I. Rybinskaya), (401) 155
- Cyclopentadienyl-ruthenium t-butyl thiol complexes, synthesis of (P.M. Treichel, R.A. Crane, K.N. Haller), (401) 173
- Ruthenium complexes for use in Langmuir-Blodgett films, preparation of (S.G. Davies, A.J. Smallridge, R. Colbrook, T. Richardson, G.G. Roberts), (401) 181
- Trinuclear ruthenium clusters, coordination of carbamates, hydrazides, guanidines and isoureas to; molecular structure of ( $\mu_2$ -H)Ru<sub>3</sub>(CO)<sub>9</sub>[ $\mu_3$ - $\eta^2$ -(1*R*,2*S*,5*R*)NETCOOC<sub>16</sub>H<sub>23</sub>] (T. Jenke, H. Stoekli-Evans, U. Bodensieck, G. Süss-Fink), (401) 347
- Ruthenium, penta- and heptanuclear  $\pi$ -toluene derivatives of (K.J. Edwards, J.S. Field, R.J. Haines, F. Mulla), (402) 113
- Mononuclear ruthenium(II) complex containing two 3e-donating  $\sigma$ -C,  $\sigma$ -N coordinated 1-aza-1,3-dien-4-yl ligands, preparation and spectroscopic characterisation of (W.P. Mul, C.J. Elsevier, J. Spaans), (402) 125
- Ru( $\eta^4$ -C<sub>8</sub>H<sub>12</sub>) $\times$ ( $\eta^6$ -C<sub>8</sub>H<sub>10</sub>), thermal decomposition of (D. Labroue, R. Pince, R. Quéau), (402) 363
- Heteronuclear and heterovalent diphenylphosphido-bridged complexes of rhodium, iridium and ruthenium; crystal structure of [(C<sub>5</sub>Me<sub>5</sub>)ClRh( $\mu$ -Cl)( $\mu$ -PPh<sub>2</sub>)Rh(COD)] $\cdot$  $\frac{1}{2}$ Et<sub>2</sub>O (COD = 1,5-cyclooctadiene) (M. Esteban, A. Pequerul, D. Carmona, F.J. Lahoz, A. Martin, L.A. Oro), (402) 421
- Ruthenium  $\alpha$ -phenylethynyl-*trans*- $\beta$ -styryl complexes, reactions with isonitriles; hemi-labile alkyne coordination (A.F. Hill, R.P. Melling, A.R. Thompsett), (402) C8

- Cobalt, ruthenium, and rhodium complexes as catalysts in the reaction of terminal olefins with a hydrosilane and with carbon monoxide (Y. Seki, K. Kawamoto, N. Chatani, A. Hidaka, N. Sonoda, K. Ohe, Y. Kawasaki, S. Murai), (403) 73
- Substituted ferrocenes and ruthenocenes, dynamic NMR studies of ring rotation in (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šík), (403) 195
- Ruthenium-catalysed oxidation of primary alcohols with sodium bromate (A. Behr, K. Eusterwiemann), (403) 209
- Ruthenium-catalysed oxidation of primary alcohols with the oxidation system *N*-methylmorpholine/hydrogen peroxide (A. Behr, K. Eusterwiemann), (403) 215
- 4,4'-Bipyridyl-bridged diruthenium(II) carbonyl catecholate complex, ligand-localised oxidation of (C. Carriero, N.G. Connelly), (403) 359
- Substituted ruthenium(II)  $\eta^5$ -cyclopentadienyl  $\eta^5$ - or  $\eta^6$ -arene salts, synthesis and nonlinear optical properties of (M. Kimura, H. Abdel-Halim, D.W. Robinson, D.O. Cowan), (403) 365
- Six-coordinate ruthenium(II) complexes containing both formate and alkenyl ligands; crystal structure of Ru(O<sub>2</sub>CH)(CO)(HC=CHPh)(PPh<sub>3</sub>)<sub>2</sub> (H. Loumrhari, L. Matas, J. Ros, M.R. Torres, A. Perales), (403) 373
- Ruthenium-containing precatalysts in the cyclotrimerisation of alkynes (E. Lindner, H. Kühbauch), (403) C9
- Annual Survey of ruthenium and osmium for the year 1988 (M.G. Richmond), (404) 213
- [CpM(CO)<sub>2</sub>R] long-chain alkyl complexes (Cp =  $\eta^5$ -C<sub>5</sub>H<sub>5</sub>; M=Fe or Ru; R=n-C<sub>6</sub>H<sub>13</sub> to n-C<sub>12</sub>H<sub>25</sub>), synthesis, characterisation and properties of (A. Emeran, M.A. Gafoor, J.K.I. Goslett, Y.-H. Liao, L. Pimble, J.R. Moss), (405) 237
- [(-)-Norphos]Ru<sup>II</sup>( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)I diastereomers, synthesis and structure of (H. Nishiyama, H. Brunner, P.G. Jones), (405) 247
- HRu<sub>3</sub>[ $\mu_3$ , $\eta^2$ -P(NEt<sub>2</sub>)<sub>2</sub>](CO)<sub>9</sub>, a compound containing a five-membered cage-like Ru<sub>3</sub>PN skeleton; crystal structure of (L. Qiwang, H. Xiang, W. Binfang, L. Shutang), (405) 257
- Metallocene-containing cryptands, NMR studies on metal complexes of (C.D. Hall, N.W. Sharpe), (405) 365
- Platinum(II) chloride complexes of 1,1'-bis(methylthio)ferrocene, 1,1'-bis(methylseleno)ferrocene and 1,1'-bis(methylthio)ruthenocene, synthesis and stereodynamics of (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne, V. Šík), (405) 375
- Chirally modified ruthenium clusters in the catalytic hydrogenation of racemic mixtures of terpenes; kinetic resolution of the enantiomers in (T. Jenke, G. Süss-Fink), (405) 383
- Polyolefinic ruthenium(0) complexes, synthesis from ruthenium(II) precursors (J.G. Toerien), (405) C43
- Bis(dimethylgermyl)alkaneruthenium tetracarbonyls, reactions with unsaturated compounds (J. Barrau, N. Ben Hamida, J. Satgé), (406) 123
- Ru<sup>II</sup>-alkyl-butadiene complexes, formation from dihalogeno-Ru<sup>IV</sup>-allyl precursors by the reaction with BrMg(CH<sub>2</sub>)<sub>4</sub>MgBr; decomposition pathway for ruthena(IV)cyclopentane intermediates (H. Nagashima, Y. Michino, K. Ara, T. Fukahori, K. Itoh), (406) 189
- Transition metal hydride complexes as Lewis bases; preparation and crystal structure of (RR,SS)-{[( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Ru(PPh<sub>3</sub>)(CN<sup>1</sup>Bu)]<sub>2</sub>( $\mu$ -H)}PF<sub>6</sub>·CH<sub>2</sub>Cl<sub>2</sub> (F.M. Conroy-Lewis, S.J. Simpson, L. Brammer, A.G. Orpen), (406) 197
- Triosmium and triruthenium clusters, reactions with the isomers methoxyallene and methyl propargyl ether; crystal structure of the methoxyallene cluster [Os<sub>3</sub>(CO)<sub>11</sub>( $\mu$ , $\eta^1$ , $\eta^3$ -CH<sub>2</sub>CCHOMe)] (A.J. Arce, Y. De Sanctis, A.J. Deeming, K.I. Hardcastle, R. Lee), (406) 209
- Ruthenium catalysed reactions of ethylene glycol with primary amines; steric factors and selectivity control (J.A. Marsella), (407) 97
- 1,1'-(1,4,10,13-Tetraoxa-7,16-diazacyclooctadecane-7,16-diyl)dicarbonyl)ruthenocene, crystal structure of (C.D. Hall, A.W. Parkins, S.C. Nyburg, N.W. Sharpe), (407) 107
- cis-Arylhydridoruthenium(II) chelate complexes as catalysts for the dimerisation of 1-alkynes to linear 1-en-3-yne; molecular structure of (pp<sub>3</sub>)RuH(C<sub>6</sub>H<sub>4</sub>CF<sub>3</sub>)<sub>3</sub> (pp<sub>3</sub> = P(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>)<sub>3</sub>) (L. Dahlenburg, K.-M. Frosin, S. Kerstan, D. Werner), (407) 115
- $\mu_3$ -Phosphinidene-bridged iron and ruthenium clusters, selective hydrogenation of diphenylacetylene and isomerisation of cis-stilbene in the presence of (M. Castiglioni, R. Giordano, E. Sappa), (407) 377
- [Ru<sub>10</sub>CCO)<sub>24</sub>]ppn<sub>2</sub>, reaction with carbon monoxide to yield [Ru<sub>3</sub>(CO)<sub>12</sub>] and [Ru<sub>6</sub>C(CO)<sub>16</sub>]<sup>2-</sup> (T. Coston, J. Lewis, D. Wilkinson, B.F.G. Johnson), (407) C13

- Ruthenium(II)-alkenyl bonds, formation of alkenedithiocarboxylate and alkenecarboxylate ligands by insertion of CS<sub>2</sub> and CO<sub>2</sub> into (H. Loumrhari, J. Ros, R. Yáñez, M.R. Torres), (408) 233
- 1-*para*-Benzozquinonyl-2,3,4,5-tetraphenylcyclopentadienyl ruthenium(II) carbonyl complexes, preparation of (S.B. Colbran, D.C. Craig, W.M. Harrison, A.E. Grimley), (408) C33
- (Cyclopentadienyl)ruthenium(1+) complexes of aromatic diterpenoids (R.C. Cambie, L.G. Mackay, P.S. Rutledge, M. Tercel, P.D. Woodgate), (409) 263
- Rigid rod, iron-metal containing polyne polymers, synthesis of (B.F.G. Johnson, A.K. Kakkar, M.S. Khan, J. Lewis), (409) C12
- [{(C<sub>5</sub>Me<sub>4</sub>R)Ru}<sub>3</sub>Ru(η<sup>3</sup>-As<sub>3</sub>)(μ<sub>3</sub>,η<sup>3</sup>-As<sub>3</sub>)(μ<sub>3</sub>-As)<sub>3</sub>] tetranuclear clusters, synthesis and crystal structure of (O.J. Scherer, C. Blath, G. Heckmann, G. Wolmershäuser), (409) C15
- Binuclear (η<sup>6</sup>-benzene)ruthenium(II) complexes with one or two bridging N-donor ligands, synthesis and structural characterisation of (W.S. Sheldrick, H.-S. Hagen-Eckhard), (410) 73
- Fe<sub>2</sub>Ru(μ<sub>3</sub>-Se)(μ<sub>3</sub>-Te)(CO)<sub>9</sub> and (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)CoFe<sub>2</sub>(μ<sub>3</sub>-Se)(μ<sub>3</sub>-Te)(CO)<sub>6</sub>, synthesis and characterisation of (D. Chakrabarty, Md.M. Hossain, R.K. Kumar, P. Mathur), (410) 143
- Ru<sub>5</sub>(μ<sub>5</sub>-C<sub>2</sub>PPh<sub>2</sub>)(μ-PPh<sub>2</sub>)(CO)<sub>13</sub>, substitution of CO by P(OEt)<sub>3</sub> in; X-ray structures of two isomers of Ru<sub>5</sub>(μ<sub>5</sub>-C<sub>2</sub>PPh<sub>2</sub>)(μ-PPh<sub>2</sub>)(CO)<sub>12</sub>{P(OEt)<sub>3</sub>} and of Ru<sub>5</sub>(μ<sub>5</sub>-C<sub>2</sub>PPh<sub>2</sub>)(μ-PPh<sub>2</sub>)(CO)<sub>11</sub>{P(OEt)<sub>3</sub>} (M.I. Bruce, M.J. Liddell, E.R.T. Tiekkink, B.K. Nicholson, B.W. Skelton, A.H. White), (410) 211
- Triosmium and triruthenium carbon radicals, reactivity toward fluoro- and hexafluorobenzene (N. Hadj-Bagheri, D.S. Strickland, S.R. Wilson, J.R. Shapley), (410) 231
- Ruthenaborane cluster HRu<sub>4</sub>(CO)<sub>12</sub>BH<sub>2</sub>, insertion of 1-phenyl-1-propyne into; isolation and molecular structure of 1-phenyl-2,3-dimethylazulene (S.M. Draper, C.E. Housecroft, A.K. Keep, D.M. Matthews, B.S. Haggerty, A.L. Rheingold), (410) C44
- Alkenyl-ruthenium(II) Ru(CO)Cl(RC = CH'R')(PPh<sub>3</sub>)<sub>2</sub> complexes, reaction with CO; formation of dicarbonyl complexes or η<sup>2</sup>-acyl complexes depending on the R and R' groups (H. Loumrhari, J. Ros, M.R. Torres, A. Santos, A.M. Echavarren), (411) 255
- (Dialkylsulfide)(arene)ruthenium(II) derivatives (M. Gaye, B. Demerseman, P.H. Dixneuf), (411) 263
- Bis(diazadiene)ruthenium, isomerisation, hydrogenation and metalation; structure of a potassium(tmeda)<sub>2</sub>ruthenate(0) derivative (V. Rosenberger, G. Fendesak, H. tom Dieck), (411) 445
- L<sub>4</sub>Ru<sup>0</sup> complex cores, inter- and intramolecular cleavage of sp<sup>2</sup>- and sp<sup>3</sup>-C-H bonds with P(o-C<sub>6</sub>H<sub>4</sub>PM<sub>2</sub>e)<sub>3</sub>, N(CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>e)<sub>3</sub>, MeP(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>e)<sub>2</sub>/PR<sub>3</sub> and MeSi(CH<sub>2</sub>PM<sub>2</sub>e)<sub>3</sub>/PR<sub>3</sub> (R = Me, OMe) as supporting ligands (L. Dahlenburg, S. Kerstan, D. Werner), (411) 457
- [(C<sub>5</sub>H<sub>5</sub>)Ru(dppm)(CH<sub>2</sub>=SO<sub>2</sub>)]<sup>+</sup>, a strong organometallic electrophile (W.A. Schenk, P. Urban), (411) C27
- Bis(tetraphenylcyclopentadienyl)ruthenium(II), synthesis, molecular structure, and <sup>1</sup>H NMR analysis of (R.J. Hoobler, J.V. Adams, M.A. Hutton, T.W. Francisco, B.S. Haggerty, A.L. Rheingold, M.P. Castellani), (412) 157
- Hydridoruthenium raft clusters H<sub>2</sub>Ru<sub>6</sub>(CO)<sub>15</sub>(L)(C<sub>6</sub>H<sub>4</sub>O) (L = CO, P(OMe)<sub>3</sub>), solution and crystal structures of (S. Bhaduri, K. Sharma, H. Khwaja, P.G. Jones), (412) 169
- Ditertiary phosphine and disphosphazane ligand-bridged derivatives of di-iron and diruthenium nonacarbonyl, electrochemical behaviour of (J.S. Field, A.M.A. Francis, R.J. Haines, S.F. Woollam), (412) 383
- Pyrazine bridged bis-allyl ruthenium(IV) complex, synthesis and characterisation of; crystal structure of [{(η<sup>3</sup>:η<sup>3</sup>-C<sub>10</sub>H<sub>16</sub>)RuCl<sub>2</sub>}\_2(μ-C<sub>4</sub>H<sub>4</sub>N<sub>2</sub>)·2CHCl<sub>3</sub>] (J.W. Steed, D.A. Tocher), (412) C34
- [(η<sup>6</sup>-C<sub>6</sub>Me<sub>6</sub>)Ru(η<sup>6</sup>-[2.2]paracyclophane)][BF<sub>4</sub>]<sub>2</sub>, reactions with nucleophiles (J.W. Steed, D.A. Tocher), (412) C37
- Ru(η<sup>6</sup>-naphthalene)(η<sup>4</sup>-1,5-cyclooctadiene)/acetonitrile system, homogeneous catalytic precursor for fast isomerization of 1,5-cyclooctadiene and 1-hexene (P. Pertici, G. Uccello Barretta, F. Burzaglio, P. Salvadori, M.A. Bennett), (413) 303
- [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)L<sub>2</sub>Ru=C=C(Me)Ph]<sup>+</sup> (L<sub>2</sub> = (PM<sub>2</sub>e)<sub>2</sub> or (PPh<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>), stereoselective nucleophilic addition of cyanide to (S.G. Davies, A.J. Smallridge), (413) 313
- Tetranuclear rhodium-triruthenium carbonyl clusters containing phosphorus(III) ligands; crystal and molecular structure of [RhRu<sub>3</sub>(μ-H)<sub>2</sub>(μ-CO)(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)] (J.-L. Le Grand, W.E. Lindsell, K.J. McCullough), (413) 321
- Three-membered metallacycle Cp<sub>3</sub>Rh<sub>3</sub>(μ-CO)<sub>3</sub>, cluster and benzenoid system in reactions with 12-electron particles PPh<sub>3</sub>M<sup>+</sup> (M = Cu, Ag, Au) and CpM<sup>+</sup> (M = Fe, Ru) (M.I. Rybinskaya, A.R. Kudinov, D.V. Muratov, N.P. Gambaryan, I.V. Stankevich, A.L. Chistyakov), (413) 419

- $[(\eta\text{-C}_5\text{R}_5)\text{M}^+]$  fragments ( $\text{M} = \text{Fe}$ ,  $\text{R} = \text{H}$ ;  $\text{M} = \text{Ru}$ ,  $\text{R} = \text{H}$ ,  $\text{Me}$ ) in synthesis of cationic clusters; crystal structure of  $[\text{RuRh}_3(\eta\text{-C}_5\text{Me}_5)(\eta\text{-C}_5\text{H}_5)_3(\mu_3\text{-CO})_3]\text{PF}_6$  (A.R. Kudinov, D.V. Muratov, M.I. Rybinskaya, P.V. Petrovskii, A.V. Mironov, T.V. Timofeeva, Yu.L. Slovokhotov, Yu.T. Struchkov), (414) 97
- Ruthenium(II) complexes as promoters of phenylacetylene dimerisation (A.M. Echavarren, J. López, A. Santos, J. Montoya), (414) 393
- Dodecacarbonyltriruthenium, reaction with hydrazones; preparation and crystal structure of  $(\mu_2\text{-H})\text{Ru}_3(\text{CO})_{10}\mu\text{-}(\text{R})\text{-NHN}=\text{CCH}_2\text{CH}(\text{CMe}_2)\text{CH}_2\text{CH}_2\text{CMe}$  (T. Jenke, U. Bodensieck, H. Stoeckli-Evans, G. Süss-Fink), (414) C28
- Mononuclear and oligomeric ruthenium(II) acetylides, synthesis of (S.J. Davies, B.F.G. Johnson, J. Lewis, P.R. Raithby), (414) C51
- $\text{RuCl}_2(\text{PPh}_3)_2$ , reactions of cyclic acetals with alcohols in the presence of (M. Bartók, K. Feldföldi, G.B. Bartók), (415) 127
- Hydridoruthenium(II) carboxylates, insertion reactions with acetylenes (S.S. Deshpande, S. Gopinathan, C. Gopinathan), (415) 265
- Anionic ruthenium iodocarbonyl complexes as selective dehydroxylation catalysts in aqueous solution (G. Braca, A.M. Raspolli Galletti, G. Sbrana), (417) 41
- Metal complexed indoles, chemistry and synthetic utility of (U.S. Gill, R.M. Moriarty, Y.Y. Ku, I.R. Butler), (417) 313
- $\text{Ru}_3(\mu_3\text{-C}_2(\text{CO}_2\text{Me})_2)(\mu\text{-CO})(\text{CO})_2(\text{PMe}_2\text{Ph})_2$ , preparation and X-ray structure of (M.I. Bruce, P.A. Humphrey, H. Miyamae, A.H. White), (417) 431
- $\text{Ru}_3(\text{CO})_{12}/\text{Ru}(\text{CO})_5/\text{CO}$  equilibrium, thermodynamic study of (R. Koelliker, G. Bor), (417) 439
- Iron and ruthenium complexes with 1,1-difluoroethene (W. Keim, W. Fischer), (417) C20
- $[\text{RuH}(\text{PPh}_3)_2(9,10\text{-dimethylanthracene})]^+$ , homogeneous catalysis of the hydrogenation of 9,10-dimethylanthracene to 1,2-dihydro-9,10-dimethylanthracene by (Z. Lin, J. Halpern), (417) C24
- Diastereoisomeric  $(\eta^6\text{-arene})\text{ruthenium(II)}$  chiral Schiff base complexes; crystal structure of a triphenylphosphine adduct (S.K. Mandal, A.R. Chakravarty), (417) C59
- Unsaturated mixed metal clusters  $\text{Fe}_2\text{Ru}_2(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})(\mu_4\text{-Te})$  and  $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})(\mu_4\text{-Te})$ , construction of (P. Mathur, D. Chakrabarty, Md. M. Hossain), (418) 415
- Selenium and tellurium atoms from selenophene and tellurophene by reaction with trinuclear iron, ruthenium, and osmium clusters, extrusion of; crystal structures of  $[\text{Os}_6(\mu\text{-H})(\mu_3\text{-Se})(\mu_4\text{-C}_4\text{H}_3)(\text{CO})_{20}]$  and of  $[\text{Ru}_4(\mu_3\text{-Se})\mu\text{-C}_4\text{H}_4](\text{CO})_{11}$  (A.J. Arce, R. Machado, C. Rivas, Y. De Santis and A.J. Deeming), (419) 63
- $[\text{RuCo}_3\text{H}(\text{CO})_{12}]$ , disordered crystal structure of (J. Pursiainen, P. Hirva and T.A. Pakkanen), (419) 193
- Trisulphur- and triselenium-bridged [3]metallocenophanes, comparison of the energy barriers to bridge reversal in (E.W. Abel, N.J. Long, K.G. Orrell, A.G. Osborne and V. Šík), (419) 375
- Ultraviolet vapour-phase absorption spectra of  $d^6$  metallocenes (S.Y. Ketkov and G.A. Domrachev), (420) 67
- Mixed chalcogen carbonyl compounds. IV. Reactivity of  $\text{Fe}_2(\text{CO})_6(\mu\text{-STe})$  and  $\text{Fe}_3(\text{CO})_9(\mu_3\text{-S})(\mu_3\text{-Te})$  towards coordinatively unsaturated species (P. Mathur, D. Chakrabarty, M.M. Hossain and R.S. Rashid), (420) 79
- Cluster chemistry. LXXI. Isomerisation of a pentanuclear ruthenium cluster by elimination of phenylphosphinidene from a tertiary phosphine. X-Ray structure of  $\text{Ru}_5(\mu_5\text{-C}_2\text{Ph})(\mu_4\text{-PPPh}_2)(\text{CO})_{13}$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (420) 87
- Cluster chemistry. LXXXIII. Preparation and X-ray structure of the hexanuclear cobalt-ruthenium cluster,  $\text{CoRu}_5(\mu_4\text{-PPPh}_2)\mu_4\text{-C}_2\text{Ph})(\mu\text{-PPPh}_2)(\text{CO})_{12}(\eta\text{-C}_5\text{H}_5)$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (420) 95
- Cluster chemistry. LXXIV. Reactions between  $\text{Ru}_5(\mu_5\text{-C}_2\text{PPPh}_2)\text{-}(\mu\text{-PPPh}_2)(\text{CO})_{13}$  and allyl halides: cluster cleavage and three-component addition reactions. Crystal structures of  $\text{Ru}_4\{\mu_4\text{-C}_2\text{PPPh}_2\}\text{Ru}(\text{CO})_2(\eta\text{-C}_3\text{H}_5)\text{-}(\mu\text{-Cl})(\mu\text{-PPPh}_2)(\text{CO})_{10}$  and  $\text{Ru}_5(\mu_5\text{-CCC(O)CH}_2\text{CH=CH}_2)(\mu\text{-PPPh}_2)_2(\mu\text{-Br})(\text{CO})_{11}$  (C.J. Adams, M.I. Bruce, M.J. Liddell and B.K. Nicholson), (420) 105
- Organometallic cumulenes: allenylidene- and alkanyl allenylidene-ruthenium complexes (A. Wolinska, D. Touchard, P.H. Dixneuf and A. Romero), (420) 217
- Ligand addition to  $[\text{Cp}^\star\text{RuCl}_4]$ —molecular structure of  $[\text{Cp}^\star\text{Ru}(\text{C}_2\text{H}_4)\text{Cl}]_2$  (U. Koelle, B.-S. Kang and U. Englert), (420) 227
- Ketone, aldehyde and aldimine complexes of cyclopentadienylruthenium organometallic Lewis acids (J.W. Faller, Y. Ma, C.J. Smart and M.J. DiVerdi), (420) 237

- Cyclopentadienyl-ruthenium and -osmium chemistry. XXXVI. Oligomerisation of phenylacetylide residues on ruthenium. Crystal structures of  $\{\text{Ru}(\text{PPh}_3\text{X}\eta\text{-C}_5\text{H}_5)\}_2(\mu\text{-C}_8\text{Ph}_4)$  and  $\{\text{Ru}(\text{PPh}_3\text{X}\eta\text{-C}_5\text{H}_5)\}_2(\mu\text{-C}_{10}\text{Ph}_4\text{C}_6\text{H}_4)$  (M.I. Bruce, G.A. Koutsantonis, M.J. Liddell and E.R.T. Tieckink), (420) 253
- Cyclopentadienyl-ruthenium and -osmium chemistry. XXXVII. Oligomerisation of  $\text{C}_2(\text{CO}_2\text{Me})_2$  at a cyclopentadienyl-ruthenium centre: X-ray structures of  $\text{Ru}[\{\eta^4\text{-CH}(\text{CO}_2\text{Me})=\text{C}(\text{CO}_2\text{Me})\text{C}(\text{CO}_2\text{Me})=\text{CH}(\text{CO}_2\text{Me})\}(\eta\text{-C}_5\text{H}_5)]$ ,  $\text{Ru}(\eta^5\text{-C}_5\text{H}_5)_2(\eta^5\text{-C}_6\text{H}(\text{CO}_2\text{Me})_6)$  and  $\text{Ru}(\eta^5\text{-C}_5\text{H}_5)_2(\eta^5\text{-C}_6\text{H}[\text{C}(\text{CO}_2\text{Me})=\text{CH}(\text{CO}_2\text{Me})](\text{CO}_2\text{Me})_6)$  (M.I. Bruce, G.A. Koutsantonis, E.R.T. Tieckink and B.K. Nicholson), (420) 271
- Pericyclic organometallic reactions. Cycloaddition reactions of  $(\eta^4\text{-cycloheptatriene})\text{Ru}(\text{CO})_3$ . Crystal structure of tricarbonyl[(2,3,4,9- $\eta$ )-bicyclo[4.2.1]non-2-ene-4,9-diy]-7,7,8,8-tetracarbonitrile]ruthenium (Z. Goldschmidt, E. Genizi, H.E. Gottlieb, D. Hezroni-Langermann, H. Berke, H.W. Bosch and J. Takats), (420) 419
- Mercury-ruthenium mixed-metal carbonyl clusters containing 2-amido-6-methylpyridine (ampy) as a  $\mu_3\eta^2$ -ligand. Crystal structures of  $[\text{Ru}_6(\mu_4\text{-Hg})(\mu_3\text{-ampy})_2(\text{CO})_{18}]\cdot 2\text{C}_4\text{H}_8\text{O}$  and  $[\text{Ru}_3(\mu\text{-HgBr})(\mu_3\text{-ampy})(\text{CO})_9]$  (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, C. Bois and Y. Jeannin), (420) 431
- The cleavage of a cluster carbonyl C–O bond by trifluoromethanesulphonic anhydride (P.J. Bailey), (420) C21
- Synthesis of slipped triple- and tetra-decker cationic ruthenium complexes with the  $\mu,\eta^5;\eta^6$ -indenyl ligand. X-ray structure of  $[(\eta\text{-C}_5\text{H}_5)\text{Ru}(\mu,\eta^5;\eta^6\text{-C}_9\text{H}_7)\text{-Ru}(\eta\text{-C}_5\text{Me}_5)]\text{PF}_6^-$  (A.R. Kudinov, P.V. Petrovskii, Yu.T. Struchkov, A.I. Yanovskii and M.I. Rubinskaya), (421) 91
- Arenecarbonylcarbide ruthenium clusters ( $\pi$ -arene)  $\text{Ru}_6\text{C}(\text{CO})_4$  with functional substituents in arene ring (v.s. Kaganovich, Z.A. Kerzina, T. Asunta, K. Wiekström and M.I. Rybinskaya), (421) 117
- Heterometallic analogues of  $[\{\text{M}(\text{CO})_2(\eta\text{-C}_5\text{H}_5)\}_2]$  ( $\text{M} = \text{Fe}, \text{Ru}, \text{Os}$ ). The synthesis of the tetracarbonylcyclopentadienylcycloheptatrienyl complexes  $[\text{M}(\text{CO})_2(\eta\text{-C}_5\text{H}_5)\text{M}'(\text{CO})_2(\eta\text{-C}_7\text{H}_7)]$  ( $\text{M} = \text{Fe}, \text{Ru}; \text{M}' = \text{Mo}, \text{W}$ ). Crystal structures of  $[\text{Ru}(\text{CO})_2(\eta\text{-C}_5\text{H}_5)\text{Mo}(\text{CO})_2(\eta\text{-C}_7\text{H}_7)]$  (Ru–Mo) and  $[\text{Mo}(\text{CO})_3(\mu\text{-}\eta^6,\eta^1\text{-C}_7\text{H}_7)\text{Ru}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)]$  (R.L. Beddoes, E.S. Davies, M. Helliwell and M.W. Whittley), (421) 285
- The chemical and electrochemical oxidation of pyridonate-bridged ruthenium(I) dimers. X-Ray structure of  $[\text{Ru}_2(\mu\text{-pyO})_2(\text{CO})_4(\text{pyOH})_2]$  ( $\text{pyOH} = 2\text{-pyridone}$ ) (P.L. Andreu, J.A. Cabeza, G.A. Carriero, V. Riera, S. García-Granda, J.F. Van der Maelen and G. Mori), (421) 305
- Metal hydride oxidation chemistry: addition of a triptyl cation to the cyclopentadienyl ring in  $\text{CpRu}(\text{PPh}_3)_2\text{H}$  via initial electron transfer (O.B. Ryan, K.-T. Smith and M. Tilset), (421) 315
- Cluster chemistry. LXXV. Tetra-, penta- and hepta-nuclear clusters derived from  $\{\text{Ru}_4(\mu\text{-H})_4(\text{CO})_{11}\}_2(\mu\text{-dppa})$  [dppa =  $\text{C}_2(\text{PPh}_2)_2$ ]: X-ray structures of  $\text{Ru}_4(\mu\text{-H})[\mu_3\text{-HC}_2\text{PPh}_2\{\text{Ru}_4(\mu\text{-H})_4(\text{CO})_{11}\}(\mu\text{-PPh}_2)(\mu\text{-CO})(\text{CO})_9]$  and  $\text{Ru}_7(\mu\text{-H})_3(\mu_5\text{-HC}_2\text{PPh}_2)(\mu\text{-PPh}_2)(\text{CO})_{16}$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (423) 83
- Cluster chemistry. LXXVI. A new  $\text{C}_2$  complex from reactions of  $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)(\mu\text{-PPh}_2)(\text{CO})_{13}$  with pyridine. X-Ray structure of  $\text{Ru}_5(\mu_5\text{-C}_2\text{X}\mu\text{-PPh}_2)_2(\text{CO})_{11}(\text{py})_2$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (423) 97
- Cluster chemistry. LXXVII. Carbide clusters from  $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2\text{X}\mu\text{-PPh}_2)(\text{CO})_{13}$  and  $\text{Fe}_2(\text{CO})_9$ : X-ray structures of  $\text{MRu}_4(\mu_5\text{-C})(\mu\text{-dppm})(\text{CO})_{13}\cdot 0.5\text{CH}_2\text{Cl}_2$  ( $\text{M} = \text{Fe}$  and  $\text{Ru}$ ) and  $\text{FeRu}_5(\mu_6\text{-C})(\mu\text{-PPh}_2)_2(\text{CO})_{13}(\text{PMMe}_2\text{Ph})$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (423) 105
- Cluster chemistry. LXXVIII. Coupling of allene dimer and acetylidyne on an  $\text{Ru}_5$  cluster. X-Ray structure of  $\text{Ru}_5(\mu_5\text{-PPh}_2\text{C}_2\text{C}_6\text{H}_8)(\mu\text{-PPh}_2)(\text{CO})_{10}(\text{P}(\text{OMe})_3)$  (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (423) 119
- $\text{Ru}_3(\text{CO})_9\text{BH}_5$  and  $[\text{Ru}_3(\text{CO})_9\text{BH}_4]^-$  as precursors to higher nuclearity homo- and heterometallic clusters: molecular structure of a second isomer of  $\text{HRu}_6(\text{CO})_{17}\text{B}$  (S.M. Draper, C.E. Housecroft, A.K. Keep, D.M. Matthews, X. Song and A.L. Rheingold), (423) 241
- Ruthenium complex-catalyzed coupling of vinyl halides with olefins (T.-a. Mitsudo, M. Takagi, S.-W. Zhang and Y. Watanabe), (423) 405
- Reaktionen von  $[\text{Cp}^*\text{RuOMe}]_2$ . XI. Austausch mit Thiolen und Molekülstruktur von  $[\text{C}_5\text{Me}_4\text{EtRuSEt}]_2$  (U. Koelle, C. Rietmann and U. Englert), (423) C20
- Complexes with functional phosphines. XVII. Synthesis and characterization of ruthenium clusters with ketophosphine ligands. Crystal structure of the cluster  $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^1\text{(O)}\text{:}\eta^2\text{(P,O)})\text{O}(\omega\text{-CPh})\text{-CHPPh}_2](\text{CO})_9$  containing a bridging phosphinoenolate ligand (P. Braunstein, S. Coco Cea, M.I. Bruce, B.W. Skelton and A.H. White), (423) C38

- Octahedral bis(acetylidyne) and bis(diacetylidyne) complexes of ruthenium(II) with linear  $C_2MC_2$  and  $C_4MC_4$  chains: the first X-ray structures of the all *trans* bis(acetylides)  $Ru(CO)_2(C\equiv CPh)_2(PEt_3)_2$  and  $Ru(CO)_2(C\equiv CH)_2(PEt_3)_2$  (Y. Sun, N.J. Taylor and A.J. Cartt), (423) C43
- A novel bifunctional bridging phosphinomethyl-alkoxycarbene ligand from (diphenylphosphinoacetylene)ruthenium complexes (M. Gaye, B. Demerseman and P.H. Dixneuf), (424) 65
- Use of  $[Me_2SSMe]BF_4$  to prepare thiolate complexes of iron and ruthenium (P.M. Treichel and E.K. Rublein), (424) 71
- Reactions of acyl(carbonyl)ruthenium(II) and acyl(carbonyl)iron(II) complexes with amines. Preparation of novel propionyl(carbamoyl)ruthenium(II) complexes (F. Ozawa, S. Chikaoka, X.-Z. Cao and A. Yamamoto), (424) 173
- Propargyl complexes of ruthenium (C.E. Shuchart, R.R. Willis and A. Wojciecki), (424) 185
- Enantioselective reduction of benzoyl derivatives of ferrocene and ruthenocene; optically active carbinols and carbenium ions (M.N. Nefedova, I.A. Mamedyarova, P.P. Petrovski and V.I. Sokolov), (425) 125
- Synthesis and reactivity of hydrido, halogeno, and  $\sigma$ -organyl ruthenatetraboranes: Crystal structure of  $[RuH(B_3H_8)(CO)(PPh_3)_2]$  (I.D. Burns, A.F. Hill, A.R. Thompsett, N.W. Alcock and K.S. Claire), (425) C8

## **Samarium**

- Tricyclopentadienyllanthanides/sodium hydride, reaction with organic halides (C. Qian, D. Zhu, Y. Gu), (401) 23
- $Cp_2^*Sm(THF)_2$ , reactions with 1,4-diazadienes and cyclooctatetraene (A. Recknagel, M. Noltemeyer, F.T. Edelmann), (410) 53
- Monomeric organosamarium(III) chalcogenolates, synthesis by reductive cleavage of E-E bonds (E = S, Se, Te) (A. Recknagel, M. Noltemeyer, D. Stalke, U. Pieper, H.-G. Schmidt, F.T. Edelmann), (411) 347
- Samarocene and ytterbocene hydroxides, synthesis and crystal structures of  $[(SmCp_2''(\mu-OH))_2]$  and  $[(YbCp_2''(\mu-OH))_2][Cp'' = \eta^5-C_5H_3(SiMe_3)_2-1,3; Cp' = \eta^5-C_5H_4SiMe_3]$  (P.B. Hitchcock, M.F. LapPERT, S. Prashar), (413) 79
- Coordinatively oversaturated metallocene derivatives, crystal and molecular structure of  $[(\eta^5-C_5H_5)_2Sm(\mu_3-H)]_2[(\mu_2-H)_2AlH \cdot N(C_2H_5)_3]_2$  and  $[(\eta^5-C_5H_4^+Bu)_2Sm]_2 \cdot (\mu_2-H)\mu \cdot [(\mu_3-H)_2Al(\mu_2-H)_2 \cdot Me_2NC_2H_4NMe_2]$  complexes (V.K. Belsky, Y.K. Gun'ko, B.M. Bulychev and G.L. Soloveichik), (419) 299
- Crystal and molecular structures of the octanuclear aluminohydride samarium complex ( $\eta^5-C_5H_3^+Bu_2$ ) $Sm[(\mu_2-H)_2(\mu_3-H)_2Al(Me_2NC_2H_4NMe_2)]_2$   $[(\eta^5-C_5H_3^+Bu_2)SmH]_2[(\mu_2-H)_3Al(\mu_2-H)Al(\mu_2-H)_3]_2$   $[(\mu_3-H)_2Sm(\eta^5-C_5H_3^+Bu_2)_2]$  (V.K. Belsky, Y.K. Gun'ko, B.M. Bulychev and G.L. Soloveichik), (420) 43
- Unsolvated lanthanidocene hydrides and borohydrides. X-Ray crystal structure of  $[(\eta^5-C_5H_3^+Bu_2)_2Ln(\mu-H)]_2$  ( $Ln = Ce, Sm$ ) and  $[(\eta^5-C_5H_3^+Bu_2)_2Sm(\mu-BH_4)]_2$  (Y.K. Gun'ko, B.M. Bulychev, G.L. Soloveichik and V.K. Belsky), (424) 289

## **Scandium**

- Mono(pentamethylcyclopentadienyl)alkoxyscandium alkyl derivatives, ( $\eta^5-C_5Me_5)(OR)ScR'$ , synthesis and characterisation of (W.E. Piers, E.E. Bunel, J.E. Bercaw), (407) 51
- Permethylscandocene complex, investigations of the magnitude of steric and  $\alpha$  deuterium kinetic isotope effects in a carbon–carbon bond-forming reaction of (W.D. Cotter, J.E. Bercaw), (417) C1

## **Selenium**

- $(CO)_6Fe_2(\mu_3-Se)(\mu_3-Te)Pt(PPh_3)_2$  and  $Fe_3(CO)_7(PPh_3)_2(\mu_3-Se)(\mu_3-Te)$ , synthesis and characterisation of (P. Mathur, D. Chakrabarty, Md.M. Hossain), (401) 167
- 3-Thia-7,7-dimethyl-7-germabicyclo[3.3.0]oct-1(5)-ene and its selenium analogue, synthesis and reactivity of (P. Mazerolles, C. Laurent), (402) 35
- Manganese cyclopentadienyldicarbonyl complexes of 1,2,3-selenadiazoles and thiadiazoles; structural comparison of free and complexed 1,2,3-benzothiadiazole and 4-phenyl-1,2,3-thiadiazole (A.J. Mayr, B. Carrasco-Flores, F. Cervantes-Lee, K.H. Pannell, L. Párkányi, K. RaghuVeer), (405) 309

- Titanocene polyselenide complexes, reactions with methyldichloroarsane; synthesis and properties of  $(CH_3C_5H_4)_2Ti(\mu_2-Se_2)_2AsCH_3$  and of  $(CH_3As)_2Se_3$  (B. Holz, R. Steudel), (406) 133
- $Fe_2Ru(\mu_3-Se)(\mu_3-Te)(CO)_9$  and  $(\eta^5-C_5H_5)CoFe_2(\mu_3-Se)(\mu_3-Te)(CO)_6$ , synthesis and characterisation (D. Chakrabarty, Md.M. Hossain, R.K. Kumar, P. Mathur), (410) 143
- Cyclopentadienyliron bis(phosphine) complexes with dimethylchalcogenide ligands; synthesis, reactivity and crystal structure of  $[C_5H_5Fe(P(OCH_3)_3)_2(S(CH_3)_2)]PF_6$  (H. Schumann, J.W. Ziller), (410) 365
- 1,5,9,13-Tetraselenacyclohexadecane selenium coronand; syntheses,  $^{77}Se$  CP-MAS solid state NMR spectra and crystal structures of adducts with copper(I) trifluoromethanesulfonate and mercury(II) cyanide (R.J. Batchelor, F.W.B. Einstein, I.D. Gay, J.-H. Gu, B.M. Pinto), (411) 147
- $Te_2S$ ,  $Te_2Se$  and  $Te_3$  bridged compounds, energies of; CNDO/2 investigation of the mechanism of the bridge reversal process (E.W. Abel, K.G. Orrell, A.G. Osborne, V. Šík, W. Guoxiong), (411) 239
- Monomeric organosamarium(III) chalcogenolates, synthesis by reductive cleavage of E-E bonds (E = S, Se, Te) (A. Recknagel, M. Noltemeyer, D. Stalke, U. Pieper, H.-G. Schmidt, F.T. Edelmann), (411) 347
- Oligonuclear ferrocene derivatives with chalcogene bridges (M. Herberhold, H.-D. Brendel, O. Nuyken, T. Pöhlmann), (413) 65
- Unsaturated mixed metal clusters  $Fe_2Ru_2(CO)_{10}(\mu-CO)(\mu_4-Se)(\mu_4-Te)$  and  $Fe_3Ru(CO)_{10}(\mu-CO)(\mu_4-Se)(\mu^4-Te)$ , construction of (P. Mathur, D. Chakrabarty, Md. M. Hossain), (418) 415
- Selenium and tellurium atoms from selenophene and tellurophene by reaction with trinuclear iron, ruthenium, and osmium clusters, extrusion of; crystal structures of  $[Os_6(\mu-H)(\mu_3-Se)(\mu_4-C_4H_3)-(CO)_{20}]$  and of  $[Ru_4(\mu_3-Se)\mu-C_4H_4](CO)_{11}$  (A.J. Arce, R. Machado, C. Rivas, Y. De Sanctis and A.J. Deeming), (419) 63
- Crystal structure and reactivity of  $[(pp_3)Rh(H)(SeCH_3)]BPh_4$ . An organometallic complex from metal-assisted Se-H activation (M. Di Vaira, D. Rovai, P. Stoppioni and M. Peruzzini), (420) 135
- Sulphur compounds. CXLV. Synthesis and properties of  $Cp_2TiS_2SiR_2$ ,  $Cp_2TiS_2GeR_2$ , and  $Cp'_2TiS_2SnR_2$  ( $Cp = \eta^5-C_5H_5$ ,  $Cp' = \eta^5-C_5H_4CH_3$ , and  $R = C_6H_5$ ) (J. Albertsen and R. Stedel), (424) 153

## Silicon

- Titanocene-catalysed hydromagnesation of silylacetylenes, reversibility of (M.A. Djadchenko, K.K. Pivnitsky, J. Spanig, H. Schick), (401) 1
- Group IVB organometallic radicals, regioselective addition to substituted benzo[2,1,3]thiadiazoles; an ESR investigation (A. Alberti, M. Benaglia, D.F. Dal Monte, M. Lucarini, G.F. Pedulli), (401) 13
- $\{PhSiO\}_6(\mu_4-O)_2(\mu_3-O)_4[Ni_8(\mu_3-O)_2](\mu_3-O)_4(\mu_4-O)_2[PhSiO\}_6 \cdot 14n-BuOH \cdot 10H_2O \cdot 2Me_2CO$  sandwich-type nickel oxide-siloxanolate complex; synthesis and crystal structure of (M.M. Levitsky, O.I. Schegolikhina, A.A. Zhdanov, V.A. Igonin, Yu.E. Ovchinnikov, V.E. Shklover, Yu.T. Struchkov), (401) 199
- Si-substituted silacyclobutanes, thermochemical properties of (M.G. Voronkov, V.A. Klyuchnikov, E.V. Sokolova, T.F. Danilova, G.N. Shvets, A.N. Korchagina, L.E. Gussel'nikov, V.V. Volkova), (401) 245
- Si-H bonds, difluorocarbene insertion into; preparation and properties of difluoromethylsilanes (H. Bürger, R. Eujen, P. Moritz), (401) 249
- 9,10-Dihydrosilaanthracenes with substituents *ortho* to the silicon and methylene bridges, synthesis of (T.C. Bedard, J.Y. Corey, L.D. Lange, N.P. Rath), (401) 261
- Chiral metal complexes of 1-dimethylphosphino-3-dimethylamines, homogeneous asymmetric catalysis by means of (C. Döbler, A. Kinting), (401) C23
- Platinum-catalysed elimination-addition reactions of trisilylethenes (T. Suzuki, P.Y. Lo), (402) 27
- $(\eta^3\text{-Allyl})Fe(CO)_4^+$ , reactions with allyl nucleophiles; approaches to selective isoprenologation (Z. Li, K.M. Nicholas), (402) 105
- 5,5'-(1,1,3,3-Tetramethyldisiloxane-1,3-diyl)bisbicyclo[2.2.1]heptane-2,3-dicarboxylic anhydride, a disiloxane equilibration approach to the preparation and characterisation of (S.A. Swint, M.A. Buese), (402) 145
- Organyltrichlorosilanes, acetolysis with acetic anhydride (K. Käppler, A. Porzel, U. Scheim, K. Rühlmann), (402) 155
- $(\eta^5-C_5H_5)_2Fe(CO)_2Si(Ph)(SiMe_3)_2$  bulky iron-substituted trisilane, X-ray structural study of (R. West, E.K. Pham), (402) 215
- 2-Silatranyl- and 2-(3,7,10-trimethyl)silatranylacetaldehydes, synthesis of (M. Nasim, L.I. Livantsova, D.P. Krut'ko, G.S. Zaitseva, V.S. Petrosyan, J. Lorberth), (402) 313

- 1-Mesityl-3-phenyl-1,2-bis(trimethylsilyl)-1-silacyclopentene and 2-mesityl-2-(phenylethylnyl)hexamethyl-trisilane, thermal isomerisation of (M. Ishikawa, Y. Yuzuriha, T. Horio, A. Kunai), (402) C20
- Substituted polysilanes, synthesis via trifluoromethanesulphonic acid (W. Uhlig), (402) C45
- Racemic acetyl(t-butyl)methylphenylsilane, stereoselective microbial reduction by *Trigonopsis variabilis* and *Corynebacterium dioxydans* (R. Tacke, S. Brakmann, F. Wuttke, J. Fooladi, C. Syldatk, D. Schomburg), (403) 29
- (t-Bu<sub>2</sub>SiOH)<sub>2</sub>, crystal structure of (R. West, E.K. Pham), (403) 43
- Trimethylsilyl ethylene oxide, reaction with  $\alpha$ -sulfonyl anions and  $\alpha,\alpha$ -sulfonyl dianions; stereocontrolled synthesis of (*E*)- and (*Z*)-allylic alcohols (P. Jankowski, S. Marczak, M. Masnyk, J. Wicha), (403) 49
- Bis(2,3-naphthalenediolato)[2-(pyrrolidinio)ethyl]silicate; synthesis and structural characterisation of a zwitterionic  $\lambda^5$ -spirosilicate (C. Strohmann, R. Tacke, G. Mattern, W.F. Kuhs), (403) 63
- Cobalt, ruthenium, and rhodium complexes as catalysts in the reaction of terminal olefins with a hydrosilane and with carbon monoxide (Y. Seki, K. Kawamoto, N. Chatani, A. Hidaka, N. Sonoda, K. Ohe, Y. Kawasaki, S. Murai), (403) 73
- Zirconium and hafnium hexakis(trimethylsilyl)metallocene dichlorides, hindered cyclopentadienyl rotation in; evidence for complete rotation of the tris(trimethylsilyl)cyclopentadienyl ligands (C.H. Winter, D.A. Dobbs, X.-X. Zhou), (403) 145
- Carbamylsilane (Me<sub>3</sub>Si)<sub>3</sub>SiC(O)NMe<sub>2</sub>, preparation and crystal structure of (S.S. Al-Juaid, Y. Derouiche, P.B. Hitchcock, P.D. Lickiss, A.G. Brook), (403) 293
- Chlorinated trimethylsilyl  $\alpha$ -allenyl alcohols, regioselective synthesis via an organoaluminium compound (D. Mesnard, L. Miginiac), (403) 299
- Trimethylsilyl group, 1,2-migration in free radicals (J.M. Harris, I. MacInnes, J.C. Walton, B. Maillard), (403) C25
- [(HOMe<sub>2</sub>Si)(Me<sub>3</sub>Si)<sub>2</sub>C]<sub>2</sub>Zn, [(MeOMe<sub>2</sub>Si)(Me<sub>3</sub>Si)<sub>2</sub>Cl]<sub>2</sub>Zn, and [(MeOMe<sub>2</sub>Si)(Me<sub>3</sub>Si)<sub>2</sub>C]<sub>2</sub>Hg, crystal structures of (F.I. Aigbirhio, S.S. Al-Juaid, C. Eaborn, A. Habtemariam, P.B. Hitchcock, J.D. Smith), (405) 149
- Silicinium ions as intermediates in reactions in solution (C. Eaborn), (405) 173
- Siloxanylpropoxy- and sodium sulfopropoxy- $\alpha,\alpha,\omega$ -trihydroperfluoroalkenes, synthesis of (G. Sonnek, C. Rabe, G. Schmaucks, R. Kaden, I. Lehms), (405) 179
- Phosphasilene, synthesis and reactivity of; an improved route to 1-(2,4,6-tri-tert-butylphenyl)-2-tert-butyl-2-(2,4,6-tri-isopropylphenyl)phosphasilene (Y. van den Winkel, H.M.M. Bastiaans, F. Bickelhaupt), (405) 183
- (Ph<sub>3</sub>P)<sub>2</sub>Pt(H)SiR<sub>3</sub> hydridosilyl complexes, base-induced (Ph<sub>3</sub>P)<sub>2</sub>Pt release from (C. Müller, U. Schubert), (405) C1
- Carbonyl carbene complex of molybdenum, preparation of naphthoquinones and indanones by annellation of (K.H. Dötz, H. Larbig), (405) C38
- 1-Aza-2-silacyclopentane derivatives, synthesis of (M.G. Voronkov, S.V. Kirpichenko, A.T. Abrosimova, A.I. Albanov), (406) 87
- Perorganyloligocyclosiloxanes, thermochemical properties of (M.G. Voronkov, V.A. Klyuchnikov, E.V. Mironenko, G.N. Shvets, T.F. Danilova, Yu.I. Khudobin), (406) 91
- Organosilicon compounds with a Si–N bond, thermochemical properties of (M.G. Voronkov, V.A. Klyuchnikov, L.I. Marenkova, T.F. Danilova, G.N. Shvets, S.I. Tsvetnitskaya, Yu.I. Khudobin), (406) 99
- Silazane derivatives of digallane(4) and diindane(4) with Ga–Ga and In–In bonds, respectively (M. Veith, F. Goffing, S. Becker, V. Huch), (406) 105
- 2,3-Bis(bromomethyl)-1,4-dibromo-2-butene, reduction in the presence of organosilicon and organogermainium dihalides (P. Mazerolles, C. Laurent), (406) 119
- Hexaethylsilsilthiane and bis(trimethylsilylalkyl)sulfides, photoelectron CTC-tetracyanoethylene spectra of (M. Yu. Maroshina, N.N. Vlasova, M.G. Voronkov), (406) 279
- Trimethylsilylquinolines and their methiodides, synthesis and <sup>1</sup>H, <sup>13</sup>C, <sup>14</sup>N, <sup>15</sup>N, <sup>29</sup>Si NMR study of (E. Lukevics, E. Liepiņš, I. Segal, M. Fleisher), (406) 283
- Silylation products of acetohydroxamic acid, structure of (J. Schraml, H.-M. Boldhaus, F. Erdt, E.W. Krahé, C. Bliefert), (406) 299
- spiro*-Germanium compounds, synthesis by 1,4-cycloaddition of cyclic diaminogermyles to 1,3-butadienes and hetero-1,3-butadienes (G. Billeb, K. Bootz, W.P. Neumann, G. Steinhoff), (406) 303
- Silicon–hydrogen bonds, nucleophilic activation of; silicon–nitrogen bond formation by (R.J.P. Corriu, D. Leclercq, P.H. Mutin, J.M. Planeix, A. Vioux), (406) C1

- Bisalkynyltitanocene complexes as bidentate chelating ligands (H. Lang, L. Zsolnai), (406) C5  
 Di-t-butylsilylene and tetra-t-butyldisilene, [4 + 1]- versus [4 + 2]-cycloaddition to 1,4-diazabutadiene (M. Weidenbruch, A. Lesch, K. Peters), (407) 31  
 Tris(trimethylsilyl)silyl silicon derivatives and related tetrasilylsilanes, preparation and reactions of (Y. Derouiche, P.D. Lickiss), (407) 41  
 Silacycloprenes, tungsten-catalysed reactions of (J. Ohshita, M. Ishikawa), (407) 157  
 Mixed trialkyltin compounds, synthesis via trialkylstannyl derivatives of sulphonmaleic and 4-sulphophthalic anhydrides (V.I. Shcherbakov, I.P. Malysheva, O.N. Druzhkova, T.N. Chulkova), (407) 181  
 Palladium-catalysed addition of tin-tin and tin-silicon bonds to allenes (T.N. Mitchell, U. Schneider), (407) 319  
 Lithiumbis(di-tert-butylfluorosilyl)phosphide, a cyclic zwitterion; synthesis and X-ray structure of (U. Klingebiel, M. Meyer, U. Pieper, D. Stalke), (408) 19  
 Eight-membered cyclosilazoxanes, rate studies of the acid-catalysed solvolysis of (Z. Lasocki, M. Witekowa), (408) 27  
 Bis(trimethylsilyl)- and bis(trimethylgermyl)carbodiimide, electron diffraction study of the gas-phase structure of (A. Hammel, H.V. Volden, A. Haaland, J. Weidlein, R. Reischmann), (408) 35  
 Metal-bridging versus chelating  $\text{Me}_2\text{Si}(\text{C}_5\text{H}_4)_2$ -ligands in binuclear complexes; crystal structures of non-sublimed  $[\text{Me}_2\text{Si}(\text{C}_5\text{H}_4)_2\text{Yb}(\mu\text{-Br})]_2$  and  $[(\text{C}_5\text{H}_5)_2\text{Yb}(\mu\text{-Cl})]_2$  (T. Akhnoukh, J. Müller, K. Qiao, X.-F. Li, R.D. Fischer), (408) 47  
 Silylenes, their role in the direct synthesis of methylchlorosilanes (M.P. Clarke, I.M.T. Davidson), (408) 149  
 Silyl-bridged di- to tetranuclear ferrocenes; synthesis, cyclic voltammetry and oxidation of (H. Atzkern, J. Hiermeier, F.H. Köhler, A. Steck), (408) 281  
 Hydrosilylation of the C=C double bond employing homogeneous catalysis with transition metals (R. Skoda-Földes, L. Kollár, B. Heil), (408) 297  
 $[(\text{Me}_3\text{SiC}_5\text{H}_4)_3\text{U}]_2[\mu\text{-O}]$  oxo-bridged bimetallic organouranium complex, synthesis and crystal structure of (J.-C. Berthet, J.-F. Le Maréchal, M. Nierlich, M. Lance, J. Vigner, M. Ephritikhine), (408) 335  
 $(\text{Me}_3\text{Si})_3\text{SiSH}$ , a new radical-based reducing agent (M. Ballestri, C. Chatgilialoglu, G. Seconi), (408) C1  
 Allylic organopotassium and organoboron compounds in the hydroxymethylation and formylation of cycloalkenes (M. Zajlewicz), (409) 103  
 Silyl-substituted phosphinomethanide ligands; preparation of dimeric Al heterocycles; crystal structure of  $\{\text{Me}_2\text{Al}[\text{C}(\text{PMe}_2)(\text{SiMe}_3)_2]\}_2$  (H.H. Karsch, K. Zellner, J. Lachmann, G. Müller), (409) 109  
 Triphenylsilanol, molecular and crystal structure of (H. Puff, K. Braun, H. Reuter), (409) 119  
 Group 14 triflates, synthesis of (W. Uhlig), (409) 377  
 Bis(alkynyl) titanocene complexes of  $\text{FeCl}_2$  and  $\text{CuCl}$  (H. Lang, M. Herres, L. Zsolnai, W. Imhof), (409) C7  
 $\alpha, \omega$ -Diphenylpermethylated oligosilanes, dearylation with triflic acid (K.E. Ruehl, K. Matyjaszewski), (410) 1  
 Trimethyliodosilane, reactions with mono-, di-, and trioxacycloalkanes (M.G. Voronkov, E.I. Dubinskaya), (410) 13  
 Hydrosilylation of carbonyl compounds catalysed by alkali metal fluorides in the presence of crown ethers (Yu. Goldberg, E. Abele, M. Shymanska, E. Lukevics), (410) 127  
 $[5,6:7,8]\text{Dibenzo-2-silabicyclo[2.2.2]octanes}$ , photochemical reactions of (B.R. Yoo, M.E. Lee, I.N. Jung), (410) 33  
 Cyclic tellurides; synthesis, reaction and ligand properties of 2,2,6,6-tetramethyl-1-oxa-4-tellura-2,6-disilacyclohexane ( $\text{C}_6\text{H}_{16}\text{OSi}_2\text{Te}$ ); crystal structure of  $\text{C}_6\text{H}_{16}\text{OSi}_2\text{TeI}_2$  (A.Z. Al-Rubaie, S. Uemura, H. Masuda), (410) 309  
 Disilanes, dehydrogenative polymerisation catalysed by titanocene and zirconocene complexes; novel route to oligo- and poly-silanes (E. Hengge, M. Weinberger, Ch. Jammegg), (410) C1  
 Bis[bis(trimethylsilyl)methyl]tin(II), examination of the reaction with calcium; molecular and crystal structure of benzyl-tris[bis(trimethylsilyl)methyl]stannane (M. Westerhausen, T. Hildenbrand), (411) 1  
 Dimethylindium- and bis(trimethylsilylmethyl)indium-dimesylamide, synthesis of; crystal structure of  $[(\text{Me}_3\text{SiCH}_2)_2\text{InN}(\text{SO}_2\text{Me})_2]_2$  (A. Blaschette, A. Michalides, P.G. Jones), (411) 57  
 $\alpha$ -Functionalised silanes, regioselective synthesis via alkylation of lithiated trimethylsilylacetonitrile (B. Mauzé, L. Miginac), (411) 69

- cis*-Fe(CO)<sub>4</sub>(SiCl<sub>3-n</sub>Me<sub>n</sub>)<sub>2</sub> (*n* = 1–3), conversion with phosphines; competition between CO substitution, SiR<sub>3</sub> elimination and formation of SiR<sub>2</sub> bridged complexes (U. Schubert, M. Knorr, C. Straßer), (411) 75
- (CO)<sub>5-n</sub>(R'<sub>3</sub>P)<sub>n</sub>M(H)SiR<sub>3</sub> (*n* = 2,3), synthesis of phosphane substituted hydridosilyl complexes of chromium, molybdenum and tungsten; synthesis of (CO)<sub>3</sub>(L–L)M(H)Cl by conversion of (CO)<sub>3</sub>(L–L)M(NCR') with HSiClR<sub>2</sub> (L–L = tmeda, bipy, dppe, (PPh<sub>3</sub>)<sub>2</sub>) (H. Piana, U. Schubert), (411) 303
- L<sub>4</sub>Ru<sup>0</sup> complex cores, inter- and intramolecular cleavage of *sp*<sup>2</sup>- and *sp*<sup>3</sup>-C–H bonds with P(o-C<sub>6</sub>H<sub>4</sub>PM<sub>2</sub>)<sub>3</sub>, N(CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>)<sub>3</sub>, MeP(CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>PM<sub>2</sub>)<sub>2</sub>/PR<sub>3</sub> and MeSi(CH<sub>2</sub>PM<sub>2</sub>)<sub>3</sub>/PR<sub>3</sub> (R = Me, OMe) as supporting ligands (L. Dahlenburg, S. Kerstan, D. Werner), (411) 457
- Polycyclic silacyclobutanes, Rh-catalyzed synthesis from norbornadiene and diphenylsilane (H. Brunner, F. Prester), (411) C1
- PhMe<sub>2</sub>E–E'Me<sub>3</sub> (E, E' = Si and Ge), aryl-substituted catenates of group 4B elements; photochemical reactions, formation of a radical pair (K. Mochida, H. Kikkawa, Y. Nakadaira), (412) 9
- Iminosilanes, facile synthesis of; crystal structure of (Me<sub>3</sub>C)<sub>2</sub>(THF)Si=N–SiMe(CMe<sub>3</sub>)<sub>2</sub> (S. Walter, U. Klingebiel, D. Schmidt-Bäse), (412) 319
- Vinylphenyl-substituted silanes, metathesis of (B. Marciniec, C. Pietraszuk), (412) C1
- (C<sub>5</sub>H<sub>5</sub>)Fe(CO)(SiHPh<sub>2</sub>)<sub>2</sub>H, a catalytically active intermediate in the hydrosilylation of acetophenone by diphenylsilane; synthesis of (H. Brunner, K. Fisch), (412) C11
- 1,2,5,6-Tetrasilacycloocta-3,7-dynes, thermal polymerisation of (M. Ishikawa, T. Hatano, T. Horio, A. Kunai), (412) C31
- Methyl- and ethylrhodium oxides, synthesis and structural aspects (W.A. Herrmann, C.C. Romao, P. Kiprof, J. Behm, M.R. Cook, M. Taillefer), (413) 11
- Organic osmium oxide, pyridine complexes of; synthesis, spectra and crystal structure of R<sub>2</sub>OsO<sub>2</sub>·(L) and R<sub>2</sub>OsO<sub>2</sub>·(L)<sub>2</sub> (W.A. Herrmann, S.J. Eder, P. Kiprof), (413) 27
- Pentamethylcyclopentadienyliridium complexes, ring-methyl activation in; synthesis of [(C<sub>5</sub>Me<sub>4</sub>CH<sub>2</sub>–E)IrL<sub>n</sub>] (E = SiMe<sub>3</sub>, PhCH<sub>2</sub>, CH<sub>2</sub>=CHCH<sub>2</sub>, PPh<sub>2</sub>, Pt(PEt<sub>3</sub>)<sub>2</sub>Cl); synthesis and crystal structure of [(1,3-C<sub>5</sub>Me<sub>3</sub>(CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>)Ir(CO)Me(Cl)] (J.A. Miguel-Garcia, H. Adams, N.A. Bailey, P.M. Maitlis), (413) 427
- Platinum–indium and cobalt–gallium complexes, synthesis of; crystal structure of [Cy<sub>2</sub>P(CH<sub>2</sub>CH<sub>2</sub>)PCy<sub>2</sub>]-Pt(InR<sub>2</sub>)X(R) (Cy = cyclo-C<sub>6</sub>H<sub>11</sub>) (R.A. Fischer, J. Behm), (413) C10
- Tolyl- and xylolpentamethyldisilanes, photolysis of (M. Ishikawa, H. Sakamoto), (414) 1
- Trimethylsilylmethyl ketones and acylsilanes, selective formation of alkenes from (A. Fürstner, G. Kollegger, H. Weidmann), (414) 295
- Six-membered fluorosilyl-functionalised hydrazines, preparation of (C. Drost, U. Klingebiel, M. Noltemeyer), (414) 307
- Rhodium-catalysed transsilylation reactions between bis(trimethylsilyl)ethyne and hexyldimethylsilane (T. Suzuki, I. Mita), (414) 311
- Lithium, samarium and ytterbium diiminophosphinates; crystal structures of Li[Ph<sub>2</sub>P(NSiMe<sub>3</sub>)<sub>2</sub>](THF)<sub>2</sub> and [Ph<sub>2</sub>P(NSiMe<sub>3</sub>)<sub>2</sub>]<sub>2</sub>Sm(μ-I)<sub>2</sub>Li(THF)<sub>2</sub> (A. Recknagel, A. Steiner, M. Noltemeyer, S. Brooker, D. Stalke, F.T. Edelmann), (414) 327
- Complexed triphosphanes, preparation and characterisation of; crystal structure of HP[PPh<sub>2</sub>Cr(CO)<sub>5</sub>]<sub>2</sub> (M. Scheer, S. Gremler, E. Herrmann, P.G. Jones), (414) 337
- Enantioselective Co-catalysed homo Diels–Alder reaction of norbornadiene with acetylenes (H. Brunner, F. Prester), (414) 401
- Pentaphenylboracyclotrisiloxane, BSi<sub>2</sub>O<sub>3</sub>Ph<sub>5</sub>, highly strained heterocyclosiloxane; synthesis and crystal structure of (D.A. Foucher, A.J. Lough, I. Manners), (414) C1
- Lithium-coordinated phosphinomethanides, influence of Co-ligands upon; crystal structure of {(THF)Li[C(SiMe<sub>3</sub>)<sub>2</sub>(PM<sub>2</sub>)])<sub>2</sub>} (H.H. Karsch, K. Zellner, S. Gamper, G. Müller), (414) C39
- Substituted  $\eta^4$ -cyclooctadiene-1,5- $\eta^5$ -cyclopentadienyl-cobalt complexes, preparation by reaction of cobaltocenes with alkene cobaltates(1–) (H.-U. Jürgens, J. Alm), (414) C61
- Organothallium(III) dimesylamides (Me<sub>3</sub>SiCH<sub>2</sub>)<sub>2</sub>TIN(SO<sub>2</sub>Me)<sub>2</sub> and (Me<sub>3</sub>SiCH<sub>2</sub>)<sub>2</sub>TIN(SO<sub>2</sub>Me)<sub>2</sub>·H<sub>2</sub>O; synthesis and X-ray structural study (A. Blaschette, P.G. Jones, A. Michalides, M. Naveke), (415) 25
- rac*-Dimethylphenyl[1-(phenylacetamido)ethyl]silane, synthesis and enantioselective enzymatic hydrolysis (H. Hengelsberg, R. Tacke, K. Fritzsche, C. Syldatk, F. Wagner), (415) 39
- $\mu$ -Methylene bis[chlorobis(trimethylsilyl)methylalane] [(Me<sub>3</sub>Si)<sub>2</sub>CH(Cl)Al–CH<sub>2</sub>–Al(Cl)CH(SiMe<sub>3</sub>)<sub>2</sub>] having a heteroadamantane structure (W. Uhl, M. Layh), (415) 181

- Alkoxy silyl amide derivatives of neodymium, samarium, and ytterbium; 4-membered anorganic ring systems in the chemistry of lanthanoids (A. Rechnagel, A. Steiner, S. Brooker, D. Stalke, F.T. Edelmann), (415) 315
- Silicon—the silicon–carbon bond; Annual Survey for the year 1988 (G.L. Larson), (416) 1
- Cyclohexyl(4-fluorophenyl) (3-piperidinopropyl)silanol(*p*-fluorohexahydro-sila-difenol, *p*-F-HHSiD) and derivatives; synthesis and antimuscarinic properties (R. Tacke, K. Mahner, C. Strohmann, B. Forth, E. Mutschler, T. Fribe, G. Lambrecht), (417) 339
- Some silicon, germanium and tin acyloxyderivatives;  $^{13}\text{C}$ ,  $^{17}\text{O}$ ,  $^{29}\text{Si}$  NMR study of (I. Ziemane, E. Liepiņš, L.M. Ignatovich, E. Lukevics), (417) 355
- Neodymium(III) complexes with bulky cyclopentadienyl and benzamidine ligands, a comparison (A. Recknagel, F. Knösel, H. Gornitzka, M. Noltemeyer, F.T. Edelmann, U. Behrens), (417) 363
- Alkynyl substituted chlorosilanes as cluster units (H. Lang, U. Lay, L. Zsolnai), (417) 377
- Zirconocene complexes, olefin hydrosilation catalyzed by (M.R. Kesti, M. Abdulrahman, R.M. Waymouth), (417) C12
- $\text{FeCl}_3$ -doped poly(alkynylsilane)s, electrical conductivity of (R.J.P. Corriu, W.E. Douglas, Z.-X. Yang, F. Garnier, A. Yasser), (417) C50
- Pyridinyloxazoline cobalt(I) complexes as catalysts, enantioselective hydrosilylierung of acetophenone (H. Brunner, K. Amberger), (417) C63
- Vinyltriethoxysilane, unusual reaction with triethoxysilane catalyzed by nickel acetylacetone (B. Marciniec, H. Maciejewski, J. Mirecki), (418) 61
- ( $\text{Me}_3\text{SiCH}_2$ )<sub>x</sub> $\text{SiH}_{4-x}$  ( $x = 1$ –4), mono-, bis-, tris- and tetrakis(trimethylsilylmethyl)silanes; synthesis and characterization of (C.K. Whitmarsh, L.V. Interrante), (418) 69
- Phenylethynyl chlorosilanes, complex-bonded; reactivity of (U. Lay, H. Lang), (418) 79
- Silatrane skeleton, effect of ring substitution on the geometry of; crystal structures of 1-phenyl-3,7,10-trimethylsilatrane and 1-*p*-tolyl-carbasilatrane (L. Párkányi, V. Fülöp, P. Hencsei, I. Kovács), (418) 173
- (=O → Si) (aryloxymethyl)trifluorosilanes, vibrational spectra of (G.A. Gavrilova, N.N. Chipanina, Yu.L. Frolov, L.I. Gubanova, M.G. Voronkov), (418) 291
- 1,1-Dimethyl-6-[1,1-dimethyl-2-(trimethylsilyl)ethyl]-4-(pentamethyldisilanyl)silepin, thermal behaviour of (H. Sakamoto, M. Ishikawa), (418) 305
- (dppe) $\text{Pt}(\text{Me}_2\text{SiCH}_2\text{CH}_2\text{SiMe}_2)$ , synthesis by exchange of silyl groups (U. Schubert, C. Müller), (418) C6
- Silatrane from reactions of chloromethylsilatrane with chlorosilanes and magnesium in tetrahydrofuran (V. Gevorgyan, L. Borisova, E. Lukevics), (418) C21
- Alkyne bridged  $\text{Co}_2\text{Mo}_2$  clusters with two triply-bridged sulphur atoms, synthesis of; crystal structure of  $[\text{Co}_2\text{Mo}_2(\eta^5\text{-Cp})_2(\mu_3\text{-S})_2(\mu_4\text{-CF}_3\text{C}_2\text{Me})(\text{CO})_4]$  (F. Robin, R. Rumin, F.Y. Pétillon, K. Foley, K.W. Muir), (418) C33
- Heterocumulenes, aminosilylation of; intramolecular decomposition of their silyl-functionalized adducts (R.J.P. Corriu, G.F. Lanneau and V.D. Mehta), (419) 9
- Aminosilanes, HeI and HeII photoelectron spectroscopic investigation of substituent effects in (A. Nagy, J.C. Green, L. Szepes and L. Zanathy), (419) 27
- Opening of cluster frameworks, increase in skeletal electrons via amino substituents (B. Eber, G. Huttner, C. Emmerich, J.C. Daran, B. Heim and Y. Jeannin), (419) 43
- Silylphosphinites and their complexation to manganese, crystal structure of *cis*- $\text{MnBr}(\text{CO})_4[\text{P}(4\text{-CH}_3\text{C}_6\text{H}_4)_2\text{OSiMe}_3]$  (K.M. Cooke, T.P. Kee, A.L. Langton and M. Thornton-Pett), (419) 171
- Bis(trimethylsilyl)methylamine, handy preparation of (J.-P. Picard, S. Grelier, J. Dunoguès, J.-M. Aizpurua and C. Palomo), (419) C1
- On carbon-13 spectra of thio- and dithiocarboxylic acids triorganogroup 14 metal esters (S. Kato, A. Hori, M. Mizuta, T. Katada, H. Ishihara, K. Fujieda and Y. Ikebe), (420) 13
- Alcynylsilanes fonctionnels. III. Synthèse régiosélective par voie organozincique de 1-triméthylsilyl-4-éne-1-ynes fonctionnels (D. Mesnard and L. Miginiac), (420) 163
- Acetylen-Komplexe von Di( $\eta^5$ -cyclopentadienyl)molybdän: Synthesen und Protonierung. Struktur von  $[\text{Cp}_2\text{MoH}(\eta^2\text{-C}_2\text{SiMe}_3)_2]\text{PF}_6$  (G.E. Herberich, U. Englert and W. Faßbender), (420) 303
- Hydrosilylation of olefins with monosilane catalyzed by transition metal complexes (M. Itoh, K. Iwata, R. Takeuchi and M. Kobayashi), (420) C5
- Photo-induced electron transfer reaction between hexamethyldisilane and quinones as studied by a CIDNP technique (M. Igarashi, T. Ueda, M. Wakasa and Y. Sakaguchi), (421) 9

- Reactions at highly sterically hindered organosilicon centres. Chemistry of  $(Me_3Si)_3CSiPhHI$ ,  $(Me_3Si)_3CSiPh(OMe)I$ , and related species (Z.H. Aiube and C. Eaborn), (421) 159
- The recycling of organosilyl protecting species used in organic synthesis and the water binding ability of the silanol  $^1BuMe_2SiOH$  (P.D. Lickiss and K.M. Stubbs), (421) 171
- Zum gezielten Aufbau oligomerer Strukturen der Elemente der 14. Gruppe mittels ihrer Triflatderivate (W. Uhlig), (421) 189
- Silicon—the silicon–carbon bond; Annual Survey for the year 1989 (G.L. Larson), (422) 1
- The crystal structure of  $(Me_3Si)_3CSi(OH)_2(OCOCF_3)$ . Novel hydrogen bonding (S.S. Al-Juaid, C. Eaborn and P.B. Hitchcock), (423) 5
- Zur Chemie der Silylene: Cycloadditionen von Methoxymethylsilylen mit Heterodienen (J. Heinicke und B. Gehrhus), (423) 13
- Synthesis of bis(pentachlorophenyl)silanediol and related compounds. Crystal structure of diethoxybis(pentachlorophenyl)silane (J. Carilla, L. Fajará, L. Juliá, J. Riera, J. Lloveras, N. Verdaguer and I. Fita), (423) 163
- Pyrrolyl compounds of main group elements. Synthesis of group 14 5-metallated pyrrole-2-carbaldehydes (F. Denat, H. Gaspard-Iloughmane and J. Dubac), (423) 173
- Siliciumverbindungen mit starken intramolekularen sterischen Wechselwirkungen. XLV. Reaktionen von Tetra-t-butyldisilen mit CN-Mehrfachbindungen: Cycloadditionen und Umlagerungen (M. Weidenbruch, A. Lesch, K. Peters and H.G. von Schnerring), (423) 329
- Darstellung und Struktur von  $(EtO)_2P(O)CH_2Si(Me)_2CH_2SnMe_2Cl$ —ein Sechsringchelat mit Seselkonformation und  $P=O \cdots Sn(Cl)Me_2CH_2$ -trigonaler Bipyramide am Lewis-aciden Zinn (U. Kolb, M. Dräger, E. Fischer and K. Jurkschat), (423) 339
- Novel class of transition metal coordination compounds with macrocyclic organosiloxanolate ligands; their synthesis and crystal structure (V.A. Igonin, O.I. Shchegolikhina, S.V. Lindeman, M.M. Levitsky, Yu.T. Struchkov and A.A. Zhdanov), (423) 351
- Synthesis of new vinylsilanes containing an asymmetric silicon via platinum catalyzed hydrosilylation of acetylene and monosubstituted alkynes (V. Gevorgyan, L. Borisova, J. Popelis, E. Lukevics, Z. Poltynowic, J. Gulinski and B. Marciniac), (424) 15
- Sulphur compounds. CXLV. Synthesis and properties of  $Cp_2TiS_2SiR_2$ ,  $Cp_2TiS_2GeR_2$ , and  $Cp'_2TiS_2SnR_2$  ( $Cp = \eta^5-C_5H_5$ ,  $Cp' = \eta^5-C_5H_4CH_3$ , and  $R = C_6H_5$ ) (J. Albertsen and R. Steudel), (424) 153
- Die Reaktion von Silyl-bis(trifluormethansulfonaten) mit Diazoessigestern und Diazomethyl-ketonen (A. Fronda, F. Krebs, B. Daucher, T. Werle und G. Maas), (424) 253
- Zur Stereochemie der mikrobiellen Reduktion von *rac*-Acetyl(t-butyl)methylphenylsilan mit *Trigonopsis variabilis* (DSM 70714) und *Corynebacterium dioxydans* (ATCC 21766): Aufklärung der absoluten Konfiguration der Biotransformationsprodukte (*Si,R,CR*)- und (*Si,S,CR*)-t-Butyl(1-hydroxyethyl)methylphenylsilan (R. Tacke, F. Wuttke und H. Henke), (424) 273
- Bifunctional catalysis in the condensation of silanols and alkoxy silanes (H.-K. Chu, R.P. Cross and D.I. Crossan), (425) 9
- Zur Kenntnis funktionalisierter Zinn-Phosphorverbindungen: Darstellung und Reaktionen von Chlorstannylphosphinen (D. Hänsgen, E. Stahlhut, H. Aldenhoven and A. Dörr), (425) 19

## Silver

- Silver–tin complex salts; a dimeric tin-complex anion with a double hydroxyl bridge; crystal structure of  $[Ag(AsPh_3)_4]_2[Sn_2Ph_4(NO_3)_4(OH)_2 \cdot 2CH_3CN$  (A. Bonardi, A. Cantoni, C. Pelizzi, G. Pelizzi, P. Tarasconi), (402) 281
- Platinum–silver cluster compound,  $[Pt_3Ag(O_3SCF_3)\mu\text{-CO}]_3(PCy_3)_2(dppp)]$  ( $Cy = C_6H_{11}$ ,  $dppp = ((C_6H_5)_2P)_2C_3H_6$ ), with a distorted tetrahedral geometry; synthesis and crystal structure of (A.D. Burrows, J.G. Jeffrey, J.C. Machell, D.M.P. Mingos), (406) 399
- Group IB metal heteronuclear cluster compounds in the solid state, high-resolution  $^{31}P\{-^1H\}$  nuclear magnetic resonance studies of (S.S.D. Brown, I.D. Salter, D.J. Smith, N.J. Clayden, C.M. Dobson), (408) 439
- $[(Co_2(CO)_6PhC_2)_2]$  acetylene-linked dimeric cluster, synthesis and structural characterisation of (B.F.G. Johnson, J. Lewis, P.R. Raithby, D.A. Wilkinson), (408) C9
- Three-membered metallacycle  $Cp_3Rh_3(\mu\text{-CO})_3$ , cluster and benzenoid system in reactions with 12-electron particles  $PPh_3M^+$  ( $M = Cu$ , Ag, Au) and  $CpM^+$  ( $M = Fe$ , Ru) (M.I. Rybinskaya, A.R. Kudinov, D.V. Muratov, N.P. Gambaryan, I.V. Stankevich, A.L. Chistyakov), (413) 419
- Acetylenes and diazoketones in organic synthesis (F. Serratosa), (413) 445

## Sodium

- Sodium alkoxides, standard enthalpies of formation (J.P. Leal, A. Pires de Matos, J.A. Martinho Simões), (403) 1
- Sodium and potassium pentamethylcyclopentadienyl pyridine solvates, preparation and crystal structures of (G. Rabe, H.W. Roesky, D. Stalke, F. Pauer, G.M. Sheldrick), (403) 11
- Allylic organopotassium and organoboron compounds in the hydroxymethylation and formylation of cycloalkenes (M. Zajdlewicz), (409) 103
- Allyl alkali metals, comparative theoretical study of (N.J.R. van Eikema Hommes, M. Bühl, P.v.R. Schleyer, Y.-D. Wu), (409) 307
- Organonatrium complexes with benzyl, xylyl, diphenylmethyl, and 1-phenyl ethyl carbanions (S. Corbeil, N.P. Lorenzen, J. Kopf, E. Weiss), (415) 293
- Alkali metal derivatives of sulfinimidamides, preparation and crystal structures (F. Pauer, D. Stalke), (418) 127
- Novel class of transition metal coordination compounds with macrocyclic organosiloxanolate ligands; their synthesis and crystal structure (V.A. Igonin, O.I. Shchegolikhina, S.V. Lindeman, M.M. Levitsky, Yu.T. Struchkov and A.A. Zhdanov), (423) 351

## Sulphur

- Reaction of norbornadiene with sulfur and dodecacarbonyltriiron; crystal structure of one of the isomeric products (A.I. Nekhaev, S.D. Alekseeva, B.I. Kolobkov, G.G. Aleksandrov, M.T. Toshev, H.B. Dustov, N.A. Parpiev), (401) 65
- $\text{Fe}_3(\eta^4\text{-C}_6\text{H}_8)(\mu\text{-CO})_2(\text{CO})_6(\mu_3\text{-S})$  trinuclear cluster, synthesis and structure of (A.I. Nekhaev, S.D. Alekseeva, B.I. Kolobkov, G.G. Aleksandrov, M.T. Toshev, H.B. Dustov), (401) 75
- Thiolato bridged-iron carbonyl complexes, synthesis of (P.M. Treichel, R.A. Crane, R. Matthews, K.R. Bonnin, D. Powell), (402) 233
- Triorganostannate esters of dicarboxylic acids, synthesis of; crystal structure of dicyclohexylammonium 2,6-pyridinedicarboxylatotributylstannate (S.W. Ng, V.G. Kumar Das, E.R.T. Tiekkink), (403) 111
- $\eta^5\text{-Cyclopentadienyl}(\text{dicarbonyl})[\eta^2\text{-diethylamino(methylthio)carbene}]$ tungsten, preparation by thiomethylation of  $\eta^5\text{-cyclopentadienyl}(\text{dicarbonyl})\text{diethylaminocarbintungsten}$  (F.R. Kreißl, N. Ullrich, H. Keller), (403) C29
- $[\text{Fe}_2(\mu\text{-C}(\text{CN})\text{SM}\text{e}_2)_2(-\text{CO})(\text{CO})_2(\text{Cp})_2]\text{SO}_3\text{CF}_3$  ( $\text{Cp} = \eta\text{-C}_5\text{H}_5$ ), reactions with alcohols (L. Busetto, M.C. Cassani, V. Zanotti, V.G. Albano, D. Braga), (415) 395

## Tantalum

- $\text{Cp}_2\text{MM}'(\text{Cp})(\text{CO})_3$  ( $\text{M} = \text{Nb}, \text{Ta}; \text{M}' = \text{Mo}, \text{W}$ ), heteronuclear complexes with bridging and semi-bridging carbonyl groups; synthesis and dynamic behaviour (D. Perrey, J.C. Leblanc, C. Moïse, J. Martin-Gil), (412) 363
- $(\text{C}_5\text{Me}_5\text{Ta})\text{PM}\text{e}_3)_2$ , ring- and phosphine-metallated forms of competitive ligand C—H bond activations (V.C. Gibson, T.P. Kee, S.T. Carter, R.D. Sanner, W. Clegg), (418) 197
- $d^2$  Tantalum complexes  $\text{Cp}_2\text{Ta}(\text{L})\text{H}$ , protonation of; synthesis and structural studies of cationic dihydride complexes (J.-F. Leboeuf, O. Lavastre, J.-C. Leblanc, C. Moïse), (418) 359

## Technetium

- $(\text{C}_5\text{Me}_5)_2\text{Tc}(\text{CO})_3$ , photolysis products of; synthesis of  $(\text{C}_5\text{Me}_5)_2\text{Tc}_2(\text{CO})_5$  and  $(\text{C}_5\text{Me}_5)_2\text{Tc}_2(\text{CO})_3$ ; X-ray structure analysis of tri- $\mu$ -carbonyl-bis[ $\eta^5$ -pentamethylcyclopentadienyl]technetium ( $\text{Tc} \equiv \text{Tc}$ ) (K. Raptis, B. Kanellakopulos, B. Nuber, M.L. Ziegler), (405) 323
- $\text{LTc}(\text{CO})_3$  compounds ( $\text{L} = \text{C}_5\text{Me}_5, \text{C}_5\text{Me}_4\text{Et}, \text{C}_9\text{H}_7$ ), preparation and characterisation of; crystal structures of  $(\eta^5\text{-C}_5\text{Me}_5)\text{M}(\text{CO})_3$  ( $\text{M} = \text{Tc}, \text{Re}$ ),  $(\eta^5\text{-C}_5\text{Me}_4\text{Et})\text{Tc}(\text{CO})_3$  and  $(\eta^5\text{-C}_9\text{H}_7)\text{Tc}(\text{CO})_3$  (K. Raptis, E. Dornberger, B. Kanellakopulos, B. Nuber, M.L. Ziegler), (408) 61
- $\text{Cp}'\text{Tc}(\text{CO})_3$  derivatives; synthesis, reactions and structure of (H.H. Knight Castro, A. Meetsma, J.H. Teuben, W. Vaalburg, K. Panek, G. Ensing), (410) 63
- Tris(cyclopentadienyl)technetium(III),  $(\eta^5\text{-C}_5\text{H}_5)_2\text{Tc}(\eta^1\text{-C}_5\text{H}_5)$ , X-ray diffraction study of; dipole moment and charge distribution of  $(\text{C}_5\text{H}_5)_3\text{M}$  ( $\text{M} = \text{V}, \text{Tc}, \text{Re}$ ) (C. Apostolidis, B. Kanellakopulos, R. Maier, J. Rebizant, M.L. Ziegler), (411) 171
- Methyl- and ethylrhodium oxides, synthesis and structural aspects (W.A. Herrmann, C.C. Romao, P. Kiprof, J. Behm, M.R. Cook, M. Taillefer), (413) 11
- Manganese, technetium, and rhenium; Annual Survey covering the year 1987 (P.M. Treichel), (416) 291

**Tellurium**

- (CO)<sub>6</sub>Fe<sub>2</sub>(μ<sub>3</sub>-Se)(μ<sub>3</sub>-Te)Pt(PPh<sub>3</sub>)<sub>2</sub> and Fe<sub>3</sub>(CO)<sub>7</sub>(PPh<sub>3</sub>)<sub>2</sub>(μ<sub>3</sub>-Se)(μ<sub>3</sub>-Te), synthesis and characterisation of (P. Mathur, D. Chakrabarty, Md.M. Hossain), (401) 167
- Fe<sub>3</sub>(CO)<sub>9</sub>(μ<sub>3</sub>-Te)<sub>2</sub>, addition to methylene groups to (P. Mathur, V.D. Reddy, R. Bohra), (401) 339
- ortho*-Tellurated azomethines with intramolecular N → Te coordination, molecular and crystal structure of (V.I. Minkin, I.D. Sadekov, A.A. Maksimenko, O.E. Kompan, Yu.T. Struchkov), (402) 331
- [{μ-CH<sub>3</sub>Te}Fe<sub>2</sub>(CO)<sub>6</sub>]<sub>2</sub>μ-Te(CH<sub>2</sub>)Te-μ] methylene-bridged double butterfly shaped complex, synthesis and structural characterisation of (P. Mathur, V.D. Reddy, K. Das, U.C. Sinha), (409) 255
- Fe<sub>2</sub>Ru(μ<sub>3</sub>-Se)(μ<sub>3</sub>-Te)(CO)<sub>9</sub> and (η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)CoFe<sub>2</sub>(μ<sub>3</sub>-Se)(μ<sub>3</sub>-Te)(CO)<sub>6</sub>, synthesis and characterisation of (D. Chakrabarty, Md.M. Hossain, R.K. Kumar, P. Mathur), (410) 143
- Cyclic tellurides; synthesis, reaction and ligand properties of 2,2,6,6-tetramethyl-1-oxa-4-tellura-2,6-disilacyclohexane (C<sub>6</sub>H<sub>16</sub>OSi<sub>2</sub>Te); crystal structure of C<sub>6</sub>H<sub>16</sub>OSi<sub>2</sub>TeI<sub>2</sub> (A.Z. Al-Rubaie, S. Uemura, H. Masuda), (410) 309
- Cyclopentadienyliron bis(phosphine) complexes with dimethylchalcogenide ligands; synthesis, reactivity and crystal structure of [C<sub>5</sub>H<sub>5</sub>Fe(P(OCH<sub>3</sub>)<sub>3</sub>)<sub>2</sub>(S(CH<sub>3</sub>)<sub>2</sub>)]PF<sub>6</sub> (H. Schumann, J.W. Ziller), (410) 365
- Te<sub>2</sub>S, Te<sub>2</sub>Se and Te<sub>3</sub> bridged compounds, energies of; CNDO/2 investigation of the mechanism of the bridge reversal process (E.W. Abel, K.G. Orrell, A.G. Osborne, V. Sik, W. Guoxiong), (411) 239
- Monomeric organosamarium(III) chalcogenolates, synthesis by reductive cleavage of E-E bonds (E = S, Se, Te) (A. Recknagel, M. Noltemeyer, D. Stalke, U. Pieper, H.-G. Schmidt, F.T. Edelmann), (411) 347
- Oligonuclear ferrocene derivatives with chalcogene bridges (M. Herberhold, H.-D. Brendel, O. Nuyken, T. Pöhlmann), (413) 65
- Pentacarbonyl(tellurobenzaldehyde)tungsten; synthesis, thermolysis and reactions with conjugated diened (H. Fischer, A. Fruh, C. Troll), (415) 211
- Compounds containing short Te-N bonds, syntheses and crystal structures of (H.W. Roesky, J. Münenberg, R. Bohra, M. Noltemeyer), (418) 339
- Aryltellurolato-bridged dinuclear complexes of platinum(II), preparation and characterization of (V.K. Jain, S. Kannan), (418) 349
- Trichloromethylisocyanide complexes, P-functionalized isocyanide; synthesis of (S. Ahn, G. Beck, W.P. Fehlhammer), (418) 365
- Unsaturated mixed metal clusters Fe<sub>2</sub>Ru<sub>2</sub>(CO)<sub>10</sub>(μ-CO)(μ<sub>4</sub>-Se)(μ<sub>4</sub>-Te) and Fe<sub>3</sub>Ru(CO)<sub>10</sub>(μ-CO)(μ<sub>4</sub>-Se)(μ<sub>4</sub>-Te), construction of (P. Mathur, D. Chakrabarty, Md. M. Hossain), (418) 415
- Selenium and tellurium atoms from selenophene and tellurophene by reaction with trinuclear iron, ruthenium, and osmium clusters, extrusion of; crystal structures of [Os<sub>6</sub>(μ-H)(μ<sub>3</sub>-Se)(μ<sub>4</sub>-C<sub>4</sub>H<sub>3</sub>)-(CO)<sub>20</sub>] and of [Ru<sub>4</sub>(μ<sub>3</sub>-Se)μ-C<sub>4</sub>H<sub>4</sub>](CO)<sub>11</sub>] (A.J. Arce, R. Machado, C. Rivas, Y. De Sanctis and A.J. Deeming), (419) 63
- TeCl<sub>4</sub> in organic synthesis. Elimination reactions (M. Albeck and T. Tamary), (420) 35
- Mixed chalcogen carbonyl compounds. IV. Reactivity of Fe<sub>2</sub>(CO)<sub>6</sub>(μ-STe) and Fe<sub>3</sub>(CO)<sub>9</sub>(μ<sub>3</sub>-S)(μ<sub>3</sub>-Te) towards coordinatively unsaturated species (P. Mathur, D. Chakrabarty, M.M. Hossain and R.S. Rashid), (420) 79
- The crystal structures of 2-(2'-pyridyl)phenyltellurium(II) bromide and of the inclusion compound bis[2-(2'-pyridyl)phenyltellurium(II) chloride]·p-ethoxyphenyl-mercury(II) chloride (M.R. Greaves, T.A. Hamor, B.J. Howlin, T.S. Lobana, S.A. Mbogo, W.R. McWhinnie and D.C. Povey), (420) 327
- Styryl coupling vs. styryl acetate formation in reactions of styryl tellurides with palladium(II) salts (S.U.H. Takahashi and K. Ohe), (423) C9
- Crystal structures of triphenyltellurium(IV) methylxanthate and isobutylxanthate — dependence on the steric influence of the alkyl group of the xanthate system (A.K. Singh, J.K. Basumatary, T.P. Singh and B. Padmanabhan), (424) 33

**Terbium**

- Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente. XXX. Kristall- und Molekülstrukturen ungewöhnlich gut kristallisierender Addukte des Grundkörpers Tris(η<sup>5</sup>-cyclopentadienyl)-lanthanoid(III) (H. Schulz, H. Schultze, H. Reddmann, M. Link und H.-D. Amberger), (424) 139

## Thallium

- Metalloporphyrins with metal–metal  $\sigma$ -bonds; synthesis, spectroscopic characterization, and electrochemistry of (P)MRe(CO)<sub>5</sub> where P is the dianion of octaethylporphyrin (OEP) or tetraphenylporphyrin (TPP) and M = Al, Ga, In or Tl (R. Guillard, A. Zrineh, A. Tabard, L. Courthaudon, B. Han, M. Ferhat, K.M. Kadish), (401) 227
- Titanium, iron and thallium vinylcyclopentadienyl derivatives, synthesis and structure of (M. Ogasa, D.T. Mallin, D.W. Macomber, M.D. Rausch, R.D. Rogers, A.N. Rollins), (405) 41
- Trifluoromethyl gallium, indium, and thallium compounds, syntheses of; mechanism of polar trifluoromethyl group transfer (D. Naumann, W. Strauß, W. Tyrra), (407) 1
- Tetrahydrofurfurylcyclopentadiene and its thallium, manganese, and rhodium complexes, synthesis of; low temperature matrix photolysis of the manganese complex (T.E. Bitterwolf, K.A. Lott, A.J. Rest), (408) 137
- [TlAu(C<sub>5</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>]<sub>2</sub> tetrametallic complex, synthesis and structure of (G.K. Anderson, N.P. Rath), (414) 129
- Organothallium(III) dimesylamides (Me<sub>3</sub>SiCH<sub>2</sub>)<sub>2</sub>TlN(SO<sub>2</sub>Me)<sub>2</sub> and (Me<sub>3</sub>SiCH<sub>2</sub>)<sub>2</sub>TlN(SO<sub>2</sub>Me)<sub>2</sub>·H<sub>2</sub>O; synthesis and X-ray structural study (A. Blaschette, P.G. Jones, A. Michalides, M. Naveke), (415) 25
- Group IIIB organometallic peroxides, synthesis and reactivity of (Yu.A. Alexandrov, N.V. Chikinova), (418) 1

## Tin

- Group IVB organometallic radicals, regioselective addition to substituted benzo[2,1,3]thiadiazoles; an ESR investigation (A. Alberti, M. Benaglia, D.F. Dal Monte, M. Lucarini, G.F. Pedulli), (401) 13
- Methyl 4,6-O-benzylidene-3-deoxy-3-triphenylstannyl- $\alpha$ -D-altropyranoside, structure and reactivity of (S.M.S.V. Doidge-Harrison, I.W. Nowell, P.J. Cox, R.A. Howie, O.J. Taylor, J.L. Wardell), (401) 273
- Organotin hexacyanoferrates, synthesis and characterisation of (A. Bonardi, C. Carini, C. Pelizzi, G. Pelizzi, G. Predieri, P. Tarasconi, M.A. Zoroddu, K.C. Molloy), (401) 283
- Linked metal acetylide complexes for nonlinear optics, synthesis of (S.J. Davies, B.F.G. Johnson, J. Lewis, M.S. Khan), (401) C43
- Polyfunctional stannylallenes, NMR spectroscopic study of (E. Liepiņš, I. Birģe, E. Lukevics, E.T. Bogoradovsky, V.S. Zagorodny), (402) 43
- ( $\eta^3$ -Allyl)Fe(CO)<sub>4</sub><sup>+</sup>, reactions with allyl nucleophiles; approaches to selective isoprenologation (Z. Li, K.M. Nicholas), (402) 105
- Substituted 9,9-bis(n-butyl)dibenzostannoles, syntheses of (H.-A. Brune, R. Hohenadel, G. Schmidtberg, U. Ziegler), (402) 171
- Dibenzoplatinol system, photochemical and UV-spectroscopic studies of (H.-A. Brune, R. Hohenadel, G. Schmidtberg, U. Ziegler), (402) 179
- Silver–tin complex salts; a dimeric tin-complex anion with a double hydroxyl bridge; crystal structure of [Ag(AsPh<sub>3</sub>)<sub>4</sub>]<sub>2</sub>[Sn<sub>2</sub>Ph<sub>4</sub>(NO<sub>3</sub>)<sub>4</sub>(OH)<sub>2</sub>]·2CH<sub>3</sub>CN (A. Bonardi, A. Cantoni, C. Pelizzi, G. Pelizzi, P. Tarasconi), (402) 281
- Triorganostannyly esters of amic acids, synthesis and spectroscopic studies of; crystal structure of the triphenylstannyl ester of citraconic acid monopyrrolidide (A. Samuel-Lewis, P.J. Smith, J.H. Aupers, D.C. Povey), (402) 319
- Triphenyltin derivatives of isothiazol-3(2H)-one 1,1-dioxides; synthesis, Mössbauer spectra, fungitotoxicity data, and crystal structure of triphenylstannyl 1,2-benzisothiazol-3(2H)-one 1,1-dioxide·N,N-dimethylformamide (S.W. Ng, A.J. Kuthubutheen, Z. Arifin, C. Wei, V.G. Kumar Das, B. Schulze, K.C. Molloy, W.-H. Yip, T.C.W. Mak), (403) 101
- Triorganostannate esters of dicarboxylic acids, synthesis of; crystal structure of dicyclohexylammonium 2,6-pyridinedicarboxylotributylstannate (S.W. Ng, V.G. Kumar Das, E.R.T. Tiekkink), (403) 111
- Diorganotin(IV) complexes with thiophenoxyacetic acid, synthesis and spectroscopic properties of; crystal structures of [<sup>n</sup>Pr<sub>2</sub>Sn(O<sub>2</sub>CCH<sub>2</sub>SPh)<sub>2</sub>] and {[R<sub>2</sub>Sn(O<sub>2</sub>CCH<sub>2</sub>SPh)]<sub>2</sub>O}<sub>2</sub>, R = <sup>n</sup>Pr, <sup>n</sup>Bu (G.K. Sandhu, N. Sharma, E.R.T. Tiekkink), (403) 119
- Trityltin acetate, <sup>2</sup>J(<sup>119</sup>Sn–O–<sup>13</sup>CO) coupling in (H.-V. Pham, B. Rusterholz, E. Pretsch, W. Simon), (403) 311
- Tris(3,5-dimethylpyrazolyl)borates, preparation and properties of (G. Gioia Lobbia, F. Bonati, P. Cecchi, A. Lorenzotti, C. Pettinari), (403) 317

- C-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325
- Organotin derivatives of 4-acyl-5-pyrazolones; crystal structure of *trans*-di(*t*-butyl)bis(1-phenyl-3-methyl-4-benzoyl-pyrazolone-5-ato)tin(IV) (C. Pettinari, G. Rafaiani, G. Gioia Lobbia, A. Lorenzotti, F. Bonati, B. Bovio), (405) 75
- Palladium catalysed and free radical hydrostannation of allenes, a comparison (T.N. Mitchell, U. Schneider), (405) 195
- Colourless C<sub>5</sub>H<sub>5</sub>SbCl<sub>2</sub>, intra- and intermolecular antimony–Cp  $\pi$ -interactions in (W. Frank), (406) 331
- Palladium-catalysed dimerisation of isoprene with carbon dioxide in the presence of organotin ethoxide and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (H. Hoberg, M. Minato), (406) C25
- Trimethyltin diazoacetic ester, Me<sub>3</sub>SnC(N<sub>2</sub>)CO<sub>2</sub>Et, crystal structure of (J. Lorberth, S.-H. Shin, H. Donath, S. Wocadlo, W. Massa), (407) 167
- n-Butyltinoxo carboxylates, [<sup>n</sup>BuSn(O)(O<sub>2</sub>CR')]<sub>6</sub> (R' = <sup>i</sup>Pr or <sup>t</sup>Bu), synthesis and characterisation of; crystal structure of [<sup>n</sup>BuSn(O)(O<sub>2</sub>C<sup>t</sup>Bu)]<sub>6</sub> (V.B. Mokal, V.K. Jain, E.R.T. Tiekkink), (407) 173
- Mixed trialkyltin compounds, synthesis via trialkylstannyl derivatives of sulphomaleic and 4-sulphophthalic anhydrides (V.I. Shcherbakov, I.P. Malysheva, O.N. Druzhkova, T.N. Chulkova), (407) 181
- Divalent organotin compounds, oxidative-addition reactions of molecular diiodine and dibromine to; crystal structures of bis[8-(dimethylamino)-1-naphthyl]tin(IV)dibromide and {2,6-[bis(dimethylamino)methyl]phenyl}(4-tolyl)tin(IV) diode (J.T.B.H. Jastrzebski, P.A. van der Schaaf, J. Boersma, G. van Koten, M. de Wit, Y. Wang, D. Heijdenrijk, C.H. Stam), (407) 301
- {Me<sub>2</sub>NCH[(EtO)<sub>2</sub>P=O]<sub>2</sub>·Me<sub>2</sub>SnCl<sub>2</sub>} dimer; crystal structure of a binuclear bis(chelate) complex of tin(IV) with *cis*-linked octahedra (J. Lorberth, S.-H. Shin, M. Otto, S. Wocadlo, W. Massa, N.S. Yashina), (407) 313
- Palladium-catalysed addition of tin–tin and tin–silicon bonds to allenes (T.N. Mitchell, U. Schneider), (407) 319
- {[R<sub>2</sub>Sn(O<sub>2</sub>C<sup>t</sup>Bu)]<sub>2</sub>O}<sub>2</sub> (R = Me, Et, <sup>n</sup>Pr and <sup>n</sup>Bu), synthesis and characterisation of; X-ray crystal structures of {[R<sub>2</sub>Sn(O<sub>2</sub>C<sup>t</sup>Bu)]<sub>2</sub>O}<sub>2</sub> (R = Me, Et) (C. Vatsa, V.K. Jain, T. Kesavadas, E.R.T. Tiekkink) (408) 157
- Triorganyltin fluorides, solid-state <sup>119</sup>Sn NMR studies of coordination in (H. Bai, R.K. Harris, H. Reuter), (408) 167
- Dimethyltin dibenzoate, X-ray crystal structure of (E.R.T. Tiekkink), (408) 323
- (Ph<sub>3</sub>Sn)<sub>2</sub>Yb(THF)<sub>4</sub>, a compound with an ytterbium–tin bond; synthesis and structure of (M.N. Bochkarev, V.V. Khramenkov, Yu.F. Rad'kov, L.N. Zakharov, Yu.T. Struchkov), (408) 329
- Group 14 metals, NQR studies of intramolecular complexes in organic compounds of (V.P. Feshin, G.A. Polygalova), (409) 1
- Allyl- and benzyl-type organoalkali intermediates, preparation via organotin compounds (O. Desponds, M. Schlosser), (409) 93
- Organotin esters of dithiocarbamylacetic acids (S.W. Ng, V.G. Kumar Das), (409) 143
- [2-(4,4-Dimethyl-2-oxazoline)-5-methylphenyl]methylphenyltin bromide, synthesis and structure of a triorganotin halide having a configurationally stable chiral tin center (J.T.B.H. Jastrzebski, E. Wehman, J. Boersma, G. van Koten, K. Goubitz, D. Heijdenrijk), (409) 157
- Triorganylstannyl diazo acetates, reactions with manganese pentacarbonyl bromide (H. Kandler, H.W. Bosch, V. Shklover, H. Berke), (409) 233
- Diorgano(chloro)tin(IV) oxinates and thioxinates, multinuclear NMR study of (A. Lyčka, J. Holeček, A. Sebald, I. Tkáč), (409) 331
- Group 14 triflates, synthesis of (W. Uhlig), (409) 377
- Rigid rod, iron-metal containing polyyne polymers, synthesis of (B.F.G. Johnson, A.K. Kakkar, M.S. Khan, J. Lewis), (409) C12
- {1-Methyl-1-aza-5-stannabicyclo[3.3.0]<sup>1,5</sup>octane}sulfide dimers and trimers; crystal structure of the dimer; study of dimer–trimer equilibrium by <sup>119</sup>Sn-NMR and FD mass spectrometry (B.M. Schmidt, M. Dräger, K. Jurkschat), (410) 43
- Diorganotin(IV) carboxylates containing 2-thiophene- or 2-furan-carboxylic acid, synthesis and characterisation of; crystal structure of [Et<sub>2</sub>Sn(O<sub>2</sub>CC<sub>4</sub>H<sub>3</sub>S)<sub>2</sub>] (C. Vatsa, V.K. Jain, T. Kesavadas, E.R.T. Tiekkink), (410) 135
- Diphenylead(IV) dichloride, molecular electronic structure of; comparison with diphenyltin(IV) dichloride by the SCF-MS molecular-orbital method (E.M. Berksoy, M.A. Whitehead), (410) 293

- Bis[bis(trimethylsilyl)methyl]tin(II), examination of the reaction with calcium; molecular and crystal structure of benzyl-tris[bis(trimethylsilyl)methyl]stannane (M. Westerhausen, T. Hildenbrand), (411) 1
- Diorganotin(IV) complexes of picolinic acid and picolinic acid *N*-oxide, preparation and tin-119m Mössbauer spectra (G.K. Sandhu, N.S. Boparoy), (411) 89
- 2-Chloro-2-n-butyl-1,3-dithia-2-stannacyclopentane, synthesis of; crystal structure of its 1,10-phenanthroline complex (G. Bandoli, U. Casellato, V. Peruzzo, G. Plazzogna), (411) 99
- 2,3-Bis(trimethylstannyl)-1-alkene moiety, Lewis acid-catalysed reaction with aldehydes (T.N. Mitchell, U. Schneider, K. Heesche-Wagner), (411) 107
- Dicyclohexylammonium tri-n-butylin 2-sulfobenzoate,  $[(c\text{-C}_6\text{H}_{11})_2\text{NH}_2]^+\text{[}^n\text{Bu}_3\text{Sn(O}_2\text{CC}_6\text{H}_4\text{-2-SO}_3\text{)}]$ , crystal structure of (S.W. Ng, V.G. Kumar Das, E.R.T. Tiekkink), (411) 121
- $[(\text{Me}_3\text{Sn}^{IV})_4\text{Fe}^{II}(\text{CN})_6 \cdot 4\text{H}_2\text{O}]_x$ , an organotin Berlin blue analogue with coordinative O-H ··· N hydrogen bonds (U. Behrens, A.K. Brimah, R.D. Fischer), (411) 325
- $[\text{O}(\text{Sn}(\text{C}_6\text{H}_3\text{Et}_2\text{-2,6})_2)_3]$ , synthesis of a four-membered ring compound containing three tin atoms and one oxygen; crystal structure of (C.J. Cardin, D.J. Cardin, M.A. Convery, M.M. Devereux), (411) C3
- 3-Ureidopropionic acid, organotin esters of; crystal structure of triphenyltin(IV) 3-ureidopropionate,  $(\text{C}_6\text{H}_5)_3\text{SnOCO}(\text{CH}_2)_2\text{NHCONH}_2$  (K.-M. Lo, V.G. Kumar Das, W.-H. Yip, T.C.W. Mak), (412) 21
- Monochloroacetyl-L-phenylalanine, diorganotin(IV) complexes of; crystal structure of  $[^n\text{Bu}_2\text{Sn(O}_2\text{C-(CH}_2\text{Ph)C(H)N(H)C(O)CH}_2\text{Cl}_2]$  (G.K. Sandhu, R. Hundal, E.R.T. Tiekkink), (412) 31
- Tetrabutylbis(3-benzoylpropionato)distannoxane dimer,  $[(\text{C}_4\text{H}_9)_2\text{SnO}_2\text{CCH}_2\text{CH}_2\text{C(O)C}_6\text{H}_5]_2\text{O}]_2$ , crystal structure of (S.W. Ng, C. Wei, V.G. Kumar Das), (412) 39
- Organotin(IV) compounds 2-(Me<sub>2</sub>NCHR)C<sub>6</sub>H<sub>4</sub>(SnR'R'Br), fixed *s-cis* arrangement of the chiral carbon and tin centers; crystal structure of 2-(Me<sub>2</sub>NCHBu<sup>1</sup>)C<sub>6</sub>H<sub>4</sub>SnMePhBr (J.T.B.H. Jastrzebski, J. Boersma, G. van Koten), (413) 43
- Distannane with very long Sn-Sn bond distance and synperiplanar conformation (M. Weidenbruch, J. Schlaefke, K. Peters, H.G. von Schnering), (414) 319
- $[\text{Bu}_4\text{N}][\text{Me}_2\text{SnCl(DMIT)}]$ , crystal structure of (S.M.S.V. Doidge-Harrison, R.A. Howie, J.T.S. Irvine, G. Spencer, J.L. Wardell), (414) C5
- Bromoheptakis(2,6-diethylphenyl)cyclotetrasiloxane; synthesis, structure and reactivity of (C.J. Cardin, D.J. Cardin, M.A. Convery, M.M. Devereux, N.B. Kelly), (414) C9
- Alkylidene dinitrosyl molybdenum complexes; synthesis and characterization of ethylidene complexes of the type  $[(\text{Et}_3\text{Sn}\cdot\text{AlCl}_3)_2\text{Mo}(\text{NO})_2(=\text{CHMe})]$  (A. Keller), (415) 97
- Bis(*o*-, *m*-, and *p*-aminobenzoate-*o*,*o'*) di-n-butyltin(IV) compounds; structures in the solid state and in solution (S.P. Narula, S.K. Bharadwaj, H.K. Sharma, Y. Sharda, G. Mairesse), (415) 203
- $[1,2\text{-Benzenedithiolate-(2-)S,S'}]_2$ -bis(methyl-1,3-dithia-benzo[c]-2-stannolanyl)[(CH<sub>3</sub>Sn)<sub>2</sub>(C<sub>6</sub>H<sub>4</sub>S<sub>2</sub>)<sub>3</sub>], synthesis and crystal structure of (V. Peruzzo, G. Plazzogna, G. Bandoli), (415) 335
- Stannacyclobutene, derivative from a sterically congested chloro(mesityl)stannane (M. Weidenbruch, K. Schäfers, J. Schlaefke, K. Peters, H.G. von Schnering), (415) 343
- p*-Tol<sub>6</sub>Sn<sub>2</sub>, *p*-Tol<sub>6</sub>PbSn, and *p*-Tol<sub>6</sub>Pb<sub>2</sub>, hexa-*p*-tolylethane derivatives; synthesis, homoeotype structures, NMR couplings, and valence vibrations (C. Schneider, M. Dräger), (415) 349
- $\{(\mu\text{-OH})(\text{Me}_3\text{Sn})_2\}_2\{(\mu\text{-NC})_2\text{Ni}(\text{CN})_2\}_2^{2-}$ , a planar heterobimetallic macrocyclic anion; stabilization of the binuclear organotin(IV) cation  $[(\mu\text{-OH})(\text{Me}_3\text{Sn})_2]^+$  within (T.M. Soliman, S.E.H. Etaiw, G. Fendesak, R.D. Fischer), (415) C1
- Some silicon, germanium and tin acyloxyderivatives; <sup>13</sup>C, <sup>17</sup>O, <sup>29</sup>Si NMR study of (I. Zicmane, E. Liepiņš, L.M. Ignatovich, E. Lukevics), (417) 355
- Diorganotin(IV) bis(8-quinolinolate) and bis(8-quinolinethiolate) complexes, stereochemistry in solution studied by NOE-difference spectroscopy (I. Tkáč, J. Holeček, A. Lyčka), (418) 311
- Dibisyl(fluorenylidene)stannane, evidence of its formation (G. Anselme, C. Couret, J. Escudié, S. Richelme, J. Satgé), (418) 321
- Dihydroxytetraorganodistannoxanes, synthesis and characterization by IR and NMR (<sup>1</sup>H, <sup>13</sup>C and <sup>119</sup>Sn) spectroscopy; crystal structure of  $\{[^n\text{Bu}_2\text{Sn(OC}_6\text{H}_4\text{OMe-2)}]_2\text{O}\}_2$  (C. Vatsa, V.K. Jain, T.K. Das, E.R.T. Tiekkink), (418) 329
- Heterodinuclear group 6 metallacarbonanes; synthesis and characterization of [*cis*-3,3,3-(CO)<sub>3</sub>-3-SnPh<sub>3</sub>-3,1,2-MC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>+</sup> (M = Cr, Mo, W) (J. Kim, Y. Do, Y.S. Sohn, C.B. Knobler, M.F. Hawthorne), (418) C1
- Me<sub>2</sub>SnCl<sub>2</sub>·(2-MeOC<sub>6</sub>H<sub>4</sub>N=CHC<sub>5</sub>H<sub>4</sub>N-2), synthesis and crystal structure of (T.K. Chattopadhyay, A.K. Kumar, A. Roy, A.S. Batsanov, E.B. Shamuratov and Yu.T. Struchkov), (419) 277

- On carbon-13 spectra of thio- and dithiocarboxylic acids triorgano group 14 metal esters (S. Kato, A. Hori, M. Mizuta, T. Katada, H. Ishihara, K. Fujieda and Y. Ikebe), (420) 13
- Preparation and  $^{119m}\text{Sn}$  Mössbauer studies of diorganotin(IV) derivatives of nicotinic acid and nicotinic acid *N*-oxide (G.K. Sandhu and N.S. Boparoy), (420) 23
- "Head-to-head" dimerization and dehydrodimerization of vinyl ketones catalyzed by modified rhodium(I) complexes (I.P. Kovalev, Y.N. Kolmogorov, Y.A. Strelenko, A.V. Ignatenko, M.G. Vinogradov and G.I. Nikishin), (420) 125
- Organozinnverbindungen. XXXIV. 2-(Chlordimethylstannyl)-4-methoxy-2',4'-dinitroazobenzol: Ein ungewöhnliches Produkt bei Azokupplungen an 3-Trimethylstannylanisol. Intramolekulare Komplexierung der Azogruppe durch die Stannylgruppe unter deren Halodemethylierung (W.P. Neumann and C. Wicenec), (420) 171
- Crystal structure of bis(*O*-isopropylidithiocarbonato)diphenyltin(IV) (N. Donoghue and E.R.T. Tiekkink), (420) 179
- Synthèse et réactivité d'un nouvel hydrure organostannique à "queue polaire": le dibutyl-4,7,10-trioxaundécylstannane (F. Ferkous, D. Messadi, B. De Jeso, M. Degueil-Castaing and B. Maillard), (420) 315
- Some new organotin(IV), aluminium(III)- $\mu$ -oxoisopropoxides and their benzoylacetone derivatives (P.N. Kapoor, A.K. Bhagi, R.N. Kapoor and H.K. Sharma), (420) 321
- Structural chemistry of organotin carboxylates. XVI. Structural variation in dicarboxylato tetraorganostannoxanes: Crystal structure of  $[(\text{Me}_2\text{Sn}(\text{O}_2\text{CC}_4\text{H}_3\text{S})_2\text{O})_2$  (C. Vatsa, V.K. Jain, T.K. Das and E.R.T. Tiekkink), (421) 21
- Interaction of naphthaleneytterbium with tetraphenyltin. Molecular structure of  $\text{Ph}_3\text{SnYb}(\text{THF})_2(\mu\text{-Ph})_3\text{Yb}(\text{THF})_3$  (M.N. Bochkarev, V.V. Khramenkov, Y.F. Rad'kov, L.N. Zakharov and Yu.T. Struchkov), (421) 29
- Synthese von Phenylbrom-, Phenyliod- und Phenylwasserstoff-trisilanen (K. Hassler und U. Katzenbeisser), (421) 151
- Trimethylsilyl-substituierte Derivate des Dimethylzinks — Synthese, spektroskopische Charakterisierung und Struktur (M. Westerhausen, B. Rademacher und W. Poll), (421) 175
- Zum gezielten Aufbau oligomerer Strukturen der Elemente der 14. Gruppe mittels ihrer Triflatderivate (W. Uhlig), (421) 189
- A simplified, high-yield synthesis, and crystal structure of [tris[bis(2,4,6-triisopropylphenyl)tin]] (F.J. Brady, C.J. Cardin, D.J. Cardin, M.A. Convery, M.M. Devereux and G.A. Lawless), (421) 199
- Reactivity of Main-Group-transition-metal bonds. IX. The kinetics of iodination of compounds containing two or more tin-transition-metal bonds (J.R. Chipperfield, S. Clark, D.E. Webster and H. Yusof), (421) 205
- Pyrrolyl compounds of main group elements. Synthesis of group 14 5-metallated pyrrole-2-carbaldehydes (F. Denat, H. Gaspard-Ioughmane and J. Dubac), (423) 173
- Preparation and  $^{119m}\text{Sn}$  Mössbauer spectral studies of diorganotin(IV) derivatives of isonicotinic acid and isonicotinic acid *N*-oxide (G.K. Sandhu and N.S. Boparoy), (423) 183
- Darstellung und Struktur von  $(\text{EtO})_2\text{P}(\text{O})\text{CH}_2\text{Si}(\text{Me})_2\text{CH}_2\text{SnMe}_2\text{Cl}$ —ein Sechsringchelat mit Sesquikonformation und  $\text{P}=\text{O} \cdots \text{Sn}(\text{Cl})\text{Me}_2\text{CH}_2$ -trigonaler Bipyramide am Lewis-aciden Zinn (U. Kolb, M. Dräger, E. Fischer und K. Jurkschat), (423) 339
- Organostannyl mediated synthesis of 1-alkyl- and 1-sulfonyl-2-trifluoromethylbenzimidazole derivatives (N. Narayanan and T.R. Balasubramanian), (423) 361
- Isolierung und Röntgenstrukturanalyse des *fac*- und *mer*-Isomeren des  $^{\text{i}}\text{PrSnCl}_3 \cdot 2\text{DMF}$  (H. Reuter und H. Puff), (424) 23
- Triphenylstannyl derivatives of isothiazol-3(2*H*)-one 1,1-dioxides. Crystal structures of 2-triphenylstannyl 1,2-benzisothiazol-3(2*H*)-one 1,1-dioxide  $[(\text{C}_6\text{H}_5)_3\text{SnNC(O)C}_6\text{H}_4\text{SO}_2]$  and 2-triphenylstannyl 4,5-dimethylisothiazol-3(2*H*)-one 1,1-dioxide  $[(\text{C}_6\text{H}_5)_3\text{SnNC(O)C(CH}_3\text{)C(CH}_3\text{)SO}_2]$  (S.W. Ng, V.G. Kumar Das, Z.-Y. Zhou and T.C.W. Mak), (424) 133
- Sulphur compounds. CXLV. Synthesis and properties of  $\text{Cp}_2\text{TiS}_2\text{SiR}_2$ ,  $\text{Cp}_2\text{TiS}_2\text{GeR}_2$ , and  $\text{Cp}'_2\text{TiS}_2\text{SnR}_2$  ( $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ ,  $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{CH}_3$ , and  $\text{R} = \text{C}_6\text{H}_5$ ) (J. Albertsen and R. Steudel), (424) 153
- Synthesis, characterisation and halogen-cleavage reactions of  $\text{Ph}_3\text{SnCH}_2\text{SR}$  ( $\text{R} = \text{benzothiazole, benzoxazole, 1-methylimidazole, pyrimidine}$ ) (K.C. Molloy and P.C. Waterfield), (424) 281
- Carbonyl olefination with  $\alpha$ -stannyl ester enolates: a new synthesis of  $\alpha,\beta$ -unsaturated esters (A. Zapata, B.M. Núñez and F.J. Ferrer), (424) C9

Zur Kenntnis funktionalisierter Zinn-Phosphorverbindungen: Darstellung und Reaktionen von Chlorstannylphosphinen (D. Hänssgen, E. Stahlhut, H. Aldenhoven and A. Dörr), (425) 19  
 Synthesis and characterization of five-coordinate platinum(II) complexes [Pt(2,9-dimethyl-1,10-phenanthroline)(SnR<sub>n</sub>X<sub>3-n</sub>)X(olefin)] (X = Cl or Br; R = Ph or Me). Molecular structure of [Pt(2,9-dimethyl-1,10-phenanthroline)(SnPh<sub>2</sub>Cl)Cl(ethylene)] (V.G. Albano, C. Castellari, V. De Felice, A. Panunzi and F. Ruffo), (425) 177

## Titanium

- Titanocene-catalysed hydromagnesation of silylacetylenes, reversibility of (M.A. Djadchenko, K.K. Pivnitsky, J. Spanig, H. Schick), (401) 1  
*ansa*-Titanocene and zirconocene complexes with asymmetrically substituted carbon bridges (G.S. Herrmann, H.G. Alt, M.D. Rausch), (401) C5  
 2,4-Di(*t*-butyl)pentadienyl anion and its Ti(II), Cr(II), and Zn(II) complexes; synthesis of (R.D. Ernst, J.W. Freeman, P.N. Sweenston, D.R. Wilson), (402) 17  
 [(C<sub>5</sub>HMe<sub>4</sub>)TiBr(μ-O)]<sub>4</sub> and [(C<sub>5</sub>Me<sub>5</sub>)TiBr(μ-O)]<sub>3</sub>, crystal and molecular structure of (S.I. Troyanov, V. Varga, K. Mach), (402) 201  
 Cationic titanocene bishexafluoroarsenate complexes (P. Gowik, T. Klapötke), (402) 349  
 [(η<sup>5</sup>-(Dimethylphosphino)cyclopentadienyl]·[η<sup>7</sup>-(dimethylphosphino)cycloheptatrienyl]titanium, heterobimetallic compounds derived from (M. Ogasa, M.D. Rausch, R.D. Rogers), (403) 279  
 Low-valent organotitanium species; their applications to organic synthesis (Y. Dang, H.J. Geise), (405) 1  
 Titanium, iron and thallium vinylcyclopentadienyl derivatives, synthesis and structure of (M. Ogasa, D.T. Mallin, D.W. Macomber, M.D. Rausch, R.D. Rogers, A.N. Rollins), (405) 41  
 Bis(η<sup>5</sup>-cyclopentadienyl)methyl(alkoxy)- and bis(η<sup>5</sup>-cyclopentadienyl)methyl(acyloxy)titanium(IV) complexes, synthesis and reactions of (R. Schobert), (405) 201  
 Stable titanium-cobalt complex with non-bridged Ti-Co bond, synthesis and characterisation of (D. Selent, R. Beckhaus, T. Bartik), (405) C15  
 [(C<sub>5</sub>Me<sub>4</sub>R)TiAs<sub>3</sub>O<sub>6</sub>], a molecule with an AB<sub>3</sub>X<sub>6</sub> adamantine skeleton (B. Bettinger, O.J. Scherer, G. Heckmann), (405) C19  
 Titanocene polyselenide complexes, reactions with methyldichloroarsane; synthesis and properties of (CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Ti(μ<sub>2</sub>-Se<sub>2</sub>)<sub>2</sub>AsCH<sub>3</sub> and of (CH<sub>3</sub>As)<sub>2</sub>Se<sub>3</sub> (B. Holz, R. Steudel), (406) 133  
 Bisalkynyltitanocene complexes as bidentate chelating ligands (H. Lang, L. Zsolnai), (406) C5  
 Allyltitanium complexes, steric effect of allyl substituent on the molecular structures of (J. Chen, Y. Kai, N. Kasai, H. Yasuda, H. Yamamoto, A. Nakamura), (407) 191  
 Etheno-bridged *ansa*-titanocene derivatives, syntheses by the McMurry reaction (P. Burger, H.H. Brintzinger), (407) 207  
 Homogeneous metal complex catalysts in the synthesis of cyclobutane and cyclopentane compounds (U.M. Dzhemilev, R.I. Khusnutdinov, G.A. Tolstikov), (409) 15  
 Bis(alkinyl) titanocene complexes of FeCl<sub>2</sub> and CuCl (H. Lang, M. Herres, L. Zsolnai, W. Imhof), (409) C7  
 Di-μ-chloro-tetrachlorobis(diethyl *o*-phthalate)dititanium(III), a new intermediate for the synthesis of titanium(III) compounds; crystal structure and properties of (P. Sobota, J. Ejfler, J. Utko, T. Lis), (410) 149  
 (μ-Alkylideneamido)titanocene complex [(C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>TiCl]<sub>2</sub>[μ-{N=C(CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>)C(CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>)=N}], preparation and crystal structure of (F. Rehbaum, K.-H. Thiele, S.I. Troyanov), (410) 327  
 Disilanes, dehydrogenative polymerisation catalysed by titanocene and zirconocene complexes; novel route to oligo- and poly-silanes (E. Hengge, M. Weinberger, Ch. Jammegg), (410) C1  
 Titanium(IV) complexes containing highly phenylated π-cyclopentadienyl ligands; structure of (π-C<sub>5</sub>H<sub>5</sub>)(π-C<sub>5</sub>Ph<sub>5</sub>)TiCl<sub>2</sub> (U. Thewalt, G. Schmid), (412) 343  
 Methyl- and ethylrhodium oxides, synthesis and structural aspects (W.A. Herrmann, C.C. Romao, P. Kiprof, J. Behm, M.R. Cook, M. Taillefer), (413) 11  
 Titanium and zirconium half-sandwich η-cycloheptatrienyl and η-toluene derivatives, one-pot synthesis of; crystal structure of [Zr(η<sup>6</sup>-C<sub>6</sub>H<sub>5</sub>Me)(PMe<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>] (G.M. Diamond, M.L.H. Green, N.M. Walker), (413) C1  
 (η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)<sub>2</sub>(Me)Ti-OC(Me)=CH<sub>2</sub>: titanium enolate or titanium alkoxide? (C.P. Gibson, D.S. Bem), (414) 23

- 1,1-Dibromo-2,3,4,5,6,7-hexaphenyltitanacycloheptatriene, preparation and molecular structure of (K. Mach, S.I. Troyanov), (414) C15
- 3,3'-Di-tert-butyltitanocene dichloride, tetramethylethanobridged; stereoselectivity in the synthesis of (M.S. Erickson, F.R. Fronczeck, M.L. McLaughlin), (415) 75
- Titanium complexes with fulvene and dimethylenecyclopentenyl ligands; formation from bis( $\eta^5$ -tetramethylcyclopentadienyl)titanium(IV) precursors (K. Mach, V. Varga, V. Hanus, P. Sedmera), (415) 87
- Aryl-substituted titanocene and zirconocene complexes with interannular tetramethylethano- or dimethylsilano-bridges; crystal structures of representative examples (P. Burger, K. Hortmann, J. Diebold, H.-H. Brintzinger), (417) 9
- Hafnium and titanium catalysts, comparative behaviour in copolymerization of  $\alpha$ -olefins (A. Altomare, R. Solaro, F. Ciardelli, L. Barazzoni, F. Menconi, S. Malquori, F. Masi), (417) 29
- Di- $\mu$ -[bis(2-phenylethyl) m-phthalate]octachlorodititanium(IV) bis(dichloromethane), crystal structure and properties (P. Sobota, J. Utko, T. Lis), (417) 389
- Group 4 1,1'-bis(trimethylsilyl)metallocenes, binuclear benzyl-1,2,4,5-tetrathiolato complexes of; synthesis and structure of [ $(\eta^5\text{-Me}_3\text{SiC}_5\text{H}_4)_2\text{TiS}_2\text{J}_2\text{TiS}_2\text{C}_6\text{H}_2$  and  $(\eta^5\text{-Me}_3\text{SiC}_5\text{H}_4)_2\text{HfS}_2\text{J}_2\text{C}_6\text{H}_2$ ] (H. Balz, H. Köpf, J. Pickardt), (417) 397
- $\text{Cp}_2\text{Ti}(\mu_2\text{-}\mu_1\text{-SO}_4)\text{Ti}(\text{H}_2\text{O})\text{Cp}_2$ , synthesis and structure of (U. Thewalt, P. Schinnerling), (418) 191
- Monopentamethylcyclopentadienyltitanium(IV) halo-alkoxides, alkyl-alkoxides and acetylacetones (P. Gómez-Sal, A. Martín, M. Mena, P. Royo and R. Serrano), (419) 77
- Darstellung und Struktur von  $\text{Cp}_2\text{Ti}(\text{SO}_3\text{CF}_3)_2(\text{THF})$  (K. Berhalter and U. Thewalt), (420) 53
- Oxidative alkylation of  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiR}$  ( $\text{R} = \text{Cl}, \text{Me}, \text{Et}, \text{CH}=\text{CH}_2, \text{Ph}, \text{OMe}, \text{N}=\text{C}(\text{H})^{\text{i}}\text{Bu}$ ) to  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}(\text{Me})\text{R}$  by group 12 organometallic compounds  $\text{MMe}_2$  (G.A. Luinstra and J.H. Teuben), (420) 337
- [ $\text{Ph}_4\text{P}][\text{CpTi}(1,2\text{-S}_2\text{C}_6\text{H}_4)_2]$ : Synthese und Molekülstruktur eines anionischen, tetragonal-pyramidal koordinierten Benzol-1,2-dithiolato-Komplexes des Titans (H. Köpf, K. Lange and J. Pickardt), (420) 345
- Preparation of metallacyclic titanocene hydrocarbyl complexes and their use in propene polymerization reactions (G. Erker, U. Korek, R. Petrenz and A.L. Rheingold), (421) 215
- $\text{Ti}^{\text{IV}}$  and  $\text{Mo}^{\text{V}}$  complexes with diesters. The crystal structure of [ $\text{o-C}_6\text{H}_4(\text{CO}_2\text{CH}_2\text{CH}_2\text{Ph})_2\text{Cl}_4\text{Ti}$ ], [ $\text{C}_2\text{O}_4(\text{CH}_2\text{CH}_2\text{Ph})_2\text{Cl}_4\text{Ti}$ ] and [ $\text{o-C}_6\text{H}_4(\text{CO}_2\text{Et})_2\text{Cl}_3\text{MoO}\cdot\text{C}_6\text{H}_6$ ] (P. Sobota, S. Szafert, J. Utko and T. Lis), (423) 195
- Sulphur compounds. CXLV. Synthesis and properties of  $\text{Cp}_2\text{TiS}_2\text{SiR}_2$ ,  $\text{Cp}_2\text{TiS}_2\text{GeR}_2$ , and  $\text{Cp}'_2\text{TiS}_2\text{SnR}_2$  ( $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$ ,  $\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{CH}_3$ , and  $\text{R} = \text{C}_6\text{H}_5$ ) (J. Albertsen and R. Steudel), (424) 153
- Synthese von  $(\eta^5\text{-Cyclopentadienyl})\text{dichloro-s-alkoxititan(IV)-Komplexen}$ . Eine Bestandsaufnahme der Reaktion von Titanocendichlorid mit Alkoholen (U. Höhlein und R. Schobert), (424) 301
- Synthesis, characterizations, and structures of group IV metal-chromium complexes bridged by the benzyloxide group (H.-M. Gau, C.-T. Chen and C.-C. Schei), (424) 307
- Cationic titanium alkyls as alkene polymerisation catalysts: solvent and anion dependence (M. Bochmann and A.J. Jaggar), (424) C5
- Frontier occupied orbitals in methyl-substituted fulvene and dimethylenecyclopentenyltitanium complexes by UV-photoelectron spectroscopy and EHT calculations (T. Vondrák, K. Mach, V. Varga and A. Terpstra), (425) 27
- Carbonylation of alkyltitanium compounds with allylic derivatives (Y. Tsuji and T. Ishii), (425) 41

## Transition Metals general

- Transition metals in organic synthesis; hydroformylation, reduction, and oxidation. Annual Survey covering the year 1989 (L. Markó), (404) 325
- Organic reactions of selected  $\pi$ -complexes; Annual Survey covering the year 1989 (G. Marr, B.W. Rockett), (416) 399
- Transition metals in organic synthesis; Annual Survey covering the year 1990 (L.S. Hegedus), (422) 301
- Organometallics of groups 4 and 5: oxidation states II and lower (F. Calderazzo and G. Pampaloni), (423) 307

## Tungsten

- Metal carbonyls, rate of O-atom transfer to; effect of substituents on amine *N*-oxides on (J.-K. Shen, Y.-C. Gao, Q.-Z. Shi, F. Basolo), (401) 295
- Alkynylalkoxycarbene metal (Cr, W) complexes, conjugate addition of alcohols to (F. Camps, A. Llebaria, J.M. Moretó, S. Ricart, J.M. Viñas, J. Ros, R. Yáñez), (401) C17
- Electron-rich tungsten diethylaminocarbyne complexes, synthesis from  $\text{Cp}(\text{Br})_2(\text{CO})\text{W}\equiv\text{CNEt}_2$  (A.C. Filippou, W. Grünleitner, E.O. Fischer), (401) C37
- ( $\eta^4$ -Norbornadiene)tetracarbonylchromium and metal VIb hexacarbonyls, chemical shift anisotropy of carbon-13 nuclei in carbonyl groups of (A. Gryff-Keller, H. Krawczyk, P. Szczęciński), (402) 77
- Molybdenum(0) and tungsten(0) ( $\eta^4$ -1-oxa-1,3-diene) complexes; compounds with *R*-(+)-pulegone ligands (T. Schmidt, C. Krüger, P. Betz), (402) 97
- Molybdenum and tungsten cycloheptadienyl complexes, interconversion of  $\eta^5$  and  $\eta^3$  hapticities in (J.R. Hinchliffe, M.W. Whiteley), (402) C50
- $\eta^5$ -Cyclopentadienyl(dicarbonyl)[ $\eta^2$ -diethylamino(methylthio)carbene]tungsten, preparation by thiomethylation of  $\eta^5$ -cyclopentadienyl(dicarbonyl)diethylaminocarbintungsten (F.R. Kreißl, N. Ullrich, H. Keller), (403) C29
- Molybdenum(0) and tungsten(0) tricarbonylphosphine complexes with chelating uninegative (X,Y)-donor ligands (X, Y = S, N, O) (A.E. Sánchez-Peláez, M.F. Perpiñán), (405) 101
- Cationic mono(but-2-yne) bis(phosphine) complexes  $[\text{W}(\text{CO})_2(\text{S}_2\text{CX})(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$  (X =  $\text{NC}_4\text{H}_8$ ,  $\text{NMe}_2$ ,  $\text{NET}_2$ ,  $\text{N}(\text{CH}_2\text{Ph})_2$  or  $\text{OEt}$ ), synthesis and spectral properties of (P.K. Baker, K.R. Flower), (405) 299
- Manganese cyclopentadienyldicarbonyl complexes of 1,2,3-selenadiazoles and thiadiazoles; structural comparison of free and complexed 1,2,3-benzothiadiazole and 4-phenyl-1,2,3-thiadiazole (A.J. Mayr, B. Carrasco-Flores, F. Cervantes-Lee, K.H. Pannell, L. Párkányi, K. RaghuVeer), (405) 309
- Tricarbonyl- $\eta^5$ -cyclohexadienyl tungstate and tricarbonyl- $\eta^5$ -cyclohexadienylmethyltungsten, synthesis of (C.G. Kreiter, T. Hellmann), (405) C6
- Four-membered ring complex  $[\text{W}(\text{CO})_5\text{PPhC(OEt)}\equiv\text{CPhCHPh}]$ , insertion reactions of  $[\text{Pt}(\text{PPh}_3)_2]$  fragments into; crystal and molecular structures of  $[\text{Pt}(\text{dppe})\text{PPh}(\text{W}(\text{CO})_5)\text{C(OEt)}\equiv\text{CPhCHPh}]$  and  $[\text{Pt}(\text{PPh}_3)_2((\text{W}(\text{CO})_5)\text{PPh=C(OEt)}\text{CPh=CHPh})(\eta^2\text{-}(\text{P}=\text{C}))$  (F.A. Ajulu, S.S. Al-Juaid, D. Carmichael, P.B. Hitchcock, M.F. Meidine, J.F. Nixon, F. Mathey, N.H. Tran Huy), (406) C20
- Electron-rich isocyanide-substituted tungsten diethylaminocarbyne complexes with a pentamethylcyclopentadienyl ligand (A.C. Filippou, W. Grünleitner), (407) 61
- Silacycloprenes, tungsten-catalysed reactions of (J. Ohshita, M. Ishikawa), (407) 157
- Heterometallic *n,π*-complexes of transition metals based on mesitylacetonitrile (V.S. Kaganovich, M.I. Rybinskaya), (407) 215
- $[\text{CpM}(\text{CO})_3\text{PPh}_2]$  ( $\text{Cp} = \eta\text{-C}_5\text{H}_5$ , M = Mo, W) metallophosphines, reactivity towards alkynes; X-ray crystal structure of  $[\text{CpMo}(\text{CO})_2(\text{Ph}_2\text{PCH=C(Ph)}\text{CO})]$  (H. Adams, N.A. Bailey, A.N. Day, M.J. Morris, M.M. Harrison), (407) 247
- Chromium and tungsten thiocarbene complexes, preparation of; insertion and annulation reactions of (K.H. Dötz, V. Leue), (407) 337
- Unidentate and heterobimetallic 1,1'-bis(diphenylphosphino)ferrocene complexes, X-ray photoelectron spectroscopic differentiation of chemically distinct phosphorus environments in (H.S.O. Chan, T.S.A. Hor, L.-T. Phang, K.L. Tan), (407) 353
- $\text{M}^+[\text{M}'\text{M}''(\text{CO})_9\text{L}]^-$  ( $\text{M}^+ = \text{Na}^+$ ,  $\text{PPN}^+$ ; M' = Cr, W; M'' = Mn, Re; L = CO,  $\text{PR}_3$ ), kinetic studies of the cleavage of (Y.K. Park, S.J. Kim, J.H. Kim, I.S. Han, C.H. Lee, H.S. Choi), (408) 193
- Iron(II), tungsten(II) and molybdenum(II) ( $\eta^5$ -iodocyclopentadienyl)metallo methyl derivatives, methyl-iodine exchange in the presence of palladium (C. Lo Sterzo), (408) 253
- $[(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_3\text{M(HgX)}]$  (M = Cr, Mo, W; X = Cl, Br, I,  $\text{N}_3$ , SCN) complexes, interaction with M'( $\text{CO}$ )<sub>3</sub>(bipy) (M' = Mo, W) fragments; trimetallic compounds containing M-Hg-M'-X arrays (J. Granifo, M.E. Vargas), (408) 357
- Trimethylphosphine-substituted  $\eta^2$ -thiocarbene complexes of tungsten, preparation and reactions of (W. Schütt, N. Ullrich, F.R. Kreißl), (408) C5
- ( $\eta^2$ -TCNE)M( $\text{CO}$ )<sub>5</sub> (M = Cr; W; TCNE = tetracyanoethene), resonance Raman study of; evidence for an olefin-complex → metallacyclopropane transition upon low-energy metal-to-ligand charge-transfer excitation (D.J. Stufkens, T.L. Snoeck, W. Kaim, T. Roth, B. Olbrich-Deussner), (409) 189
- ( $\eta^5\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_{13}\text{H}_9$ )M( $\text{CO}$ )<sub>3</sub>Me complexes (M = Mo, W), photoinduced reactions with acetylenes (H.G. Alt, J.S. Han, H.E. Maisel), (409) 197

- [WI<sub>2</sub>(CO)(NCMe)(η<sup>2</sup>-RC<sub>2</sub>R)<sub>2</sub>] (R = Me or Ph) and [WI(CO)(NCMe){Ph<sub>2</sub>P(CH<sub>2</sub>)PPh<sub>2</sub>}](η<sup>2</sup>-MeC<sub>2</sub>Me)][BF<sub>4</sub>], acetonitrile exchange reactions with thiophene-3-acetonitrile; crystal structure of [WI<sub>2</sub>(CO)(NCCH<sub>2</sub>(3-C<sub>4</sub>H<sub>3</sub>S))(η<sup>2</sup>-MeC<sub>2</sub>Me)<sub>2</sub>] (P.K. Baker, M.G.B. Drew, S. Edge, S.D. Riyard), (409) 207
- Seven-coordinate molybdenum(II) and tungsten(II) organophosphine carbonyl iodide derivatives, NMR study of (P.K. Baker, M.A. Beckett, L.M. Severs), (409) 213
- Molybdenum and tungsten monoalkyne complexes, syntheses and NMR studies of (J.L. Davidson, F. Sence), (409) 219
- Tungsten carbonyl complexes derived from the dinitrogen complex *trans*-[W(N<sub>2</sub>)<sub>2</sub>(Ph<sub>2</sub>PCH<sub>2</sub>CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub> (T. Ishida, Y. Mizobe, T. Tanase, M. Hidai), (409) 355
- [(OC)<sub>3</sub>M(μ-Cl)(μ-CO)(μ-dppm)Rh(NBD)] (M = Mo, W) tribridged derivatives, formation by ring opening of [(OC)<sub>4</sub>M(dppm-PP)]; crystal structure of [(OC)<sub>3</sub>Mo(μ-Cl)(μ-CO)(μ-dppm)Rh(NBD)] (M. Cano, J.V. Heras, P. Ovejero, E. Pinilla, A. Monge), (410) 101
- Carbonylchromium, carbonylmolybdenum and carbonyltungsten complexes of the asymmetrically substituted phosphino sulfur diimides <sup>1</sup>Bu<sub>2</sub>P(NSN)PPh<sub>2</sub> and <sup>1</sup>Bu<sub>2</sub>P(NSN)P(<sup>1</sup>Bu)Ph; <sup>1</sup>H, <sup>13</sup>C and <sup>31</sup>P NMR spectroscopic characterisation of (M. Herberhold, S.M. Frank, B. Wrackmeyer), (410) 159
- Tungsten(0) anionic ethylisocyanide complex, alkylation of; synthesis of alkyl-, η<sup>2</sup>-iminoacyl-, 1-azaallyl- and dialkylaminocarbene complexes of tungsten (A.C. Filippou, W. Grünleitner, P. Kiprof), (410) 175
- CH<sub>3</sub>C[CH<sub>2</sub>NC-M(CO)<sub>x</sub>]<sub>3</sub> complexes (M = Cr, W, x = 5; M = Fe, x = 4), synthesis of; crystal structure of CH<sub>3</sub>C[CH<sub>2</sub>NC-W(CO)<sub>3</sub>]<sub>3</sub> (F.E. Hahn, M. Tamm), (410) C9
- CpMH=C(NHEt)Ph)(CO)<sub>2</sub> (M = Mo, W) system, hydride to carbene migration in; crystal structure of CpMo(η<sup>2</sup>-CH(Ph)NHEt)(CO)<sub>2</sub> (H. Adams, N.A. Bailey, M.J. Winter, S. Woodward), (410) C21
- Molybdenum(II) and tungsten(II) tricyclohexylphosphinecarbonylsulphide seven-coordinate complexes (P.K. Baker, D. ap Kendrick), (411) 215
- (CO)<sub>5-n</sub>(R'<sub>3</sub>P)<sub>n</sub>M(H)SiR<sub>3</sub> (n = 2,3), synthesis of phosphane substituted hydridosilyl complexes of chromium, molybdenum and tungsten; synthesis of (CO)<sub>3</sub>(L-LM(H)Cl by conversion of (CO)<sub>3</sub>(L-L)M(NCR') with HSiClR<sub>2</sub> (L-L = tmeda, bipy, (PPh<sub>3</sub>)<sub>2</sub>) (H. Piana, U. Schubert), (411) 303
- Tungsten(0)-diethylaminocomplex, stepwise preparation from ethylisocyanide precursors (A.C. Filippou, W. Grünleitner, E.O. Fischer), (411) C21
- Binuclear oxo complexes of molybdenum and tungsten with chalcogenes (O, S, Se, Te) as bridging ligands; synthesis and characterisation of (M. Gorzellik, H. Bock, L. Gang, B. Nuber, M.L. Ziegler), (412) 95
- Mechanism of the W(CO)<sub>6</sub>-mediated C-S bond cleavage reaction (C.-T. Ng, T.-Y. Luh), (412) 121
- Cp<sub>2</sub>MM'(Cp)(CO)<sub>3</sub> (M = Nb, Ta; M' = Mo, W), heteronuclear complexes with bridging and semi-bridging carbonyl groups; synthesis and dynamic behaviour (D. Perrey, J.C. Leblanc, C. Moise, J. Martin-Gil), (412) 363
- [M(η<sup>3</sup>-C(CF<sub>3</sub>)C(CF<sub>3</sub>)SR)(CF<sub>3</sub>≡CCF<sub>3</sub>)(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)], molybdenum and tungsten η<sup>2</sup>-C,C-vinyl complexes; reactions with alkynes leading to novel metallacycles (L. Carlton, N.M. Agh-Atabay, J.L. Davidson), (413) 205
- [NEt<sub>4</sub>][WFe<sub>2</sub>{μ<sub>3</sub>-η<sup>3</sup>,η<sup>5</sup>-CH=C(<sup>1</sup>Bu)C(O)C<sub>2</sub>B<sub>9</sub>H<sub>8</sub>Me<sub>2</sub>}(CO)<sub>8</sub>], a tungstendiiiron cluster derived from [Fe<sub>2</sub>(CO)<sub>9</sub>] and [NEt<sub>4</sub>][W(≡CC<sup>1</sup>Bu)(CO)<sub>2</sub>(η<sup>5</sup>-C<sub>2</sub>B<sub>9</sub>H<sub>9</sub>Me<sub>2</sub>)] (N. Carr, S.J. Dossett, F.G.A. Stone), (413) 223
- M-Hg-M (M = Cr, Mo, W) structural unit, unexpected metal–metal bond formation in; synthesis and characterisation of [η<sup>5</sup>-RC<sub>5</sub>H<sub>4</sub>(CO)<sub>3</sub>M]<sub>2</sub>Hg complexes (R = H, M = W; R = CH<sub>3</sub>CO, M = Cr, Mo, W) (L.-C. Song, H. Yang, Q. Dong, Q.-M. Hu), (414) 137
- W(CO)<sub>3</sub>(PMTA) (PMTA = MeN(CH<sub>2</sub>CH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub>) as a starting material for syntheses of W(CO)<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub>, W(CO)<sub>3</sub>(η<sup>6</sup>-arene), and protonated W(H)(CO)<sub>3</sub>(PR<sub>3</sub>)<sub>3</sub><sup>+</sup> complexes (V. Zanotti, V. Rutar, R.J. Angelici), (414) 177
- Bis(η<sup>5</sup>-cyclopentadienyl)dihydridotungsten, insertion reactions with activated olefins and with acetylenes; preparation of tungstenocene alkyl hydrides, olefin and acetylene complexes (G.E. Herberich, W. Barlage, K. Linn), (414) 193
- Complexed triphosphanes, preparation and characterisation of; crystal structure of HP[PPh<sub>2</sub>Cr(CO)<sub>5</sub>]<sub>2</sub> (M. Scheer, S. Gremler, E. Herrmann, P.G. Jones), (414) 337
- [W<sub>2</sub>(CO)<sub>10</sub>]<sup>2-</sup> and [Mo<sub>2</sub>(CO)<sub>8</sub>(μ-H)(μ-L)]<sup>-</sup> (L = bidentate phosphines) anions, their use in the synthesis of mercury-transition metal clusters (R. Reina, O. Rossell, M. Seco, A. Perales), (415) 101

- Pentacarbonyl(tellurobenzaldehyde)tungsten; synthesis, thermolysis and reactions with conjugated diened (H. Fischer, A. Fruh, C. Troll), (415) 211
- $\eta^2$ -iminoacyl and  $\eta^3$ -1-azaallyl complexes of tungsten, four- and five-membered metallaheterocycles; selective synthesis via C-C bond-forming reactions with alkyl isocyanides (A.C. Filippou, C. Völkl, P. Kiprof), (415) 375
- Tungsten and molybdenum alkylidyne complexes [ $M(=CR)(CO)_2(\eta\text{-}C_5H_5)$ ], reactions with the bidentate secondary phosphine  $C_6H_4(PH_2)_2$ ; crystal structure of the phospha-allyl complex [ $W(\sigma, \eta^3\text{-}C_6H_4(PH_2)\text{PC(OH)}CH(C_6H_4\text{Me-4})\text{-}1,2)(CO)(\eta\text{-}C_5H_5)$ ] (J.C. Jeffery, D.W.I. Sams, K.D.V. Weerasura), (418) 395
- Bis(cyclopentadienyl)(diphenylacetato)(2,2-diphenylethenyl)tungsten, unexpected product of the reaction of bis(cyclopentadienyl)dihydridotungsten and diphenylketene; synthesis and structure of (G.E. Herberich, K. Linn), (418) 409
- Heterodinuclear group 6 metallacarboranes; synthesis and characterization of [*cis*-3,3,3-(CO)<sub>3</sub>-SnPh<sub>3</sub>-3,1,2-MC<sub>2</sub>B<sub>9</sub>H<sub>11</sub>]<sup>-</sup> ( $M = Cr, Mo, W$ ) (J. Kim, Y. Do, Y.S. Sohn, C.B. Knobler, M.F. Hawthorne), (418) C1
- CpMH=C(OEt)Ph](CO)<sub>2</sub> ( $M = Mo, W$ ) system, hydride to carbene migration and subsequent rearrangement of the consequent products to  $\eta^3$ -benzyl complexes; crystal structure of the  $\eta^3$ -benzyl CpW(CO)<sub>2</sub>{ $\eta^3$ -C<sub>6</sub>H<sub>5</sub>CH(OEt)}
- (H. Adams, N.A. Bailey, M.J. Winter, S. Woodward), (418) C39
- Group VI metal carbonyls, ( $\eta^6$ -C<sub>6</sub>H<sub>6</sub>)Cr(CO)<sub>3</sub>, ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Mn(CO)<sub>3</sub>, ( $\eta^5$ -CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Mn(CO)<sub>3</sub> and ( $\eta^5$ -C<sub>5</sub>H<sub>5</sub>)Re(CO)<sub>3</sub>, photolysis in Nujol at 77 K (T.E. Bitterwolf, K.A. Lott, A.J. Rest and J. Mascetti), (419) 113
- Alkyne, cyclobutadiene and cyclopentadienone complexes of molybdenum and tungsten (J.L. Davidson), (419) 137
- W(CO)<sub>6</sub>, photochemical reaction with CCl<sub>4</sub>; matrix-isolation studies of intermediates in (T. Szymbańska-Buzar, A.J. Downs, T.M. Greene and M.A.K. Wood), (419) 151
- Reactivity of Main-Group-transition-metal bonds. IX. The kinetics of iodination of compounds containing two or more tin-transition-metal bonds (J.R. Chipperfield, S. Clark, D.E. Webster and H. Yusof), (421) 205
- Preparation and spectroscopic properties of a series of new bimetallic phosphine-bridged seven-coordinate complexes of the type [M<sub>2</sub>I<sub>4</sub>(CO)<sub>6</sub>L<sub>2</sub>( $\mu$ -L<sub>2</sub>)] { $M = Mo$  or  $W$ ; L = PPh<sub>3</sub>, AsPh<sub>3</sub> or SbPh<sub>3</sub>; L<sub>2</sub> = Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub> ( $n = 1, 2$  or 4); L<sub>2</sub> = [Fe( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>PPh<sub>2</sub>)<sub>2</sub>]} (P.K. Baker, M. van Kampen and D. ap Kendrick), (421) 241
- Spectroscopic properties of conjugated metal–carbon multiple bonds: synthesis and absorption spectra of the “dialkylidynes” (RO)<sub>3</sub>W≡C–C≡W(OR)<sub>3</sub> (OR = OCMe<sub>3</sub>, OCMe<sub>2</sub>CF<sub>3</sub>, OCMe<sub>2</sub>Et) (T.M. Gilbert and R.D. Rogers), (421) C1
- Preparation and properties of molybdenum and tungsten dinitrogen complexes. XXXIV. Nucleophilic addition reactions of hydrazido(2-) complexes with diphenylketene: characterization of new acylhydrazido(2-) complexes of molybdenum and tungsten (T. Aoshima, Y. Mizobe, M. Hidai and J. Tsuchiya), (423) 39
- Electrochemical generation of 17-electron cation radicals from arene–M(CO)<sub>3</sub> ( $M = Cr, Mo, W$ ) and thiophene–Cr(CO)<sub>3</sub> complexes in MeCN: Conventional cyclic voltammetric studies with digital simulation and microelectrode voltammetry in the absence of supporting electrolyte (F. Rourke, R. Gash and J.A. Crayton), (423) 223
- Rapid [SR]<sup>+</sup> exchange between metal-thiolate and metal-organic disulphide complexes (P.M. Treichel and E.K. Rublein), (423) 391
- Photochemische Reaktion von Pentacarbonyl(1-ethoxy-2-methyl-2-propenyliden)wolfram und Dekacarbonyldirhenium (C.G. Kreiter, S. Schufft und G. Heckmann), (424) 163
- Organometallic chemistry of diphosphazanes. IV. Reactions of the cyclodiphosphazane, *cis*-[<sup>1</sup>BuNP(OPh)]<sub>2</sub> with M(CO)<sub>6</sub> ( $M = Cr, Mo, W$ ), rhodium(I), palladium(II) and platinum(II) derivatives (M.S. Balakrishna and S.S. Krishnamurthy), (424) 243
- Bis( $\eta^5$ -cyclopentadienyl)tungsten chemistry: hydration reactions of functional alkyl hydrides and olefin complexes (G.E. Herberich and K. Linn), (425) C4

## Uranium

- [ $(Me_3SiC_5H_4)_3U_2[\mu\text{-O}]$ ] oxo-bridged bimetallic organouranium complex, synthesis and crystal structure of (J.-C. Berthet, J.-F. Le Maréchal, M. Nierlich, M. Lance, J. Vigner, M. Ephritikhine), (408) 335

- Actinide complexes with tripod ligands, synthesis and structure of  $[\text{CpCo}(\text{P}(\text{O})(\text{OEt})_2)_3]\text{UCl}_3(\text{THF})$  (M. Wedler, J.W. Gilje, M. Noltemeyer, F.T. Edelmann), (411) 271
- Stable biscyclopentadienyl uranium(IV) dichloride containing two  $\text{C}_5\text{H}_4$ -rings interlinked by a metal-coordinated 2,6-bis(methylene)pyridyl unit; synthesis, spectroscopy, and crystal structure of (G. Paolucci, R.D. Fischer, F. Benetollo, R. Seraglia, G. Bombieri), (412) 327
- Tri-tert-butyl methoxide uranium(IV) complexes, influence of electronic factors on the structure and stability of (C. Baudin, D. Baudry, M. Ephritikhine, M. Lance, A. Navaza, M. Nierlich, J. Vigner), (415) 59
- Uraniumindenyl compound, uranium(III)-tetrahydrofuran bond disruption enthalpy in (X. Jemine, J. Goffart, S. Bettomville, J. Fuger), (415) 363
- Tricyclopentadienyluranium azide complexes (J.-C. Berthet, M. Lance, M. Nierlich, J. Vigner and M. Ephritikhine), (420) C9

## Vanadium

- $\text{VO}(\text{OR})\text{Cl}_2$ -induced cyclisation of diketene via ring opening (T. Hirao, T. Fujii, Y. Ohshiro), (407) C1
- $\text{Cp}_2\text{VCl}(\text{AsF}_6)$  and other vanadocene(IV) hexafluoropnicogenate species, synthesis and EPR spectroscopic characterisation of; HF-catalysed oxidation to cationic vanadocenium(V) complexes; structure of  $[\text{Cp}_2\text{VCl}_2]^+[\text{AsF}_6]^-$  (F.H. Görlitz, P.K. Gowik, T.M. Klapötke, D. Wang, R. Meier, J. v. Welzen), (408) 343
- Naphthaleneytterbium, reactions with bis(cyclopentadienyl)complexes of cobalt, nickel, chromium and vanadium; crystal structure of the triple-decker compound  $\text{CpVC}_{10}\text{H}_8\text{VCp}$  (M.N. Bochkarev, I.L. Fedushkin, H. Schumann, J. Loebel), (410) 321
- $[\text{V}_2\text{Cp}_2(\text{CO})_4(\mu-\text{SMe})_2]$ , ligand substitution by isocyanides or trimethyl phosphite; synthesis and spectroscopic characterisation of the mono- and di-substituted derivatives; electrochemical behaviour of the parent complex (F.Y. Pétillon, P. Schollhammer, J. Talarmin), (411) 159
- Tris(cyclopentadienyl)technetium(III),  $(\eta^5\text{-C}_5\text{H}_5)_2\text{Tc}(\eta^1\text{-C}_5\text{H}_5)$ , X-ray diffraction study of; dipole moment and charge distribution of  $(\text{C}_5\text{H}_5)_3\text{M}$  ( $\text{M} = \text{V}, \text{Tc}, \text{Re}$ ) (C. Apostolidis, B. Kanellakopoulos, R. Maier, J. Rebantz, M.L. Ziegler), (411) 171
- Alkylvanadium complexes, synthesis and characterisation of; molecular structure of  $\text{trans-}[V(\text{CO})_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$  (D. Rehder, F. Süßmilch, W. Priebisch, M. Fornalczuk), (411) 357
- $[\text{V}(\eta^6\text{-1,3,5-Me}_3\text{C}_6\text{H}_3)_2]\text{AlCl}_4$ , primary product of the reduction of  $\text{VCl}_3$  by  $\text{Al}/\text{AlCl}_3/1,3\text{-Me}_3\text{C}_6\text{H}_3$ , crystal structure of; further reduction to  $\text{V}(\eta^6\text{-1,3,5-Me}_3\text{C}_6\text{H}_3)_2$  (F. Calderazzo, G. Pampaloni, L. Rocchi, F. Marchetti), (417) C16
- Binuclear pentamethylcyclopentadienylvanadium complexes with tellurium bridges; use of  $^{51}\text{V}$  NMR spectroscopy for identification of new vanadium complexes with different chalcogenes in the bridging system (M. Herberhold and M. Schrepfermann), (419) 85

## Ytterbium

- $[(\eta\text{-C}_5\text{H}_4\text{CH}_3)_2\text{Yb}^{\text{III}}(\text{OC}_4\text{H}_8)]_2(\mu_2\text{-O})$ , a binuclear lanthanoid complex with strictly linear Ln–O–Ln skeleton (M. Adam, G. Massarweh, R.D. Fischer), (405) C33
- Bis(t-butylcyclopentadienyl)lanthanoid chloride complexes, syntheses of; crystal structures of bis(t-butylcyclopentadienyl)chloro(bistetrahydrofuran)praseodymium and bis(t-butylcyclopentadienyl)chlorotetrahydrofuranytterbium (Q. Shen, M. Qi, J. Guan, Y. Lin), (406) 353
- Metal-bridging versus chelating  $\text{Me}_2\text{Si}(\text{C}_5\text{H}_4)_2$ -ligands in binuclear complexes; crystal structures of non-sublimed  $[\text{Me}_2\text{Si}(\text{C}_5\text{H}_4)_2\text{Yb}(\mu\text{-Br})]_2$  and  $[(\text{C}_5\text{H}_5)_2\text{Yb}(\mu\text{-Cl})]_2$  (T. Akhnoukh, J. Müller, K. Qiao, X.-F. Li, R.D. Fischer), (408) 47
- $(\text{Ph}_3\text{Sn})_2\text{Yb}(\text{THF})_4$ , a compound with an ytterbium–tin bond; synthesis and structure of (M.N. Bochkarev, V.V. Khramenkov, Yu.F. Rad'kov, L.N. Zakharov, Yu.T. Struchkov), (408) 329
- Naphthaleneytterbium, reactions with bis(cyclopentadienyl)complexes of cobalt, nickel, chromium and vanadium; crystal structure of the triple-decker compound  $\text{CpVC}_{10}\text{H}_8\text{VCp}$  (M.N. Bochkarev, I.L. Fedushkin, H. Schumann, J. Loebel), (410) 321
- Organolanthanide hydrides,  $[\text{O}(\text{CH}_2\text{CH}_2\text{C}_5\text{H}_4)_2\text{LnH}]_2$  ( $\text{Ln} = \text{Gd}, \text{Er}, \text{Yb}, \text{Lu}, \text{Y}$ ); synthesis and reactivity towards alkenes, alkynes and organic halides (Z. Xie, C. Qian, Y. Huang), (412) 61
- $(\eta^5\text{-C}_5\text{H}_5)_2\text{Yb}\cdot\text{DME}$ , synthesis and structure of (J. Jin, S. Jin, W. Chen), (412) 71
- Samarocene and ytterbocene hydroxides, synthesis and crystal structures of  $[(\text{SmCp}_2''(\mu\text{-OH}))_2]$  and  $[(\text{YbCp}_2(\mu\text{-OH}))_2][\text{Cp}'' = \eta^5\text{-C}_5\text{H}_3(\text{SiMe}_3)_2\text{-1,3}; \text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{SiMe}_3]$  (P.B. Hitchcock, M.F. LapPERT, S. Prashar), (413) 79

Alumohydride group, change of bonding mode in biscyclopentadienylhydrido REM complexes from heterometallic to homometallic; crystal and molecular structures of  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Yb}(\mu_3\text{-H})_2](\mu_2\text{-H})\text{AlH}_2\cdot\text{N}(\text{C}_2\text{H}_5)_3)_2\cdot\text{C}_6\text{H}_6$ ,  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Lu}(\mu_2\text{-H})_2](\mu_2\text{-H})_2\text{AlH}\cdot\text{N}(\text{C}_2\text{H}_5)_3)_2\cdot\text{C}_6\text{H}_6$  and  $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Lu}]_3(\mu_2\text{-H})_2(\mu_3\text{-H})$  (S.Ya. Knjazhanskij, B.M. Bulychev, O.K. Kireeva, V.K. Belsky, G.L. Soloveichik), (414) 11

Interaction of naphthalenenytterbium with tetraphenyltin. Molecular structure of  $\text{Ph}_3\text{SnYb}(\text{THF})_2(\mu\text{-Ph})_3\text{Yb}(\text{THF})_3$  (M.N. Bochkarev, V.V. Khramenkov, Y.F. Rad'kov, L.N. Zakharov and Yu.T. Struchkov), (421) 29

## **Yttrium**

Tricyclopentadienyllanthanides/sodium hydride, reaction with organic halides (C. Qian, D. Zhu, Y. Gu), (401) 23

Organic rare earth hydrides, synthesis and reactivity of (C. Ye, C. Qian, X. Yang), (407) 329

Dicyclopentadienyltytrium chloride, reaction with acyl chlorides in tetrahydrofuran; acylative cleavage of the Cp-Y π-bond and tetrahydrofuran ring (C. Qian, A. Qiu, Y. Huang, W. Chen), (412) 53

Organolanthanide hydrides,  $[\text{O}(\text{CH}_2\text{CH}_2\text{C}_5\text{H}_4)_2\text{LnH}]_2$  ( $\text{Ln} = \text{Gd}, \text{Er}, \text{Yb}, \text{Lu}, \text{Y}$ ); synthesis and reactivity towards alkenes, alkynes and organic halides (Z. Xie, C. Qian, Y. Huang), (412) 61

## **Zinc**

Nickel-catalysed electrochemical Reformatsky reaction with methyl chlorodifluoroacetate (S. Mcharek, S. Sibille, J.-Y. Nédélec, J. Périchon), (401) 211

2,4-Di(t-butyl)cyclopentadienyl anion and its Ti(II), Cr(II), and Zn(II) complexes; synthesis of (R.D. Ernst, J.W. Freeman, P.N. Swepston, D.R. Wilson), (402) 17

Allylic organozinc derivatives, condensation with nitriles; synthesis of β,γ-unsaturated ketones (P. Marceau, L. Gautreau, F. Béguin, G. Guillaumet), (403) 21

C-Nitroso compounds and their coordination complexes, NO bond length/stretching frequency relationships in (M. Cameron, B.G. Gowenlock, G. Vasapollo), (403) 325

$[(\text{HOMe}_2\text{Si})(\text{Me}_3\text{Si})_2\text{Cl}_2\text{Zn}$ ,  $[(\text{MeOMe}_2\text{Si})(\text{Me}_3\text{Si})_2\text{Cl}_2\text{Zn}$ , and  $[(\text{MeOMe}_2\text{Si})(\text{Me}_3\text{Si})_2\text{Cl}_2\text{Hg}$ , crystal structures of (F.I. Aigbirhio, S.S. Al-Juaid, C. Eaborn, A. Habtemariam, P.B. Hitchcock, J.D. Smith), (405) 149

Trialkylindium compounds, reaction with (cyclopentadienyl)(tricarbonyl)metal radicals (D.L. Thorn), (405) 161

Organozinc derivatives of functionalised allylic compounds, preparation and properties of (F. Lambert, B. Kirschleger, J. Villiéras), (405) 273

Organozinc derivatives of β-functionalised γ-substituted allylic compounds, reactivity of; preparation of β- and γ-substituted α-methylene γ-butyrolactones (F. Lambert, B. Kirschleger, J. Villiéras), (406) 71

Organolithium and -magnesium compounds, conjugated addition to α-ethylenic mesitylketones (F. Barbot, D. N'Goma, Ph. Miginiac), (410) 277

Methyl- and ethylrhodium oxides, synthesis and structural aspects (W.A. Herrmann, C.C. Romao, P. Kiprof, J. Behm, M.R. Cook, M. Taillefer), (413) 11

Synthèse en une étape par voie organométallique d'amines primaires β-acétyléniques α-ramifiées (L. Lebouret, G. Courtois and L. Miginiac), (420) 155

Alcynylsilanes fonctionnels. III. Synthèse régiosélective par voie organozincique de 1-triméthylsilyl-4-éne-1-yne fonctionnels (D. Mesnard and L. Miginiac), (420) 163

Oxidative alkylation of  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{TiR}$  ( $\text{R} = \text{Cl}, \text{Me}, \text{Et}, \text{CH}=\text{CH}_2, \text{Ph}, \text{OMe}, \text{N}=\text{C}(\text{H})^{\prime}\text{Bu}$ ) to  $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}(\text{Me})\text{R}$  by group 12 organometallic compounds  $\text{MMe}_2$  (G.A. Lijnstra and J.H. Teuben), (420) 337

Trimethylsilyl-substituierte Derivate des Dimethylzinks — Synthese, spektroskopische Charakterisierung und Struktur (M. Westerhausen, B. Rademacher und W. Poll), (421) 175

## **Zirconium**

ansa-Titanocene and zirconocene complexes with asymmetrically substituted carbon bridges (G.S. Herrmann, H.G. Alt, M.D. Rausch), (401) C5

Dinuclear ( $\eta^2$ -formaldehyde)zirconocene dimer t-butylisothiocyanate complexes, structure of (G. Erker, M. Mena, C. Krüger, R. Noe), (402) 67

- Hydrozirconation of (*E*)-3-methoxy-1-phenyl-1-propene and (*E*)-3-phenyl-2-propenol (S. Karlsson, A. Hallberg, S. Gronowitz), (403) 133
- Zirconium and hafnium hexakis(trimethylsilyl)metallocene dichlorides, hindered cyclopentadienyl rotation in; evidence for complete rotation of the tris(trimethylsilyl)cyclopentadienyl ligands (C.H. Winter, D.A. Dobbs, X.-X. Zhou), (403) 145
- Zirconocene dichloride/methylalumoxane systems, catalysts for olefin polymerization; IR and NMR studies of (L.A. Nekhaeva, G.N. Bondarenko, S.V. Rykov, A.I. Nekhaev, B.A. Krentsel, V.P. Mar'in, L.I. Vyshinskaya, I.M. Krapova, A.V. Polonskii, N.N. Korneev), (406) 139
- ( $\eta^2$ -Phenyltrimethylsilylacetylene)zirconocene generated *in situ*, reactions of (G. Erker, R. Zwettler), (409) 179
- Disilanes, dehydrogenative polymerisation catalysed by titanocene and zirconocene complexes; novel route to oligo- and poly-silanes (E. Hengge, M. Weinberger, Ch. Jammegg), (410) C1
- Heterobimetallic Zr,Hf( $\mu$ -formaldehyde) complex from [(CH<sub>2</sub>O)ZrCp<sub>2</sub>] dimer and hafnocene dichloride (G. Erker, M. Mena, M. Bendix), (410) C5
- Bis( $\eta^5$ -pentamethylcyclopentadienyl)zirconium(IV) dialkyls, autoxidations of (J.M. Atkinson, P.B. Brindley), (411) 131
- Titanium and zirconium half-sandwich  $\eta$ -cycloheptatrienyl and  $\eta$ -toluene derivatives, one-pot synthesis of; crystal structure of [Zr( $\eta^6$ -C<sub>6</sub>H<sub>5</sub>Me)(PMe<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>] (G.M. Diamond, M.L.H. Green, N.M. Walker), (413) C1
- Group 4 1,1'-bis(trimethylsilyl)metallocenes, binuclear benzyl-1,2,4,5-tetrathiolato complexes of; synthesis and structure of [( $\eta^5$ -Me<sub>3</sub>SiC<sub>5</sub>H<sub>4</sub>)<sub>2</sub>TiS<sub>2</sub>]<sub>2</sub>TiS<sub>2</sub>C<sub>6</sub>H<sub>2</sub> and [( $\eta^5$ -Me<sub>3</sub>SiC<sub>5</sub>H<sub>4</sub>)<sub>2</sub>HfS<sub>2</sub>]<sub>2</sub>C<sub>6</sub>H<sub>2</sub> (H. Balz, H. Köpf, J. Pickardt), (417) 397
- Homogeneous Cp<sub>2</sub>ZrCl<sub>2</sub>/methylaluminoxane Ziegler-Natta catalysts, reversible and irreversible deactivation of propene polymerization (D. Fischer, R. Mühlaupt), (417) C7
- Zirconocene complexes, olefin hydrosilation catalyzed by (M.R. Kesti, M. Abdulrahman, R.M. Waymouth), (417) C12
- A binuclear zirconium fulvalene tetraallyl complex, preparation and molecular structure of [ $\eta^5$ : $\eta^5$ -C<sub>10</sub>H<sub>8</sub>][CpZr( $\eta^1$ -CH<sub>2</sub>CHCH<sub>2</sub>) $\chi$ ( $\eta^3$ -CH<sub>2</sub>CHCH<sub>2</sub>)]<sup>2</sup> (Y. Wielstra, R. Duchateau, S. Gambarotta, C. Bensimon, E. Gabe), (418) 183
- Influence of the modification of the ligands on hex-1-ene hydroformylation catalyzed by [HRh(P(OPh)<sub>3</sub>)<sub>4</sub>] and [HRh(CO)(P(OPh)<sub>3</sub>)<sub>3</sub>]. Catalytic activity of the system [HRh(P(OPh)<sub>3</sub>)<sub>4</sub>]+Cp<sub>2</sub>Zr(CH<sub>2</sub>PPh<sub>2</sub>)<sub>2</sub> (A.M. Trzeciak, J.J. Ziolkowski and R. Choukroun), (420) 353
- Preparation and some properties of chiral *ansa*-mono( $\eta^5$ -fluorenyl)zirconium(IV) complexes (B. Rieger), (420) C17
- Molekulares Design von Übergangsmetallkomplexverbindungen zur Aktivierung kleiner Moleküle. II. Zur Carbonylierung von Bis( $\pi$ -pentamethylcyclopentadienyl)metall-1,3-Dien-Komplexen Cp<sup>\*</sup><sub>2</sub>M (1,3-Dien) (M = Zr, Hf)—Bildung einkerniger Metall-Endiolate (R. Beckhaus, D. Wilbrandt, S. Flatau und W.-H. Böhmer), (423) 211
- Synthesis, characterizations, and structures of group IV metal-chromium complexes bridged by the benzyloxide group (H.-M. Gau, C.-T. Chen and C.-C. Schei), (424) 307

## Book reviews

- Gmelin handbook of inorganic chemistry, 8th edition, index. Formula index, 2nd supplement (K.R. Seddon), (402) C37
- Dictionary of organometallic compounds. Cumulative structure index to volumes 1–5, edited by J.E. Macintyre (C. Eaborn), (402) C38
- Tin-based antitumour drugs, edited by M. Gielen (C. Eaborn), (402) C38
- A guide for the perplexed organic experimentalist, 2nd edition, by H.J. Loewenthal (C. Eaborn), (402) C63
- Advances in physical organic chemistry, vol. 25, edited by D. Bethell (C. Eaborn), (403) C32
- Gmelin handbook of inorganic chemistry, 8th edition, Sn. Organotin compounds. Part 17 (C. Eaborn), (403) C32
- The chemistry of the metal–carbon bond, volume 5, organometallic compounds in organic and biological syntheses, edited by F.R. Hartley (P.A. Chaloner), (403) C33
- Photocatalysis: fundamentals and applications, edited by N. Serpone and E. Pelizzetti (K.R. Seddon), (405) C11

- Gmelin handbook of inorganic chemistry, 8th edition, Pb. Organolead compounds, part 2 (C. Eaborn), (405) C12
- Nomenclature of organic chemistry: recommendations 1990, edited by G.J. Leigh (C. Eaborn), (405) C48
- Stereochemistry of organometallic and inorganic compounds, vol. 3: chemical bonds — better ways to make them and break them, edited by I. Bernal (P.A. Chaloner), (406) C9
- Gmelin handbook of inorganic chemistry, 8th edition, Ge—Organogermanium compounds, part 3 (C. Eaborn), (406) C10
- Gmelin handbook of inorganic chemistry, 8th edition, Fe—Organoiron compounds, part A9: ferrocene 9 (K.R. Seddon), (406) C11
- Inorganic reactions and methods, vol. 3. The formation of bonds to halogens (part 1), edited by A.P. Hagen (C. Eaborn), (406) C11
- Gmelin handbook of inorganic chemistry, 8th edition, W—Tungsten, supplement volume A3: physical properties (continued) (K.R. Seddon), (406) C12
- Atomic and molecular clusters, edited by E.R. Bernstein (A.J. Stace), (407) C5
- Transition metals in total synthesis, by P.J. Harrington (J.R. Hanson), (407) C6
- Gmelin handbook of inorganic chemistry, 8th edition, Mn—coordination compounds of manganese (G.J. Leigh), (407) C7
- Advances in sonochemistry, vol. 1, edited by T.J. Mason (P.D. Lickiss), (407) C16
- The chemistry of functional groups. The chemistry of enols, edited by Z. Rappoport (J.R. Hanson), (407) C16
- Steroids made it possible, by C. Djerassi (J.R. Hanson), (407) C34
- Directory of graduate research 1989 (C. Eaborn), (408) C22
- Gmelin handbook of inorganic chemistry, 8th edition, organotitanium compounds, part 5 (G.J. Leigh), (408) C22
- Inorganic synthesis vol. 26, editor-in-chief H.D. Kaesz (A.C. Sullivan), (409) C5
- Dictionary of chemistry and chemical technology: English—German, edited by H. Gross (J.D. Smith), (412) C27
- Reagents for transition metal complex and organometallic syntheses, inorganic syntheses, Vol. 28, edited by R.J. Angelici (J.D. Smith), (414) C33
- The chemistry of organophosphorus compounds vol. 1, primary, secondary and tertiary phosphines, polyphosphines and heterocyclic organophosphorus(III) compounds, edited by F.R. Hartley (J.D. Smith), (415) C6
- Sonochemistry: The Uses of Ultrasound in Chemistry, edited by T.J. Mason (P.D. Lickiss), (417) C66
- Enzymatic and Model Carboxylation and Reduction Reactions for Carbon Dioxide Utilization, edited by M. Aresta and T.V. Schloss (G.L. Leigh), (417) C67
- Biocoordination chemistry: coordination equilibria in biological systems, by K. Burger (G.J. Leigh) (418) C17
- Electroorganic synthesis; best synthetic methods, Tatsuya Shono (P.A. Chaloner), (418) C17
- Inorganic synthesis, vol. 27, editor-in-chief A.P. Ginsberg (A.C. Sullivan), (423) C29
- Advances in silicon chemistry, vol. 1, G.L. Larson (Ed.) (C. Eaborn), (423) C31
- Heteroatom chemistry, E. Block (Ed.) (C. Eaborn), (423) C33
- Gmelin handbook of inorganic chemistry, 8th edition, Sb—organoantimony compounds, part V, compounds of pentavalent antimony with two, three, and one Sb—C bonds (C. Eaborn), (424) C1