

JOURNAL OF ORGANOMETALLIC CHEMISTRY, VOL. 435 (1992)

SUBJECT INDEX

Actinides

Molekülstruktur und Ladungsverteilung der 4f- und 5f-Elemente. III. Dipolmomente und Ladungsverteilung der 1:1-Addukte der Tris(cyclopentadienyl)lanthanide(III) bzw. -actinoide(III) mit Tetrahydrofuran; Röntgenstrukturanalyse von $(C_5H_5)_3Dy \cdot THF$ (R. Maier, B. Kanellakopulos, C. Apostolidis und B. Nuber), (435) 275

Boron

A simple convenient preparation of ω -aminoboronic acids and esters (J.M. Jego, B. Carboni and M. Vaultier), (435) 1

Synthesis and solution properties of the boron-containing clusters $HM_4(CO)_{12}BAu_2(dppf)$ ($M = Fe$ or Ru) and the competitive formation of $\{HM_4(CO)_{12}BHAu\}_2(dppf)$ for $M = Ru$ ($dppf = 1,1'$ -bis(diphenylphosphino)ferrocene). Molecular structure of $HRu_4(CO)_{12}BAu_2(dppf) \cdot 2 CHCl_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9

Chromium

Symmetric *versus* unsymmetric ligand orientations in (pentadienyl)Cr(I or II)(L)₂(L') species, and a spontaneous reduction of Cr^{II} to Cr^I (T.D. Newbound, A.M. Arif, D.R. Wilson, A.L. Rheingold and R.D. Ernst), (435) 73

Germanium

Amination of vinylsilanes and vinylgermanes containing an activated double bond (V. Dirnens, D.I. Barabanov, E. Liepiņš, L.M. Ignatovich and E. Lukevics), (435) 257

Gold

Asymmetrical monocationic or neutral gold(II) complexes: X-ray crystal structure of $[(C_6F_5)Au(CH_2 \cdot PPh_2CH_2)_2AuPPh_3]ClO_4 \cdot 0.5CH_2Cl_2$ (A. Laguna, M. Laguna, J. Jiménez, F.J. Lahoz and E. Olmos), (435) 235

Hafnium

Preparation and crystal structures of the complexes $(\eta^5-C_5H_4CMe_2 \cdot \eta^5-C_{13}H_8)MCl_2$ ($M = Zr, Hf$) and their role in the catalytic formation of syndiotactic polypropylene (A. Razavi and J. Ferrara), (435) 299

Iridium

Ligand redistribution reactions of some organometallic rhodium and iridium complexes (P.J. Stang and Y.-H. Huang), (435) 185

Surface organometallic chemistry: formation of $[Ir_8(CO)_{22}]^{2-}$ on magnesium oxide (S.D. Maloney, M.J. Kelley and B.C. Gates), (435) 377

Iron

Synthesis and solution properties of the boron-containing clusters $HM_4(CO)_{12}BAu_2(dppf)$ ($M = Fe$ or Ru) and the competitive formation of $\{HM_4(CO)_{12}BHAu\}_2(dppf)$ for $M = Ru$ ($dppf = 1,1'$ -bis(diphenylphosphino)ferrocene). Molecular structure of $HRu_4(CO)_{12}BAu_2(dppf) \cdot 2 CHCl_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9

Synthesis and characterization of Group IV metal–iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $\text{Cp}_2\text{Ti}[(\mu\text{-O}_2\text{CC}_5\text{H}_4)\text{Fe}(\text{CO})_2(\text{CH}_2\text{C}_6\text{H}_5)]_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

P_4 -Liganden mit maximaler Elektronendonorfähigkeit II. $[\{\text{Fe}(\text{CO})_4(\mu\text{-P}_2)\}_2\{\mu\text{-Fe}_2(\text{CO})_6\}_2]$ —ein Komplex mit einer ungewöhnlich stabilisierten P_4 -Einheit (M. Scheer, M. Dargatz, K. Schenzel und P.G. Jones), (435) 123

Electrochemistry of acetylide complexes of iron (L.D. Field, A.V. George, F. Laschi, E.Y. Malouf and P. Zanello), (435) 347

An X-ray crystallographic and high field NMR study of [7-dehydrocholesteryl acetate] $\text{Fe}(\text{CO})_3$: synthesis of a B-ring cyclohexadienyl cationic complex (R.E. Perrier, C.S. Frampton and M.J. McGlinchey), (435) 357

Lanthanides

Molekülstruktur und Ladungsverteilung der 4f- und 5f-Elemente. III. Dipolmomente und Ladungsverteilung der 1:1-Addukte der Tris(cyclopentadienyl)lanthanoid(III) bzw. -actinoide(III) mit Tetrahydrofuran; Röntgenstrukturanalyse von $(\text{C}_5\text{H}_5)_3\text{Dy}\cdot\text{THF}$ (R. Maier, B. Kanellakopulos, C. Apostolidis und B. Nuber), (435) 275

Manganese

The synthesis of alkylmanganese(III) complexes. Crystal structure of $\text{MnMe}(\text{2-Me}_2\text{NCH}_2\text{C}_6\text{H}_4)_2$ (J.L. Latten, R.S. Dickson, G.B. Deacon, B.O. West and E.R.T. Tiekkari), (435) 101

Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109

Metallocenes

Synthesis and solution properties of the boron-containing clusters $\text{HM}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf})$ ($\text{M} = \text{Fe}$ or Ru) and the competitive formation of $[\text{HM}_4(\text{CO})_{12}\text{BHAu}_2(\text{dppf})]$ for $\text{M} = \text{Ru}$ ($\text{dppf} = 1,1'\text{-bis(diphenylphosphino)ferrocene}$). Molecular structure of $\text{HRu}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf}) \cdot 2 \text{CHCl}_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9

ansa-Metallocene derivatives of Ti^{IV} and Zr^{IV} with the shortest $-\text{C}(\text{CH}_3)_2-$ bridge (I.E. Nifant'ev, A.V. Churakov, I.F. Urazowski, S.G. Mkoyan and L.O. Atovmyan), (435) 37

Synthesis and characterization of Group IV metal–iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $\text{Cp}_2\text{Ti}[(\mu\text{-O}_2\text{CC}_5\text{H}_4)\text{Fe}(\text{CO})_2(\text{CH}_2\text{C}_6\text{H}_5)]_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

Early-transition-metal ketene complexes: synthesis, reactivity and structure of ketene complexes of bis(trimethylsilyl)niobocene, X-ray structure of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Br}(\text{Ph}_2\text{C}=\text{C=O-C}_6\text{H}_4\text{O})]$ (A. Antinolo, A. Otero, M. Fajardo, C. Lopez-Mardomingo, D. Lucas, Y. Mugnier, M. Lanfranchi and M.A. Pellinghelli), (435) 55

Darstellung und Eigenschaften von und Reaktionen mit metallhaltigen Heterocyclen. LXXVIII. Das Verhalten von monomeren und dimeren Thiophosphinitonickel-Komplexen gegenüber Alkinen und Phosphaalkinen (E. Lindner, A. Nothdurft, R. Fawzi und C. Maichle), (435) 213

Synthese und Kristallstruktur von $\text{Cp}'_2\text{Zr}[\text{P}(\text{SiMe}_3)_2]_2$ ($\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Me}$) — Erste röntgenstrukturanalytische Untersuchung eines Zirkonocen-Bisposphido-Komplexes (F. Lindenberg und E. Hey-Hawkins), (435) 291

Preparation and structure of half-open vanadocene, $\text{V}[2,4-(\text{CH}_3)_2\text{C}_5\text{H}_5](\text{C}_5\text{H}_5)\text{CO}$ (Y. Wang, Y. Liang, J. Liu, S. Jin and Y. Lin), (435) 311

Electrochemical studies on organometallic compounds. XI. Electrogeneration and ESR characterization of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Cl}(\text{P}(\text{OMe})_3)]^+$ (D. Lucas, Y. Mugnier, A. Antinolo, A. Otero and M. Fajardo), (435) C3

Molybdenum

Preparation and properties of molybdenum and tungsten dinitrogen complexes. XXXV. Synthesis of diazoalkane and hydrazido(2-) complexes of tungsten containing a π -acceptor ligand (T. Aoshima, T. Tamura, Y. Mizobe and M. Hidai), (435) 85

- Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109
 Structural properties of cationic molybdenum and tungsten allyl derivatives (B.J. Brisdon, M. Cartwright, A.G.W. Hodson, M.F. Mahon and K.C. Molloy), (435) 319

Nickel

- Reactivity of the $[(\text{Ni}(\text{C}_6\text{F}_5)_2(\mu\text{-OH}))_2]^{2+}$ ion towards β -diketones, 8-hydroxiquinoline and heterocyclic thiones. Crystal structure of $[\text{NBu}_4][\text{Ni}(\text{C}_6\text{F}_5)_2(\text{C}_5\text{H}_4\text{NS-2})]$ (G. López, G. Sánchez, G. García, J. García, A. Martínez, J.A. Hermoso and M. Martínez-Ripoll), (435) 193
 A kinetic study of the coordination template effect in the nickel-catalysed formation of arylphosphonium salts (D.W. Allen and P.E. Cropper), (435) 203
 Darstellung und Eigenschaften von und Reaktionen mit metallhaltigen Heterocyclen. LXXVIII. Das Verhalten von monomeren und dimeren Thiophosphinitonickel-Komplexen gegenüber Alkinen und Phosphaalkinen (E. Lindner, A. Nothdurft, R. Fawzi und C. Maichle), (435) 213

Niobium

- Early-transition-metal ketene complexes: synthesis, reactivity and structure of ketene complexes of bis(trimethylsilyl)niobocene. X-ray structure of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Br}(\text{Ph}_3\text{C}\equiv\text{O}-\text{C},\text{O})]$ (A. Antinolo, A. Otero, M. Fajardo, C. Lopez-Mardomingo, D. Lucas, Y. Mugnier, M. Lanfranchi and M.A. Pellinghelli), (435) 55
 Electrochemical studies on organometallic compounds. XLJ. Electrogeneration and ESR characterization of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Cl}\{\text{P}(\text{OMe})_3\}]^+$ (D. Lucas, Y. Mugnier, A. Antinolo, A. Otero and M. Fajardo), (435) C3

Osmium

- Synthesis, characterization and hindered rotation of triosmium clusters ($\mu\text{-H}\text{Os}_3(\text{CO})_{10}\text{L}[\text{ON}(\text{CF}_3)_2]$ [$\text{L} = \text{CO, CH}_3\text{CN}$ and PhCN]) (H.G. Ang, C.H. Koh and W.L. Kwik), (435) 149
 Triosmium clusters containing bridging sulphur, hydrido, and diphosphine ligands: a study of hydride site preferences and mobility (K.A. Azam, S.E. Kabir, A. Miah, M.W. Day, K.I. Hardcastle, E. Rosenberg and A.J. Deeming), (435) 157

Palladium

- Pallada(IV)cyclopentane chemistry, including the synthesis of tris(pyrazol-1-yl)borate complexes (A.J. Carty and P.R. Traill), (435) C8

Phosphorus

- Neutrale und kationische Rhodiumkomplexe mit den potentiell zweizähnigen Phosphanliganden ${}^1\text{Pr}_2\text{PCH}_2\text{CH}_2\text{OMe}$, ${}^1\text{Pr}_2\text{PCH}_2\text{CH}_2\text{NMe}_2$ und ${}^1\text{Pr}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{NMe}_2$ (H. Werner, A. Hampp und B. Windmüller), (435) 169
 A kinetic study of the coordination template effect in the nickel-catalysed formation of arylphosphonium salts (D.W. Allen and P.E. Cropper), (435) 203
 Darstellung und Eigenschaften von und Reaktionen mit metallhaltigen Heterocyclen. LXXVIII. Das Verhalten von monomeren und dimeren Thiophosphinitonickel-Komplexen gegenüber Alkinen und Phosphaalkinen (E. Lindner, A. Nothdurft, R. Fawzi und C. Maichle), (435) 213

Platinum

- Electron transfer reactions involving $\text{trans-}[\text{PtH}_2(\text{PCy}_3)_2]$ and fluorinated benzonitriles (S. Hintermann, P.S. Pregosin, H. Rüegger and H.C. Clark), (435) 225

Rhenium

- Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109

Rhodium

Neutrale und kationische Rhodiumkomplexe mit den potentiell zweizähnigen Phosphanliganden $^i\text{Pr}_2\text{PCH}_2\text{CH}_2\text{OMe}$, $^i\text{Pr}_2\text{PCH}_2\text{CH}_2\text{NMe}_2$ und $^i\text{Pr}_2\text{PCH}_2\text{CH}_2\text{CH}_2\text{NMe}_2$ (H. Werner, A. Hampp und B. Windmüller), (435) 169

Ligand redistribution reactions of some organometallic rhodium and iridium complexes (P.J. Stang and Y.-H. Huang), (435) 185

Ruthenium

Synthesis and solution properties of the boron-containing clusters $\text{HM}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf})$ ($\text{M} = \text{Fe}$ or Ru) and the competitive formation of $(\text{HM}_4(\text{CO})_{12}\text{BHAu})_2(\text{dppf})$ for $\text{M} = \text{Ru}$ ($\text{dppf} = 1,1'\text{-bis(diphenylphosphino)ferrocene}$). Molecular structure of $\text{HRu}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf}) \cdot 2 \text{CHCl}_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9

Asymmetric hydrogenation of prochiral carboxylic acids and functionalized carbonyl compounds catalysed by ruthenium(II)-binap complexes with aryl nitriles (binap = (*R*)- or (*S*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) (L. Shao, K. Takeuchi, M. Ikemoto, T. Kawai, M. Ogasawara, H. Takeuchi, H. Kawano and M. Saburi), (435) 133

Silicon

Polycyclopentadienyls: synthesis of arylsilylcyclopentadienyl compounds and their η^5 -complexes with tungsten (H. Plenio), (435) 21

Silicon–carbon unsaturated compounds. XL. Photolysis of 1,4-bis(2-phenyltetramethylsilanyl)benzene (M. Ishikawa, K. Watanabe, H. Sakamoto and A. Kunai), (435) 249

Amination of vinylsilanes and vinylgermanes containing an activated double bond (V. Dirnens, D.I. Barabanov, E. Liepiņš, L.M. Ignatovich and E. Lukevics), (435) 257

Synthese und Kristallstruktur von $\text{Cp}'_2\text{Zr}[\text{PSiMe}_3]_2$ ($\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Me}$) — Erste röntgenstrukturanalytische Untersuchung eines Zirkonocen–Bispophosphido-Komplexes (F. Lindenberg und E. Hey-Hawkins), (435) 291

Tin

Complexes of dibutyldihalotin(IV) with imidazole and pyrazole. The crystal structure of dibutylchlorobis(pyrazole)tin(IV) (A. Sánchez González, J.S. Casas, J. Sordo and G. Valle), (435) 29

Sterically hindered organotin compounds. II. Synthesis of the organotin halides $(\text{mesityl})_2\text{SnX}_n\text{Y}_{2-n}$, $(\text{mesityl})\text{SnX}_n\text{Y}_{3-n}$ and $\text{R}_4\text{Sn}_2\text{X}_2$ ($\text{R} = 2,4,6\text{-}^i\text{Pr}_3\text{C}_6\text{H}_2$, $\text{X} = \text{Cl}$, $\text{Y} = \text{Br}$). Crystal structure of the ditin species $(\text{R}_2\text{BrSn})_2$ (P. Brown, M.F. Mahon and K.C. Molloy), (435) 265

Titanium

ansa-Metallocene derivatives of Ti^{IV} and Zr^{IV} with the shortest $-\text{C}(\text{CH}_3)_2-$ bridge (I.E. Nifant'ev, A.V. Churakov, I.F. Urazowski, S.G. Mkoyan and L.O. Atovmyan), (435) 37

Synthesis and characterization of Group IV metal–iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $\text{Cp}_2\text{Ti}[(\mu\text{-O}_2\text{CC}_5\text{H}_4)\text{Fe}(\text{CO})_2(\text{CH}_2\text{C}_6\text{H}_5)]_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

Tungsten

Polycyclopentadienyls: synthesis of arylsilylcyclopentadienyl compounds and their η^5 -complexes with tungsten (H. Plenio), (435) 21

Preparation and properties of molybdenum and tungsten dinitrogen complexes. XXXV. Synthesis of diazoalkane and hydrazido(2^-) complexes of tungsten containing a π -acceptor ligand (T. Aoshima, T. Tamura, Y. Mizobe and M. Hidai), (435) 85

Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109

Structural properties of cationic molybdenum and tungsten allyl derivatives (B.J. Brisdon, M. Cartwright, A.G.W. Hodson, M.F. Mahon and K.C. Molloy), (435) 319

EPR studies of photochemical reactions of hexacarbonylbis($\eta^5\text{-}2,4\text{-cyclopentadien-1-yl}$)ditungsten, $[\text{CpW}(\text{CO})_3]_2$, with quinones (M. Hanaya and M. Iwaizumi), (435) 337

Vanadium

Preparation and structure of half-open vanadocene, V[2,4-(CH₃)₂C₅H₅](C₅H₅)CO (Y. Wang, Y. Liang, J. Liu, S. Jin and Y. Lin), (435) 311

Zirconium

ansa-Metallocene derivatives of Ti^{IV} and Zr^{IV} with the shortest -C(CH₃)₂- bridge (I.E. Nifant'ev, A.V. Churakov, I.F. Urazowski, S.G. Mkoyan and L.O. Atovmyan), (435) 37

Synthesis and characterization of Group IV metal-iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of Cp₂Ti[μ -O₂CC₅H₄]Fe(CO)₂(CH₂C₆H₅)₂ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

Synthese und Kristallstruktur von Cp'₂Zr{P(SiMe₃)₂}₂ (Cp' = η⁵-C₅H₄Me) — Erste röntgenstrukturanalytische Untersuchung eines Zirkonocen-Bisphosphido-Komplexes (F. Lindenberg und E. Hey-Hawkins), (435) 291

Preparation and crystal structures of the complexes (η^5 -C₅H₄CMe₂- η^5 -C₁₃H₈)MCl₂ (M = Zr, Hf) and their role in the catalytic formation of syndiotactic polypropylene (A. Razavi and I. Ferrara), (435) 299

Book review

Reaction Mechanisms of Inorganic and Organometallic Systems, R.B. Jordan (D.A. Brown), (435) C1