

## Subject Index of Volume 449

---

### Aluminium

The systems  $\text{LiAlH}_4/\text{NH}_4\text{X}$  and  $\text{N}_2\text{H}_5\text{Cl}$  as precursor sources for AlN (J.F. Janik and R.T. Paine), 39

Tertiary phosphine adducts of alane and gallane (F.M. Elms, M.G. Gardiner, G.A. Koutsantonis, C.L. Raston, J.L. Atwood and K.D. Robinson), 45

Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido [ $(^t\text{Bu})\text{PH}$ ]<sup>-</sup> and arsenido [ $(^t\text{Bu})\text{AsH}$ ]<sup>-</sup> units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), 61

### Antimony

Primary and secondary trimethylsilylmethylstibines. Synthesis, characterization and chemical vapor deposition properties (D.G. Hendershot and A.D. Berry), 119

Antimony sources for MOCVD. The use of  $\text{Et}_4\text{Sb}_2$  as a p-type dopant for  $\text{Hg}_{1-x}\text{Cd}_x\text{Te}$  and crystal structure of the adduct  $[\text{Et}_4\text{Sb}_2 \cdot 2\text{CdI}_2]_n$  (R.S. Dickson, K.D. Heazle, G.N. Pain, G.B. Deacon, B.O. West, G.D. Fallon, R.S. Rowe, P.W. Leech and M. Faith), 131

### Arsenic

Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido [ $(^t\text{Bu})\text{PH}$ ]<sup>-</sup> and arsenido [ $(^t\text{Bu})\text{AsH}$ ]<sup>-</sup> units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), 61

Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of  $[\text{R}(\text{Me}_3\text{SiCH}_2)\text{InE}(\text{SiMe}_3)_2]_2$  ( $\text{R} = \text{Ph}, \text{E} = \text{As}; \text{R} = \text{Me}, \text{E} = \text{As}, \text{P}; \text{R} = \text{Cl}, \text{E} = \text{P}$ ) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), 85

Aspects of the chemistry of vinylarsinic compounds. The preparation of divinylarsinic acid, tetravinyldiarsine oxide, and tetravinyldiarsine, and the X-ray crystal structure of the helical divinylarsinic acid (V.L. Goedken, L.F. Brough and W.S. Rees, Jr.), 125

### Boron

Synthesis and second-order nonlinear optical properties of three coordinate organoboranes with diphenylphosphino and ferrocenyl groups as electron donors: crystal and molecular structures of (*E*)- $\text{D}-\text{CH}=\text{CH}-\text{B}(\text{mes})_2$  and  $\text{D}-\text{C}\equiv\text{C}-\text{B}(\text{mes})_2$  [ $\text{D} = \text{P}(\text{C}_6\text{H}_5)_2$ , ( $\eta$ - $\text{C}_5\text{H}_5$ ) $\text{Fe}(\eta\text{-C}_5\text{H}_4)$ ; mes = 2,4,6-( $\text{CH}_3)_3\text{C}_6\text{H}_2$ ] (Z. Yuan, N.J. Taylor, Y. Sun, T.B. Marder, I.D. Williams and L.-T. Cheng), 27

### Cadmium

Structural studies of some Group 12 metal alkyl adducts: the X-ray crystal structures of  $\text{Me}_2\text{Zn}[\text{Me}_2\text{N}(\text{CH}_2)_2\text{NMe}_2]$ ,  $\text{Me}_2\text{Cd}[\text{Me}_2\text{N}(\text{CH}_2)_2\text{NMe}_2]$ ,  $(\text{Me}_3\text{CCH}_2)_2\text{Zn}[\text{Me}_2\text{N}(\text{CH}_2)_2\text{NMe}_2]$  and  $(\text{Me}_3\text{CCH}_2)_2\text{Cd}[\text{Me}_2\text{N}(\text{CH}_2)_2\text{NMe}_2]$  (P. O'Brien, M.B. Hursthouse, M. Motavalli, J.R. Walsh and A.C. Jones), 1

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors  $\text{M}^{\text{II}}\text{M}^{\text{IV}}\text{P}_2$  where  $\text{M}^{\text{II}} = \text{Zn}$  and  $\text{Cd}$ , and

$\text{M}^{\text{IV}} = \text{Ge}$  and  $\text{Sn}$  (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conradi), 9

Synthesis and characterization of sterically encumbered Li, Na, and K aryl tellurolates, and some Pt(II), Ir(I), and Cd(II) derivatives (Ph.J. Bonasia and J. Arnold), 147

### Chalcogenides

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), 167

### Cobalt

Formation of a high surface area, regular porous solid from the cluster of clusters,  $\text{Zn}_4^{\text{II}}\text{O}[(\text{CO})_9\text{Co}_3\text{CCO}_2]_6$  (W. Cen, B. Ladna, T.P. Fehlner, A.E. Miller and D. Yue), 19

Study of CoGa deposition from the single source precursor  $(\text{CO})_4\text{CoGaCl}_2$  (THF) (F. Maury, L. Brandt and H.D. Kaesz), 159

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), 167

### Copper

Chemistry of copper(I)  $\beta$ -diketonate complexes. VI. Synthesis, characterization and chemical vapor deposition of 2,2-dimethyl-6,6,7,7,8,8-heptafluoro-3,5-octanedione (fod), copper(I)L complexes and the solid state structure of (fod) $\text{Cu}(\text{PMMe}_3)$  (K.M. Chi, T.S. Corbitt, M.J. Hampden-Smith, T.T. Kodas and E.N. Duesler), 181

A comparison of heterometallic alkoxide molecules containing copper(I) or copper(II) and zirconium (B.A. Vaartstra, J.A. Samuels, E.H. Barash, J.D. Martin, W.E. Streib, C. Gasser and K.G. Caulton), 191

### Gallium

Tertiary phosphine adducts of alane and gallane (F.M. Elms, M.G. Gardiner, G.A. Koutsantonis, C.L. Raston, J.L. Atwood and K.D. Robinson), 45

Synthesis, characterization, and reactivity of new alkylgallium acetylides (K.E. Lee and K.T. Higa), 53

Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido [ $(^t\text{Bu})\text{PH}$ ]<sup>-</sup> and arsenido [ $(^t\text{Bu})\text{AsH}$ ]<sup>-</sup> units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), 61

Gallium and indium compounds containing three different substituents. Crystal and molecular structure of  $[(\text{Me}_3\text{CCH}_2)\text{Cl}-\text{GaPP}_2\text{H}_3](\text{O.T. Beachley, Jr., J.D. Maloney and R.D. Rogers})$ , 69

Study of CoGa deposition from the single source precursor  $(\text{CO})_4\text{CoGaCl}_2$  (THF) (F. Maury, L. Brandt and H.D. Kaesz), 159

## Germanium

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors  $M^{II}M^{IV}P_2$  where  $M^{II}$  = Zn and Cd, and  $M^{IV}$  = Ge and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conradi), 9

## Gold

Monomeric to polymeric gold(I) complexes from bifunctional ligands  $R_2PC\equiv CH$  (G. Jia, R.J. Puddephatt and J.J. Vittal), 211

## Indium

Gallium and indium compounds containing three different substituents. Crystal and molecular structure of  $[(Me_3CCH_2)ClGaPPPh_2]_3$  (O.T. Beachley, Jr., J.D. Maloney and R.D. Rogers), 69

Raman spectroscopic studies of gas phase  $Me_3In$ ,  $Et_3In$  and  $EtMe_2In$  (Z.S. Huang, C. Park and T.J. Anderson), 77

Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of  $[R(Me_3SiCH_2)InE(SiMe_3)_2]_2$  ( $R = Ph, E = As; R = Me, E = As, P; R = Cl, E = P$ ) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), 85

Indium tert-butylthiolates as single source precursors for indium sulfide thin films: is molecular design enough? (A.N. MacInnes, M.B. Power, A.F. Hepp and A.R. Barron), 95

An indium-phosphorus heterocubane as a new type of indium phosphide precursor (D.A. Atwood, A.H. Cowley, R.A. Jones and M.A. Mardones), C1

## Iridium

Synthesis and characterization of sterically encumbered Li, Na, and K aryl tellurolates, and some Pt(II), Ir(I), and Cd(II) derivatives (Ph.J. Bonasia and J. Arnold), 147

## Iron

Synthesis and second-order nonlinear optical properties of three coordinate organoboranes with diphenylphosphino and ferrocenyl groups as electron donors: crystal and molecular structures of (*E*)-D-CH=CH-B(mes)<sub>2</sub> and D-C≡C-B(mes)<sub>2</sub> [D = P(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>, ( $\eta$ -C<sub>5</sub>H<sub>5</sub>)Fe( $\eta$ -C<sub>5</sub>H<sub>4</sub>); mes = 2,4,6-(CH<sub>3</sub>)<sub>3</sub>C<sub>6</sub>H<sub>2</sub>] (Z. Yuan, N.J. Taylor, Y. Sun, T.B. Marder, I.D. Williams and L.-T. Cheng), 27

## Lithium

The systems LiAlH<sub>4</sub>/NH<sub>4</sub>X and N<sub>2</sub>H<sub>5</sub>Cl as precursor sources for AlN (J.F. Janik and R.T. Paine), 39

## Metallocenes

Synthesis and second-order nonlinear optical properties of three coordinate organoboranes with diphenylphosphino and ferrocenyl groups as electron donors: crystal and molecular structures of (*E*)-D-CH=CH-B(mes)<sub>2</sub> and D-C≡C-B(mes)<sub>2</sub> [D = P(C<sub>6</sub>H<sub>5</sub>)<sub>2</sub>, ( $\eta$ -C<sub>5</sub>H<sub>5</sub>)Fe( $\eta$ -C<sub>5</sub>H<sub>4</sub>); mes = 2,4,6-(CH<sub>3</sub>)<sub>3</sub>C<sub>6</sub>H<sub>2</sub>] (Z. Yuan, N.J. Taylor, Y. Sun, T.B. Marder, I.D. Williams and L.-T. Cheng), 27

## Phosphorus

Tertiary phosphine adducts of alane and gallane (F.M. Elms, M.G. Gardiner, G.A. Koutsantonis, C.L. Raston, J.L. Atwood and K.D. Robinson), 45

Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido [<sup>t</sup>BuPH]<sup>-</sup> and arsenido [<sup>t</sup>BuAsH]<sup>-</sup> units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), 61

Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of  $[R(Me_3SiCH_2)InE(SiMe_3)_2]_2$  ( $R = Ph, E = As; R = Me, E = As, P; R = Cl, E = P$ ) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), 85

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), 167

An indium-phosphorus heterocubane as a new type of indium phosphide precursor (D.A. Atwood, A.H. Cowley, R.A. Jones and M.A. Mardones), C1

Monomeric to polymeric gold(I) complexes from bifunctional ligands  $R_2PC\equiv CH$  (G. Jia, R.J. Puddephatt and J.J. Vittal), 211

## Platinum

Synthesis and characterization of sterically encumbered Li, Na, and K aryl tellurolates, and some Pt(II), Ir(I), and Cd(II) derivatives (Ph.J. Bonasia and J. Arnold), 147

## Rhodium

Studies of the thermal and photochemical lability of olefin complexes of rhodium(I) to form Rh and Rh<sub>2</sub>O<sub>3</sub> (Z. Duan, M.J. Hampden-Smith and A.P. Sylvester), 173

## Selenium

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), 167

## Silicon

Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of  $[R(Me_3SiCH_2)InE(SiMe_3)_2]_2$  ( $R = Ph, E = As; R = Me, E = As, P; R = Cl, E = P$ ) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), 85

Measurement of surface tension for polysilanes (T. Fujisaka, R. West and C. Murray), 105

Pyrolysis of poly(silylene)diacetylenes: direct evidence between their morphology and thermal behavior (R. Corriu, Ph. Gerbier, C. Guérin, B. Henner and R. Fourcade), 111

## Silver

Crystal structure of  $[CF_3CF=C(CF_3)AgI_4]$  and its use as a chemical vapor deposition precursor for silver films (P.M. Jeffries, S.R. Wilson and G.S. Girolami), 203

## Sulphur

Indium tert-butylthiolates as single source precursors for indium sulfide thin films: is molecular design enough? (A.N. MacInnes, M.B. Power, A.F. Hepp and A.R. Barron), 95

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), 167

## Tellurium

Evidence for a free radical mechanism in the decomposition of bis(but-2-enyl)tellurium (J. Stevenson, W. Bell, J. Ferry, D.J. Cole-Hamilton and J.E. Hails), 141

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), 167

**Tin**

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors  $M^{II}M^{IV}P_2$  where  $M^{II}$  = Zn and Cd, and  $M^{IV}$  = Ge and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conradi), 9

**Zinc**

Structural studies of some Group 12 metal alkyl adducts: the X-ray crystal structures of  $Me_2Zn[Me_2N(CH_2)_2NMe_2]$ ,  $Me_2Cd[Me_2N(CH_2)_2NMe_2]$ ,  $(Me_3CCH_2)_2Zn[Me_2N(CH_2)_2NMe_2]$  and  $(Me_3CCH_2)_2Cd[Me_2N(CH_2)_2NMe_2]$  (P. O'Brien, M.B. Hursthouse, M. Mottevalli, J.R. Walsh and A.C. Jones), 1

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors  $M^{II}M^{IV}P_2$  where  $M^{II}$  = Zn and Cd, and  $M^{IV}$  = Ge and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conradi), 9

Formation of a high surface area, regular porous solid from the cluster of clusters,  $Zn_4^{II}O[(CO)_9Co_3CCO_2]_6$  (W. Cen, B. Ladna, T.P. Fehlner, A.E. Miller and D. Yue), 19

**Zirconium**

A comparison of heterometallic alkoxide molecules containing copper(I) or copper(II) and zirconium (B.A. Vaartstra, J.A. Samuels, E.H. Barash, J.D. Martin, W.E. Streib, C. Gasser and K.G. Caulton), 191