

Cumulative Subject Index of Volumes 426 (1992)–450 (1993)

Actinides

Molekülstruktur und Ladungsverteilung der 4f- und 5f-Elemente. III. Dipolmomente und Ladungsverteilung der 1:1-Addukte der Tris(cyclopentadienyl)lanthanide(III) bzw. -actinoide(III) mit Tetrahydrofuran; Röntgenstrukturanalyse von $(C_5H_5)_3Dy \cdot THF$ (R. Maier, B. Kanellakopulos, C. Apostolidis und B. Nuber), (435) 275

Aluminium

Die Reaktion des Tri(isobutyl)alans mit Kalium—eine Neuuntersuchung; Kristallstruktur des Kalium[tetra(isobutyl)alanats] (W. Uhl und J. Wagner), (427) 151

Planar tetracoordinate carbon stabilized in a dimetallic hafnium/aluminium compound: formation and crystal structure of $Cp_2Hf[\mu-\eta^1:\eta^2-MeCC(C_6H_{11})][\mu-CC(C_6H_{11})]AlMe_2$ (M. Albrecht, G. Erker, M. Nolte und C. Krüger), (427) C21

Methylation of bis(oxazolinyl)pyridinerhodium(III) chlorides with trimethylaluminum (H. Nishiyama, T. Hirai und K. Itoh), (431) 227

Synthesis and characterization of alkylaluminum and -gallium derivatives of thiomorpholine. Molecular structures of $Me_3Al \cdot N(H)C_4H_8S$ and $[Et_2Al(\mu-NC_4H_8S)]_2$ (M. Taghiof, D.G. Hendershot, M. Barber und J.P. Oliver), (431) 271

Alkylidenedinitrosylmolybdenum complexes. Preparation, characteristics and metathesis activity of dicarboxylatoethylidenedinitrosylmolybdenum complexes (A. Keller), (436) 199

Reactions of trimethylaluminium with 2-[methyl-bis(trimethylsiloxy)silyl]but-2-ene-1,4-diol: synthesis and structure of $[Al(CH_3)]_2[OCH_2(SiO_2Me(OSiMe_3)_2)C=C(H)CH_2O_2]_2[Al(CH_3)_2]_2$ (S. Pasynkiewicz und W. Ziemkowska), (437) 99

Stereoselective opening of chiral α -stannyacetals with organometallic reagents (J.-L. Parrain, J.-C. Cintrat und J.-P. Quintard), (437) C19

Kinetically stable adduct of samarocene with aluminium deuteride ($\eta-C_5H_5^1Bu_2)_2Sm(\mu_2-D)_2AlD \cdot Me_2NC_2H_4NMe_2$ (Y.K. Gun'ko, B.M. Bulychev, V.K. Belsky und G.L. Soloveichik), (440) 47

Etude de la fixation du groupe tertiaire *gem*-diméthylpropynyle $HC\equiv CC(CH_3)_2$ par voie organométallique; applications (F. Barbot und P. Miginiac), (440) 249

Allénysilanes fonctionnels. II. Synthèse par voie organoaluminique du 2-triméthylsilylbuta-2,3-diénal et application à la synthèse d'alcools secondaires α -alléniques triméthylsilylés (D. Mesnard und L. Miginiac), (440) 277

Die Reaktion des Tetrakis[bis(trimethylsilyl)methyl]dialans(4) mit Schwefelkohlenstoff: Molekülstrukturen mit Al-S-Al- und Al-CS₂-Al-Fragment (W. Uhl, A. Vester und W. Hiller), (443) 9

Reaction of Me_3Ga and Me_3Al with the tertiary-tetraaza analog of adamantine, hexamethylenetetramine (N_4 -Ada). Syntheses and molecular structures of $[(Me_3M)_n \cdot N_4-Ada]$ ($n = 1-4$, M = Ga or Al) (J.B. Hill, S.J. Eng, W.T. Pennington und G.H. Robinson), (445) 11

Dimethylsiloxy-aluminiumdihalide dimers. Molecular structures of $[Me_2HSiOAlX_2]_2$ (X = Cl, I) and $[Me_3SiOAl(BH_4)_2]_2$ (P. Bissinger, P. Mikulcik, J. Riede, A. Schier und H. Schmidbaur), (446) 37

The crystal structure and properties of $[1,3,5-C_6H_3(CO_2Me)_3 \cdot 3AlCl_3]$ (J. Utko, M. Wróblewska, T. Lis und P. Sobota), (447) 159

The systems $LiAlH_4/NH_4X$ and N_2H_5Cl as precursor sources for AlN (J.F. Janik und R.T. Paine), (449) 39

Tertiary phosphine adducts of alane and gallane (F.M. Elms, M.G. Gardiner, G.A. Koutsantonis, C.L. Raston, J.L. Atwood und K.D. Robinson), (449) 45

Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido [$(^1Bu)PH$]⁻ and arsenido [$(^1Bu)AsH$]⁻ units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder und C.M. Nunn), (449) 61 Über dimere Dimethylmetall-bis(trimethylgermyl)amide des Aluminiums, Galliums und Indiums (M. Trapp, H.-D. Hausen, G. Weckler und J. Weidlein), (450) 53

Intramolekulare Basenkoordination je nach Bedarf: Der 1-(Dimethylaminoethyl)-2,3,4,5-tetramethylcyclopentadienyl-Ligand in der Komplexchemie von s-, p-, d- und f-Block-Elementen (P. Jutzi, J. Dahlhaus und M.O. Kristen), (450) C1

Antimony

Strukturen von $[Me_4Sb]_2[MeSbI_4]$, $MeSbI_2$, und $[Me_4Sb]I$. Darstellung von $Me_3Sb \cdot MeSbI_2$ und Farbwechsel bei $Me_4Sb_2 \cdot Me_2SbBr$ (H.J. Breunig, K.H. Ebert, S. Gülec, M. Dräger, D.B. Sowerby und M.J. Begley), (427) 39

Primary and secondary trimethylsilylmethylstibines. Synthesis, characterization and chemical vapor deposition properties (D.G. Hendershot und A.D. Berry), (449) 119

Antimony sources for MOCVD. The use of Et_4Sb_2 as a p-type dopant for $Hg_{1-x}Cd_xTe$ and crystal structure of the adduct $[Et_4Sb_2 \cdot 2CdI_2]_n$ (R.S. Dickson, K.D. Heazle, G.N. Pain, G.B. Deacon, B.O. West, G.D. Fallon, R.S. Rowe, P.W. Leech und M. Faith), (449) 131

Arsenic

Metallorganische Verbindungen der Lanthanoide. LXIX. Intramolekulare Stabilisierung der Lanthanoid-Kohlenstoff- σ -Bindung in Mono(cyclopentadienyl)lanthanoidkomplexen (H. Schumann, J.A. Meese-Markscheffel, A. Dietrich und J. Pickardt), (433) 241

η^1 - und η^2 -Komplexe von Cp^* -substituierten Diphenophenen und Arsaphenphenen: Synthese, Struktur und Umlagerung unter Cp^* -Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammel), (439) 279

Synthesis of the first diarsadigallatane (D.A. Atwood, A.H. Cowley, R.A. Jones und M.A. Mardones), (439) C33

- Synthesen und photochemische Untersuchungen von *ortho*- und *meta*-substituierten Bis(phenyl)[η^2 -1,2-bis(diphenylarsino)ethan bzw. -ethen]platin(II)-Verbindungen (H.-A. Brune, E. Hupfer, G. Schmidberg und U. Ziegler), (440) 303
- Bis(diisopropylgalliumdiisopropylphosphid) und -arsenid. Kristallstruktur des Phosphidderivats—ein “angegedeuter” butterfly (G.G. Hoffmann, R. Fischer, U. Schubert und B. Hirle), (441) 7
- Stereoselektive Additionsreaktionen von Komplex-gebundenen Phosphanen und Arsanen an Alkine (H. Lang und U. Lay), (441) 389
- Additionsreaktionen von $(CO)_5CrEPhH_2$ ($E = P, As$) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), (444) 53
- Reactions of coordinated phosphines and arsines: stereoselective reactions at the chlorophosphine-*P* stereocentre in the complex $[(R^*), (R^*, R^*)](\pm)[(\eta^5-C_5H_5)(1,2-C_6H_4(PMePh)_2)-Fe(PCl-MePh)]PF_6 \cdot CH_2Cl_2$ (G. Salem, G.B. Shaw, A.C. Willis and S.B. Wild), (444) 147
- Atropisomerism in (\pm) -7-methyl- and -phenyl-substituted dinaphtho[2,1-*b*;1',2'-*d*]phospholes and dinaphth[2,1-*b*;1',2'-*d*]-arsoles (A.A. Watson, A.C. Willis and S.B. Wild), (445) 71
- Verbrückte Diarsinidenkomplexe $[(Cp'(CO)_2Mn)_2As]_2E$ ($E = O, S, Se$) und deren nukleophile Spaltung (C. Emmerich, G. Huttner und A. Asam), (447) 71
- Substitution, nukleophile und oxidative Addition an Arsinidenkomplexe. Zur Reaktion von $[Cp'(CO)_2Mn]_2AsCl$ mit PhSeH und verwandten Chalkogenverbindungen (C. Emmerich und G. Huttner), (447) 81
- Reaktionen von $[Cp'(CO)_2Mn]_2AsCl$ mit Kohlenstoff-Nukleophilen (C. Emmerich und G. Huttner), (448) 83
- Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido [$(^1Bu)PH^-$] and arsenido [$(^1Bu)AsH^-$] units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), (449) 61
- Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of $[R(Me_3SiCH_2)InE(SiMe_3)]_2$ ($R = Ph, E = As; R = Me, E = As, P; R = Cl, E = P$) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), (449) 85
- Aspects of the chemistry of vinylarsenic compounds. The preparation of divinylarsinic acid, tetravinyl diarsine oxide, and tetravinyl diarsine, and the X-ray crystal structure of the helical divinylarsinic acid (V.L. Goedken, L.F. Brough and W.S. Rees, Jr.), (449) 125
- Reaktive Arsen-Heterocyclen. II. 3-Arsolene als Arsiniden-Quelle (W.A. Schenk und M. Stubbe), (450) C4
- Barium**
- Reaction of organocalcium and -barium complexes with pyrazines. X-ray structural characterization of $[(Me_5C_5)_2Ba]_2(\mu-C_4H_4N_2)$ (R.A. Williams, T.P. Hanusa and J.C. Huffman), (429) 143
- Bismuth**
- Manganese carbonyl complexes of 2,5-dimethylbismolyl. The crystal and molecular structure of (η^1 -2,5-dimethylbismolyl) manganese pentacarbonyl (A.J. Ashe, III, J.W. Kampf and D.B. Puranik), (447) 197
- Boron**
- Polyazolyl chelate chemistry. IV. Dihydrobis(benzotriazolyl)borato complexes of ruthenium(II) (J. Cartwright and A.F. Hill), (429) 229
- Sterically induced opening of a *clos* carbametallaborane: synthesis and characterisation of 1,2- Ph_2 -3-(η - C_5Me_3)-3,1,2-*pseudoclos*- $RhC_2B_9H_9$ (Z.G. Lewis and A.J. Welch), (430) C45
- Derivatives of tetrahedral boronic acids (M. Biedrzycki, W.H. Scouten and Z. Biedrzycka), (431) 255
- Carbonylation of $>B-H$ -boranes; evidence for a Fischer-Tropsch type reaction (M. Yalpani and R. Köster), (434) 133
- A simple convenient preparation of ω -aminoboronic acids and esters (J.M. Jego, B. Carboni and M. Vaultier), (435) 1
- Synthesis and solution properties of the boron-containing clusters $HM_4(CO)_{12}BAu_2(dppf)$ ($M = Fe$ or Ru) and the competitive formation of $\{HM_4(CO)_{12}BHAu_2\}_2(dppf)$ for $M = Ru$ ($dppf = 1,1'$ -bis(diphenylphosphino)ferrocene). Molecular structure of $HRu_4(CO)_{12}BAu_2(dppf) \cdot 2 CHCl_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9
- Synthesis and structural characterization of cycloborasiloxanes: the X-ray crystal structures of *cyclo*-1,3,3,5,5-pentaphenyl-1-bora-3,5-disiloxane and *cyclo*-1,3,3,5,7,7-hexaphenyl-1,5-dibora-3,7-disiloxane (B.J. Brisdon, M.F. Mahon, K.C. Molloy and P.J. Schofield), (436) 11
- Steric and metal-centre control of two-, three- and four-orbital carborane-metal interaction in tin and platinum complexes of the $\{C_2B_8H_9X\}$ ligand, where $X = H$ and Ph (J.D. Kennedy, K. Nestor, B. Šíbr, M. Thornton-Pett and G.S.A. Zammit), (437) C1
- Indenyl carbametallaboranes. III. The synthesis and characterization of novel rhodium carbaborane compounds with $3-(C_2B_9)-8-\{(C_9H_6)Rh(C_9H_7)\}-3,1,2$ -*clos*- RhC_2B_9 , $10-\{(C_9H_6)Rh(C_9H_7)\}-7,8$ -*nido*- C_2B_9 , $9-\{(C_9H_6)Rh(C_9H_7)\}-7,8$ -*nido*- C_2B_9 and $3-(\eta-C_9H_7)-3,1,2$ -*clos*- RhC_2B_9 structures (Z.G. Lewis and A.J. Welch), (438) 353
- Methylphenylborohydride salts and methylphenylborane: new reagents for preparing methylphenylborane derivatives (Z. Wang, C. Narayana, P.P. Wadgaonkar and G.W. Kabalka), (440) 243
- Sodium tetraphenylborate as a phenylating reagent in the palladium-catalyzed phenylation of alkenes and acid chlorides (C.S. Cho, K. Itotani and S. Uemura), (443) 253
- Asymmetric catalysis. LXXX. An optically-active tetrakispyrazolylborate: Synthesis and use in Cu-catalysed enantioselective cyclopropanation (H. Brunner, U.P. Singh, T. Boeck, S. Altmann, T. Scheck and B. Wrackmeyer), (443) C16
- A new type of stabilized carbonucleophiles with a carboranyl group (S.C. Shim, J.G. Shim, S.Y. Chae, S.Y. Lee and V.N. Kalinin), (443) C22
- B-C bond formation by ene-type reactions of dimethylaminobis(trifluoromethyl)borane with nitriles and carbonyl compounds. Crystal structure of $(NCCH_2)_2CF_3)_2B \cdot NHMe_2$ (A. Ansorge, D.J. Brauer, H. Bürger, T. Hagen and G. Pawelke), (444) 5
- Trisubstituted alkenes from alkynes and aryl iodides by boron-to-zinc transmetallation and palladium(0) catalyzed cross coupling (K.A. Agrios and M. Srebnik), (444) 15
- Photochemistry of tris(dimethylpyrazolyl)-boratorrhodium dicarbonyl and bis(dimethylpyrazolyl)-boratorrhodium dicarbonyl complexes in low temperature media at 12–298 K: some insights into C–H activation processes (P.E. Bloyce, J. Mascetti and A.J. Rest), (444) 223
- Stereoselective synthesis of allylic boronates via palladium-catalyzed cross-coupling reaction of Knochel's (dialkoxyboryl)methylzinc reagents with 1-halo-1-alkenes (T. Watanabe, N. Miyaura and A. Suzuki), (444) C1
- Formation and instability of an α -acetoxyborane (J.L. Hubbard), (445) 7
- Synthesis, molecular structure and NMR behaviour of $[9,9-(PMe_2Ph)_2arachno-9,6,8-PtS_2B_7H_7]$: Direct evidence for the ten-vertex borane vertex-flip mechanism of isomerization (J.H. Jones, X.L.R. Fontaine, N.N. Greenwood, J.D. Kennedy, M. Thornton-Pett, B. Šíbr and H. Langhoff), (445) C15

The ammine complex of tris(trifluoromethyl)borane, and some N-acyl derivatives. Crystal structure of $(CF_3)_3B \cdot NH_3$ and $(CF_3)_3B \cdot NH=CHN(CH_3)_2$ (A. Ansorge, D.J. Brauer, H. Bürger, B. Krumm and G. Pawelke), (446) 25

Synthesis and second-order nonlinear optical properties of three coordinate organoboranes with diphenylphosphino and ferrocenyl groups as electron donors: crystal and molecular structures of (*E*)-D-CH=CH-B(mes)₂ and D-C≡C-B(mes)₂ [D = P(C₆H₅)₂, (η -C₅H₅)Fe(η -C₅H₄); mes = 2,4,6-(CH₃)₃C₆H₂] (Z. Yuan, N.J. Taylor, Y. Sun, T.B. Marder, I.D. Williams and L.-T. Cheng), (449) 27

Convenient method for the preparation of catecholborane and promotion of the formation of alkenyl catecholborane using BH₃ complexes (Y. Suseela and M. Periasamy), (450) 47

Crystal structure of Fe(η^6 -C₅H₅BCMe₃)₂, a "problem structure" (M.R. Churchill and R.F. See), (450) 171

Cadmium

Reactions of sterically hindered organozinc and organocadmium compounds containing functional silicon centres. Crystal structures of Zn[CSiMe₃)₂(SiMe₂OCOCF₃)₂ and Cd[CSiMe₃)₂(SiMe₂OMe)₂] (S.S. Al-Juaid, C. Eaborn, A. Habtemariam, P.B. Hitchcock and J.D. Smith), (437) 41

Chiral cyclopentadiene-cyclopentadienyl complexes of cobalt (V.P. Maryin and P.V. Petrovskii), (441) 125

Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241

Synthesis of phenyl(perfluoroorgano)mercury derivatives C₆H₅Hg(R_f) [R_f = C_nF_{2n+1} (n = 1–4 or 6) or C₆F₅] by the reactions of bis(perfluoroorgano)cadmium compounds with phenylmercury chloride (H. Layeghi, D. Naumann and W. Tyrra), (441) 355

Structural studies of some Group 12 metal alkyl adducts: the X-ray crystal structures of Me₂Zn[Me₂N(CH₂)₂NMe₂], Me₂Cd[Me₂N(CH₂)₂NMe₂], (Me₃CCH₂)₂Zn[Me₂N(CH₂)₂NMe₂] and (Me₃CCH₂)₂Cd[Me₂N(CH₂)₂NMe₂] (P. O'Brien, M.B. Hursthorne, M. Mottevalli, J.R. Walsh and A.C. Jones), (449) 1

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors M^{II}M^{IV}P₂ where M^{II} = Zn and Cd, and M^{IV} = Ge and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conradi), (449) 9

Synthesis and characterization of sterically encumbered Li, Na, and K aryl tellurolates, and some Pt(II), Ir(I), and Cd(II) derivatives (Ph.J. Bonasia and J. Arnold), (449) 147

Calcium

Reaction of organocalcium and -barium complexes with pyrazines. X-ray structural characterization of [(Me₅C₅)₂Ba]₂(μ -C₄H₄N₂) (R.A. Williams, T.P. Hanusa and J.C. Huffman), (429) 143

Characterization of 1:1 heterobimetallic complexes of (η^5 -C₅Me₅)Ir(CO)₂ with zinc, cadmium and mercury(II) chloride and the X-ray structure determination of the 1:2 mercury complex (F.W.B. Einstein, X. Yan, X. Zhang and D. Sutton), (439) 221

Intramolekulare Basenkoordination je nach Bedarf: Der 1-(Dimethylaminoethyl)-2,3,4,5-tetramethylcyclopentadienyl-Ligand in der Komplexchemie von s-, p-, d- und f-Block-Elementen (P. Jutzi, J. Dahlhaus and M.O. Kristen), (450) C1

Cerium

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), (444) C15

Cesium

The preference of 1-methylallyl polar organometallics and carbanions for *cis* rather than for *trans* geometries (P. von R. Schleyer, J. Kaneti, Y.-D. Wu and J. Chandrasekhar), (426) 143

Chalcogenides

Zur Addition von Schwefel und Selen an Dicarbonyl(η^5 -cyclopentadienyl)[η^2 -(methylthio)methylcarben]wolfram-tetrafluoroborat (F.R. Kreißl and N. Ullrich), (426) C8

Structure and dynamics of [C₅H₅Fe(CO)₂(ECH₃)₂]BF₄ (E = S, Se, Te) complexes studied by ¹³C solid state NMR spectroscopy (A.P.M. Kentgens, H. Karrenbeld, E. De Boer and H. Schumann), (429) 99

Diferrocenyldichalcogenide: Synthese und Struktur von Bis[2-(dimethylaminomethyl)ferrocenyl]diselenid (H. Gornitzka, S. Besser, R. Herbst-Irmer, U. Kilimann, F.T. Edelmann und K. Jacob), (437) 299

Trigon-planar koordiniertes Selen bzw. Tellur in [Cp'(CO)₂Mn]₂E-Mes⁺ (E = Se, Te): Verbindungen vom "Iniden"-Typ (P. Lau, G. Huttner und L. Zsolnai), (440) 41

Cyclopentadienyl chromium chalcogenide complexes. Synthesis of [CpCr(EPh)₂]₂E (E = S, Se) and their Cr(CO)₅ adducts. Crystal structures of [CpCr(SePh)₂]Se · 1/2C₆H₆ and [CpCr(SePh)₂]Se [Cr(CO)₅] (L.Y. Goh, M.S. Tay, Y.Y. Lim, W. Chen, Z.-Y. Zhou and T.C.W. Mak), (441) 51

Umsetzungen der lithisierten Halbsandwich-Chalkogenole, (CO)₃MC₅H₄-ELi (M = Mn, Re; E = S, Se, Te) (M. Herberhold und M. Biersack), (443) 1

Stereochemistry of the compounds [Fe₃(CO)₉XY]. Crystal and molecular structures of the mixed triiron carbonyl compounds [Fe₃(CO)₉XY] (X, Y = S, Se, or Te) (G. Gervasio), (445) 147

Verbrückte Diarsinidenkomplexe {[Cp'(CO)₂Mn]₂As]₂E (E = O, S, Se) und deren nukleophile Spaltung (C. Emmerich, G. Huttner und A. Asam), (447) 71

Substitution, nukleophile und oxidative Addition an Arsinidenkomplexe. Zur Reaktion von [Cp'(CO)₂Mn]₂AsCl mit PhSeH und verwandten Chalkogenverbindungen (C. Emmerich und G. Huttner), (447) 81

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), (449) 167

Chromium

Metall- π -Komplexe von Benzolderivaten. XXXIX. Di[1,4-bis(diphenylphosphano)- η^6 -benzol]chrom: ein Baustein für polynukleare Bimetallkomplexe? (Ch. Elschenbroich, J. Sebbach, B. Metz and G. Heikenfeld), (426) 173

Synthesis of isomeric benchetroenic alkenes by Ti-induced carbonyl coupling reaction (J. Besançon, J. Szymoniak and C. Moise), (426) 325

A study of polarities, polarisabilities and infrared spectra of LM(CO)₅ complexes (M = Cr, Mo or W; L = trimethylamine, quinuclidine, pyridine, *p*-methyl- or *p*-*t*-butylpyridine) and of the X-ray crystal structure of quinuclidine-Cr(CO)₅: evidence for π -acceptor behaviour of coordinated pyridine (M.J. Aroney, R.M. Clarkson, T.W. Hambley and R.K. Pierens), (426) 331

Metallorganische Lewis-Säuren. XLIV. Rheniooxy(organyl)carben-Komplexe von Chrom, Molybdän und Wolfram (P.M. Fritz, J. Breimair, B. Wagner and W. Beck), (426) 343

Synthese und Kristallstrukturanalyse von Alkinylcarbenkomplex-verbrückten Dicobalthexacarbonylkomplexen (A. Wienand, H.-U. Reissig, H. Fischer, D. Pflumm and C. Troll), (427) C9

- Reaktionen von Nitrosylkomplexen. XI. Überführung von μ -NO- in μ -NH₂-Liganden durch katalytische Hydrierung am Beispiel von [Cp'Co(μ -NO)]₂ sowie von [CpCr(μ -NO)NO]₂ (J. Müller, I. Sonn und M. Strampfer), (427) C15**
- Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XXXIX. Synthese und dynamisches Verhalten von Tetracarbonyl- η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien-metall(0)- und Dicarbonyl-bis(η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien)metall(0)-Komplexe der Elemente Chrom, Molybdän und Wolfram (S. Özkar, N.K. Tunali and C.G. Kreiter), (428) 345**
- Synthèse de chalcones à base de tétrathiafulvalène: RC(=O)CH=C(CH₃)TTF, TTFC(=O)CH=C(CH₃)R (R = TTF: C₆H₃S₄; Bct: C₆H₅Cr(CO)₃; Fc: C₅H₄FeC₅H₅). Propriété conductrice d'un nouveau complexe de transfert de charge TTFC(=O)CH=C(CH₃)TTF · TCNQ (J. Besançon, A. Radecki-Sudre, J. Szymoniak and J. Padiou), (429) 335**
- Vapor-phase electronic absorption spectrum of (η^5 -cyclopentadienyl)(η^5 -cycloheptatrienyl)chromium (S.Y. Ketkov), (429) C38**
- Synthese und strukturelle Charakterisierung von bcpm-Metallcarbonyl-Dimeren (bcpm = Biscyclopentadienylmethan) mit Metallen der 6. Nebengruppe (H. Brunner, R. Graßl, J. Wachter, B. Nuber and M.L. Ziegler), (431) 143**
- Untersuchungen zum Einfluß des Biscyclopentadienylmethanliganden auf die Bildung von Cyclopentadienylchromsulfiden (H. Brunner, R. Graßl and J. Wachter), (431) 151**
- Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XL. Synthese und dynamisches Verhalten von Tricarbonyl- η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien-trimethylphosphit-metall(0)- und Dicarbonyl- η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien-bis(trimethylphosphit)metall(0)-Komplexen der Elemente Chrom, Molybdän und Wolfram (C.G. Kreiter, M. Kotzian, S. Özkar and I. Abu-Mour), (431) 159**
- Dihydropyridine-induced reductions of methyl(ethoxy)carbene complexes of chromium and tungsten: formation and X-ray crystal structure of the adduct between pyridine and the corresponding ethylidene carbene complexes (F. Cohen, R. Goumont, H. Rudler, J.C. Daran and R.A. Toscano), (431) C6**
- Mono et dinitrosation en position benzylque des complexes du benzène chrometricarbonyle. Synthèse directe d'oximes et d'hydroximates (M.-C. Senechal-Tocquer, D. Senechal, J.-Y. Le Bihan, D. Gentric, B. Caro, M. Gruselle et G. Jaouen), (433) 261**
- Homoleptische metallorganische Innerkomplexe des 8-Dimethylaminonaphthyl-1-Anions (H. Drews), (433) C1**
- Einfluß verschiedenartig substituierter Cyclopentadienylliganden auf die Chemie schwefelreicher Organometallkomplexe mit Metallen der 6. Nebengruppe (H. Brunner, R. Graßl, W. Meier, J. Wachter, B. Nuber und M.L. Ziegler), (434) 63**
- Übergangsmetall-Stannyl-Komplexe. V. Darstellung der anionischen Stannyl-Komplexe K[(π -Aromat)(CO)₂CrSnPh₃]⁻ und Reaktion mit Organozinnhalogeniden zu Bis(stannyl)-Komplexen (π -Aromat)(CO)₂Cr(SnPh₃)(SnR₃) (π -Aromat = C₆H_{6-n}Me_n) (U. Schubert und J. Schubert), (434) 169**
- Synthese und Charakterisierung von Hydridotrispyrazolylborat-Heterobimetall-Komplexen mit Titan und Molybdän (J. Ipaktschi und W. Sulzbach), (434) 287**
- Symmetric versus unsymmetric ligand orientations in (pentadienyl)Cr(I or II)(L)₂(L') species, and a spontaneous reduction of Cr^{II} to Cr^I (T.D. Newbound, A.M. Arif, D.R. Wilson, A.L. Rheingold and R.D. Ernst), (435) 73**
- Pentacarbonylchromium complexes containing the fluorinated isocyanide ligands H₂FCNC and HF₂CNC (D. Lentz and D. Preugschat), (436) 185**
- Reactions of silyl enol ethers with [ethoxy(phenylethylnyl) carbene]chromium and -tungsten complexes (F. Camps, L. Jordi, J.M. Moretó, S. Ricart, A.M. Castaño and A.M. Echavarren), (436) 189**
- Oxidative addition reactions of [Os₃(CO)₁₀(MeCN)₂] with 2-imine-substituted derivatives of furan, thiophene, selenophene and pyrrole (A.J. Arce, Y. De Sanctis, L. Hernandez, M. Marquez and A.J. Deeming), (436) 351**
- Static and dynamic structures of pentacarbonyl-chromium(0) and -tungsten(0) complexes of dithioether ligands. I. Symmetrical dithioether complexes and X-ray crystal structure of [W(CO)₅(¹BuSCH₂S^tBu)] (E.W. Abel, K.G. Orrell, H. Rahoo, V. Šík, M.A. Mazid and M.B. Hursthouse), (437) 191**
- { η^6 -aren(CO)₂Cr} als metallorganische 16-Elektronenkomplexfragmente: Synthese und Eigenschaften von Stibinidenkomplexen [{ η^6 -aren(CO)₂Cr}₂SbX] (F. Bringewski, G. Huttner, W. Imhof und L. Zsolnai), (439) 33**
- Reaktionen von Komplexliganden. LI. Sechsgliedrige Aminocarbon-Chelatkomplexe: Synthese, Alkin/Carben-Verknüpfung und Carbenanellierung (K.H. Dötz, A. Rau und K. Harms), (439) 263**
- η^1 - und η^2 -Komplexe von Cp*-substituierten Diphosphinen und Arsaphosphinen: Synthese, Struktur und Umlagerung unter Cp*-Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammmer), (439) 279**
- Studies on olefin-coordinating transition metal carbene complexes. XIX. Synthesis and structural characterization of the cycloalkene ligated carbonyl-chromium and -tungsten carbene complexes (J. Chen and B. Wang), (440) 67**
- Conjugate additions to alkynylalkoxycarbenemetal (Cr or W) complexes (A. Llebaria, J.M. Moretó, S. Ricart, J. Ros, J.M. Viñas and R. Yáñez), (440) 79**
- Wittig and Reformatsky chemistry of the formyl and acetyl derivatives of (η^5 -C₅H₅)Mn(CO)₃, (η^5 -C₅H₅)Re(CO)₃ and (η^6 -C₆H₆)Cr(CO)₃ (T.E. Bitterwolf and X. Dai), (440) 103**
- Electron transfer catalyzed substitution in carbonyl complexes. VI. Highly variable rates of substitution by tetracyanoethylene (B. Schwederski, W. Kaim, B. Olbrich-Deussner and T. Roth), (440) 145**
- P₄-Liganden mit maximaler Elektronendonorfähigkeit. III. Zum Reaktionsverhalten von Phosphankomplexen des Platin mit P₄-Phosphor (M. Scheer, M. Dargatz und A. Rufińska), (440) 327**
- Cyclopentadienyl chromium chalcogenide complexes. Synthesis of [CpCr(EPh)₂]E (E = S, Se) and their Cr(CO)₅ adducts. Crystal structures of [CpCr(SePh)₂]Se · 1/2C₆H₆ and [CpCr(SePh)₂]Se [Cr(CO)₅] (L.Y. Goh, M.S. Tay, Y.Y. Lim, W. Chen, Z.-Y. Zhou and T.C.W. Mak), (441) 51**
- Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241**
- Static and dynamic structures of pentacarbonyl-chromium(0) and -tungsten(0) complexes of dithioether ligands. II. Unsymmetrical dithioether ligand complexes of general type [M(CO)₅(MeSCH₂SR)] (E.W. Abel, K.G. Orrell, H. Rahoo and V. Šík), (441) 255**
- Stereoselektive Additionsreaktionen von Komplex-gebundenen Phosphanen und Arsanen an Alkine (H. Lang und U. Lay), (441) 389**
- Kohlenwasserstoffverbrückte Komplexe. XXVI. Reaktionen von anionischen Aromatenkomplexen [(OC)₃Cr(Ar)]⁻ mit metallorganischen Elektrophilen. Darstellung von (OC)₃Cr(μ - η^6 : η^6 -C₆H₅-C₇H₇)M(CO)₃ (M = Cr, Mo), (OC)₃Cr(μ - η^6 : η^5 -C₆H₅-C(O)C₅H₄)Fe(η^5 -C₅H₅), (OC)₃Cr(μ - η^6 : η^6 -C₆H₅S-C₇H₇)Mo(CO)₃, (OC)₃Cr(μ - η^6 : η^6 -C₉H₇-C₇H₇)Mo(CO)₃, (OC)₃Cr(μ - η^6 : η^6 -C₁₃H₉-C₇H₇)M(CO)₃ und (OC)₃Cr(μ - η^6 : η^6 -C₁₃H₁₁-C₇H₇)M(CO)₃ (M = Cr, Mo, W) (J. Breimair, M. Wieser und W. Beck), (441) 429**
- Static and dynamic structures of pentacarbonyl-chromium(0) and tungsten(0) complexes of dithioether ligands. III. 1,3-Metallo-**

- tropic shifts of sulphur coordination sites (E.W. Abel, K.G. Orrell, H. Rahoo and V. Šik), (441) 441
- Antiferromagnetic complexes with metal–metal bonds. XXVI. Synthesis, molecular structure and magnetic properties of mixed-metal triangular clusters, $\text{Cp}_2\text{Cr}(\mu\text{-SR})(\mu_3\text{-S})_2\text{FeCp}$ ($\text{R} = \text{CMe}_3$ or Ph), and linear mixed-metal chain $\text{CpCr}(\mu\text{-SPh})_3\text{Fe}(\mu\text{-SPh})_3\text{CrCp}$ (S.E. Nefedov, A.A. Pasynskii, I.L. Eremenko, G.S. Gasanov, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky and Y.T. Struchkov), (443) 101
- Synthesis, structure, and chemistry of heterobimetallic phosphido-bridged complexes $\text{CpFe}(\text{CO})_2(\mu\text{-PPPh}_2)\text{M}(\text{CO})_5$ ($\text{M} = \text{Cr, Mo, W}$) (S.-G. Shyu, P.-J. Lin and Y.-S. Wen), (443) 115
- Übergangsmetall-substituierte Diphosphene. XXX. η^1 - und η^2 -Komplexe der Metallodiphosphene ($\eta^5\text{-L}(\text{CO})_2\text{Fe-P=P-Mes}^*$ ($\text{L} = \text{C}_5\text{Me}_5$, $\text{C}_5\text{Me}_4\text{Et}$, $\text{C}_5\text{Me}_4^n\text{Bu}$, $1,3\text{-}^1\text{Bu}_2\text{C}_5\text{H}_3$; $\text{Mes}^* = 2,4,6\text{-}^1\text{Bu}_3\text{C}_6\text{H}_2$). Röntgenstrukturanalyse von $\text{Pt}(\eta^2\text{-}[(\eta^5\text{-C}_5\text{Me}_4\text{Et})(\text{CO})_2\text{Fe-P=P-Mes}^*]\text{PPh}_3)_2$ (L. Weber, I. Schumann, H.-G. Stammmer and B. Neumann), (443) 175
- Synthesis and characterization of some new cobaloximes, including the first members of a new class of heterobimetallic cobaloxime complexes. Crystal structure of $[(\eta^6\text{-C}_6\text{H}_5\text{CH}_2\text{Co}(\text{DH})_2(\text{py}))\text{Cr}(\text{CO})_3]$ (M.R. Domingo, A. Irving, Y.-H. Liao, J.R. Moss and A. Nash), (443) 233
- Additionsreaktionen von $(\text{CO})_5\text{CrEPhH}_2$ ($\text{E} = \text{P, As}$) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), (444) 53
- Subvalent Group 14 metal compounds. XV. Germanium and tin amides $\text{M}(\text{NR}_2)_2$ as ligands in carbonylchromium(0) chemistry. The crystal structure of the compounds $cis\text{-}[\text{Cr}(\text{CO})_4\{\text{Sn}(\text{NR}'_2)_2\}_2]$ [$\text{M} = \text{Ge or Sn}$, $\text{R}' = \text{SiMe}_3$ or $\text{NR}'_2 = \text{NCMe}_2(\text{CH}_2)_3\text{CMe}_2$] (S.L. Ellis, P.B. Hitchcock, S.A. Holmes, M.F. Lappert and M.J. Slade), (444) 95
- Antiferromagnetic complexes with metal–metal bonds. XXVII. Synthesis, molecular structures and magnetic properties of Cr,Re-clusters $\text{Cp}_2\text{Cr}(\mu\text{-SCMe}_3)_2(\mu_3\text{-S})\text{Re}_2(\text{CO})_9$ and $\text{Cp}_2\text{Cr}(\mu_3\text{-SCMe}_3)_2(\mu_3\text{-S})_2\text{Re}_2(\text{CO})_6$ (A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, B. Orzasakhatov, A.A. Zharkikh, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky and Y.T. Struchkov), (444) 101
- Umsetzungen von Metallorganylen mit dem capto-dativ substituierten Ethylen $\text{H}_2\text{C=C}(\text{CN})(\text{S}^1\text{Butyl})$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammmer und M.L. Ziegler), (444) 107
- Trapping of a carbanionic intermediate in the reaction of alkynyl-alkoxy carbene metal (Cr) complexes and ethyl diethoxyacrylate (L. Jordi, J.M. Moretó, S. Ricart, J.M. Viñas, E. Molins and C. Miravittles), (444) C28
- Dichromastibacumulenium-Ionen $[\eta^6\text{-aren} = (\text{CO})_2\text{Cr=Sb=Cr}(\text{CO})_2\text{aren-}\eta^6]^{+}$: Stabilität in Abhängigkeit von η^6 -gebundenen Arenliganden am Chromatom (F. Bringewski, G. Huttner und L. Zsolnai), (447) 233
- ^{13}C and ^{19}F NMR study on the structure and conformation of tricarbonylchromium complexes of biphenyl derivatives (P. Szczeciński and J. Zachara), (447) 241
- Group IV metal–chromium complexes bridged by a benzoate group (H.-M. Gau, C.-T. Chen, T.-T. Jong and M.-Y. Chien), (448) 99
- Cobalt**
- Neuartige Synthesen monosubstituierter Nickelocene als Zugang zu mehrkernigen Metallocensystemen (H. Schottenberger, G. Ingram and D. Obendorf), (426) 109
- Selective homolytic reactions of alkylcobaloxime with nitrogen monoxide (M. Kijima, H. Yamashita and T. Sato), (426) 399
- Synthese und Kristallstrukturanalyse von Alkinylcarbenkomplexver-
- brückten Dicobalthexacarbonylkomplexen (A. Wienand, H.-U. Reissig, H. Fischer, D. Pflumm and C. Troll), (427) C9
- Reaktionen von Nitrosylkomplexen. XI. Überführung von $\mu\text{-NO}$ in $\mu\text{-NH}_2$ -Liganden durch katalytische Hydrierung am Beispiel von $[\text{Cp}'\text{Co}(\mu\text{-NO})_2]$ sowie von $[\text{CpCr}(\mu\text{-NO})\text{NO}]_2$ (J. Müller, I. Sonn and M. Strampfer), (427) C15
- Charge-transfer salts of carbonylmetalates as outer-sphere ion pairs in photochemical and thermal electron transfer (C.-H. Wei, T.M. Bockman and J.K. Kochi), (428) 85
- Preparation and properties of inclusion compounds of cobalt cluster complexes with γ -cyclodextrin (M. Shimada, A. Harada and S. Takahashi), (428) 199
- Two-dimensional nutation ^{59}Co NMR of solid $\text{Co}_4(\text{CO})_{12}$ (T. Eguchi, H. Nakayama, H. Ohki, S. Takeda, N. Nakamura, S. Kernaghan and B.T. Heaton), (428) 207
- Rupture of the triple $\text{C}\equiv\text{C}$ and $\text{C}\equiv\text{O}$ bonds in the mixed-metal clusters with Ni_2Mo_2 and Mo_4Co_3 cores. Synthesis and molecular structures of $\text{Cp}_4\text{Ni}_2\text{Mo}_2(\mu_3\text{-CPh})_2(\mu_3\text{-CO})_2$ and $\text{Cp}_4\text{Mo}_4(\mu_3\text{-OH})(\mu_3\text{-CPh})_2(\mu_6\text{-C})\text{Co}_3(\mu\text{-CO})_3(\text{CO})_3$ (A.D. Shaposhnikova, M.V. Drab, G.L. Kamalov, A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, Y.T. Struchkov and A.I. Yanovsky), (429) 109
- Organoerdmetallkomplexe von *d*-Block-Elementen. IV. "Ein-Topf-Synthese" flüchtiger Organogalliocobalt-Komplexe, Molekülstruktur von $\{\mu\text{-[3-(Diethylamino)propyl]gallio}\text{-[bis(tetracarbonylcobalt)]}$ (R.A. Fischer, J. Behm and T. Priermeier), (429) 275
- Metal–metal bonds extended over a porphyrin ring. III. Syntheses and spectral studies of new metal–metal bonded indium porphyrin complexes, (TBPP) In-ML and (SFP) In-ML (S. Takagi, Y. Kato, H. Furuta, S. Onaka and T.K. Miyamoto), (429) 287
- Synthesis, characterization, crystal structure, and solution conformation of $(-\text{)}_{436}(\text{S}_{\text{Co}}\text{S}_{\text{C}})(\eta^5\text{-C}_9\text{H}_7)\text{Co}^*(\text{C}_3\text{F}_7)(\text{Ph}_2\text{PNHC}^*\text{H-(CH}_3\text{)Ph})$, an easily resolvable chiral-at-metal complex (C. Jablonski, Z. Zhou and J.N. Bridson), (429) 379
- Photochemical and thermal disproportionation of $[(\text{CO})_4\text{CoM}(\text{CO})_3(\text{LL})]$ ($\text{M} = \text{Mn, Re}$; $\text{LL} = 2,2'\text{-bipyridine, 2-pyridinecarbaldehyde N-isopropylimine}$) complexes (H. Knoll, W.J.G. De Lange, H. Hennig, D.J. Stufkens and A. Oskam), (430) 123
- Sulphur dioxide insertion into organocobaloximes: a true insertion or a case of radical chain process? (B.D. Gupta, M. Roy, M. Oberoi and V. Dixit), (430) 197
- Further synthetic and structural studies on cobalt carbonyl containing antimony complexes (N.C. Norman, P.M. Webster and L.J. Farrugia), (430) 205
- S_2CPR_3 adducts as bridging ligands in tricobalt clusters. X-Ray structure of $[\text{Co}_3(\text{CO})_7(\mu_3\text{-CH})(\mu_2\text{-S}_2\text{CPCy}_3)]$ (B. Alvarez, D. Miguel, V. Riera and S. García-Granda), (430) C19
- Adduct between quadricyclane and $(\eta^5\text{-cyclopentadienyl})(1,2\text{-benzenedithiolato})\text{cobalt(III)}$: formation and structure (M. Kajitani, H. Hatano, T. Fujita, T. Okumachi, H. Nagao, T. Akiyama and A. Sugimori), (430) C64
- Optically active complexes of transition metals (Rh^{I} , Ru^{II} , Co^{II} and Ni^{II}) with 2-aminocarbonylpyrrolidine ligands. Selective catalysts for hydrogenation of prochiral olefins (A. Corma, M. Iglesias, C. Del Pino and F. Sánchez), (431) 233
- Use of metal complexation in the synthesis of medium-ring acetylenic lactones (S.D. Najdi, M.M. Olmstead and N.E. Schore), (431) 335
- Camphor-derived alcohols as chiral auxiliaries for asymmetric Paason-Khand bicyclizations. Enantioselective synthesis of α -methoxyenones (X. Verdaguera, A. Moyano, M.A. Pericàs, A. Riera, A.E. Greene, J.F. Piniella and A. Alvarez-Larena), (433) 305

- Homoleptische metallorganische Innerkomplexe des 8-Dimethylaminonaphthal-1-Anions (H. Dreys), (433) C1
- Reactivity of main-group–transition-metal bonds. X. The kinetics of iodination of compounds containing one, two, three or four tin–cobalt bonds (J.R. Chipperfield, D.E. Webster and H. Yusof), (434) 53
- Mercury-bridged transition-metal clusters. Synthesis of pentanuclear Ru_3HgM ($M = \text{Mo}$, W , or Co) clusters and X-ray structure of $[(\text{Ru}_3(\mu_3,\eta^2\text{-ampy})(\text{CO})_9)\text{-}(\mu_3\text{-Hg})\text{Co}(\text{CO})_4]$ (Hampy = 2-amino-6-methylpyridine) (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, S. García-Granda and J.F. Van der Maelen), (434) 123
- Reactions of 1,3,5-trithiacyclohexane with tetrahedral ruthenium–cobalt clusters (S. Rossi, J. Pursiainen and T.A. Pakkanen), (436) 55
- Crystal and molecular structure of $[\text{Co}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ (M.Y. Antipin, Y.T. Struchkov, A.N. Cherneva, M.F. Meidine and J.F. Nixon), (436) 79
- The reaction of diphenylphosphine, PPPh_2H , with an alkyne-bridged dicobalt carbonyl complex; the synthesis crystal structure and reactivity of the complex $[\text{Co}_2(\mu\text{-C}(\text{CO}_2\text{Me})=\text{CHCO}_2\text{Me})(\mu\text{-PPPh}_2\text{H})(\text{CO})_4]$ (A.J.M. Caffyn, M.J. Mays, G. Conole, M. McPartlin and H.R. Powell), (436) 83
- Cobalcarbonyl-Komplexe von Alkinyl-Silanen (H. Lang, U. Lay und M. Weinmann), (436) 265
- Alkylcobalamins: formation by enantioselective alkylation of cob(I)alamin, ^1H NMR spectra, and conformational analysis of the alkyl group (R.J. Anderson, R.M. Dixon and B.T. Golding), (437) 227
- Synthese, Charakterisierung und Reaktivität heterodinukleärer Komplexe: $[(\text{CO})_4\text{Fe}(\mu\text{-P}^t\text{Bu}_2)\text{Rh}(\text{CO})\text{L}](\text{Fe}-\text{Rh})$ ($L = \text{CO}$, HP^tBu_2), $[(\text{CO})_3\text{Co}(\mu\text{-H})(\mu\text{-P}^t\text{Bu}_2)\text{Rh}(\text{CO})(\text{HP}^t\text{Bu}_2)](\text{Co}-\text{Rh})$ (B. Walther, H.-C. Böttcher, M. Scheer, G. Fischer, D. Fenske und G. Süss-Fink), (437) 307
- Syntheses and characterization of tri-metal cluster compounds; X-ray structure determination of $[\text{Co}_2\text{Mo}(\mu_3\text{-CC}_6\text{H}_5)\text{Br}(\text{CO})_8(\text{bipy})]$ ($\text{bipy} = \alpha,\alpha\text{-dipyridyl}$) (Y. Yin, H. Sun, D. Jin and Y. Zhu), (439) 45
- The synthesis of dimetallic thioalkyne complexes of cobalt and molybdenum (S.C. Bennett, A. Gelling and M.J. Went), (439) 189
- Synthesis and characterization of cobalt and molybdenum complexes derived from linear conjugated diynes, triynedienes and tetracynetrienes (W.E. Lindsell, P.N. Preston and P.J. Tomb), (439) 201
- Cycloadditionsreaktionen von Organometall-Komplexen. XV. [2 + 3]-und [2 + 1]-Cycloadditionsreaktionen von Carbonylrhodium- und Carbonylcobalt-Komplexen mit Nitriloxiden und Aziden. Die Kristall- und Molekülstruktur von $\text{C}_5\text{H}_5(\text{P}^t\text{Pr}_3)\text{RhC}(\text{O})\text{ONC-C}_6\text{H}_5$ (H. Werner, U. Brekau, O. Nürnberg und B. Zeier), (440) 389
- The preparation and structures of two Ge–Co–Fe anionic clusters; $[\text{GeCoFe}_3(\text{CO})_{14}]^-$ and $[\text{Ge}_2\text{Co}_5\text{Fe}_2(\text{CO})_{22}]^-$ (S.G. Anema, G.C. Barris, K.M. Mackay and B.K. Nicholson), (441) 35
- Chiral cyclopentadiene–cyclopentadienyl complexes of cobalt (V.P. Maryin and P.V. Petrovskii), (441) 125
- Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241
- Stereoselektive Additionsreaktionen von Komplex-gebundenen Phosphanen und Arsanen an Alkine (H. Lang und U. Lay), (441) 389
- A low-valence cobalt cluster formed via capture of sulphur atom from ethanedithiol – synthesis and crystal structure of $[\text{Co}_7(\mu_4\text{-S})_3(\mu_3\text{-S})_3(\text{PPh}_3)_6\text{Br}]\cdot\text{CH}_3\text{OH}$ (F. Jiang, X. Lei, M. Hong, Z. Huang, B. Kang and H. Liu), (443) 229
- Synthesis and characterization of some new cobaloximes, including the first members of a new class of heterobimetallic cobaloxime complexes. Crystal structure of $[(\eta^6\text{-C}_6\text{H}_5\text{CH}_2\text{Co}(\text{DH})_2(\text{py})\text{Cr}(\text{CO})_3)]$ (M.R. Domingo, A. Irving, Y.-H. Liao, J.R. Moss and A. Nash), (443) 233
- Preparation and properties of cyclodextrin–metal carbonyl inclusion compounds (M. Shimada, Y. Morimoto and S. Takahashi), (443) C8
- Synthesis and X-ray structure of a highly-hindered N -functionalized alkyl-cobalt(II) complex $\text{trans-}[\text{Co}(\text{C}(\text{SiMe}_3)_2\text{C}_5\text{H}_4\text{N}-2)]$ (W.-P. Leung, H.-K. Lee, Z.-Y. Zhou and T.C.W. Mak), (443) C39
- Umsetzungen von Metallorganylen mit dem capto-dativ substituierten Ethylen $\text{H}_2\text{C}=\text{C}(\text{CN})(\text{S}^t\text{Butyl})$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammel und M.L. Ziegler), (444) 107
- Ein- bis dreikernige Metall- π -Derivate von Hericen mit $(\text{Me}_3\text{P})_3\text{Fe}$ und CpCo -Fragmenten (F.H. Köhler und A. Steck), (444) 165
- Formation of $(\eta^2\text{-O}_2)\text{Co}(\text{CO})$ and $(\eta^2\text{-O}_2)\text{Co}(\text{CO})_2$ by photolysis of $\text{Co}_2(\text{CO})_8$ in dioxygen matrices at 20 K (M.J. Almond and R.H. Orrin), (444) 199
- Synthesis and crystal structure of $[\text{Co}_2(\text{CO})_4(\text{PhC}\equiv\text{CC}(\text{O})\text{CH}_3)_3]$. Its role in the cyclotrimerization of 1-phenylbut-1-yn-3-one to 1,3,5-triphenyltris(carboxymethyl)benzene (G. Gervasio, E. Sappa and L. Markó), (444) 203
- Synthesis of the square-bipyramidal cluster $[\text{Co}_4(\mu_4\text{-SiMe}_2)_2(\text{CO})_{11}]$ by two routes and its reaction with GeMe_2H_2 . The crystal structures of $[\text{Co}_4(\mu_4\text{-SiMe}_2)_2(\text{CO})_{11}]$ and $[\text{Co}_4(\mu_4\text{-SiMe}_2)_2(\mu\text{-GeMe}_2)_2(\text{CO})_{10}]$ (S.G. Anema, S. Kim Lee, K.M. Mackay and B.K. Nicholson), (444) 211
- Ligand exchange in cyclopentadienylcobalt diselenolanes. Isolation and characterization of the Se_3 -chelate, $\text{C}_5\text{H}_5\text{Co}[(\text{SeC}_8\text{H}_{12})_2\text{Se}]$ (C.P. Morley and R.R. Vaughan), (444) 219
- Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), (444) C15
- 1,1'-Bis(diphenylphosphino)ferrocene ligand substitution in the benzylidyne-capped cluster $\text{PhCCo}_3(\text{CO})_9$. Synthesis, X-ray structure, and redox reactivity of $\text{PhCCo}_3(\text{CO})_7(\text{dpfp})$ (W.H. Watson, A. Nagl, S. Hwang and M.G. Richmond), (445) 163
- Coordination of a tripodal phosphine sulphide P_2S towards various transition metal ions (H.-E. Wang, M.-C. Cheng, G.-H. Lee, S.-M. Peng and S.-T. Liu), (445) 171
- Synthetic pathways for selective introduction of a dicobalt hexacarbonyl cluster into a polyfunctional molecule: methotrexate (M. Salman and G. Jaouen), (445) 237
- Sterically crowded $\text{PhCCo}_3(\text{CO})_9$ derivatives: X-ray structures of the mono- and bis-diphenylphosphinomethane substitution products (A.J. Downard, B.H. Robinson and J. Simpson), (447) 281
- Relative metal-ion affinities of alkylamines in the gas-phase (L.-Z. Chen and J.M. Miller), (448) 225
- Formation of a high surface area, regular porous solid from the cluster of clusters, $\text{Zn}_4^{\text{II}}\text{O}[(\text{CO})_9\text{Co}_3\text{CCO}_2]_6$ (W. Cen, B. Ladna, T.P. Fehlner, A.E. Miller and D. Yue), (449) 19
- Study of CoGa deposition from the single source precursor $(\text{CO})_4\text{CoGaCl}_2$ (THF) (F. Maury, L. Brandt and H.D. Kaesz), (449) 159
- The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), (449) 167
- Reactivity of tri- and tetra-nuclear ethylidyne cyclopentadienylcobalt⁺ clusters towards protonic acids (H. Wadeohl and H. Pritzlow), (450) 9
- Formation of the unusual alkynyl carboxamide complex, $\text{C}_2\text{H}_5\text{N}(\text{H})\text{C}(\text{O})\text{C}\equiv\text{QCo}_2(\text{CO})_6\text{CCo}_3(\text{CO})_9$ (G.H. Worth, B.H. Robinson and J. Simpson), (450) 219

Hydroformylation and hydrocarbethoxylation of 1,2-dicarbethoxy-1,2,3,6-tetrahydropyridazine and 1,2-dicarbethoxy-1,2,3,4-tetrahydropyridazine (G. Menchi, S. Paganelli, U. Matteoli, A. Scrivanti and C. Botteghi), (450) 229

Copper

Substitution of the acetoxy groups of dialkoxyacetates by organometallic reagents: a route to allyl-, propargyl-, homoallyl-, homopropargyl- and α -stannylacetals (I. Beaudet, A. Duchêne, J.-L. Parrain and J.-P. Quintard), (427) 201

Fast-atom bombardment (FAB) mass spectra of coordination complexes: aggregation processes in the mass spectra of Group 11 alkynyl complexes (M.I. Bruce and M.J. Liddell), (427) 263

Chemie polyfunktioneller Moleküle. CIX. Synthese, Molekül- und Kristallstruktur des über ($\text{NH} \cdots \text{O}$)- und ($\text{OH} \cdots \text{NC}$)-Wasserstoffbrückenbindungen verknüpften tetraedrischen Chelatkomplexes $[\text{Cu}(\text{CN})(\text{Ph}_2\text{P}-\text{NH}-\text{PPH}_2)\text{PPh}_3] \cdot \text{CH}_3\text{OH}$ (J. Ellermann, F.A. Knoch, K.J. Meier and M. Moll), (428) C44

Trimethylsilylcyclooctadiene-transition metal complexes: metal-catalysed protodesilylation of cyclic vinylsilanes, and transfer hydrogenation promoted by the displaced silyl group (B.S. Bandodkar and G. Nagendrappa), (430) 373

Palladium- and light-enhanced ring-opening of oxiranes by copper chloride (J. Muzart and A. Riahi), (433) 323

Ethylnlfrocene durch direkte Einführung von Acetylenen (M. Buchmeiser und H. Schottenberger), (436) 223

Chemistry of iridium carbonyl cluster complexes. Synthesis, chemical characterization and solid state structure of the mixed-metal carbonyl cluster $[\text{PPh}_4]^-\text{Ir}_6(\mu\text{-CO})_3(\text{CO})_{12}(\mu_3\text{-Cu}(\text{NCMe})) \cdot \text{CH}_2\text{Cl}_2$ and its conversion to the dianion $[\text{Ir}_{12}(\text{CO})_{26}]^{2-}$ (R. Della Pergola, L. Garlaschelli, F. Demartin, M. Manassero and N. Masciocchi), (436) 241

Stereoselective opening of chiral α -stannylacetals with organometallic reagents (J.-L. Parrain, J.-C. Cintrat and J.-P. Quintard), (437) C19

Synthesis, electrochemistry and complexation studies of new redox active bisferrocene acyclic and macrocyclic thioethers (P.D. Beer, J.E. Nation, M.E. Harman and M.B. Hursthouse), (441) 465

Asymmetric catalysis. LXXX. An optically-active tetrakispyrazolylborate: Synthesis and use in Cu-catalysed enantioselective cyclopropanation (H. Brunner, U.P. Singh, T. Boeck, S. Altmann, T. Scheck and B. Wrackmeyer), (443) C16

Exclusive β -hydrosilylation of acrylates catalyzed by copper-tetramethylethylenediamine (P. Boudjouk, S. Kloos and A.B. Rajkumar), (443) C41

A novel high nuclearity bimetallic cluster: the crystal structure of $[\text{N}(\text{PPh}_3)_2^-\text{Ag}_6\text{Cu}_7(\text{C}_2\text{Ph})_{14}]$ (M.S. Hussain and O.M. Abu-Salah), (445) 295

Chemistry of copper(I) β -diketonate complexes. VI. Synthesis, characterization and chemical vapor deposition of 2,2-dimethyl-6,6,7,7,8,8,8-heptafluoro-3,5-octanedione (fod), copper(I)L complexes and the solid state structure of (fod) $\text{Cu}(\text{PMe}_3)$ (K.M. Chi, T.S. Corbitt, M.J. Hampden-Smith, T.T. Kodas and E.N. Duesler), (449) 181

A comparison of heterometallic alkoxide molecules containing copper(I) or copper(II) and zirconium (B.A. Vaartstra, J.A. Samuels, E.H. Barash, J.D. Martin, W.E. Streib, C. Gasser and K.G. Caulton), (449) 191

Erbium

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente. XXXIII. Esteraddukte des Grundkörpers Tris(η^5 -cyclopentadienyl)-lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Europium

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

The first examples of intermolecular weak (agostic) γ -methyl–metal interactions in organolanthanide complexes: the synthesis and X-ray structures of $[(\text{Yb}(\eta\text{-Cp})_2)_2]$ and $[(\text{Eu}(\eta\text{-Cp})_2)_2]$ [$\text{Cp} = \text{C}_5\text{H}_3(\text{SiMe}_3)_2\text{-}1,3$] (P.B. Hitchcock, J.A.K. Howard, M.F. LapPERT and S. Prashar), (437) 177

Gadolinium

Synthesis and crystal structure of $(\text{C}_9\text{H}_7\text{GdCl}_2 \cdot 3\text{THF})\text{THF}$ (F. Gao, G. Wei, Z. Jin and W. Chen), (438) 289

Gallium

Organoerdmetallkomplexe von d-Block-Elementen. IV. “Ein-Topf-Synthese” flüchtiger Organogalliocobalt-Komplexe, Molekülstruktur von $\{\mu\text{-[3-(Diethylamino)propyl]gallio}\text{-[bis(tetracarbonylcobalt)]}$ (R.A. Fischer, J. Behm and T. Priermeier), (429) 275

Unusual intramolecular hydrogen bonding in $\text{Cy}_2\text{GaBrNH}_2\text{Ph}$. The first structurally characterized dialkyl gallium halide primary amine adduct (D.A. Atwood, A.H. Cowley and R.A. Jones), (430) C29

Synthesis and characterization of alkylaluminum and -gallium derivatives of thiomorpholine. Molecular structures of $\text{Me}_3\text{Al} \cdot \text{N}(\text{H})\text{C}_4\text{H}_8\text{S}$ and $[\text{Et}_2\text{Al}(\mu\text{-NC}_4\text{H}_8\text{S})_2]$ (M. Taghiof, D.G. Hendershot, M. Barber and J.P. Oliver), (431) 271

Ligand redistribution reactions of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ (O.T. Beachley, Jr., T.L. Royster, Jr. and J.R. Arhar), (434) 11

Primary amide and amine complexes of gallium and indium: X-ray crystal structures of $[\text{Me}_2\text{Ga}(\mu\text{-NH}(\text{'Bu}))_2]$, $\text{Me}_3\text{Ga} \cdot \text{NH}_2(\text{'Bu})$ and $\text{Me}_3\text{In} \cdot \text{NH}_2(\text{'Bu})$ (D.A. Atwood, R.A. Jones, A.H. Cowley, S.G. Bott and J.L. Atwood), (434) 143

Organoerdmetall-Komplexe von d-Block-Elementen. V. Selektive Reduktion von Bis[carbonyl(η^5 -cyclopentadienyl)nickel] zum Carbonyl(η^5 -cyclopentadienyl)nickel-Anion durch C_8K . Synthese und Struktur von $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})\text{Ni} \cdot \text{Ga}[\text{CH}_2\text{C}(\text{CH}_3)_3]_2(\text{OC}_4\text{H}_8)$ (R.A. Fischer, J. Behm, E. Herdtweck und C. Kronseder), (437) C29

Organometallic precursors to the formation of GaN by MOCVD: structural characterisation of $\text{Me}_3\text{Ga} \cdot \text{NH}_3$ by gas-phase electron diffraction (M.J. Almond, C.E. Jenkins, D.A. Rice and K. Hagen), (439) 251

Synthesis of the first diarsadigallatane (D.A. Atwood, A.H. Cowley, R.A. Jones and M.A. Mardones), (439) C33

Bis(diisopropylgalliumdiisopropylphosphid) und -arsenid. Kristallstruktur des Phosphidderivats—ein “angedeuteter” butterfly (G.G. Hoffmann, R. Fischer, U. Schubert und B. Hirle), (441) 7

Synthesis and properties of pentafluorophenylgallium derivatives. The preparation of tris(pentafluorophenyl)gallium and its adducts, and of bis(pentafluorophenyl)gallium bromide, pentafluorophenylgallium dibromide, and tetrabutylammonium tetrakis(pentafluorophenyl)gallate (K. Ludovici, W. Tyrra and D. Naumann), (441) 363

Organometallic precursors for the formation of gallium nitride by metal organic chemical vapour deposition. Thermal decomposition of trineopentyl gallium and of its adducts with ammonia, dimethylamine and t-butylamine (M.J. Almond, C.E. Jenkins and D.A. Rice), (443) 137

Darstellung und Eigenschaften der t-Butylgalliumbromide und -iodide. Struktur von Di(t-butyl)galliumiodid (G.G. Hoffmann, D. Hellert und M. Warren), (444) 21

- Reaction of Me_3Ga and Me_3Al with the tertiary-tetraaza analog of adamantane, hexamethylenetetramine ($\text{N}_4\text{-Ada}$). Syntheses and molecular structures of $[(\text{Me}_3\text{M})_n \cdot \text{N}_4\text{-Ada}]$ ($n = 1\text{--}4$, M = Ga or Al) (J.B. Hill, S.J. Eng, W.T. Pennington and G.H. Robinson), (445) 11
- Tertiary phosphine adducts of alane and gallane (F.M. Elms, M.G. Gardiner, G.A. Koutsantonis, C.L. Raston, J.L. Atwood and K.D. Robinson), (449) 45
- Synthesis, characterization, and reactivity of new alkylgallium acetylides (K.E. Lee and K.T. Higa), (449) 53
- Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido ($[(^t\text{Bu})\text{PH}]^-$) and arsenido ($[(^t\text{Bu})\text{AsH}]^-$) units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), (449) 61
- Gallium and indium compounds containing three different substituents. Crystal and molecular structure of $[(\text{Me}_3\text{CCH}_2)\text{ClGa-PPPh}_2]_3$ (O.T. Beachley, Jr., J.D. Maloney and R.D. Rogers), (449) 69
- Study of CoGa deposition from the single source precursor $(\text{CO})_4\text{CoGaCl}_2$ (THF) (F. Maury, L. Brandt and H.D. Kaesz), (449) 159
- Über dimere Dimethylmetall-bis(trimethylgermyl)amide des Aluminiums, Galliums und Indiums (M. Trapp, H.-D. Hausen, G. Weckler und J. Weidlein), (450) 53
- ### Germanium
- Fast atom bombardment mass spectra of silatrane and silocanes (I. Mažeika, S. Grinberga, A.P. Gaukman, G.I. Zelchan and E. Lukevics), (426) 41
- Tetra-*p*-tolyl-Verbindungen *p*-Tol₄Si und *p*-Tol₄Ge: ein Beitrag zur Konfiguration der Tetraaryl-Methan-Analoga Ar₄M (M = C, Si, Ge, Sn, Pb) (M. Charissé, S. Roller and M. Dräger), (427) 23
- Synthesis and reactivity of germylliron complexes $[\text{Ph}_2\text{HGeFe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ and $[\text{RH}_2\text{GeFe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ (R = Ph or 2,4,6-Me₃C₆H₂) (A. Castel, P. Rivière, M. Ahbala, J. Satgé, M. Soufiaoui and N. Knouzi), (427) 91
- Reactions of pentafluorophenyl germanium hydrides with nickelocene (L.V. Pankratov, V.I. Nevodchikov, L.N. Zakharov, M.N. Bochkarev, I.V. Zdanovich, V.N. Latyaeva, A.N. Lineva, A.S. Batsanov and Y.T. Struchkov), (429) 13
- Sur la stabilisation d'une germa-imine par effet stérique et mésomérie (M. Rivière-Baudet and A. Morère), (431) 17
- Si-, Ge- and Sn-diine derivatives in 1,3-dipolar cycloaddition reactions (A.D. Jurayev, A.G. Makhsumov, K. Yuldasheva and M.A. Tolipova), (431) 129
- Gas phase ion-molecule reactions in methylgermane/oxygen, /ammonia, and /unsaturated hydrocarbon systems (L. Operti, M. Splendore, G.A. Vaglio, P. Volpe, M. Speranza and G. Occhipi), (433) 35
- Nouveaux arylidihydrogermyllithium (A. Castel, P. Rivière, J. Satgé et D. Desor), (433) 49
- Multinuclear NMR spectroscopic studies of some organometallic N-substituted anilines (B.K. Barr, A.J. Herman, L.K. Myers, P.I. Young, C.D. Schaeffer, Jr., H.J. Eppley, J.C. Otter and C.H. Yoder), (434) 45
- 3,5-Di-*tert*-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151
- Amination of vinylsilanes and vinylgermanes containing an activated double bond (V. Dirmens, D.I. Barabanov, E. Liepiņš, L.M. Ignatovich and E. Lukevics), (435) 257
- Addition of sulphenyl halides to alkenyl-metal compounds. III. Arenesulphenyl chloride additions to triethoxy- and triphenylvinylsilanes and to triphenylvinylgermanium. Crystal structure of $(4\text{-Me-2-O}_2\text{NC}_6\text{H}_3\text{S})\text{CH}(\text{SiPh}_3)\text{CH}_2\text{Cl}$ (R.A. Howie, G.M. Spencer, J.L. Wardell and J.N. Low), (437) 111
- Excimer laser photolysis of tetramethylgermane (J. Pola and R. Taylor), (437) 271
- Verbindungen des Germaniums und Zinns. XII. Ein Telluratrigerman durch Tellur-Insertion in ein Cyclotrigerman (M. Weidenbruch, A. Ritschl, K. Peters und H.G. von Schnerring), (437) C25
- Organoerdmetall-Komplexe von *d*-Block-Elementen. V. Selektive Reduktion von Bis[carbonyl(η^5 -cyclopentadienyl)nickel] zum Carbonyl(η^5 -cyclopentadienyl)nickel-Anion durch C₈K. Synthese und Struktur von $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})\text{Ni-Ga}[\text{CH}_2\text{C}(\text{CH}_3)_3]_2(\text{OC}_4\text{H}_8)$ (R.A. Fischer, J. Behm, E. Herdtweck und C. Kronseder), (437) C29
- Verbindungen des Germaniums und Zinns. XI. Neue viergliedrige Ringe: Thiatrigerman, Selenatrigerman und *N*-Phenyltrigermacyclobutanimin (M. Weidenbruch, A. Ritschl, K. Peters und H.G. von Schnerring), (438) 39
- Bis(4-fluorophenyl)methyl(1*H*-1,2,4-triazol-1-yl-methyl)germane, a germanium analogue of the agricultural fungicide flusilazole: synthesis and biological properties (R. Tacke, B. Becker, D. Berg, W. Brandes, S. Dutzmann and K. Schaller), (438) 45
- Synthesis of polymetallacycloalkanes (P. Mazerolles and C. Laurent), (439) 115
- ¹³C NMR studies of silyl and germyl complexes of rhodium and iridium containing carbonyl ligands (E.A.V. Ebsworth and S. Moreton), (439) 213
- The preparation and structures of two Ge-Co-Fe anionic clusters; $[\text{GeCoFe}_3(\text{CO})_{14}]^-$ and $[\text{Ge}_2\text{Co}_5\text{Fe}_2(\text{CO})_{22}]^-$ (S.G. Anema, G.C. Barris, K.M. Mackay and B.K. Nicholson), (441) 35
- Cleavage of Si-C and Ge-C bonds in heterysilanes and -germanes by organolithium reagents (V. Gevorgyan, L. Borisova and E. Lukevics), (441) 381
- Synthesis of stanna- and germa-cycloheptatrienes (Y. Nakadaira, R. Sato and H. Sakurai), (441) 411
- Etude comparative des réactions par transfert monoélectronique entre les germylamines primaires, secondaires et tertiaires et la 3,5-di-*tert*-butylorthoquinone (M. Riviere-Baudet, A. Morere et A. Khallaayoun), (443) 43
- Etude du transfert monoélectronique entre des germylanions et des piègeurs de spin dia et paramagnétiques (P. Rivière, A. Castel, D. Desor et C. Abdennadher), (443) 51
- Sila-Riechstoffe und Riechstoffisostere. XII. Geruchsvergleiche homologer Organoelementverbindungen der vierten Hauptgruppe (C, Si, Ge, Sn) (U. Wannagat, V. Damrath, V. Huch, M. Veith und U. Harder), (443) 153
- Subvalent Group 14 metal compounds. XV. Germanium and tin amides M(NR₂)₂ as ligands in carbonylchromium(0) chemistry. The crystal structure of the compounds *cis*-[Cr(CO)₄{Sn(NR'₂)₂}₂] [M = Ge or Sn, R' = SiMe₃ or NR₂ = NCMe₂(CH₂)₃CMe₂] (S.L. Ellis, P.B. Hitchcock, S.A. Holmes, M.F. Lappert and M.J. Slade), (444) 95
- Synthesis of the square-bipyramidal cluster $[\text{Co}_4(\mu_4\text{-SiMe})_2(\text{CO})_{11}]$ by two routes and its reaction with GeMe₂H₂. The crystal structures of $[\text{Co}_4(\mu_4\text{-SiMe})_2(\text{CO})_{11}]$ and $[\text{Co}_4(\mu_4\text{-SiMe})_2\mu\text{-GeMe}_2](\text{CO})_{10}$ (S.G. Anema, S. Kim Lee, K.M. Mackay and B.K. Nicholson), (444) 211
- Über gemischte Gruppe 14-Gruppe 14-Bindungen. V. Sechs Perphenylierte Propan-Analoga Ph₃M-M'Ph₂-M''Ph₃ (M, M', M'' = Si/Ge): ein Beitrag zum Elektronegativitätsunterschied zwischen Silicium und Germanium (M. Charissé, M. Mathes, D. Simon und M. Dräger), (445) 39
- Chemistry of heavy carbene analogues R₂M (M = Si, Ge, Sn). XVII. Reactions of free dimethylgermylene with H-acidic compounds and donors (B. Klein, W.P. Neumann, M.P. Weisbeck and S. Wienken), (446) 149
- Complexes du fer hydro-digermaniés: $[\text{R}_2\text{HGeGeR}_2\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2]$ (R = Et, Ph) — étude photolytique (A. Castel, P.

Rivière, M. Ahbala, J. Satgé, M. Soufiaoui et N. Knouzi), (447) 123

Sila et germacyclopentanes C-bifonctionnels. I. 1-Sila et 1-germacyclopentan-2,3 et 3,4-diols (G. Manuel et R. Boukherroub), (447) 167

Über gemischte Gruppe 14–Gruppe 14-Bindungen. VII. Ein Inkrementensystem für ^{13}C -NMR chemische Verschiebungen bei arylierten Gruppe 14-Verbindungen (C. Schneider-Koglin, B. Mathiasch und M. Dräger), (448) 39

Substitution of fluorine in $\text{C}_6\text{F}_5\text{X}$ ($\text{X} = \text{COOMe}$, CF_3 , CN , NO_2) and pentafluoropyridine by the triethylgermyl group (V.V. Bardin, L.S. Pressman and G.G. Furin), (448) 55

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors $\text{M}^{\text{II}}\text{M}^{\text{IV}}\text{P}_2$ where $\text{M}^{\text{II}} = \text{Zn}$ and Cd , and $\text{M}^{\text{IV}} = \text{Ge}$ and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conrad), (449) 9

Über dimere Dimethylmetall-bis(trimethylgermyl)amide des Aluminiums, Galliums und Indiums (M. Trapp, H.-D. Hausen, G. Weckler und J. Weidlein), (450) 53

Chemie der schweren Carben-Analogen R_2M , $\text{M} = \text{Si}$, Ge , Sn . XVIII. Eine unerwartete Ge-C-Spaltung des als Dimethylgermylen-Speicher dienenden 7,7-Dimethyl-7-germanorbornadiens (B. Klein und W.P. Neumann), (450) 91

Gold

Metal-ion dependent reactivity of 2-(2'-thienyl)pyridine (Hthpy) (E.C. Constable and L.R. Sousa), (427) 125

Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphan-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe $(\text{CO})_3(\text{R}'_3\text{P})(\text{R}_3\text{Si})\text{Fe}-\text{ML}_n$ ($\text{M} = \text{Cu}$, Ag , Au , Hg) (G. Reinhard, B. Hirle and U. Schubert), (427) 173

Fast-atom bombardment (FAB) mass spectra of coordination complexes: aggregation processes in the mass spectra of Group 11 alkynyl complexes (M.I. Bruce and M.J. Liddell), (427) 263

Chemistry of hexaruthenium carbidocarbonyl cluster complexes. Syntheses and X-ray crystal structures of $\text{Ru}_6(\text{C}(\text{CO}))_{15}(\mu\text{-SePh})(\mu\text{-AuPPh}_3)$, $\text{Ru}_6(\text{C}(\text{CO}))_{14}(\mu\text{-SePh})_2$, and $\text{Ru}_6(\text{C}(\text{CO}))_{14}(\mu\text{-SePh})(\mu,\eta^3\text{-C}_3\text{H}_5)$ (T. Chihera and H. Yamazaki), (428) 169

Synthesis, structure and properties of dimethyl(alkoxycarbonyl)gold(III) complexes having a triphenylphosphine ligand (S. Komiya, T. Sone, S. Ozaki, M. Ishikawa and N. Kasuga), (428) 303

The gold(I)-catalyzed aldol reaction utilizing chiral ferrocenylamine ligands: synthesis of *N*-benzyl-substituted ligands (S.D. Pastor, R. Kesselring and A. Togni), (429) 415

Stereoselective formation and reactions of tetraalkylaurates(III) (S. Komiya, S. Ozaki, I. Endo, K. Inoue, N. Kasuga and Y. Ishizaki), (433) 337

Asymmetrical monocationic or neutral gold(II) complexes: X-ray crystal structure of $[(\text{C}_6\text{F}_5)\text{Au}(\text{CH}_2\text{PPh}_2\text{CH}_2)_2\text{AuPPh}_3]\text{ClO}_4 \cdot 0.5\text{CH}_2\text{Cl}_2$ (A. Laguna, M. Laguna, J. Jiménez, F.J. Lahoz and E. Olmos), (435) 235

The synthesis and characterization of σ -bonded gold carbaborane compounds: *closو*-carbaboranes as good σ -donor ligands (B.D. Reid and A.J. Welch), (438) 371

The use of bis(diphenylphosphinoacetylene) and its digold derivative as linking groups in osmium cluster chemistry. Crystal structures of $[(\text{Os}_3(\text{CO}))_{11}(\text{dppa})]$, $[\text{Os}_3(\text{CO})_{10}(\text{dppa})]_2$ and $[\text{Os}_4\text{H}(\text{CO})_{12}\text{Au}(\text{dppa})]_2$ ($\text{dppa} = \text{Ph}_2\text{PC}\equiv\text{CPPh}_2$) (A.J. Amoroso, B.F.G. Johnson, J. Lewis, A.D. Massey, P.R. Raithby and W.T. Wong), (440) 219

Syntheses of gold–manganese and gold–rhenium clusters, $[(\text{Ph}_3\text{PAu})_3\text{M}(\text{CO})_4]$ and $[(\text{Ph}_3\text{PAu})_4\text{M}(\text{CO})_4]^+$ ($\text{M} = \text{Mn}$, Re), and the crystal structure of $[(\text{Ph}_3\text{PAu})_4\text{Mn}(\text{CO})_4]\text{BF}_4^-$ (B.K.

Nicholson, M.I. Bruce, O. Bin Shawkataly and E.R.T. Tiekkink), (440) 411

Neutral or cationic (μ -methylene)bisylide digold(III) complexes (M. Bardaji, M.C. Gimeno, J. Jiménez, A. Laguna, M. Laguna and P.G. Jones), (441) 339

Synthesis and structural characterisation of some triangular and tetrahedral mixed metal platinum–gold clusters (C.M. Hill, D.M.P. Mingos, H. Powell and M.J. Watson), (441) 499

Monomeric to polymeric gold(I) complexes from bifunctional ligands $\text{R}_2\text{PC}\equiv\text{CH}$ (G. Jia, R.J. Puddephatt and J.J. Vittal), (449) 211

Hafnium

Planar tetracoordinate carbon stabilized in a dimetallic hafnium/aluminium compound: formation and crystal structure of $\text{Cp}_2\text{Hf}(\mu\text{-}\eta^1 : \eta^2\text{-MeCC}(\text{C}_6\text{H}_{11})\text{J}\mu\text{-CC}(\text{C}_6\text{H}_{11})\text{JAlMe}_2$ (M. Albrecht, G. Erker, M. Nolte and C. Krüger), (427) C21

Agostic interaction in early transition metal alkyls and their role in catalytic activity for olefin polymerizations (K. Mashima and A. Nakamura), (428) 49

Base-free cationic 14-electron alkyls of Ti, Zr and Hf as polymerisation catalysts: a comparison (M. Bochmann and S.J. Lancaster), (434) C1

Preparation and crystal structures of the complexes $(\eta^5\text{-C}_5\text{H}_4\text{CMe}_2\text{-}\eta^5\text{-C}_{13}\text{H}_8)\text{MCl}_2$ ($\text{M} = \text{Zr}$, Hf) and their role in the catalytic formation of syndiotactic polypropylene (A. Razavi and J. Ferrara), (435) 299

Ferrozentrale Liganden in der Lanthanid-Chemie: Alkoxide und Thiolate (H. Gornitzka, F.T. Edelmann und K. Jacob), (436) 325

Condensation of primary silanes in the presence of $\text{Cp}_2\text{MCl}_2/\text{nBuLi}$ ($\text{M} = \text{Ti}$, Zr , Hf) (J.Y. Corey and X.-H. Zhu), (439) 1

Heterozyklen als Liganden. XVI. $(2,5\text{-C}_4\text{'Bu}_2\text{RHN})\text{MCl}_3$ ($\text{M} = \text{Ti}$, Zr , Hf ; $\text{R} = \text{H}$, SiMe_3)—Azacyclopentadienyl-Komplexe der Gruppe 4-Metalle (N. Kuhn, S. Stubenrauch, R. Boese und D. Bläser), (440) 289

Holmium

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Indium

Metal-metal bonds extended over a porphyrin ring. III. Syntheses and spectral studies of new metal–metal bonded indium porphyrin complexes, (TBPP)In–ML and (5FP)In–ML (S. Takagi, Y. Kato, H. Furuta, S. Onaka and T.K. Miyamoto), (429) 287

Die Molekül- und Kristallstruktur von $(\text{CH}_3)_2\text{InN}_2\text{C}_3\text{H}_3$ (H.-D. Hausen, K. Locke und J. Weidlein), (429) C27

Primary amide and amine complexes of gallium and indium: X-ray crystal structures of $[\text{Me}_2\text{Ga}(\mu\text{-NH}(\text{tBu}))_2]$, $\text{Me}_3\text{Ga} \cdot \text{NH}_2(\text{tBu})$ and $\text{Me}_3\text{In} \cdot \text{NH}_2(\text{tBu})$ (D.A. Atwood, R.A. Jones, A.H. Cowley, S.G. Bott und J.L. Atwood), (434) 143

Organometal-Komplexe von *d*-Block-Elementen. V. Selektive Reduktion von Bis[carbonyl(η^5 -cyclopentadienyl)nickel] zum Carbonyl(η^5 -cyclopentadienyl)nickel-Anion durch C_8K . Synthese und Struktur von $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})\text{Ni-Ga}[\text{CH}_2\text{C}(\text{CH}_3)_3]_2(\text{OC}_4\text{H}_8)$ (R.A. Fischer, J. Behm, E. Herdtweck und C. Kronseder), (437) C29

Synthese und Charakterisierung von ${}^1\text{Pr}_2\text{InF}$ und $[\text{K}(15\text{-Krone-5})_2\text{I}^1\text{Pr}_2\text{InCl}_2]$. Die Kristallstrukturen von $[{}^1\text{Pr}_2\text{InCl}]_x$ und $[\text{K}(15\text{-Krone-5})_2\text{I}^1\text{Pr}_2\text{InCl}_2]$ (B. Neumüller, F. Gahlmann, M. Schäfer und S. Magull), (440) 263

$(\text{CH}_3)_3\text{In} \cdot 1,4\text{-X}_2\text{C}_4\text{H}_8$ (mit $\text{X} = \text{O}$, S), zwei stabile Addukte mit Kettenstruktur (J. Blank, H.-D. Hausen, W. Schwarz und J. Weidlein), (443) 145

- Die Kristallstruktur von $[(\text{CH}_3)_2\text{InCN}]_n$ (J. Blank, H.-D. Hausen und J. Weidlein), (444) C4
- Gallium and indium compounds containing three different substituents. Crystal and molecular structure of $[(\text{Me}_3\text{CCH}_2)\text{ClGa-PPPh}_2]_3$ (O.T. Beachley, Jr., J.D. Maloney and R.D. Rogers), (449) 69
- Raman spectroscopic studies of gas phase Me_3In , Et_3In and EtMe_2In (Z.S. Huang, C. Park and T.J. Anderson), (449) 77
- Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of $[\text{R}(\text{Me}_3\text{SiCH}_2)\text{InE}(\text{SiMe}_3)_2]_2$ ($\text{R} = \text{Ph}$, $\text{E} = \text{As}$; $\text{R} = \text{Me}$, $\text{E} = \text{As}$, P ; $\text{R} = \text{Cl}$, $\text{E} = \text{P}$) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), (449) 85
- Indium tert-butylthiolates as single source precursors for indium sulfide thin films: is molecular design enough? (A.N. MacInnes, M.B. Power, A.F. Hepp and A.R. Barron), (449) 95
- An indium-phosphorus heterocubane as a new type of indium phosphide precursor (D.A. Atwood, A.H. Cowley, R.A. Jones and M.A. Mardones), (449) C1
- Über dimere Dimethylmetall-bis(trimethylgermyl)amide des Aluminiums, Galliums und Indiums (M. Trapp, H.-D. Hausen, G. Weckler und J. Weidlein), (450) 53

Iridium

- Synthesis and X-ray structure of the heteronuclear cluster, $(\mu\text{-H})_2(\eta^5\text{C}_5\text{H}_5)\text{IrOs}_3(\text{CO})_{10}$ (L.-Y. Hsu, W.-L. Hsu, D.A. McCarthy, J.A. Krause, J.-H. Chung and S.G. Shore), (426) 121
- Chemoselective asymmetric hydrogenation of α,β -unsaturated carbonyl compounds to allylic alcohols catalysed by $[\text{Ir}(\text{binap})(\text{cod})]\text{BF}_4$ -aminophosphine (K. Mashima, T. Akutagawa, X. Zhang, H. Takaya, T. Taketomi, H. Kumabayashi and S. Akutagawa), (428) 213
- Hydride addition to and reduction of $\text{Cp}^*\text{Ir}(\eta^6\text{-BT})^{2+}$ and $\text{Cp}^*\text{Ir}(\eta^6\text{-DBT})^{2+}$, where BT = benzo[*b*]thiophene and DBT = dibenzothiophene (J. Chen, Y. Su, R.A. Jacobson and R.J. Angelici), (428) 415
- Spectroscopic investigations of catalytic iridium carbonyl species in sodium chloroaluminate melts (L.J. Tortorelli, P.A. Flowers, B.L. Harward, G. Mamantov and L.N. Klatt), (429) 119
- A convenient lactonization of diols to γ - and δ -lactones catalyzed by transition metal polyhydrides (Y. Lin, X. Zhu and Y. Zhou), (429) 269
- Binuclear hydroxo-, methoxo-, and pyrazolate-bridged complexes of ruthenium(II). Molecular structure of $[(\text{p-cymene})\text{Ru}]_2(\mu\text{-pz})(\mu\text{-OH})_2\text{BF}_4$ (*p*-cymene = $\text{p-MeC}_6\text{H}_4\text{CHMe}_2$, pz = pyrazolate) (D. Carmona, A. Mendoza, J. Ferrer, F.J. Lahoz and L.A. Oro), (431) 87
- Reactivity studies of square planar platinum, iridium, and rhodium triflate complexes with alkynols (P.J. Stang and Y.-H. Huang), (431) 247
- Geometrical isomerism arising from oxidative addition of an *o*-quinone monoimide to $\text{IrCl}(\text{CO})(\text{PPh}_3)_2$ (P.J. Brothers, G.R. Clark, C.E.F. Rickard and H.W. Heine), (433) 203
- Coordination chemistry of $[\text{CH}\{\text{P}(\text{S})\text{Ph}_2\}_2]^-$: X-ray diffraction studies of *S,S*-chelate complexes of iridium and rhodium (J. Browning, G.W. Bushnell, K.R. Dixon and R.W. Hiltz), (434) 241
- Ligand redistribution reactions of some organometallic rhodium and iridium complexes (P.J. Stang and Y.-H. Huang), (435) 185
- Surface organometallic chemistry: formation of $[\text{Ir}_8(\text{CO})_{22}]^{2-}$ on magnesium oxide (S.D. Maloney, M.J. Kelley and B.C. Gates), (435) 377
- Chemistry of iridium carbonyl cluster complexes. Synthesis, chemical characterization and solid state structure of the mixed-metal

carbonyl cluster $[\text{PPh}_4]\text{Ir}_6(\mu\text{-CO})_3(\text{CO})_{12}(\mu_3\text{-Cu}(\text{NCMe})) \cdot \text{CH}_2\text{Cl}_2$ and its conversion to the dianion $[\text{Ir}_{12}(\text{CO})_{26}]^{2-}$ (R. Della Pergola, L. Garlaschelli, F. Demartin, M. Manassero and N. Masciocchi), (436) 241

Decarboxylation syntheses of transition metal organometallics. VI. Preparations of polyfluorophenyliridium(I) compounds (G.B. Deacon and S.J. Faulks), (437) 239

Carbon monoxide activation by poly(1-pyrazolyl)boratoiridium complexes (M.J. Fernandez, M.J. Rodriguez and L.A. Oro), (438) 337

Electronic structure of catalytic intermediates for production of H_2 : $(\text{C}_5\text{Me}_5)\text{Ir}(\text{bpy})$ and its conjugated acid (M. Ladwig and W. Kaim), (439) 79

^{13}C NMR studies of silyl and germyl complexes of rhodium and iridium containing carbonyl ligands (E.A.V. Ebsworth and S. Moreton), (439) 213

Characterization of 1:1 heterobimetallic complexes of $(\eta^5\text{-C}_5\text{Me}_5)\text{Ir}(\text{CO})_2$ with zinc, cadmium and mercury(II) chloride and the X-ray structure determination of the 1:2 mercury complex (F.W.B. Einstein, X. Yan, X. Zhang and D. Sutton), (439) 221

Synthesis and molecular structure of new families of iridium(III)- Cp^* and rhodium(III)- Cp^* complexes derived from 1,2-dicyanoethene-1,2-dithiolate, 2,2'-biimidazole or 2,2'-bithiazole. Single crystal structures of $[(\eta^5\text{-Me}_5\text{C}_5)\text{Ir}(\text{biimH}_2)\text{Cl}] \text{Cl}$ and $[(\eta^5\text{-Me}_5\text{C}_5)\text{Rh}(\text{dcdt})]$ (R. Ziessel, M.-T. Youinou, F. Balegroune and D. Grandjean), (441) 143

Carbon monoxide activation in $[\text{Ir}(\text{HBPz}_3)(\text{CO})_2]$: formation of carbamoyl complexes by direct reaction with amines (M.J. Fernandez, J. Modrego, M.J. Rodriguez, M.C. Santamaría and L.A. Oro), (441) 155

Synthesis and nuclear magnetic resonance studies of some ethylene and ethyltris(pyrazolyl)boratoiridium complexes (M.A. Ciriano, M.J. Fernández, J. Modrego, M.J. Rodríguez and L.A. Oro), (443) 249

Chloro-, Methyl- und Hydrido-Iridiumkomplexe mit Ph_2CN_2 und anderen Diazoalkan-Liganden (L. Brandt, J. Wolf und H. Werner), (444) 235

Synthesis, structure and hydrogenation of η^3 -benzyl diphosphine complexes of rhodium and iridium (M.D. Fryzuk, D.H. McConville and S.J. Rettig), (445) 245

$\text{IrCl}_2\text{H}(\text{P}^+\text{Pr}_3)_2$ as catalyst precursor for the reduction of unsaturated substrates (H. Werner, M. Schulz, M.A. Esteruelas and L.A. Oro), (445) 261

Synthesis and reactivity of binuclear 7-azaindolate complexes of iridium. I. Characterization of isomers by H,H-COSY NMR spectroscopy (M.A. Ciriano, J.J. Pérez-Torrente and L.A. Oro), (445) 267

Synthesis and reactivity of binuclear 7-azaindolate complexes of iridium. II. Oxidative-addition reactions of halogens and halocarbons to $[\text{Ir}(\mu\text{-aza})(\text{CO})_2]_2$ (M.A. Ciriano, J.J. Pérez-Torrente and L.A. Oro), (445) 273

The synthesis of ionic iridium(III) carbene complexes of the type $[\text{Ir}=\text{CR}-\text{NHMe}]\text{Cl}(\text{OSO}_2\text{CF}_3)_2(\text{CO})(\text{PPh}_3)_2]^+[\text{O}_3\text{SCF}_3]^-$ (B.L. Booth and A.C. Wickens), (445) 283

Dehydrogenative silation, isomerization and the control of *syn*- vs. *anti*-addition in the hydrosilation of alkynes (C.-H. Jun and R.H. Crabtree), (447) 177

Transition metal polyhydrides-catalyzed addition of activated nitriles to aldehydes and ketones via Knoevenagel condensation (Y. Lin, X. Zhu and M. Xiang), (448) 215

Synthesis and characterization of sterically encumbered Li, Na, and K aryl tellurolates, and some Pt(II), Ir(I), and Cd(II) derivatives (Ph.J. Bonasia and J. Arnold), (449) 147

Hydroformylation and hydrocarbonylation of 1,2-dicarbethoxy-1,2,3,6-tetrahydropyridazine and 1,2-dicarbethoxy-1,2,3,4-tetrahy-

- dropyridazine (G. Menchi, S. Paganelli, U. Matteoli, A. Scrivanti and C. Botteghi), (450) 229
- Ethylene hydrogenation by the hydride alkoxide species $\text{Ir}(\text{H})_2(\text{OCH}_2\text{CF}_3)(\text{P}^{\text{i}}\text{Bu}_2\text{Ph})_2$. Ethylene-induced reductive elimination of alcohol from Ir^{III} (B. Hauger and K.G. Caulton), (450) 253
- Iron**
- Synthesis and characterization of various coordination modes of 1,1'-bis(diphenylphosphino)ferrocene in iron carbonyl complexes. X-Ray crystal structures of $(\eta^1\text{-BPPF})\text{Fe}(\text{CO})_4$, $(\mu,\eta^2\text{-BPPF})\text{Fe}(\text{CO})_4)_2$, and $(\mu,\eta^2\text{-BPPF})\text{Fe}_3(\text{CO})_{10}$ (T.-J. Kim, S.-C. Kwon, Y.-H. Kim, N.H. Heo, M.M. Teeter and A. Yamano), (426) 71
- Aminsubstituenten als Elektronendonatoren in Clustern (B. Eber, G. Huttner, W. Imhof, J.C. Daran and Y. Jeannin), (426) 87
- Synthesis and evaluation of the catalytic properties of homo- and hetero-bimetallic complexes containing bridging diphenylphosphido ligands (Z. He, N. Lugan, D. Neibecker, R. Mathieu and J.-J. Bonnet), (426) 247
- Metallorganische Lewis-Säuren. XLIV. Rhenioxy(organyl)carben-Komplexe von Chrom, Molybdän und Wolfram (P.M. Fritz, J. Breimair, B. Wagner and W. Beck), (426) 343
- The thermal dissociation process of metal carbonyls on the heated Re surface: $\text{Fe}(\text{CO})_5$ and $\text{Mn}_2(\text{CO})_{10}$ (T. Fujii, K. Kakizaki and H. Ishii), (426) 361
- Electron transfer in asymmetric mixed-valence 3,1'''-(propane-1,3-diy)-1,1'-biferrocenium triiodide (T.-Y. Dong and H.-M. Lin), (426) 369
- Synthesis and reactivity of germyliiron complexes $[\text{Ph}_2\text{HGeFe}(\eta^5\text{-C}_5\text{H}_5)\text{CO}]_2$ and $[\text{RH}_2\text{GeFe}(\eta^5\text{-C}_5\text{H}_5)\text{CO}]_2$ ($\text{R} = \text{Ph}$ or 2,4,6- $\text{Me}_3\text{C}_6\text{H}_2$) (A. Castel, P. Rivière, M. Ahbala, J. Satgé, M. Soufiaoui and N. Knouzi), (427) 91
- The effect of interannular bridge on electron-transfer rate in mixed-valence biferrocenium triiodide salts (T.-Y. Dong, T.-Y. Lee and H.-M. Lin), (427) 101
- Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphan-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe $(\text{CO})_3(\text{R}'_3\text{P})(\text{R}_3\text{Si})\text{Fe}-\text{ML}_n$ ($\text{M} = \text{Cu}, \text{Ag}, \text{Au}, \text{Hg}$) (G. Reinhard, B. Hirle and U. Schubert), (427) 173
- Di(tertiobutylcyclopentadienyl)zircona[1]ferrocenophane précurseur de zirconocenes disubstitués dissymétriquement (R. Broussier, A. Da Rold and B. Gautheron), (427) 231
- Structure and Mössbauer effect study of $[\text{Et}_4\text{N}]_2[\text{Fe}_2(\text{CO})_8]$ (J.M. Cassidy, K.H. Whitmire and G.J. Long), (427) 355
- Charge-transfer salts of carbonylmetalates as outer-sphere ion pairs in photochemical and thermal electron transfer (C.-H. Wei, T.M. Bockman and J.K. Kochi), (428) 85
- Mono- and binuclear σ -aryl iron-lithium hydrides; synthesis and molecular structure (T.A. Bazhenova, L.M. Kachapina, A.E. Shilov, M.Y. Antipin and Y.T. Struchkov), (428) 107
- Photoreactions of silyliiron(II) complexes $\text{Cp}^*\text{Fe}(\text{CO})_2\text{SiMe}_3$ ($\text{Cp}^* = \eta^5\text{-C}_5\text{H}_5$, $\eta^5\text{-C}_5\text{Me}_5$) in the presence of trihydrosilanes (Y. Kawano, H. Tobita and H. Ogino), (428) 125
- Isolation of cationic cyclopentadienyl iron complexes containing phosphinite and formation of metallaphosphineoxide complexes by the Arbuzov-like dealkylation reaction (H. Nakazawa, M. Yamaguchi, K. Kubo and K. Miyoshi), (428) 145
- Synthese und Charakterisierung von 1,3-disubstituierten Decamethylcyclohexasilanen (E. Hengge and M. Eibl), (428) 335
- Cluster mit Fe_6Cd - und Fe_6Hg -Baueinheiten (W. Deck, A.K. Powell and H. Vahrenkamp), (428) 353
- Synthese, Struktur und Reaktivität der Ferriophosphonium-Salze $[\text{CpFe}(\text{CO})(\text{L})(\text{PPh}_2\text{CH}_2\text{R}')]\text{BF}_4$ ($\text{L} = \text{CO}, \text{CH}_3\text{CN}, \text{P}^{\text{n}}\text{Bu}_3$; $\text{R}' = \text{Ph}, \text{CN}, \text{COOEt}, \text{PPh}_2, \text{SiMe}_3$); Kristallstrukturen von $[\text{CpFe}(\text{CO})_2\text{PPh}_2\text{CH}_2\text{SiMe}_3]\text{BF}_4$ und $[\text{CpFe}(\text{PPh}_2\text{CH}_2\text{Ph})\text{CpFe}(\text{CO})](\mu\text{-CO})_2$ (C. Klasen, I.-P. Lorenz, S. Schmid and G. Beuter), (428) 363
- Darstellung μ_3 -Chalkogenverbrückter dreikerniger Carbonyleisencluster durch die Reaktionsfolge XCN^- , Et_3O^+ . Aufbau und Dynamik von $(\mu_3\text{-X})(\mu_3\text{-PR})\text{Fe}_3(\text{CO})_7\text{L}_2$ ($\text{X} = \text{S}, \text{Se}, \text{Te}; \text{L} = \text{CO}, \text{EtNC}, \text{P}(\text{OMe})_3$) (W. Imhof, G. Huttner, B. Eber and D. Günauer), (428) 379
- $[\text{Rh}(\text{CO})]^-$, $[\text{Rh}_3(\text{CO})_{15}]^-$, and bimetallic clusters as catalysts for the carbonylation of nitrobenzene to methyl phenylcarbamate (F. Ragagni, S. Cenini, A. Fumagalli and C. Crotti), (428) 401
- Thermolysis of $\text{Cp}(\text{CO})_n\text{M}-\text{CH}(\text{CH}_3)\text{C}_6\text{H}_5$ ($n = 2, \text{M} = \text{Fe}, \text{Ru}; n = 3, \text{M} = \text{Mo}, \text{W}$). Some vertical trends in transition metal-alkyl bond dissociation energies (C. Mancuso and J. Halpern), (428) C8
- Resolution of $[(\eta^4\text{-benzylideneacetone})\text{Fe}(\text{CO})_3]$. Structure and configurational stability of $[(p\text{S})\text{-}(benzylideneacetone})\text{Fe}(\text{CO})_2\text{L}^*]$ ($\text{L}^* = (+)\text{-neomenthyldiphenylphosphine}$) (A. Marcuzzi, A. Linden, D. Rentsch and W. Von Philipsborn), (429) 87
- Structure and dynamics of $[\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{E}(\text{CH}_3)_2)]\text{BF}_4$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$) complexes studied by ^{13}C solid state NMR spectroscopy (A.P.M. Kentgens, H. Karrenbeld, E. De Boer and H. Schumann), (429) 99
- Synthèse de chalcones à base de tétrathiafulvalène: $\text{RC}(=\text{O})\text{CH}=\text{C}(\text{CH}_3)\text{TTF}$, $\text{TTFC}(=\text{O})\text{CH}=\text{C}(\text{CH}_3)\text{R}$ ($\text{R} = \text{TTF}: \text{C}_6\text{H}_3\text{S}_4$; $\text{Bct}: \text{C}_6\text{H}_5\text{Cr}(\text{CO})_3$; $\text{Fc}: \text{C}_5\text{H}_4\text{FeC}_5\text{H}_5$). Propriété conductrice d'un nouveau complexe de transfert de charge $\text{TTFC}(=\text{O})\text{CH}=\text{C}(\text{CH}_3)\text{TTF} \cdot \text{TCNQ}$ (J. Besançon, A. Radecki-Sudre, J. Szymoniak and J. Padiou), (429) 335
- Facile conversion of $(\eta^5\text{-C}_5\text{R}_5)\text{M}(\text{CO})_2$ -halide complexes to halomethyl, alkoxyethyl, and cyanomethyl derivatives ($\text{R} = \text{H}, \text{CH}_3$; $\text{M} = \text{Fe}, \text{Ru}$; halide = Cl, Br, I) (J.L. Hubbard and W.K. McVicar), (429) 369
- The gold(I)-catalyzed aldol reaction utilizing chiral ferrocenylamine ligands: synthesis of *N*-benzyl-substituted ligands (S.D. Pastor, R. Kesselring and A. Togni), (429) 415
- Intramolecular nucleophilic attack on cationic iron(II) vinylidene complexes: synthesis and crystal structure of the alkenyl $[\text{Fe}(\text{C}=\text{CMe}^1\text{Bu})(\eta^5\text{-C}_5\text{H}_5)(\text{Ph}_2\text{P})_2\text{CH}]$ containing an unprecedented bicyclopentane ring system (M.P. Gamasa, J. Gimeno, E. Lastra, B.M. Martín, A. Aguirre, S. García-Granda and P. Pertíerra), (429) C19
- In-situ* intercalative oxidation of some pyridine derivatives within the cavity of the three-dimensional polymeric host $[(\text{Me}_3\text{Pb})_3\text{Fe}(\text{CN})_6]_\infty$ (A.M.A. Ibrahim, S.E.H. Etaiw and T.M. Soliman), (430) 87
- Silicon–silicon interaction in bis(silylene)iron, disilanyliron, and bis(silyl)iron complexes (K. Ueno, H. Tobita and H. Ogino), (430) 93
- Synthesis and some properties of diferrocenyl thionketones and dynamic behavior of some [1.1]ferrocenophane derivatives (M. Sato and M. Asai), (430) 105
- Organometalloidal derivatives of the transition metals. XXX. Mass spectrometry of transition metal substituted disilanes (A. Guerrero, J. Gómez-Lara, J. Cervantes, L. Velasco, H. Sharma and K.H. Pannell), (430) 273
- Synthesis of new *N*-substituted imidazolyl and 1*H*-1,2,4-triazolyl derivatives of $(\eta^6\text{-arene})(\eta^5\text{-cyclopentadienyl})\text{Fe}(\text{II})$ salts (R.M.G. Roberts), (430) 327

Penta- and hexaphospha ferrocenes as ligands: crystal and molecular structures of $[\text{Fe}(\eta^5\text{-P}_3\text{C}_2\text{^tBu}_2)_2\text{W}(\text{CO})_5]$, $[\text{Fe}(\eta^5\text{-P}_3\text{C}_2\text{^tBu}_2)\text{X}(\eta^5\text{-P}_2\text{C}_3\text{^tBu}_3)\text{W}(\text{CO})_5]$ and the novel triruthenium carbonyl cluster complex $[\text{Fe}(\eta^5\text{-P}_3\text{C}_2\text{^tBu}_2)_2\text{Ru}_3(\text{CO})_{10}]$ containing two inter-linked $\eta^5\text{-P}_3\text{C}_2\text{^tBu}_2$ ring systems (R. Bartsch, A. Gelessus, P.B. Hitchcock and J.F. Nixon), (430) C10

Reactions of cationic vinylidene complexes $[\text{Fe}(=\text{C=C(R}^1\text{)R}^2)(\eta\text{-C}_5\text{H}_5)\text{dppm}]^+$ [dppm = bis(diphenylphosphino)methane] with nucleophiles: stereoselective synthesis and crystal structure of the alkenyl complex (*E*)- $[\text{Fe}(\text{C(H)=C(Me)Ph})(\eta\text{-C}_5\text{H}_5)\text{dppm}]$ (M.P. Gamasa, J. Gimeno, E. Lastra, M. Lanfranchi and A. Tiripicchio), (430) C39

The synthesis and polymerisation of some new 1,1'-substituted vinyl-ferrocenes of the type $[\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{X})(\eta^5\text{-C}_5\text{H}_4(\text{CH=CH}_2))] (\text{X} = \text{Cl}, \text{Br}, \text{CN} \text{ or } \text{CONH}_2)$ (S. Afeworki, P.K. Baker, O.L. Parri and C. D'Silva), (431) 81

Synthesis and structure of $[\text{NEt}_4]\text{Fe}(\text{CO})_3(\mu\text{-SPh})_2\text{Mo}(\text{CO})_3(\text{SPh})$: a heterodinuclear compound with bridging and terminal thiolate ligands (L.D. Rosenheim and P.E. Fanwick), (431) 327

Interesting magnetic effects caused by bulky substituents in ferrocenium salts. Zero applied-field magnetic hyperfine interactions in 1,1',3,3'-tetrakis(trimethylsilyl)ferrocenium triflate (A. Houlton, R.A. Brown, J.R. Miller, R.M.G. Roberts, J. Silver and M. Thomas), (431) C17

Ein neues Verfahren zur Synthese von Metall-funktionalisierten Diphosphenen, Arsaphosphenen und Tetraphospha-bicyclobutanen (P. Jutzi and S. Opela), (431) C29

Photolysis of $\text{Fe}(\text{CO})_4^{2-}$ in an aqueous NaOH glass at 77 K. Formation of solvated electrons (H. Kunckel and A. Vogler), (431) C42

Synthesis and characterisation of a bulky chelating bis(phosphine) ligand, 1,2-bis(diⁿbutylphosphino)ethane (DBPE), and its iron metal coordinated complexes, $\text{Fe}(\text{DBPE})_2\text{Cl}_2$ and $\text{Fe}(\text{DBPE})_2(-\text{C}\equiv\text{C-C}_6\text{H}_5)_2$ (J. Lewis, M.S. Khan, A.K. Kakkar, P.R. Raithby, K. Fuhrmann and R.H. Friend), (433) 135

HPLC studies of $\text{Fe}_2(\text{CO})_6(\text{ligand})$ complexes (D. Osella, O. Gambino, C. Nervi and M. Ravera), (433) 287

Homoleptische metallorganische Innerkomplexe des 8-Dimethylaminonaphthyl-1-Anions (H. Dreys), (433) C1

One-electron reduction of the iron carbene complexes $[\text{Fe}(\text{Cp}^*)(\text{CO})_2(=\text{C}(\text{OMe})\text{R})]^+$ ($\text{R} = \text{H}$ or Me) and hydrogen-atom *vs.* alkyl-radical transfer in the 19-electron intermediates (S. Nlate, V. Guerchais and C. Lapinte), (434) 89

Functionalization of chlorobenzenes via their CpFe^+ salts: a synthesis of substituted diphenyl ethers (R.C. Cambie, S.J. Janssen, P.S. Rutledge and P.D. Woodgate), (434) 97

The constitution of a ferrocenyl diketone: solution and solid state NMR spectroscopy. Crystal structure of 1-ferrocenyl-3-hydroxybut-2-en-1-one (W. Bell, J.A. Crayston, C. Glidewell, M.A. Mazid and M.B. Hursthouse), (434) 115

Synthesis and solution properties of the boron-containing clusters $\text{HM}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf})$ ($\text{M} = \text{Fe}$ or Ru) and the competitive formation of $\{\text{HM}_4(\text{CO})_{12}\text{BHAu}_2\}_{2}(\text{dppf})$ for $\text{M} = \text{Ru}$ ($\text{dppf} = 1,1'\text{-bis(diphenylphosphino)ferrocene}$). Molecular structure of $\text{HRu}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf}) \cdot 2 \text{CHCl}_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9

Synthesis and characterization of Group IV metal–iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $\text{Cp}_2\text{Ti}[(\mu\text{-O}_2\text{CC}_5\text{H}_4)\text{Fe}(\text{CO})_2(\text{CH}_2\text{C}_6\text{H}_5)]_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

P_4 -Liganden mit maximaler Elektronendonorfähigkeit II. $[(\text{Fe}(\text{CO})_4(\mu\text{-P}_2))_2(\mu\text{-Fe}_2(\text{CO})_6)_2]$ —ein Komplex mit einer ungewöhnlich stabilisierten P_4 -Einheit (M. Scheer, M. Dargatz, K. Schenzel und P.G. Jones), (435) 123

Electrochemistry of acetylide complexes of iron (L.D. Field, A.V. George, F. Laschi, E.Y. Malouf and P. Zanello), (435) 347

An X-ray crystallographic and high field NMR study of [7-dehydrocholesteryl acetate] $\text{Fe}(\text{CO})_3$: synthesis of a B-ring cyclohexadienyl cationic complex (R.E. Perrier, C.S. Frampton and M.J. McGlinchey), (435) 357

Umsetzungen von Molybdän-Arsen-Tetrahedranen mit $\text{Fe}_2(\text{CO})_9$ und $\text{Fe}_3(\text{CO})_{12}$ (M. Gorzellik, B. Nuber und M.L. Ziegler), (436) 207

Ethynylferrocene durch direkte Einführung von Acetylenen (M. Buchmeiser und H. Schottenberger), (436) 223

Chelatstabilisierung von Lanthanid(III)-Organylen: Über 2-(Dimethylaminomethyl)ferrocenyl-Komplexe ausgewählter Sel tener Erden (K. Jacob, W. Kretschmer, K.-H. Thiele, H. Gornitzka, F.T. Edelmann, I. Pavlik, A. Lyčka und J. Holeček), (436) 231

Ferrocenaltige Liganden in der Lanthanid-Chemie: Alkoxide und Thiolate (H. Gornitzka, F.T. Edelmann und K. Jacob), (436) 325
 α -Ferrocenyl-Komplexe des Rutheniums (M. Herberhold, W. Feger und U. Kölle), (436) 333

The reactivity of bridging hydride ligands at the di-iron centre. Synthesis and reactivity towards alkynes of $[\text{Fe}_2(\text{CO})_6(\mu\text{-H})(\mu\text{-CO})(\mu\text{-dppm})][\text{BF}_4]$. X-ray crystal structure of $[\text{Fe}_2(\text{CO})_6(\mu\text{-PhC=CH}_2)(\mu\text{-dppm})][\text{BF}_4]$ (J. Boothman and G. Hogarth), (437) 201

Catalytic de-ligation of bis(arene)iron dication upon electro-reduction (Z.J. Karpinski and J.K. Kochi), (437) 211

Diferrocenyl dichalkogenide: Synthese und Struktur von Bis[2-(dimethylaminomethyl)ferrocenyl]diselenid (H. Gornitzka, S. Besser, R. Herbst-Irmer, U. Kilimann, F.T. Edelmann und K. Jacob), (437) 299

Synthese, Charakterisierung und Reaktivität heterodinuklearer Komplexe: $[(\text{CO})_4\text{Fe}(\mu\text{-P}^t\text{Bu}_2)\text{Rh}(\text{CO})\text{L}](\text{Fe-Rh})$ ($\text{L} = \text{CO}$, HP^tBu_2), $[(\text{CO})_3\text{Co}(\mu\text{-H})(\mu\text{-P}^t\text{Bu}_2)\text{Rh}(\text{CO})(\text{HP}^t\text{Bu}_2)](\text{Co-Rh})$ (B. Walther, H.-C. Böttcher, M. Scheer, G. Fischer, D. Fenske und G. Süss-Fink), (437) 307

Synthesis, structure and redox chemistry of ferrocenylsilylmethyldinetricobaltnonacarbonyl complexes, $\text{FcSi}(\text{R})_2\text{CCo}_3(\text{CO})_9, 1,1'\text{-Fc}'[\text{Si}(\text{R})_2\text{CCo}_3(\text{CO})_9]_2$ ($\text{R} = \text{Me}$, Et , Ph) and their derivatives (J. Borgdorff, E.J. Ditzel, N.W. Duffy, B.H. Robinson and J. Simpson), (437) 323

Significance of entropies and volumes of activation for the addition of 4-formylpyridine to tricarbonyl(1-5- η -diienyl)iron(II) complexes (diienyl = C_6H_7 , 2-MeOC₆H₆ and C₇H₉) (T.I. Odiaka and R. van Eldik), (438) 131

A reinvestigation of the reaction of $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_{4-n}(\text{CNR})_n]$ ($n = 1$ or 2) with strong alkylating agents (G. Cox, C. Dowling, A.R. Manning, P. McArdle and D. Cunningham), (438) 143

Electrochemical behaviour of *trans*- $[\text{FeH}(\text{CNR})(\text{dppe})_2]^+$. Kinetic parameters determined by digital simulation of cyclic voltammetry (M.A.N.D.A. Lemos and A.J.L. Pombeiro), (438) 159

Anodic oxidation of the ferrocenyl-derived β -ketophosphine $[\text{Ph}_2\text{PCH}_2\text{CO}(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)]$ and of two of its palladium complexes. Ferricinium-mediated oxidation of palladium(II)-phosphine complexes (A. Louati, M. Gross, L. Douce and D. Matt), (438) 167

The reaction between ferrocenyllithium and trimethylacetylchloride to form either trimethylacetylferrocene or 1,1'-bis(ferrocenyl)-2,2'-dimethylpropan-1-ol, an unusual hydrogen bonded dimer (H.K. Sharma, F. Cervantes-Lee and K.H. Pannell), (438) 183

First optically active ferricinium salts: a circular dichroism study (V.I. Sokolov, I.A. Mamedyarova, M.N. Nefedova and G. Snatzke), (438) C26

$\text{Fe}(\text{CO})_2[\text{P}(\text{OMe})_3]_2\text{L}$, Dicarbonylbis(trimethylphosphite)eisen-Komplexe mit den Liganden L = Ethen, Diphenylethin und SO₂ (F. Meier-Brocks, R. Albrecht und E. Weiss), (439) 65

η^1 - und η^2 -Komplexe von Cp^* -substituierten Diphosphenen und Arsaphosphenen: Synthese, Struktur und Umlagerung unter

- Cp⁺-Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammmer), (439) 279**
- Ferrocenliganden in der Lanthanid-Chemie: Chelatstabilisierung des ersten Diorganyllanthanid(III)-halogenids (H. Gornitzka, A. Steiner, D. Stalke, U. Kiliemann, F.T. Edelmann, K. Jacob und K.-H. Thiele), (439) C6**
- Phosphonium-Übergangsmetallkomplexe. XX. Stabilisierung von Mesitylphosphiniden durch ein Cp(CO)₂W- und ein Cp(CO)₂Fe-Fragment (U.-A. Hirth und W. Malisch), (439) C16**
- Protophilic isotopic hydrogen exchange of 1-ferrocenyl-2-(nitrophenyl)ethylenes (Z.V. Todres and D.S. Ermekov), (439) C28**
- A new reaction of [Cp(CO)₂Fe(PPh₂H)]PF₆ with NaBH₄ to give Cp(CO)(H)Fe(PPh₂H) via Cp(CO)₂FeH and PPh₂H (H. Nakazawa, K. Kubo, C. Kai and K. Miyoshi), (439) C42**
- High resolution EPR spectra of ferrocenyl(nitrophenyl)ethylene anion-radicals (G.F. Pedulli and Z.V. Todres), (439) C46**
- Cis/trans isomerization of β-ferrocenylacrylonitrile during protophilic isotope hydrogen exchange (Z.V. Todres, D.S. Ermekov and I.A. Garbuzova), (439) C53**
- An investigation of steric effects along a cyclopentadienyl substituent: a study by ¹H, ¹³C and ²⁹Si NMR spectroscopy of bis(trimethylsilyl)cyclopentadienyl iron complexes (D. White, L. Carlton and N.J. Coville), (440) 15**
- The Wittig reaction in the generation of organometallic compounds containing alkenes as side groups (E.J. Miller, C.A. Weigelt, J.A. Serth, R. Rusyid, J. Brenner, L.A. Luck and M. Godlewski), (440) 91**
- Methylisocyanide derivatives of molybdenocene and tungstenocene: preparation, reactivity and electronic structure: crystal structures of [(η⁵-C₅H₅)₂WBr(CNMe)]Br and [(η⁵-C₅H₅)₂MoCl(N(H)-CH₃N(H)CH₃)]BF₄]₂·CH₃CN (M.J. Calhorda, A.R. Dias, M.T. Duarte, A.M. Martins, P.M. Matias and C.C. Romão), (440) 119**
- Katalytische Cocyclisierungen von Ethin mit Nitrilen an Bis(η²-ethen)(η⁶-toluol)eisen als Katalysator (U. Schmidt und U. Zenneck), (440) 187**
- Schwefel(IV)-Verbindungen als Liganden. XVII. Übergangsmetallvermittelte Insertion von Schwerfeldioxid in die C–O-Einfachbindung (W.A. Schenk und J. Pfeffermann), (440) 341**
- The synthesis of phosphine derivatives of [Fe₄N(CO)₁₂]⁻: crystal structures of [Fe₄N(CO)₉(NO)(PPh₃)₂], [HFe₄N(CO)₁₁(PPh₃)] and [(PPh₃)₂N]Fe₄N(CO)₁₁(PM₂Ph)] (A. Gourdon and Y. Jeannin), (440) 353**
- The preparation and structures of two Ge–Co–Fe anionic clusters; [GeCoFe₃(CO)₁₄]⁻ and [Ge₂Co₅Fe₂(CO)₂₂]⁻ (S.G. Anema, G.C. Barris, K.M. Mackay and B.K. Nicholson), (441) 35**
- Das Verhalten von stereoisomeren Bistriflaten des Tetrahydrofurans gegenüber ein- und zweiwertigen Carbonylmittallen (E. Lindner, M. Pabel, R. Fawzi und M. Steimann), (441) 63**
- Coupling reactions of ynamines. Syntheses, structural characterization and electrochemistry of some *nido* and *closو ferracyclopentadiene complexes (B. Heim, J.C. Daran, Y. Jeannin, B. Eber, G. Huttner and W. Imhof), (441) 81***
- Synthesis, reactivity of the [(CO)₃(L)Fe(CO₂R)₂] dialkoxy carbonyl carbonyl iron complexes (L = CO or PPh₃; R = CH₃, C₂H₅), and an easy access to [(CO)₃Fe(CO₂Me)]⁺ (J.-Y. Salaün, G. Le Gall, P. Laurent and H. Des Abbayes), (441) 99**
- On the problem of stabilization of α-metallocenylcarbocations. Synthesis, ¹H and ¹³C NMR spectra of metallocenyldiphenylmethyl hexafluorophosphates and X-ray investigation of (C₅H₅Osc₅H₄CPh₂)⁺PF₆⁻ (U. Turpeinen, A.Z. Kreindlin, P.V. Petrovskii and M.I. Rybinskaya), (441) 109**
- Highly selective hydroformylation and dimerization reactions of 2-ferrocenylpropene (L. Kollár and B. Floris), (441) 117**
- Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241**
- Crystal and molecular structure of bis(μ₃-tellurido)-decacarbonyltri-iron, [Fe₃(CO)₁₀Te₂] (G. Gervasio), (441) 271**
- Reaction of Ru₃(CO)₁₂ with ferrocenylacetylene. Crystal and molecular structure of Ru₂(CO)₆[C₄H₂(C₅H₄FeC₅H₅)₂] (A.A. Koridze, A.I. Yanovsky and Y.T. Struchkov), (441) 277**
- Substituted cyclopentadienyl complexes. X. An absolute integrated infrared intensity study of [(η⁵-C₅H₄R)Fe(CO)(L)] (K.E. Du Plooy, T.A. Ford and N.J. Coville), (441) 285**
- Kohlenwasserstoffverbrückte Komplexe. XXVI. Reaktionen von anionischen Aromatenkomplexen [(OC)₃Cr(Ar)]⁻ mit metallorganischen Elektrophilen. Darstellung von (OC)₃Cr(μ-η⁶: η⁶-C₆H₅-C₇H₇)M(CO)₃ (M = Cr, Mo), (OC)₃Cr(μ-η⁶: η⁵-C₆H₅-C(O)C₅H₄)Fe(η⁵-C₅H₅), (OC)₃Cr(μ-η⁶: η⁶-C₈H₅-C₇H₇)Mo(CO)₃, (OC)₃Cr(μ-η⁶: η⁶-C₉H₇-C₇H₇)Mo(CO)₃, (OC)₃Cr(μ-η⁶: η⁶-C₁₃H₉-C₇H₇)M(CO)₃ und (OC)₃Cr(μ-η⁶: η⁶-C₁₃H₁₁-C₇H₇)M(CO)₃ (M = Cr, Mo, W) (J. Breimair, M. Wieser und W. Beck), (441) 429**
- Synthesis and reactivity of tricarbonyl(η⁵-4-triethylsilyl-1-methylpentadienyl)iron(+1) cation (W.A. Donaldson, P.T. Bell and M.-J. Jin), (441) 449**
- Neue Derivierungswege ethinylsubstituierter Metallocene (M. Buchmeiser und H. Schottenberger), (441) 457**
- Synthesis, electrochemistry and complexation studies of new redox active bisferrocene acyclic and macrocyclic thioethers (P.D. Beer, J.E. Nation, M.E. Harman and M.B. Hursthouse), (441) 465**
- Electronic structure of ferrocenylacrylonitrile ion radicals (Z.V. Todres, A.I. Safronov, D.S. Ermekov and R.M. Minyaev), (441) 479**
- Antiferromagnetic complexes with metal–metal bonds. XXVI. Synthesis, molecular structure and magnetic properties of mixed-metal triangular clusters, Cp₂Cr₂(μ-SR)(μ₃-S)₂FeCp (R = CMe₃ or Ph), and linear mixed-metal chain CpCr(μ-SPh)₃Fe(μ-SPh)₃CrCp (S.E. Nefedov, A.A. Pasynskii, I.L. Eremenko, G.S. Gasanov, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky and Y.T. Struchkov), (443) 101**
- Synthesis, structure, and chemistry of heterobimetallic phosphido-bridged complexes CpFe(CO)₂(μ-PPh₂)M(CO)₅ (M = Cr, Mo, W) (S.-G. Shyu, P.-J. Lin and Y.-S. Wen), (443) 115**
- Solvent, N-alkyl substituent and coordination effects on the electrochemistry of 2,11-diaza-5,8-dithia[12](1,1')ferrocenophanes (R.A. Holwerda, J.S. Kim, T.W. Robison, R.A. Bartsch and B.P. Czech), (443) 123**
- Übergangsmetall-substituierte Diphenophene. XXX. η¹- und η²-Komplexe der Metallocidophenene (η⁵-L)(CO)₂Fe-P=P-Mes* (L = C₅Me₅, C₅Me₄Et, C₅Me₄ⁿBu, 1,3-^tBu₂C₅H₃; Mes* = 2,4,6-^tBu₃C₆H₂). Röntgenstrukturanalyse von Pt[η²-(η⁵-C₅Me₄Et)(CO)₂Fe-P=P-Mes*](PPh₃)₂ (L. Weber, I. Schumann, H.-G. Stammmer and B. Neumann), (443) 175**
- The redox behaviour of ferrocene derivatives. I. 1,2,3-Tritellura[3]ferrocenophane and 1,3-ditellura[3]ferrocenophanes (P. Zanello, G. Opronolla, M. Casarin, M. Herberhold and P. Leitner), (443) 199**
- Oxidatively induced nucleophilic capture vs. degradation of cyclopentadienyl iron derivatives of simple carboxylic acids and of α-amino acids. A comparative study (C. Amiens, G. Balavoine and F. Guibé), (443) 207**
- Preparation and properties of cyclodextrin–metal carbonyl inclusion compounds (M. Shimada, Y. Morimoto and S. Takahashi), (443) C8**

- Umsetzungen von Metallorganylen mit dem capto-dativ substituierten Ethylen $H_2C=C(CN)(S^tButyl)$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammmer und M.L. Ziegler), (444) 107
- Regioselectivity of nucleophilic additions to cyclopentadienyliron complexes of substituted benzenes: a novel theoretical approach based on Boltzmann probabilities (X. Luo, G.A. Artega, C. Zhang and P.G. Mezey), (444) 131
- Stereoselectivity in the formation of tricarbonyliron complexes of some dihydrobiphenyls (B.M.R. Bandara, A.J. Birch and B. Chauncy), (444) 137
- An investigation of the barriers to trimethylenemethane (TMM) rotation in compounds of the type $\eta^4\text{-TMMFe}(\text{CO})_2\text{L}$ ($\text{L} = 'Bu\text{NC}$, AsPh_3 , tertiary phosphines); examples of coordinated tertiary phosphine rotational isomerism (L. Girard and M.C. Baird), (444) 143
- Reactions of coordinated phosphines and arsines: stereoselective reactions at the chlorophosphine-*P* stereocentre in the complex $[(R^*), (R^*, R^*)]\text{-}[(\eta^5\text{-C}_5\text{H}_5)(1,2\text{-C}_6\text{H}_4(\text{PMePh})_2)\text{-Fe}(\text{PCl-MePh})\text{PF}_6 \cdot \text{CH}_2\text{Cl}_2]$ (G. Salem, G.B. Shaw, A.C. Willis and S.B. Wild), (444) 147
- Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), (444) 155
- Ein- bis dreikernige Metall- π -Derivate von Hericen mit $(\text{Me}_3\text{P})_3\text{Fe}$ und CpCo -Fragmenten (F.H. Köhler und A. Steck), (444) 165
- The redox behaviour of ferrocene derivatives. II. 8-Ferrocenyl-2-hydroxytricyclo[7.3.1.0^{2,7}]tridecan-13-one and 2,2-dimethylpropylferrocene (A. Cinquantini, G. Oppomolla, P. Zanello and G. Giorgi), (444) 179
- Übergangsmetall-Silyl-Komplexe. XLV. Erzeugung von Fluorsilyl-Liganden durch selektive Spaltung metallgebundener Si-Ph-Gruppen (S. Gilbert und U. Schubert), (444) C12
- Synthesis of $\text{Me}_2\text{Si}\text{-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{Fe}(\text{CO})\text{-}\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2$ and a novel reaction between its Si–Si and Fe–Fe bonds (H. Sun, S. Xu, X. Zhou, H. Wang, R. Wang and X. Yao), (444) C41
- Zur Chemie des Dimesityleisens. VI. Die Struktur von Tetramesityldieisen (H. Müller, W. Seidel und H. Görts), (445) 133
- The preparation, polymerization and copolymerization of $(\eta^5\text{-vinylcyclopentadienyl})\text{dicarbonylmethyliron}$ monomer (A.K. Saha and M.M. Hossain), (445) 137
- The reaction of iron-alkyls with triethyloxonium tetrafluoroborate: formation of iron ethoxycarbene complexes (J.R. Tinder, R.R. Kohl, R.S. Bly and M.M. Hossain), (445) 143
- Stereochemistry of the compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$. Crystal and molecular structures of the mixed triiron carbonyl compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$ ($X, Y = \text{S}, \text{Se}$, or Te) (G. Gervasio), (445) 147
- Synthesis of 1-alkynyl(phthalocyaninato)iron(II) and -ruthenium(II) complexes (M. Hanack, S. Knecht and H.-J. Schulze), (445) 157
- 1,1'-Bis(diphenylphosphino)ferrocene ligand substitution in the benzylidene-capped cluster $\text{PhCCo}_3(\text{CO})_9$. Synthesis, X-ray structure, and redox reactivity of $\text{PhCCo}_3(\text{CO})_7\text{(dppf)}$ (W.H. Watson, A. Nagl, S. Hwang and M.G. Richmond), (445) 163
- Coordination of a tripodal phosphine sulphide P_2S towards various transition metal ions (H.-E. Wang, M.-C. Cheng, G.-H. Lee, S.-M. Peng and S.-T. Liu), (445) 171
- Enantiomerically pure C_2 -symmetric bridged ferrocene and titanocene derivatives (K.-L. Gibis, G. Helmchen, G. Huttner and L. Zsolnai), (445) 181
- Un nouveau type de ferrocénophane, premier exemple de structure dimère (R. Broussier, Y. Gobet, R. Amardeal, A. Da Rold, M.M. Kubicki et B. Gautheron), (445) C4
- Mixed (μ -alkane- and arenethiolato) $(\mu$ -phosphido)hexacarbonyldi-iron complexes. Synthesis and P–Cl reactivity (D. Seyerth, K.S. Brewer, G.B. Womack and C.M. Archer), (446) 167
- ^1H , ^{13}C NMR spectroscopic studies of iodoferrocenyl-ruthenocenium $^+$ salts (M. Watanabe, T. Iwamoto, H. Sano and I. Motoyama), (446) 177
- Umsetzung von Alkynylphosphanen mit Hauptgruppenelement-verbrückten Eisencarbonylclustern (W. Imhof, B. Eber, G. Huttner und C. Emmerich), (447) 21
- Photochemisch induzierte Addition von Alkinen und Phosphaalkinen an die Cluster $\text{Fe}_3(\text{CO})_9(\mu_3\text{-PR})(\mu_3\text{-X})$ ($\text{X} = \text{Se}, \text{Te}; \text{R} = ^i\text{Pr}, 'Bu$) (W. Imhof und G. Huttner), (447) 31
- Photochemical head-to-head ($\text{R} = \text{H}$) vs. head-to-tail dimerization ($\text{R} = \text{Ph}$) of iron acetylide complexes $(\eta^5\text{-C}_5\text{Me}_5)\text{Fe}(\text{CO})_2\text{-C}\equiv\text{C-R}$ leading to a dimetallacyclopentenone and a μ -vinylidene complex (M. Akita, S. Sugimoto, M. Terada and Y. Moro-oka), (447) 103
- Electron transfer in mixed-valence 1',1''-bis(*m*-bromobenzyl)biferrocenium triiodide (T.-Y. Dong and C.-C. Schei), (447) 107
- Studies on olefin-coordinating transition metal carbene complexes. XX. Synthesis of 1,3-cyclohexadiene(dicarbonyl)[ethoxy(aryl)carbene]iron complexes and their phosphine adducts. Crystal structure of $\text{C}_6\text{H}_8(\text{CO})_2\text{FeC}(\text{OC}_2\text{H}_5)\text{C}_6\text{H}_4\text{CH}_3\text{-}\sigma$ (J. Chen, Y. Yu, L. Hu and Z. Jin), (447) 113
- Complexes du fer hydro-digermaniés: $[\text{R}_2\text{HGEGeR}_2\text{Fe}(\eta^5\text{-C}_5\text{H}_5\text{XCO})_2]$ ($\text{R} = \text{Et}, \text{Ph}$) — étude photolytique (A. Castel, P. Rivière, M. Ahbala, J. Satgé, M. Soufiaoui et N. Knouzi), (447) 123
- Organometaloidal derivatives of the transition metals. XXXIII. Reactions of ferrocenylcarboxaldehyde with base and trimethylsilylcyanide. A new high yield reaction for the formation of cyanoamines or silylethers (E. Peña, M. Rivera-Claudio, R.N. Kapoor and K.H. Pannell), (447) 265
- Zur Reaktion von metallkoordinierten 1-Molybda-2-phospha-1,3-dienen mit Alkinen (H. Lang, M. Leise und L. Zsolnai), (447) C1
- Synthesis of the heterodinuclear Fe-Pt complex $[(\text{OC})_3\text{Fe}(\mu\text{-CN}(2,6\text{-xylyl}))(\mu\text{-dppm})\text{Pt}(\text{PPh}_3)]$ containing a bridging isonitrile ligand and electrophilic additions leading to μ -aminocarbyne complexes (M. Knorr, T. Faure and P. Braunstein), (447) C4
- Synthesis and solution dynamics of $[\text{Cp}(\text{CO})_2\text{Fe}]_2(\text{CH}=\text{CH}_2)^+\text{BF}_4^-$, a μ -($\eta^1 : \eta^2$) vinyl complex not containing a metal–metal bond (D.L. Tarazano, T.W. Bodnar and A.R. Cutler), (448) 139
- A rare example of three C–H activations of the methyl group: synthesis and crystal structure of $[\text{Fe}_3(\text{CO})_9(\mu_3\text{-C=N(CH}_3)_2)(\mu_3\text{-CC}_2\text{H}_5)]$ (S. Jeannin, Y. Jeannin, F. Robert and C. Rosenberger), (448) 151
- Heterometallic clusters in homogeneous catalysis. Hydrogenation of diphenylacetylene and isomerization of *cis*-stilbene in the presence of $[\text{Fe}_2\text{Ru}(\text{CO})_{12}]$, $[\text{FeRu}_2(\text{CO})_{12}]$, $[\text{H}_2\text{FeRu}_3(\text{CO})_{13}]$, $[\text{H}_2\text{Ru}_4(\text{CO})_{13}]$, and related compounds. Identification of organometallic complexes and a discussion of their role (R. Giordano and E. Sappa), (448) 157
- ^1H , ^{13}C NMR and ^{57}Fe -Mössbauer spectroscopic studies of α,α -ferrocenylruthenocenylmethylium $^+$ ion (M. Watanabe, T. Iwamoto, S. Nakashima, H. Sakai, I. Motoyama and H. Sano), (448) 167
- The complexation of Y^{3+} by 1,1'-(1,4,10-trioxa-7,13-diazacyclopentadecane-7,13-diyl)ferrocene (C.D. Hall, J.H.R. Tucker and S.Y.F. Chu), (448) 175
- Ferrocene compounds. XX. Synthesis and reactions of some ferrocene fulvenes (S. Kovač, V. Rapić and N. Filipović-Marinić), (448) 181
- Triphenylphosphanchalkogenide als selektive Reagentien zum Aufbau von μ_3 -Chalkogenverbrückten Eisencarbonylclustern (W. Imhof und G. Huttner), (448) 247
- Synthesis and second-order nonlinear optical properties of three coordinate organoboranes with diphenylphosphino and ferrocenyl groups as electron donors: crystal and molecular structures of (*E*)-D-CH=CH-B(mes)₂ and D-C≡C-B(mes)₂ [D = P(C₆H₅)₂,

$(\eta\text{-C}_5\text{H}_5)\text{Fe}(\eta\text{-C}_5\text{H}_4)$; mes = 2,4,6-(CH₃)₃C₆H₂] (Z. Yuan, N.J. Taylor, Y. Sun, T.B. Marder, I.D. Williams and L.-T. Cheng), (449) 27

A facile synthetic route to (η^5 -benzoylcyclopentadienyl)metal compounds (S.S. Jones, M.D. Rausch and T.E. Bitterwolf), (450) 27

Crystal structure of Fe(η^6 -C₅H₅BCMe₃)₂, a "problem structure" (M.R. Churchill and R.F. See), (450) 171

Ferrocene derivatives. XXVIII. The metallation of phenylazoferrrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177

Ferrocene derivatives, LXXI; Stereochemistry of metallocenes, LVI. Synthesis and structure of optically active ferrocenylaminoalcohols (H. Wally, C. Kratky, W. Weissensteiner, M. Widhalm and K. Schlägl), (450) 185

The redox behaviour of ferrocene derivatives. III. Ferrocenylacyl complexes (η^5 -C₅H₅)Fe(η^5 -C₅H₄-COER₃), E = C, Si or Ge; R = Me or Ph (P. Zanello, G. Oppromolla, L. Pardi, K.H. Pannell and H.K. Sharma), (450) 193

Intramolekulare Basenkoordination je nach Bedarf: Der 1-(Dimethylaminooethyl)-2,3,4,5-tetramethylcyclopentadienyl-Ligand in der Komplexchemie von s-, p-, d- und f-Block-Elementen (P. Jutzi, J. Dahlhaus und M.O. Kristen), (450) C1

Lanthanides

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Molekülstruktur und Ladungsverteilung der 4f- und 5f-Elemente. III. Dipolmomente und Ladungsverteilung der 1:1-Addukte der Tris(cyclopentadienyl)lanthanoide(III) bzw. -actinoide(III) mit Tetrahydrofuran; Röntgenstrukturanalyse von (C₅H₅)₃Dy · THF (R. Maier, B. Kanellakopulos, C. Apostolidis und B. Nuber), (435) 275

Chelatstabilisierung von Lanthanid(III)-Organylen: Über 2-(Dimethylaminomethyl)ferrocenyl-Komplexe ausgewählter Seltener Erden (K. Jacob, W. Kretschmer, K.-H. Thiele, H. Gornitzka, F.T. Edelmann, I. Pavlik, A. Lyčka und J. Holeček), (436) 231

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente. XXXIII. Esteraddukte des Grundkörpers Tris(η^5 -cyclopentadienyl)-lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Lanthanum

Molekülstruktur und Ladungsverteilung in metallorganischen Verbindungen der 4f- und 5f-Elemente. II. Dielektrische Untersuchungen an metallorganischen Komplexen der 4f- und Übergangselemente mit C_{3v}- und D_{3h}-Molekülsymmetrie (R. Maier, B. Kanellakopulos und C. Apostolidis), (427) 33

Studies on organolanthanide complexes. XLII. Synthesis of the unsolvated monomeric complexes (MeOCH₂CH₂C₅H₄)₃Ln (Ln = La, Pr) and X-ray structure of (MeOCH₂CH₂C₅H₄)₃Pr (C. Qian, B. Wang, D. Deng, G. Wu and P. Zheng), (427) C29

Metallorganische Verbindungen der Lanthanoide. LXXI. Synthese und Röntgenstrukturanalyse von CpLa(C₅H₄PPh₂)₂(THF), einem zweizähnigen Liganden zur Herstellung heterobimetallischer Lanthanoidkomplexe (H. Schumann, J.A. Meese-Markscheffel, B. Gorella and F.H. Görlitz), (428) C27

Metallorganische Verbindungen der Lanthanoide. LXX. Bis(cyclopentadienyl)seltenerdkomplexe: Synthese und Röntgenstrukturanalyse von dem intramolekular stabilisierten Lutetiumalkyl Cp₂Lu(CH₂)₃NMe₂ und dem monomeren basenfreien Yttriumcarboxylat Cp₂Y[η²-O₂C(CH₂)₃NMe₂] (H. Schumann, J.A. Meese-Markscheffel, A. Dietrich and F.H. Görlitz), (430) 299

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente. XXXIII. Esteraddukte des Grundkörpers Tris(η^5 -cyclopentadienyl)-lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Komplexbildung. XL. Darstellung und Kristallstruktur des Tetra(allyl)lanthanat(III)-Komplexes [Li(μ-C₄H₈O₂)_{3/2}]La(η^3 -C₃H₅)₄ eines Katalysators für die stereospezifische Butadienpolymerisation (R. Taube, H. Windisch, F.H. Görlitz und H. Schumann), (445) 85

Metallorganische Verbindungen der Lanthanoide. LXXIV. Tris(tetramethylcyclopentadienyl)komplexe der Lanthanoide (H. Schumann, M. Glanz und H. Hemling), (445) C1

Lead

Tetra-p-tolyl-Verbindungen p-Tol₄Si und p-Tol₄Ge: ein Beitrag zur Konfiguration der Tetraaryl-Methan-Analoga Ar₄M (M = C, Si, Ge, Sn, Pb) (M. Charissé, S. Roller and M. Dräger), (427) 23

Structure and Mössbauer effect study of [Et₄N]₂[Fe₂(CO)₈] (J.M. Cassidy, K.H. Whitmire and G.J. Long), (427) 355

Structure and equilibria in triorganolead halide adduct formation (H.J. Eppley, J.L. Ealy, C.H. Yoder, J.N. Spencer and A.L. Rheingold), (431) 133

3,5-Di-tert-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151

Trifluormethylsubstituierte Plumbane. Synthese, Eigenschaften und NMR-Spektren der Verbindungen (CF₃)_nPbR_{4-n} (n = 1–3) und (CF₃)_nPbR_{3-n}X (n = 1, 2; R = Me, Et; X = Hal) (R. Eujen and A. Patorra), (438) 57

Synthesis and spectroscopic properties of tetrakis(trifluoromethyl)plumbane, (CF₃)₄Pb (R. Eujen and A. Patorra), (438) C1

Untersuchungen an Arylbleitriacetaten. Röntgenstrukturanalyse von 2-Methylphenyl- und 2-Chlorphenylbleitriacetat (F. Huber, H. Preut, D. Scholz and M. Schürmann), (441) 227

Synthesis of α -alkoxyalkyltributyllead compounds via the reaction of tributylplumbyllithium with α -chloroethers, and the conjugate addition of tributylplumbyllithium to enones (I. Suzuki, T. Furuta and Y. Yamamoto), (443) C6

Über gemischte Gruppe 14–Gruppe 14-Bindungen. VI. Hexa- σ -tolylethan-Analoga o-Tol₆Sn₂, o-Tol₆PbSn und o-Tol₆Pb₂: ein Vergleich von Bindungsstärke und Polarität in der Reihung Sn-Sn, Pb-Sn, Pb-Pb (C. Schneider-Koglin, K. Behrends und M. Dräger), (448) 29

Über gemischte Gruppe 14–Gruppe 14-Bindungen. VII. Ein Inkrementensystem für ¹³C-NMR chemische Verschiebungen bei arylierten Gruppe 14-Verbindungen (C. Schneider-Koglin, B. Mathiasch und M. Dräger), (448) 39

Lithium

The preference of 1-methylallyl polar organometallics and carbanions for *cis* rather than for *trans* geometries (P. von R. Schleyer, J. Kaneti, Y.-D. Wu and J. Chandrasekhar), (426) 143

Some observations on the pyrolysis of methyl lithium (J.R. Baran, Jr. and R.J. Lagow), (427) 1

Synthesis and crystal structure of the tris(cyclopentadienyl)phenyl complex of neodymium, (Li · 3DME)[(η⁵-C₅H₅)₃NdC₆H₅] (H. Gao, Q. Shen, J. Hu, S. Jin and Y. Lin), (427) 141

Mono- and binuclear σ -aryl iron-lithium hydrides; synthesis and molecular structure (T.A. Bazhenova, L.M. Kachapina, A.E. Shilov, M.Y. Antipin and Y.T. Struchkov), (428) 107

Convenient route to di- and triorganosilyl ethyl ethers and the corresponding di- and triorganosilanes (J.M. Tour, J.A. John and E.B. Stephens), (429) 301

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. IV. Synthese von $\text{Li}_2\text{Ni}-(\text{CH}_2-\text{CH}_2\text{CH}_2\text{CH}_2)_2(\text{Et}_2\text{O})$ und seine Umsetzung mit Kohlenmonoxid zu $\text{Li}_2\text{Ni}_6(\text{CO})_{12}(\text{thf})_6$ (H.-O. Fröhlich, B. Hippler and B. Hofmann), (430) 133

Nouveaux arylidihydrogermyllithium (A. Castel, P. Rivière, J. Satgé et D. Desor), (433) 49

Stereoselective opening of chiral α -stannylacetals with organometallic reagents (J.-L. Parrain, J.-C. Cintrat and J.-P. Quintard), (437) C19

Solid-state and solution structures of three lithiumsulfinimidamides: identification of two distinct structural types (F.T. Edelmann, F. Knösel, F. Pauer, D. Stalke and W. Bauer), (438) 1

Structures of lithium salts of substituted 1-phenylpropynes: ^{13}C NMR and MNDO study (P.I. Dem'yanov, I.M. Styrov, D.P. Krut'ko, M.V. Vener and V.S. Petrosyan), (438) 265

Synthesis of polymetallacycloalkanes (P. Mazerolles and C. Laurent), (439) 115

Reactions of chelating alkali metal amides with the Grignard reagent $^n\text{BuMgCl}$: crystal structures and solution ^1H NMR spectroscopic studies of the expected mono(amido)-product $[\text{PhCH}_2(\text{Me}_2\text{N}-\text{CH}_2\text{CH}_2)\text{NMg}^n\text{Bu}]_2$ and the unexpected bis(amido)-product $[(\text{Ph}(2\text{-Pyr})\text{N})_2\text{Mg} \cdot (\text{THF})_2]$ (K.W. Henderson, R.E. Mulvey, W. Clegg and P.A. O'Neil), (439) 237

Le (chlorodiisopropyl)silyl fluorenyllithium: un équivalent synthétique du diisopropyl(fluorenylidène)silène (C. Couret, J. Escudé, G. Delpon-Lacaze et J. Satgé), (440) 233

Selectivity between deprotonation and attack on tellurium in the reaction of 1,4-thiatellurins with strongly basic reagents (H.J.M. Schoufs, A. Maercker and L. Brandsma), (441) 1

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. V. Synthese von $\text{Li}_2\text{Ni}-(\text{CH}_2-\text{CH}_2\text{CH}_2\text{CH}_2)_2(\text{Solv}_x)$ aus $\text{K}[\text{Ni}(\text{NPh}_2)_3](\text{THF})$ und $\text{LiCH}_2\text{CH}_2-\text{CH}_2\text{CH}_2\text{Li}$; Molekülstruktur von $(\text{Li}(\text{THF})_2)_2\text{Ni}^+ \text{CH}_2\text{CH}_2-\text{CH}_2\text{CH}_2)_2$ (H.-O. Fröhlich, R. Wyrwa und H. Görts), (441) 169

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. VI. Synthese und Eigenschaften von Zincacyclohexankomplexen des Typs $(\text{Li}(\text{D})_2)_2\text{Zn}(\text{CH}_2\text{CH}_2\text{CH}_2-\text{CH}_2\text{CH}_2)_2$ ($\text{D} = (\text{C}_2\text{H}_5)_2\text{O}$; $2\text{D} = (\text{CH}_3)_2\text{NC}_2\text{H}_4\text{N}(\text{CH}_3)_2$); Molekülstruktur von $\{\text{Li}(\text{CH}_3)_2\text{NC}_2\text{H}_4\text{N}(\text{CH}_3)_2\}_2\text{Zn}^+ \text{CH}_2\text{CH}_2-\text{CH}_2\text{CH}_2\text{CH}_2)_2$ (H.-O. Fröhlich, B. Kosan, B. Müller und W. Hiller), (441) 177

Metallation reactions. XVIII. Regioselective metallation of (alkylthio)benzenes by superbases (S. Cabiddu, C. Fattuoni, C. Floris, G. Gelli and S. Melis), (441) 197

Umsetzungen der lithiierten Halbsandwich-Chalkogenole, $(\text{CO})_3\text{MC}_5\text{H}_4-\text{ELi}$ ($\text{M} = \text{Mn, Re}$; $\text{E} = \text{S, Se, Te}$) (M. Herberhold und M. Biersack), (443) 1

Etude du transfert monoélectronique entre des germylanions et des piégeurs de spin dia et paramagnétiques (P. Rivière, A. Castel, D. Desor et C. Abdennadher), (443) 51

A solid state ^7Li NMR study of phenyllithium aggregates (D. Johnels), (445) 1

4-Phenylbenzylidene benzylamine: a new and convenient reagent for the titration of solutions of lithium alkyls and metal amides (L. Duhamel and J.-C. Plaquevent), (448) 1

Synthesis of α -stanny α,β -unsaturated carboxylic esters (A.J. Zapata, C. Fortoul R. and C. Acuña A.), (448) 69

The systems $\text{LiAlH}_4/\text{NH}_4\text{X}$ and $\text{N}_2\text{H}_5\text{Cl}$ as precursor sources for AlN (J.F. Janik and R.T. Paine), (449) 39

Lutetium

Metallorganische Verbindungen der Lanthanoide. LXX. Bis(cyclopentadienyl)seltenerdkomplexe: Synthese und Röntgen-strukturanalyse von dem intramolekular stabilisierten Lutetiumalkyl

$\text{Cp}_2\text{Lu}(\text{CH}_2)_3\text{NMe}_2$ und dem monomeren basenfreien Yttrium-carboxylat $\text{Cp}_2\text{Y}[\eta^2-\text{O}_2\text{C}(\text{CH}_2)_3\text{NMe}_2]$ (H. Schumann, J.A. Meese-Marktscheffel, A. Dietrich and F.H. Görlitz), (430) 299

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Metallorganische Verbindungen der Lanthanoide. LXIX. Intramolekulare Stabilisierung der Lanthanoid-Kohlenstoff- σ -Bindung in Mono(cyclopentadienyl)lanthanoidkomplexen (H. Schumann, J.A. Meese-Marktscheffel, A. Dietrich und J. Pickardt), (433) 241

Alpha-metatalated diazoalkanes: synthesis of yttrium, lutetium and ytterbium trimethylsilyldiazomethyl complexes (H. Siebald, M. Dartiguenave and Y. Dartiguenave), (438) 83

Synthesis and characterization of the lutetium naphthalene complex, $\text{CpLuC}_{10}\text{H}_8(\text{DME})$ (A.V. Protchenko, L.N. Zakharov, M.N. Bochkarev and Y.T. Struchkov), (447) 209

Magnesium

Cyclopentadienylmagnesium alkoxides derived from polyfunctional alcohols (O.N.D. Mackey and C.P. Morley), (426) 279

Substitution of the acetoxy groups of dialkoxymethylationacetates by organometallic reagents: a route to allyl-, propargyl-, homoallyl-, homopropargyl- and α -stannylacetals (I. Beaudet, A. Duchêne, J.-L. Parrain and J.-P. Quintard), (427) 201

Reactions of chelating alkali metal amides with the Grignard reagent $^n\text{BuMgCl}$: crystal structures and solution ^1H NMR spectroscopic studies of the expected mono(amido)-product $[\text{PhCH}_2(\text{Me}_2\text{N}-\text{CH}_2\text{CH}_2)\text{NMg}^n\text{Bu}]_2$ and the unexpected bis(amido)-product $[(\text{Ph}(2\text{-Pyr})\text{N})_2\text{Mg} \cdot (\text{THF})_2]$ (K.W. Henderson, R.E. Mulvey, W. Clegg and P.A. O'Neil), (439) 237

Allénysilanes fonctionnels. II. Synthèse par voie organoaluminique du 2-triméthylsilylbuta-2,3-diényal et application à la synthèse d'alcools secondaires α -alléniques triméthylsilylés (D. Mesnard et L. Miginiac), (440) 277

Synthesis of phenyl(perfluoroorgano)mercury derivatives $\text{C}_6\text{H}_5\text{Hg}(\text{R}_f)$ [$\text{R}_f = \text{C}_n\text{F}_{2n+1}$ ($n = 1-4$ or 6) or C_6F_5] by the reactions of bis(perfluoroorgano)cadmium compounds with phenylmercury chloride (H. Layeghi, D. Naumann and W. Tyrra), (441) 355

Conjugate addition of Grignard reagents to *para*-substituted nitrobenzenes (S.J. Bentley and D.J. Milner), (447) 1

Synthese und Kristallstruktur der Grignardverbindungen $\text{C}_5\text{H}_5\text{MgCl} \cdot \text{O}(\text{C}_2\text{H}_5)_2$ und $\text{C}_5\text{Me}_5\text{MgCl} \cdot \text{O}(\text{C}_2\text{H}_5)_2$ (C. Dohmeier, D. Loos, C. Robl, H. Schnöckel und D. Fenske), (448) 5

A polar substituent effect on the ring-cleavage rearrangement of 1-arylcyclobutylmethyl Grignard reagents (E.A. Hill and B. Li), (448) 9

Manganese

Reactions of η^2 -13-acyltetracarbonylmanganese complexes derived from podocarpic acid with alkenes; cyclopentaannulation of ring C (R.C. Cambie, M.R. Metzler, C.E.F. Rickard, P.S. Rutledge and P.D. Woodgate), (426) 213

The thermal dissociation process of metal carbonyls on the heated Re surface: $\text{Fe}(\text{CO})_5$ and $\text{Mn}_2(\text{CO})_{10}$ (T. Fujii, K. Kakizaki and H. Ishii), (426) 361

ansa-Carben-Komplexe durch Insertion/Cyclisierung aus kationischen Carbin-Komplexen und Cyanamiden (H. Fischer and C. Troll), (427) 77

The first examples of insertion of SnCl_2 into the Mn–Cl and Re–Cl bonds of octahedral complexes: X-ray structure of $[\text{Mn}(\text{CO})_3(\text{SnCl}_3)(\text{S}_2\text{CPCy}_3)] \cdot \text{CH}_2\text{Cl}_2$ (B. Alvarez, D. Miguel, J.A. Pérez-Martínez, V. Riera and S. García-Granda), (427) C33

Charge-transfer salts of carbonylmetalates as outer-sphere ion pairs in photochemical and thermal electron transfer (C.-H. Wei, T.M. Bockman and J.K. Kochi), (428) 85

- Coupling reactions of diterpenoid η^2 -tetracarbonylmanganese complexes with alkynes (R.C. Cambie, M.R. Metzler, P.S. Rutledge and P.D. Woodgate), (429) 41
- Reactions with alkenes of η^2 -7-oxotetracarbonylmanganese complexes derived from diterpenoids (R.C. Cambie, M.R. Metzler, P.S. Rutledge and P.D. Woodgate), (429) 59
- Metal-metal bonds extended over a porphyrin ring. III. Syntheses and spectral studies of new metal-metal bonded indium porphyrin complexes, (TBPP)In-ML and (5FP)In-ML (S. Takagi, Y. Kato, H. Furuta, S. Onaka and T.K. Miyamoto), (429) 287
- Formation of a manganese alkyl complex as an intermediate in the ring-opening hydroformylation of α -cyclopropylstyrene by HMn(CO)₅ (R.M. Bullock and B.J. Rappoli), (429) 345
- Orthomanganated arenes in synthesis. VIII. Mono- and dicyclomanganation of diacetyl benzenes (N.P. Robinson, L. Main and B.K. Nicholson), (430) 79
- Photochemical and thermal disproportionation of [(CO)₄CoM(CO)₃ (LL)] (M = Mn, Re; LL = 2,2'-bipyridine, 2-pyridinecarbaldehyde N-isopropylimine) complexes (H. Knoll, W.J.G. De Lange, H. Hennig, D.J. Stufkens and A. Oskam), (430) 123
- Reactions of η^2 -tetracarbonylmanganese complexes derived from podocarpic acid with electrophiles; functionalization of ring C (R.C. Cambie, M.R. Metzler, C.E.F. Rickard, P.S. Rutledge and P.D. Woodgate), (431) 177
- Synthesis and chemistry of 1-acetyl, 2-methylsubstituted cyclopentadienyl thallium and its chiral manganese and rhodium carbonyl derivatives. The molecular structure of [1-CH₃CO, 2-CH₃C₅H₃]Rh(CO)[PC₆H₅]₃] (T.E. Bitterwolf, T.L. Hubler and A.L. Rheingold), (431) 199
- Homoleptische metallorganische Innerkomplexe des 8-Dimethylaminonaphthyl-1-Anions (H. Drews), (433) C1
- 3,5-Di-tert-butylthiobenzoyltrifluorosilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151
- Synthesis and structure analysis of [¹⁴BuC₆H₅)Mn(CO)₃]ClO₄ and its reaction with some nucleophiles (E. Jeong and Y.K. Chung), (434) 225
- The vacuum ultraviolet spectrum of [Mn₂(CO)₁₀] (A. Márquez, J.F. Sanz, M. Gelizé and A. Dargelos), (434) 235
- Hydrogen bonds and conformations of α -carbinol derivatives of cyclopentadienyltricarbonyls of manganese and rhenium (E.S. Shubina, A.N. Krylov, T.V. Timofeeva, Y.T. Struchkov, A.G. Ginzburg, N.M. Loim and L.M. Epstein), (434) 329
- The synthesis of alkylmanganese(III) complexes. Crystal structure of MnMe(2-Me₂NCH₂C₆H₄)₂ (J.L. Latten, R.S. Dickson, G.B. Deacon, B.O. West and E.R.T. Tiekkink), (435) 101
- Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109
- Fulvenylsubstituierte Cp-Liganden. III. Komplexe des Anions [C₅H₄CH=C₉H₆]⁻ mit Molybdän, Mangan und Rhodium (P. Härtter, J. Behm und K.J. Burkert), (438) 297
- Activation of carbon–carbon bond in the Mn-mediated cycloaddition reaction between disilacyclobutene and cyclohepta-1,3-diene (H.-J. Chiang and C.-S. Liu), (438) C9
- Electron spin resonance spectroscopy of Mn(CO)₅⁺ radicals generated in the gas phase thermolysis of Mn₂(CO)₁₀ (K. Mach, J. Nováková and J.B. Raynor), (439) 341
- Alkylmanganese pentacarbonyls (J.-A.M. Andersen and J.R. Moss), (439) C25
- Trigonal-planar koordiniertes Selen bzw. Tellur in [Cp'(CO)₂Mn]₂E-Mes⁺ (E = Se, Te): Verbindungen vom "Iniden"-Typ (P. Lau, G. Huttner und L. Zsolnai), (440) 41
- The Wittig reaction in the generation of organometallic compounds containing alkenes as side groups (E.J. Miller, C.A. Weigelt, J.A. Serth, R. Rusyid, J. Brenner, L.A. Luck and M. Godlewski), (440) 91
- Wittig and Reformatsky chemistry of the formyl and acetyl derivatives of (η^5 -C₅H₅)Mn(CO)₃, (η^5 -C₅H₅)Re(CO)₃ and (η^6 -C₆H₆)Cr(CO)₃ (T.E. Bitterwolf and X. Dai), (440) 103
- Electron transfer catalyzed substitution in carbonyl complexes. VI. Highly variable rates of substitution by tetracyanoethylene (B. Schwederski, W. Kaim, B. Olbrich-Deussner and T. Roth), (440) 145
- Syntheses of gold–manganese and gold–rhenium clusters, [(Ph₃P-Au)₃M(CO)₄] and [(Ph₃PAu)₄M(CO)₄]⁺ (M = Mn, Re), and the crystal structure of [(Ph₃PAu)₄Mn(CO)₄]BF₄ (B.K. Nicholson, M.I. Bruce, O. Bin Shawkataly and E.R.T. Tiekkink), (440) 411
- Das Verhalten von stereoisomeren Bistriflaten des Tetrahydrofuran gegenüber ein- und zweiwertigen Carbonylmetallaten (E. Lindner, M. Pabel, R. Fawzi und M. Steimann), (441) 63
- Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241
- Kohlenwasserstoffverbrückte Komplexe. XXVI. Reaktionen von anionischen Aromatenkomplexen [(OC)₃Cr(Ar)]⁻ mit metallorganischen Elektrophilen. Darstellung von (OC)₃Cr(μ - η^6 : η^6 -C₆H₅-C₇H₇)M(CO)₃ (M = Cr, Mo), (OC)₃Cr(μ - η^6 : η^5 -C₆H₅-C(O)C₅H₄)Fe(η^5 -C₅H₅), (OC)₃Cr(μ - η^6 : η^6 -C₈H₅S-C₇H₇)Mo(CO)₃, (OC)₃Cr(μ - η^6 : η^6 -C₉H₇-C₇H₇)Mo(CO)₃, (OC)₃Cr(μ - η^6 : η^6 -C₁₃H₉-C₇H₇)M(CO)₃ und (OC)₃Cr(μ - η^6 : η^6 -C₁₃H₁₁-C₇H₇)M(CO)₃ (M = Cr, Mo, W) (J. Breimair, M. Wieser und W. Beck), (441) 429
- Umsetzungen der lithiierten Halbsandwich-Chalkogenole, (CO)₃MC₅H₄-ELi (M = Mn, Re; E = S, Se, Te) (M. Herberhold und M. Biersack), (443) 1
- A new preparation of tricarbonyl(η^5 -pentadienyl)manganese (K.F. McDaniel and A.B.A. Afif Abu-Baker), (443) 107
- Preparation and properties of cyclodextrin–metal carbonyl inclusion compounds (M. Shimada, Y. Morimoto and S. Takahashi), (443) C8
- X-ray structure of [Mn(η -C₆H₅Me)₂]⁺I⁻ (D. O'Hare, V. Murphy, A. Bland and P. Scott), (443) C37
- Cyclometallated complexes of Pd^{II} and Mn^I with *N,N*-terephthalylidenebis(cyclohexylamine) (J.M. Vila, M. Gayoso, M.T. Pereira, M. López, G. Alonso and J.J. Fernández), (445) 287
- Electrospray mass spectrometric studies of some cationic and anionic Group 6 and Group 7 metal carbonyl complexes (I. Ahmed, A.M. Bond, R. Colton, M. Jurcevic, J.C. Traeger and J.N. Walter), (447) 59
- Verbrückte Diarsinidenkomplexe {[Cp'(CO)₂Mn]₂As]₂E (E = O, S, Se) und deren nukleophile Spaltung (C. Emmerich, G. Huttner und A. Asam), (447) 71
- Substitution, nukleophile und oxidative Addition an Arsinidenkomplexe. Zur Reaktion von [Cp'(CO)₂Mn]₂AsCl mit PhSeH und verwandten Chalkogenverbindungen (C. Emmerich und G. Huttner), (447) 81
- Manganese carbonyl complexes of 2,5-dimethylbismolyl. The crystal and molecular structure of (η^1 -2,5-dimethylbismolyl) manganese pentacarbonyl (A.J. Ashe, III, J.W. Kampf and D.B. Puranik), (447) 197
- Reaktionen von [Cp'(CO)₂Mn]₂AsCl mit Kohlenstoff-Nukleophilen (C. Emmerich und G. Huttner), (448) 83
- Reaktivität von Tricarbonyl- $\eta^{3:2}$ -2,6-cyclononadienyl-mangan-Komplexen, photochemische Ligandabspaltung (C.G. Kreiter und K. Lehr), (448) 107
- Metallorganische Chemie des Technetiums. VIII. Technetium(I)-carbonyl-Komplexe mit Polypyrazol-1-yl-borato-Liganden im Vergleich mit seinen Mn- und Re-Homologen (J.E. Joachim, C. Apostolidis, B. Kanellakopulos, R. Maier, N. Marques, D. Meyer,

J. Müller, A. Pires de Matos, B. Nuber, J. Rebizant und M.L. Ziegler), (448) 119

A facile synthetic route to (η^5 -benzoylcyclopentadienyl)metal compounds (S.S. Jones, M.D. Rausch and T.E. Bitterwolf), (450) 27

Ferrocene derivatives. XXVIII. The metallation of phenylazoferrrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177

Mercury

Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphoran-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe $(CO)_3(R'_3P)(R_3Si)Fe-ML_n$ (M = Cu, Ag, Au, Hg) (G. Reinhard, B. Hirle and U. Schubert), (427) 173

The effect of ligands, solvent and temperature on the reactions of allyltin(IV) compounds with singlet oxygen (H.-S. Dang and A.G. Davies), (430) 287

Synthese 1,4-disubstituierter Tetrasilane durch selektive Spaltung von Si–Phenyl-Bindungen (H. Stüger), (433) 11

Mercuriation of 3,4,5-trimethoxybenzoic acid. Synthesis of derivatives of the mercuriation product and crystal and molecular structure of $[PPh_3(CH_2Ph)][HgCl(C_6H(CO_2)_3-6-(OMe)_3-2,3,4)]$ (J. Vicente, J.-A. Abad, A. Sandoval and P.G. Jones), (434) 1

Mercury-bridged transition-metal clusters. Synthesis of pentanuclear Ru_3HgM (M = Mo, W, or Co) clusters and X-ray structure of $[(Ru_3(\mu_3,\eta^2\text{-ampy})(CO)_9)\cdot(\mu_3\text{-Hg})Co(CO)_4]$ (Hampy = 2-amino-6-methylpyridine) (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, S. García-Granda and J.F. Van der Maelen), (434) 123

Synthesis, characterisation and molecular structure of $[Ru_5C(CO)_15(HgCF_3)(CF_3CO_2)]$ (B.F.G. Johnson, J. Lewis, P.R. Raithby, V.P. Saharan and W.T. Wong), (434) C10

β -Peroxyalkyl radicals and their role in the generation of epoxides during the autoxidation of alkenes. An example of the interplay of free radical, peroxide and organometallic chemistries (J.L. Courtneidge and M. Bush), (437) 57

The thermal decomposition of dibenzylmercury: a suitable model reaction for determination of substituent effects on radical stability? (R.A. Jackson), (437) 77

Preparation of polyfunctional diorganomercurials and their trans-metallation to diorganozincs. Applications to the preparation of optically active secondary alcohols (M.J. Rozema, D. Rajagopal, C.E. Tucker and P. Knochel), (438) 11

Characterization of 1:1 heterobimetallic complexes of (η^5 - C_5Me_5)Ir(CO)₂ with zinc, cadmium and mercury(II) chloride and the X-ray structure determination of the 1:2 mercury complex (F.W.B. Einstein, X. Yan, X. Zhang and D. Sutton), (439) 221

Uptake of a Hg or Hg₂ unit into the center of cage-type platinum clusters supported by diphosphines having long methylene carbon chains: X-ray structure of $[Hg_2Pt_6(dpphex)_3(RNC)_6] \cdot 2C_6H_6$ (dpphex = 1,6-bis(diphenylphosphino)hexane, R = 2,6-dimethylphenyl) (T. Tanase, T. Horiuchi, Y. Yamamoto and K. Kobayashi), (440) 1

Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241

Crown compounds for anions. A spinning top-shaped complex of cyclic pentameric perfluoroisopropylidenemercury with two chloride anions (V.B. Shur, I.A. Tikhonova, F.M. Dolgushin, A.I. Yanovsky, Y.T. Struchkov, A.Y. Volkonsky, E.V. Solodova, S.Y. Panov, P.V. Petrovskii and M.E. Vol'pin), (443) C19

Selectivity of mercury(II) salts in reactions with α,β -unsaturated stannyll esters (C. Deb and B. Basu), (443) C24

Mass spectroscopic and nuclear magnetic resonance evidence confirming the existence of methyl mercury hydride (P.J. Craig, D. Mennie, M. Needham, N. Oshah, O.F.X. Donard and F. Martin), (447) 5

Complexes derived from 4-amino-3-methyl-5-thione-1,2,4-triazole and monoorganomercury moieties. Crystal structure of (4-amino-5-mercaptop-3-methyl-1,2,4-triazolato)methylmercury(II) (J.M. Varela, A. Macías, J.S. Casas and J. Sordo), (450) 41

Darstellung und spektroskopische Charakterisierung verzweigter funktioneller Hexasilangerüste (H. Stüger und P. Lassacher), (450) 79

Metallocenes

Cyclopentadienylmagnesium alkoxides derived from polyfunctional alcohols (O.N.D. Mackey and C.P. Morley), (426) 279

Electron transfer in asymmetric mixed-valence 3,1'''-(propane-1,3-diy)-1,1'-biferrocenium triiodide (T.-Y. Dong and H.-M. Lin), (426) 369

Alkinkomplexe des Titanocens und Permethyltitanoцен ohne zusätzliche Liganden — Erste Strukturvergleiche (U. Rosenthal, H. Görls, V.V. Burlakov, V.B. Shur and M.E. Vol'pin), (426) C53

Direct proof of the molecular structure of dimeric titanocene; the X-ray structure of $\mu\cdot(\eta^5:\eta^5\text{-fulvalene})\text{-di}\cdot(\mu\text{-hydrido})\text{-bis}(\eta^5\text{-cyclopentadienyltitanium}) \cdot 1.5\text{benzene}$ (S.I. Troyanov, H. Antropiusová and K. Mach), (427) 49

ansa-Metallocene derivatives. XXV. Synthesis, crystal structure and reactions of a tetramethylmethano-bridged vanadocene dichloride, $(CH_3)_4C_2(C_5H_4)_2VCl_2$ (B. Dorer, J. Diebold, O. Weyand and H.-H. Brintzinger), (427) 245

Insertion of alkynes into zirconium–nitrogen bonds leading to 2,3-diaza metalacyclopentenes. Generation and reactivity of $Cp_2Zr(N_2Ph_2)$ (P.J. Walsh, F.J. Hollander and R.G. Bergman), (428) 13

Reactions of pentafluorophenyl germanium hydrides with nickelocene (L.V. Pankratov, V.I. Nevodchikov, L.N. Zakharov, M.N. Bochkarev, I.V. Zdanovich, V.N. Latyaeva, A.N. Lineva, A.S. Batsanov and Y.T. Struchkov), (429) 13

Synthèse de chalcones à base de tétrathiafulvalène: $RC(=O)CH=C(CH_3)TTF$, $TTFC(=O)CH=C(CH_3)R$ (R = TTF: $C_6H_3S_4$; Bct: $C_6H_5Cr(CO)_3$; Fc: $C_5H_4FeC_5H_5$). Propriété conductrice d'un nouveau complexe de transfert de charge $TTFC(=O)CH=C(CH_3)TTF \cdot TCNQ$ (J. Besançon, A. Radecki-Sudre, J. Szymoniak and J. Padiou), (429) 335

Facile synthesis of α,β -unsaturated niobium acyls via γ -hydride abstraction (A.S. Hneihen, M.C. Fermin, J.J. Maas and J.W. Bruno), (429) C33

Cyclopentadienyl- and methylcyclopentadienyl-vanadium complexes containing chalcogen bridges: dechalcogenation to pseudocubane clusters (M. Herberhold, M. Schrepfermann and J. Darkwa), (430) 61

Synthesis and some properties of diferrocenyl thioketones and dynamic behavior of some [1.1]ferrocenophane derivatives (M. Sato and M. Asai), (430) 105

The synthesis and crystal structure of 1,2,3-trithia[3]osmocenophane (E.W. Abel, N.J. Long, A.G. Osborne, M.B. Hursthouse and M.A. Mazid), (430) 117

Synthesis of new N-substituted imidazolyl and 1*H*-1,2,4-triazolyl derivatives of (η^6 -arene) \cdot (η^5 -cyclopentadienyl)iron(II) salts (R.M.G. Roberts), (430) 327

Penta- and hexaphosphorus ferrocenes as ligands: crystal and molecular structures of $[Fe(\eta^5\text{-P}_3C_2'Bu_2)_2W(CO)_5]$, $[Fe(\eta^5\text{-P}_3C_2'Bu_2)(\eta^5\text{-P}_2C_3'Bu_3)W(CO)_5]$ and the novel triruthenium carbonyl cluster complex $[Fe(\eta^5\text{-P}_3C_2'Bu_2)_2Ru_3(CO)_{10}]$ containing two inter-linked $\eta^5\text{-P}_3C_2'Bu_2$ ring systems (R. Bartsch, A. Gelessus, P.B. Hitchcock and J.F. Nixon), (430) C10

Organo-Übergangsmetall-Chemie hochfluorierter Ligand-Systeme. XIV. Vanadocenhexafluoroarsenat-Komplexe: Struktur von $Cp_2V(AsF_6)_2$ und $^{51}\text{V-NMR}$ von $[Cp_2VCl_2]^{+}[AsF_6]^{-}$ (P. Gowik, T.M. Klapötke, K. Siems and U. Thewalt), (431) 47

The synthesis and polymerisation of some new 1,1'-substituted vinyl-ferrrocenes of the type $[\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{X})(\eta^5\text{-C}_5\text{H}_4(\text{CH}=\text{CH}_2))]$ ($\text{X} = \text{Cl}, \text{Br}, \text{CN}$ or CONH_2) (S. Afeworki, P.K. Baker, O.L. Parri and C. D'Silva), (431) 81

Structure of an oxazirconacyclohexadiene synthesized by thermolysis of difurylzirconocene (G. Erker, R. Petrenz, C. Krüger and M. Nolte), (431) 297

Interesting magnetic effects caused by bulky substituents in ferrocenium salts. Zero applied-field magnetic hyperfine interactions in 1,1',3,3'-tetrakis(trimethylsilyl)ferrocenium triflate (A. Houlton, R.A. Brown, J.R. Miller, R.M.G. Roberts, J. Silver and M. Thomas), (431) C17

Mehrkernige Zirkonocen-Ethylkomplexe (H.G. Alt, C.E. Denner und R. Zenk), (433) 107

Ligand redistribution reactions of $\text{Me}_2\text{Ga}(\text{C}_5\text{H}_5)$ and $\text{MeGa}(\text{C}_5\text{H}_5)_2$ (O.T. Beachley, Jr., T.L. Royster, Jr. and J.R. Arhar), (434) 11

Functionalization of chlorobenzenes via their CpFe^+ salts: a synthesis of substituted diphenyl ethers (R.C. Cambie, S.J. Janssen, P.S. Rutledge and P.D. Woodgate), (434) 97

The constitution of a ferrocenyl diketone: solution and solid state NMR spectroscopy. Crystal structure of 1-ferrocenyl-3-hydroxybut-2-en-1-one (W. Bell, J.A. Crayston, C. Glidewell, M.A. Mazid and M.B. Hursthouse), (434) 115

Synthesis and solution properties of the boron-containing clusters $\text{HM}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf})$ ($\text{M} = \text{Fe}$ or Ru) and the competitive formation of $\{\text{HM}_4(\text{CO})_{12}\text{BHAu}_2\}_{\text{dppf}}$ for $\text{M} = \text{Ru}$ ($\text{dppf} = 1,1'\text{-bis(diphenylphosphino)ferrocene}$). Molecular structure of $\text{HRu}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf}) \cdot 2 \text{ CHCl}_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9

ansa-Metallocene derivatives of Ti^{IV} and Zr^{IV} with the shortest $-\text{C}(\text{CH}_3)_2-$ bridge (I.E. Nifant'ev, A.V. Churakov, I.F. Urazowski, S.G. Mkoyan and L.O. Atovmyan), (435) 37

Synthesis and characterization of Group IV metal-iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $\text{Cp}_2\text{Ti}(\mu\text{-O}_2\text{CC}_5\text{H}_4)\text{Fe}(\text{CO})_2(\text{CH}_2\text{C}_6\text{H}_5)_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

Early-transition-metal ketene complexes: synthesis, reactivity and structure of ketene complexes of bis(trimethylsilyl)niobocene, X-ray structure of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Br}(\text{Ph}_2\text{C}=\text{C=O-C}_6\text{O})]$ (A. Antinolo, A. Otero, M. Fajardo, C. Lopez-Mardomingo, D. Lucas, Y. Mugnier, M. Lanfranchi and M.A. Pellinghelli), (435) 55

Darstellung und Eigenschaften von und Reaktionen mit metallhaltigen Heterocyclen. LXXVIII. Das Verhalten von monomeren und dimeren Thiophosphinitonickel-Komplexen gegenüber Alkinen und Phosphaalkinen (E. Lindner, A. Nothdurft, R. Fawzi und C. Maichle), (435) 213

Synthese und Kristallstruktur von $\text{Cp}'_2\text{Zr}(\text{PSiMe}_3)_2$ ($\text{Cp}' = \eta^5\text{-C}_5\text{H}_4\text{Me}$) — Erste röntgenstrukturanalytische Untersuchung eines Zirkonocen-Bisphosphido-Komplexes (F. Lindenberg und E. Hey-Hawkins), (435) 291

Preparation and structure of half-open vanadocene, $\text{V}[2,4\text{-}(\text{CH}_3)_2\text{C}_5\text{H}_5](\text{C}_5\text{H}_5)\text{CO}$ (Y. Wang, Y. Liang, J. Liu, S. Jin and Y. Lin), (435) 311

Electrochemical studies on organometallic compounds. XLII. Electrogeneration and ESR characterization of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Cl}(\text{POMe})_3]^+$ (D. Lucas, Y. Mugnier, A. Antinolo, A. Otero and M. Fajardo), (435) C3

Zum Koordinationsverhalten der Lewis-Basen HCN und DCN gegenüber $\text{Cp}_2\text{Ti}(\text{AsF}_6)_2$ (A. Schulz und T.M. Klapötke), (436) 179

Chelatstabilisierung von Lanthanid(III)-Organylen: Über 2-(Dimethylaminomethyl)ferrocenyl-Komplexe ausgewählter Seltener Erden (K. Jacob, W. Kretschmer, K.-H. Thiele, H. Gornitzka, F.T. Edelmann, I. Pavlik, A. Lyčka und J. Holeček), (436) 231

Ferrocenhaltige Liganden in der Lanthanid-Chemie: Alkoxide und

Thiolate (H. Gornitzka, F.T. Edelmann und K. Jacob), (436) 325

α -Ferrocenyl-Komplexe des Rutheniums (M. Herberhold, W. Feger und U. Kölle), (436) 333

The first examples of intermolecular weak (agostic) γ -methyl–metal interactions in organolanthanide complexes: the synthesis and X-ray structures of $[(\text{Yb}(\eta\text{-Cp}')_2)_2]$ and $[(\text{Eu}(\eta\text{-Cp}')_2)_2]$ [$\text{Cp}' = \text{C}_5\text{H}_3(\text{SiMe}_3)_2\text{-}1,3]$ (P.B. Hitchcock, J.A.K. Howard, M.F. LapPERT and S. Prashar), (437) 177

Diferrocenyldichalkogenide: Synthese und Struktur von Bis[2-(dimethylaminomethyl)ferrocenyl]diselenid (H. Gornitzka, S. Besser, R. Herbst-Irmer, U. Kilimann, F.T. Edelmann und K. Jacob), (437) 299

Synthesis, structure and redox chemistry of ferrocenylsilylmethylylidetricobaltnonacarbonyl complexes, $\text{FcSi}(\text{R})_2\text{CCo}_3(\text{CO})_9$, $1,1'\text{-Fc}'[\text{Si}(\text{R})_2\text{CCo}_3(\text{CO})_9]_2$ ($\text{R} = \text{Me}, \text{Et}, \text{Ph}$) and their derivatives (J. Borgdorff, E.J. Ditzel, N.W. Duffy, B.H. Robinson and J. Simpson), (437) 323

The reaction between ferrocenyllithium and trimethylacetylchloride to form either trimethylacetylferrocene or 1,1'-bis(ferrocenyl)-2,2'-dimethylpropan-1-ol, an unusual hydrogen bonded dimer (H.K. Sharma, F. Cervantes-Lee and K.H. Pannell), (438) 183

First optically active ferricinium salts: a circular dichroism study (V.I. Sokolov, I.A. Mamedyarova, M.N. Nefedova and G. Snatzke), (438) C26

Catalytic activity of the titanocene complex with tolane in homogeneous hydrogenation of unsaturated hydrocarbons (V.B. Shur, V.V. Burlakov and M.E. Vol'pin), (439) 303

Ferrocenliganden in der Lanthanid-Chemie: Chelatstabilisierung des ersten Diorganyllanthanid(III)-halogenids (H. Gornitzka, A. Steiner, D. Stalke, U. Kilimann, F.T. Edelmann, K. Jacob und K.-H. Thiele), (439) C6

Protophilic isotopic hydrogen exchange of 1-ferrocenyl-2-(nitrophenyl)ethylenes (Z.V. Todres and D.S. Ermekov), (439) C28

Die Bildung von $[\text{Cp}_2\text{Ti}(\text{C}=\text{CSiMe}_3)_2]$ aus Titanocene and 1,4-Bis(trimethylsilyl)-1,3-Butadiin (U. Rosenthal and H. Görls), (439) C36

High resolution EPR spectra of ferrocenyl(nitrophenyl)ethylene anion-radicals (G.F. Pedulli and Z.V. Todres), (439) C46

Cis/trans isomerization of β -ferrocenylacrylonitrile during protophilic isotope hydrogen exchange (Z.V. Todres, D.S. Ermekov and I.A. Garbuzova), (439) C53

Kinetically stable adduct of samarcene with aluminium deuteride ($\eta\text{-C}_5\text{H}_3^+\text{Bu}_2)_2\text{Sm}(\mu\text{-D})_2\text{AlD} \cdot \text{Me}_2\text{NC}_2\text{H}_4\text{NMe}_2$ (Y.K. Gun'ko, B.M. Bulychev, V.K. Belsky and G.L. Soloveichik), (440) 47

Methylisocyanide derivatives of molybdenocene and tungstenocene: preparation, reactivity and electronic structure: crystal structures of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{WBr}(\text{CNMe})]\text{Br}$ and $[(\eta^5\text{-C}_5\text{H}_5)_2\text{MoC}(\text{NHC}_6\text{H}_4)_3\text{NHC}(\text{NHC}_6\text{H}_4)_3]\text{BF}_4)_2 \cdot \text{CH}_3\text{CN}$ (M.J. Calhorda, A.R. Dias, M.T. Duarte, A.M. Martins, P.M. Matias and C.C. Romão), (440) 119

On the problem of stabilization of α -metallocenylcarbocations. Synthesis, ^1H and ^{13}C NMR spectra of metallocenyl diphenylmethyl hexafluorophosphates and X-ray investigation of $(\text{C}_5\text{H}_5\text{OsC}_5\text{H}_4\text{CPh}_2)^+\text{PF}_6^-$ (U. Turpeinen, A.Z. Kreindlin, P.V. Petrovskii and M.I. Rybinskaya), (441) 109

Highly selective hydroformylation and dimerization reactions of 2-ferrocenylpropene (L. Kollár and B. Floris), (441) 117

Chiral cyclopentadiene–cyclopentadienyl complexes of cobalt (V.P. Maryin and P.V. Petrovskii), (441) 125

Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241

Reaction of $\text{Ru}_3(\text{CO})_{12}$ with ferrocenylacetylene. Crystal and molec-

- ular structure of $\text{Ru}_2(\text{CO})_6[\text{C}_4\text{H}_2(\text{C}_5\text{H}_4\text{FeC}_5\text{H}_5)_2]$ (A.A. Kordze, A.I. Yanovsky and Y.T. Struchkov), (441) 277
- Substituted cyclopentadienyl complexes. X. An absolute integrated infrared intensity study of $[(\eta^5\text{-C}_5\text{H}_4\text{R})\text{Fe}(\text{CO})(\text{L})\text{I}]$ (K.E. Du Plooy, T.A. Ford and N.J. Coville), (441) 285
- Mixed-valence state of 1',1''-dialkylbiruthenocenium(II,IV) salts (M. Watanabe, T. Iwamoto, H. Sano, A. Kubo and I. Motoyama), (441) 309
- Rearrangement processes of bis-(η^5 -alkylcyclopentadienyl)-zirconium dichlorides under electron impact (Y.A. Andrianov and V.P. Maryin), (441) 419
- Neue Derivierungswege ethinylsubstituierter Metallocene (M. Buchmeiser und H. Schottenberger), (441) 457
- Synthesis, electrochemistry and complexation studies of new redox active bisferrocene acyclic and macrocyclic thioethers (P.D. Beer, J.E. Nation, M.E. Harman and M.B. Hursthouse), (441) 465
- Electronic structure of ferrocenylacrylonitrile ion radicals (Z.V. Tordes, A.I. Safronov, D.S. Ermekov and R.M. Minyaev), (441) 479
- Benzylverbindungen elektronenärmer Übergangsmetalle. Molekülstrukturen von $\text{Cp}_2\text{Ti}(\text{CH}_2\text{Ph})_2$ und $\text{CpZr}(\text{CH}_2\text{Ph})_3$ (J. Scholz, F. Rehbaum, K.-H. Thiele, R. Goddard, P. Betz und C. Krüger), (443) 93
- Carbonyl adducts of open titanocenes – syntheses and structures (Y. Wang, Y.-Q. Liang, Y.-J. Liu and J.-Z. Liu), (443) 185
- The redox behaviour of ferrocene derivatives. I. 1,2,3-Tritellura[3]ferrocenophane and 1,3-ditellura[3]ferrocenophanes (P. Zanello, G. Oppomolla, M. Casarin, M. Herberhold and P. Leitner), (443) 199
- Oxidatively induced nucleophilic capture vs. degradation of cyclopentadienyl iron derivatives of simple carboxylic acids and of α -amino acids. A comparative study (C. Amiens, G. Balavoine and F. Guibé), (443) 207
- Synthesis and structure of mono-THF solvates of bis(cyclopentadienyl)samarium(II) complexes: $(\text{C}_5\text{Me}_5)_2\text{Sm}(\text{THF})$ and $[\text{C}_5\text{H}_2(\text{SiMe}_3)_3\text{I}\text{C}_5\text{H}_3(\text{SiMe}_3)_2]\text{Sm}(\text{THF})$ (W.J. Evans, G. Kociok-Köhne, S.E. Foster, J.W. Ziller and R.J. Doedens), (444) 61
- Photochemistry of $\text{Cp}_2\text{Ti}(\text{CO})_2$ and $\text{Cp}_2\text{Ti}(\text{N}_3)_2$ in solid matrices and in solution at low temperatures (M. Tacke, C. Klein, D.J. Stufkens and A. Oskam), (444) 75
- The redox behaviour of ferrocene derivatives. II. 8-Ferrocenyl-2-hydroxytricyclo[7.3.1.0^{2,7}]tridecan-13-one and 2,2-dimethylpropyl-ferrocene (A. Cinquantini, G. Oppomolla, P. Zanello and G. Giorgi), (444) 179
- Ligand exchange in cyclopentadienylcobalt diselenolenes. Isolation and characterization of the Se_3 -chelate, $\text{C}_5\text{H}_5\text{Co}(\text{SeC}_8\text{H}_{12})_2\text{Se}$ (C.P. Morley and R.R. Vaughan), (444) 219
- Nickelocene-catalyzed polymerization of acetylenes (W.E. Douglas and A.S. Overend), (444) C62
- Ketene acetal metallocene complexes. Crystal structure of $(\text{C}_5\text{H}_5)_2\text{Ti}(\text{Cl})\text{OC}(\text{OCH}_3)\text{C}(\text{CH}_3)_2$ (K. Hortmann, J. Diebold and H.-H. Brintzinger), (445) 107
- Stabilization of a catalytically syndiotactic-specific metallocene cation by trimethylphosphine. The crystal structure of $[(\eta^5\text{-C}_5\text{H}_4)\text{CMe}_2\text{-}\eta^5\text{-}(\text{C}_{13}\text{H}_8)\text{ZrMePMe}_3]^+ [\text{B}(\text{C}_6\text{F}_5)_4]^-$ (A. Razavi and U. Vargha), (445) 111
- Studies on molybdenocene derivatives: reactions of $[\text{Cp}_2\text{Mo}(\eta^2\text{-NCMe})]$ and preparation of alkyl hydride complexes. Crystal structure of $[\text{Cp}_2\text{Mo}(\text{PMe}_3)]$ (C.G. De Azevedo, M.A.A.F. De C.T. Carrondo, A.R. Dias, A.M. Martins, M.F.M. Piedade and C.C. Romão), (445) 125
- 1,1'-Bis(diphenylphosphino)ferrocene ligand substitution in the benzylidine-capped cluster $\text{PhCCo}_3(\text{CO})_9$. Synthesis, X-ray structure, and redox reactivity of $\text{PhCCo}_3(\text{CO})_7(\text{dpfp})$ (W.H. Watson, A. Nagl, S. Hwang and M.G. Richmond), (445) 163
- Enantiomerically pure C_2 -symmetric bridged ferrocene and titanocene derivatives (K.-L. Gibis, G. Helmchen, G. Huttner and L. Zsolnai), (445) 181
- Un nouveau type de ferrocénophane, premier exemple de structure dimère (R. Broussier, Y. Gobet, R. Amardeal, A. Da Rold, M.M. Kubicki et B. Gautheron), (445) C4
- Chemistry of decamethylsilicocene: oxidative addition of compounds with X-H bonds (X = F, Cl, Br, O, S) (P. Jutzi, E.-A. Bunte, U. Holtmann, B. Neumann and H.-G. Stammle), (446) 139
- ^1H , ^{13}C NMR spectroscopic studies of iodoferrocenylruthenocenium $^+$ salts (M. Watanabe, T. Iwamoto, H. Sano and I. Motoyama), (446) 177
- Fulvalene cyclopentadienyl titanium and zirconium(III) and -(IV) complexes. X-Ray crystal structure of $[(\text{Ti}(\eta^5\text{-C}_5\text{H}_5)\text{Cl})_2(\mu\text{-O})(\mu\text{-}\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)]$ (A. Cano, T. Cuena, G. Rodríguez, P. Royo, C. Cardin and D.J. Wilcock), (447) 51
- Crystal and molecular structure of bis(tetramethylcyclopentadienyl)titanium halides, $(\text{C}_5\text{HMe}_4)_2\text{TiCl}$, $(\text{C}_5\text{HMe}_4)_2\text{TiI}$ and $(\text{C}_5\text{HMe}_4)_2\text{TiCl}_2$ (S.I. Troyanov, V.B. Rybakov, U. Thewalt, V. Varga and K. Mach), (447) 221
- Organometallic derivatives of the transition metals. XXXIII. Reactions of ferrocenylcarboxaldehyde with base and trimethylsilylcyanide. A new high yield reaction for the formation of cyanamines or silyl ethers (E. Peña, M. Rivera-Claudio, R.N. Kapoor and K.H. Pannell), (447) 265
- ^1H , ^{13}C -NMR and ^{57}Fe -Mössbauer spectroscopic studies of α,α -ferrocenylruthenocenylmethylium $^+$ ion (M. Watanabe, T. Iwamoto, S. Nakashima, H. Sakai, I. Motoyama and H. Sano), (448) 167
- The complexation of Y^{3+} by 1,1'-(1,4,10-trioxa-7,13-diazacyclopentadecane-7,13-diyl)dicarbonyl)ferrocene (C.D. Hall, J.H.R. Tucker and S.Y.F. Chu), (448) 175
- Ferrocene compounds. XX. Synthesis and reactions of some ferrocene fulvenes (S. Kovač, V. Rapić and N. Filipović-Marinić), (448) 181
- Synthesis and second-order nonlinear optical properties of three coordinate organoboranes with diphenylphosphino and ferrocenyl groups as electron donors: crystal and molecular structures of (*E*)-D-CH=CH-B(*mes*)₂ and D-C≡C-B(*mes*)₂ [D = P(*C*₆*H*₅)₂, (η -C₅H₅)Fe(η -C₅H₄)₂; *mes* = 2,4,6-(CH₃)₃C₆H₂] (Z. Yuan, N.J. Taylor, Y. Sun, T.B. Marder, I.D. Williams and L.-T. Cheng), (449) 27
- Stereoselective propene polymerization at a metallocene/alumoxane catalyst derived from the chirally-substituted “meso-like” (*p*-R,*p*-S)bis[1-(neoisopinocampyl)-4,5,6,7-tetrahydroindenyl]zirconium dichloride (G. Erker, M. Aulbach, C. Krüger and S. Werner), (450) 1
- Carbonylation of 2-bornenyl(methyl)zirconocene (G. Erker, R. Noe, M. Albrecht and A.L. Rheingold), (450) 137
- Rearrangement of cyclooctyne to 1,2-cyclooctadiene within the coordination sphere of $\text{CpMn}(\text{CO})_2$ (S.M. Coughlan and G.K. Yang), (450) 151
- Ferrocene derivatives. XXVIII. The metallation of phenylazoferrrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177
- Ferrocene derivatives, LXXI; Stereochemistry of metallocenes, LVI. Synthesis and structure of optically active ferrocenylaminoalcohols (H. Wally, C. Kratky, W. Weissensteiner, M. Widhalm and K. Schlögl), (450) 185
- The redox behaviour of ferrocene derivatives. III. Ferrocenylacyl complexes ($\eta^5\text{-C}_5\text{H}_5$) Fe($\eta^5\text{-C}_5\text{H}_4\text{-COER}_3$), E = C, Si or Ge; R = Me or Ph (P. Zanello, G. Oppomolla, L. Pardi, K.H. Pannell and H.K. Sharma), (450) 193

Intramolekulare Basenkoordination je nach Bedarf: Der 1-(Dimethylaminoethyl)-2,3,4,5-tetramethylcyclopentadienyl-Ligand in der Komplexchemie von s-, p-, d- und f-Block-Elementen (P. Jutzi, J. Dahlhaus und M.O. Kristen), (450) C1

Molybdenum

Crystal and molecular structure of tetracarbonyl (6,6'-dimethyl-2,2'-bipyridine)molybdenum benzene solvate (P.N.W. Baxter, J.A. Connor, J.D. Wallis and D.C. Povey), (426) 187

Electrochemistry and bonding in biscyclopentadienyl complexes of molybdenum, tungsten, and niobium with *o*-phenanthroline and bipyridine. Crystal and molecular structures of $[Nb(\eta^5-C_5H_5)_2(bpy)]PF_6$ ₂ and $[M(\eta^5-C_5H_5)_2(o\text{-phen})]PF_6$ ₂ ($M = Mo$ or W) (M.J. Calhorda, M.A.A.F. de C.T. Carrondo, A. Bram, P.N. Olsen, A.R. Dias, A.M. Freitas, M.H. Garcia and M.F.M. Piedade), (426) 195

Synthesis and evaluation of the catalytic properties of homo- and hetero-bimetallic complexes containing bridging diphenylphosphido ligands (Z. He, N. Lugan, D. Neibecker, R. Mathieu and J.-J. Bonnet), (426) 247

A study of polarities, polarisabilities and infrared spectra of LM(CO)₅ complexes ($M = Cr$, Mo or W ; L = trimethylamine, quinuclidine, pyridine, *p*-methyl- or *p*-*t*-butylpyridine) and of the X-ray crystal structure of quinuclidine-Cr(CO)₅: evidence for π -acceptor behaviour of coordinated pyridine (M.J. Aroney, R.M. Clarkson, T.W. Hambley and R.K. Pierens), (426) 331

Metallorganische Lewis-Säuren. XLIV. Rheniooxy(organyl)carben-Komplexe von Chrom, Molybdän und Wolfram (P.M. Fritz, J. Breimair, B. Wagner and W. Beck), (426) 343

Controllable migration of acyl groups from molybdenum to phosphido ligands: synthesis of acylphosphine complexes (P. Blenkins, M.H. Lavender and M.J. Morris), (426) C28

Synthese und strukturelle Charakterisierung von Bis(pentamethylcyclopentadienylmolybdän- μ -sulfido)-Komplexen mit μ , η^1 -SSR-Liganden (H. Brunner, R. Graßl, J. Wachter, B. Nuber and M.L. Ziegler), (427) 57

Réaction du bis(dicarbonyl η^5 -cyclopentadiénymolybdène- (I)) avec l'octa-1,7-dyne. Réactivité du μ -octa-1,7-dyne bis(dicarbonyl η^5 -cyclopentadiénymolybdène(I)) vis-à-vis de réactifs spécifiques de la triple liaison carbone-carbone (N. Le Berre-Cosquer and R. Kergoat), (427) 325

The synthesis and substitution reactions of CpM(CO)₂(μ -RC≡CR)Co(CO)₃ ($M = Mo$, W ; $R = COOEt$, $COOMe$) (C.-A. Dickson and N.J. Coville), (427) 335

A novel Redox/Fission reaction of the molybdenum tetracarbonyl complex of 3-diphenylphosphino-(1*R*)-(+)-camphor dimethylhydrazone. Crystal structure of $[Mo(CO)_4(PPh_2C_{10}H_{15}NH)]$ (S.D. Perera, B.L. Shaw and M. Thornton-Pett), (428) 59

Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XXXIX. Synthese und dynamisches Verhalten von Tetracarbonyl- η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien-metall(0)- und Dicarbonyl-bis(η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien)metall(0)-Komplexen der Elemente Chrom, Molybdän und Wolfram (S. Özkar, N.K. Tunali and C.G. Kreiter), (428) 345

Thermolysis of Cp(CO)_nM-CH(CH₃)C₆H₅ ($n = 2$, $M = Fe$, Ru ; $n = 3$, $M = Mo$, W). Some vertical trends in transition metal-alkyl bond dissociation energies (C. Mancuso and J. Halpern), (428) C8

Übergangsmetall-Carbin-Komplexe. CII. Alkylierung von Na[CpMo(CO)₂(CN'Bu)]; Einfluß des Isocyanid-Substituenten auf den Reaktionsablauf (A.C. Filippou, W. Grünleitner and E.O. Fischer), (428) C37

Rupture of the triple C≡C and C≡O bonds in the mixed-metal clusters with Ni₂Mo₂ and Mo₄Co₃ cores. Synthesis and molecular structures of Cp₄Ni₂Mo₂(μ_3 -CPh)₂(μ_3 -CO)₂ and Cp₄Mo₄

(μ_3 -OH)(μ_3 -CPh)₂(μ_6 -C)Co₃(μ -CO)₃(CO)₃ (A.D. Shaposhnikova, M.V. Drab, G.L. Kamalov, A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, Y.T. Struchkov and A.I. Yanovsky), (429) 109

Komplexinduzierter Abbau von As₄S₄: Synthese von Molybdän-Schwefel-Komplexen und Molybdän-Arsen-Tetrahedranen (M. Gorzellik, B. Nuber and M.L. Ziegler), (429) 153

Organometallic complexes of the diphosphene [2,4,6-(CF₃)₃C₆H₂]₂P₂ (K.B. Dillon and H.P. Goodwin), (429) 169

Photolytisch induzierte Erzeugung und komplexchemische Stabilisierung eines Molybdän-Arsen-Dreifachbindungssystems; Röntgenstrukturanalysen der Cluster μ_3 -AsCp₃Mo₂Ni(CO)₄, As₂Cp₆Mo₄Ni₂(CO)₇ und μ_3 -AsCpMoPt₂(CO)₄(PPh₃)₂ (M. Gorzellik, B. Nuber, T. Bohn and M.L. Ziegler), (429) 173

Metal-metal bonds extended over a porphyrin ring. III. Syntheses and spectral studies of new metal-metal bonded indium porphyrin complexes, (TBPP)In-ML and (5FP)In-ML (S. Takagi, Y. Kato, H. Furuta, S. Onaka and T.K. Miyamoto), (429) 287

Tris(η^4 -1-oxa-1,3-diene)molybdenum and -tungsten: Preparation and properties of new homoleptic enone complexes (T. Schmidt and S. Neis), (430) C5

Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CX. Methylierte Nitridomolybdän(VI)-Komplexe (W.A. Herrmann, S. Bogdanović, J. Behm and M. Denk), (430) C33

Oxidation of the molybdenum hydride CpMo(CO)₂(PPh₃)H. Synthesis of *cis*- and *trans*-CpMo(CO)₂(PPh₃)(NCMe)⁺ and the kinetics of their interconversion (K.-T. Smith and M. Tilset), (431) 55

Synthese und strukturelle Charakterisierung von bcpm-Metallcarbonyl-Dimeren (bcpm = Biscyclopentadienylmethan) mit Metallen der 6. Nebengruppe (H. Brunner, R. Graßl, J. Wachter, B. Nuber and M.L. Ziegler), (431) 143

Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XL. Synthese und dynamisches Verhalten von Tricarbonyl- η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien-trimethylphosphit-metall(0)- und Dicarbonyl- η^4 -tricyclo[6.3.0.0^{2,7}]undeca-3,5-dien-bis(trimethylphosphit)metall(0)-Komplexen der Elemente Chrom, Molybdän und Wolfram (C.G. Kreiter, M. Kotzian, S. Özkar and I. Abu-Mour), (431) 159

Synthese des neuartigen organometallischen Molybdän-Antimon-Clusters (μ_4 -Sb)₂Cp₄Mo₅(CO)₁₄ (M. Gorzellik, B. Nuber and M.L. Ziegler), (431) 171

Reaktion von Hydridotrispirazolylboratomolybdän-Komplexen des Typs [TpMo(CO)₂(η^4 -dien)]⁺ mit Ein-Elektronen-Donor-Reagenzien (J. Ipaktschi and A. Hartmann), (431) 303

Molybdenum-95 nuclear magnetic resonance studies on molybdenum carbonyl complexes of isonitriles and amines (C.A. Merlic and B. Adams), (431) 313

Synthesis and structure of [NEt₄]Fe(CO)₃(μ -SPh)₂Mo(CO)₃(SPh): a heterodinuclear compound with bridging and terminal thiolate ligands (L.D. Rosenhein and P.E. Fanwick), (431) 327

Übergangsmetall-Heteroallen-Komplexe. XXVI. Synthese und Struktur von zweikernigen Molybdän-Thioketen-Komplexen (W. Ziegler, D. Wormsbächer and U. Behrens), (431) C11

X-Ray structure analyses of diphosphinidene cyclobutene and its chelate type tetracarbonylmolybdenum(0) complex (K. Toyota, K. Tashiro, M. Yoshifuji, I. Miyahara, A. Hayashi and K. Hirotsu), (431) C35

Reaktionen von Komplexliganden. XLVII. Darstellung von Pentacarbonyl-(Aryl- und Heteroarylcarben)-Komplexen des Molybdäns (K.H. Dötz and H. Larbig), (433) 115

Synthesis of molybdenum-triosmium cluster complexes: reaction of (μ -H)₂Os₃(CO)₁₀ with (η^5 -C₅H₅)₂Mo≡CC₆H₄Me (J.T. Park, J.-J. Cho, K.-M. Chun and S.-S. Yun), (433) 295

Einfluß verschiedenartig substituierter Cyclopentadienylliganden auf die Chemie schwefelreicher Organometallkomplexe mit Metallen der 6. Nebengruppe (H. Brunner, R. Graßl, W. Meier, J. Wachter, B. Nuber und M.L. Ziegler), (434) 63

Mercury-bridged transition-metal clusters. Synthesis of pentanuclear Ru_3HgM ($M = \text{Mo}, \text{W}, \text{or Co}$) clusters and X-ray structure of $[\{\text{Ru}_3(\mu_3,\eta^2\text{-ampy})(\text{CO})_9\} \cdot (\mu_3\text{-Hg})\text{Co}(\text{CO})_4]$ (Hampy = 2-amino-6-methylpyridine) (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, S. García-Granda and J.F. Van der Maelen), (434) 123

Coordinated 2-halo-1,3,2-dioxaphosphorinane ligands. III. *cis*- $\text{Mo}(\text{CO})_4\{\text{REP}(\text{OCH}_2\text{CMe}_2\text{CH}_2\text{O})_2\}$ ($E = \text{O}, \text{S}, \text{NH}; \text{R} = \text{alkyl, aryl or ER} = \text{Cl, Br}$) complexes: syntheses, spectroscopic studies and the molecular structures of the *cis*- $\text{Mo}(\text{CO})_4\{\text{t-PrEP}(\text{OCH}_2\text{CMe}_2\text{CH}_2\text{O})_2\}$ ($E = \text{O}, \text{NH}$) complexes (G.M. Gray and W. Watt), (434) 181

Carbyne–carbyne coupling at a single metal center. Synthesis and structure of bis(diethylamino)acetylene complexes of molybdenum(II) and tungsten(II) (A.C. Filippou, C. Völk, W. Grünlein and P. Kiprof), (434) 201

Synthese und Charakterisierung von Hydridotrispyrazolylborat-Heterobimetall-Komplexen mit Titan und Molybdän (J. Ipaktschi und W. Sulzbach), (434) 287

Synthese und stereokontrollierte Funktionalisierung von Hydridotrispyrazolylborat-Molybdän-Dien-Komplexen (J. Ipaktschi, A. Hartmann und R. Boese), (434) 303

Synthese und stereokontrollierte Funktionalisierung von Hydridotrispyrazolylborat-Molybdän-Dien-Komplexen (J. Ipaktschi, A. Hartmann und R. Boese), (434) 303

Synthesis and X-ray crystal structure of *trans*- $[\text{MoF}(\equiv\text{CCH}_2\text{Bu})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPH}_2)_2]\text{BF}_4$, a paramagnetic alkylidynefluorocomplex (M.A.N.D.A. Lemos, A.J.L. Pombeiro, D.L. Hughes and R.L. Richards), (434) C6

Nucleophilic attack by isocyanides, phosphines and cyclohexenesulfide on the α -carbon of “side-on” bonded μ - σ : η^2 -(4e)-vinylidenes; formation of thioketene and thioaldehyde dimolybdenum complexes (M. Bamber, S.F.T. Froom, M. Green, M. Schulz and H. Werner), (434) C19

Preparation and properties of molybdenum and tungsten dinitrogen complexes. XXXV. Synthesis of diazoalkane and hydrazido(2-) complexes of tungsten containing a π -acceptor ligand (T. Aoshima, T. Tamura, Y. Mizobe and M. Hidai), (435) 85

Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109

Structural properties of cationic molybdenum and tungsten allyl derivatives (B.J. Bridson, M. Cartwright, A.G.W. Hodson, M.F. Mahon and K.C. Molloy), (435) 319

Alkylidenedinitrosylmolybdenum complexes. Preparation, characteristics and metathesis activity of dicarboxylatoethylidenedinitrosylmolybdenum complexes (A. Keller), (436) 199

Umsetzungen von Molybdän-Arsen-Tetrahedranen mit $\text{Fe}_2(\text{CO})_9$ und $\text{Fe}_3(\text{CO})_{12}$ (M. Gorzellik, B. Nuber und M.L. Ziegler), (436) 207

Silylated diphosphite complexes of molybdenum. Crystal structures of the compounds $\text{Mo}(\text{CO})_4[\text{P}(\text{OMe})_2\text{O}]_2\text{SiR}^2\text{R}^3$ ($\text{R}^2 = \text{Ph}$ or $\text{R}^2 = \text{Me}$, $\text{R}^3 = \text{Ph}$) (V. Sum, T.P. Kee and M. Thornton-Pett), (438) 89

Organometallics of diphosphazanes. V. The different coordination behaviour of *cis*- and *trans*-cyclodiphosphazanes towards Group 6 metal carbonyl moieties (V.S. Reddy, S.S. Krishnamurthy and M. Nethaji), (438) 99

Fulvenylsubstituierte Cp-Liganden. III. Komplexe des Anions $[\text{C}_5\text{H}_4\text{CH}=\text{C}_9\text{H}_6]^-$ mit Molybdän, Mangan und Rhodium (P. Härtel, J. Behm und K.J. Burkert), (438) 297

Heterobimetallic Group 6-rhodium complexes. II. Reactions of $[\text{Rh}(\text{S}_2\text{CNEt}_2)(\text{CO})_2]$ with the complexes $[\text{M}(\text{CO})_3(\eta^2\text{-phen})(\eta^1\text{-dppm})]$ ($\text{M} = \text{Mo}$ or W) and $[\text{Mo}(\text{CO})_4(\eta^2\text{-dppm})]$ as P-donor systems (M. Cano, P. Ovejero and J.V. Heras), (438) 329

Syntheses and characterization of tri-metal cluster compounds; X-ray structure determination of $[\text{Co}_2\text{Mo}(\mu_3\text{-CC}_6\text{H}_5)\text{Br}(\text{CO})_8(\text{bipy})]$ ($\text{bipy} = \alpha,\alpha\text{-dipyridyl}$) (Y. Yin, H. Sun, D. Jin and Y. Zhu), (439) 45

Half-sandwich complexes with intramolecular amino group coordination: synthesis and structure of cationic molybdenum dienyl complexes (T.-F. Wang and Y.-S. Wen), (439) 155

The synthesis of dimetallic thioalkyne complexes of cobalt and molybdenum (S.C. Bennett, A. Gelling and M.J. Went), (439) 189

Synthesis and characterization of cobalt and molybdenum complexes derived from linear conjugated diynes, triynedienes and tetracynetrienes (W.E. Lindsell, P.N. Preston and P.J. Tomb), (439) 201

η^1 - und η^2 -Komplexe von Cp^* -substituierten Diphenophenen und Arsaphenophenen: Synthese, Struktur und Umlagerung unter Cp^* -Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammler), (439) 279

Some P–H bond substitution derivatives and fluxionality of $\text{Cp}_2\text{Mo}_2(\mu\text{-H})(\mu\text{-PPH})\text{CO}_4$ (S. Woodward and M.D. Curtis), (439) 319

The synthesis of dinuclear molybdenum and tungsten complexes with the bridging heterodifunctional ligand $\text{C}_5\text{H}_4\text{PPh}_2$: X-ray crystal structure of $\text{Mo}_2(\mu\text{-C}_5\text{H}_4\text{PPh}_2)(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_5$ (T.J. Duckworth, M.J. Mays, G. Conole and M. McPartlin), (439) 327

Cycloadditionsreaktionen von Phospheniumkomplexen. VIII. Regioselektive Cycloaddition chiraler Phosphenium-Komplexe $\text{Cp}(\text{OC})(\text{Ph}_2\text{R})\text{P}=\text{PPh}_2$ ($\text{M} = \text{Mo, W}; \text{R} = \text{H, Me}$) mit Isothiocyanaten (H. Pfister und W. Malisch), (439) C11

Phosphenium-Übergangsmetallkomplexe. XXI. Schwefel- und Sele-naddition an die Metall–Phosphor-Doppelbindung der zweikernigen Phosphinidenkomplexe $\text{Cp}(\text{CO})_2\text{M}=\text{PMes}[\text{M}(\text{CO})_3\text{Cp}]$ ($\text{M} = \text{Mo, W}$) (U.-A. Hirth, W. Malisch und H. Käb), (439) C20

Effect of pressure on intramolecular ring-closure reactions of molybdenum carbonyl complexes induced by flash photolysis (K. Bal Reddy, B.R. Brady, E.M. Eyring and R. Van Eldik), (440) 113

Methylisocyanide derivatives of molybdenocene and tungstenocene: preparation, reactivity and electronic structure: crystal structures of $[(\eta^5\text{-C}_5\text{H}_5)_2\text{WBr}(\text{CNMe})]\text{Br}$ and $[(\eta^5\text{-C}_5\text{H}_5)_2\text{MoC}(\text{N}(\text{H})=\text{CH}_3)\text{N}(\text{H})\text{C}(\text{N}(\text{H})\text{CH}_3)]\text{BF}_4$ · CH_3CN (M.J. Calhorda, A.R. Dias, M.T. Duarte, A.M. Martins, P.M. Matias and C.C. Romão), (440) 119

Elektrophile Addition des Thiomethyl-Kations an η^5 -Cyclopenta-dienyl(ethoxymethylcarben)- (nitrosyl)(trimethylphosphin)molyb-dän (F.R. Kreißl und N. Ullrich), (440) 335

Schwefel(IV)-Verbindungen als Liganden. XVII. Übergangsmetall-vermittelte Insertion von Schwefeldioxid in die C–O-Einfach-bindung (W.A. Schenck und J. Pfeffermann), (440) 341

Das Verhalten von stereoisomeren Bistriflaten des Tetrahydrofurans gegenüber ein- und zweiwertigen Carbonylmallatten (E. Lindner, M. Pabel, R. Fawzi und M. Steimann), (441) 63

Kohlenwasserstoffverbrückte Komplexe. XXVI. Reaktionen von anionischen Aromatenkomplexen $[(\text{OC})_3\text{Cr}(\text{Ar})]^-$ mit metallorganischen Elektrophilen. Darstellung von $(\text{OC})_3\text{Cr}(\mu\text{-}\eta^6:\eta^6\text{-C}_6\text{H}_5\text{-C}_7\text{H}_7)\text{M}(\text{CO})_3$ ($\text{M} = \text{Cr, Mo}$), $(\text{OC})_3\text{Cr}(\mu\text{-}\eta^6:\eta^5\text{-C}_6\text{H}_5\text{-C}(\text{O})\text{C}_5\text{H}_4)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)$, $(\text{OC})_3\text{Cr}(\mu\text{-}\eta^6:\eta^6\text{-C}_8\text{H}_5\text{S-C}_7\text{H}_7)\text{Mo}(\text{CO})_3$, $(\text{OC})_3\text{Cr}(\mu\text{-}\eta^6:\eta^6\text{-C}_9\text{H}_7\text{-C}_7\text{H}_7)\text{Mo}(\text{CO})_3$ und $(\text{OC})_3\text{Cr}(\mu\text{-}\eta^6:\eta^6\text{-C}_{13}\text{H}_9\text{-C}_7\text{H}_7)\text{M}(\text{CO})_3$ ($\text{M} = \text{Cr, Mo, W}$) (J. Breimair, M. Wieser und W. Beck), (441) 429

Synthesis, structure, and chemistry of heterobimetallic phosphido-bridged complexes $\text{CpFe}(\text{CO})_2(\mu\text{-PPH}_2)\text{M}(\text{CO})_5$ ($\text{M} = \text{Cr, Mo, W}$) (S.-G. Shyu, P.-J. Lin and Y.-S. Wen), (443) 115

Additionsreaktionen von $(\text{CO})_5\text{CrEPH}_2$ ($\text{E} = \text{P, As}$) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), (444) 53

Synthesis and X-ray structure of $(-\text{[Mo}(\eta^5\text{-C}_5\text{Me}_4\text{-3-pinanyl})\text{CO})_3\text{Me}]$ (R. Lai, L. Bousquet and A. Heumann), (444) 115

Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenyliindol (D. Garn, F. Knoch und H. Kisch), (444) 155

Molekülstruktur von Tetracarbonyl-bis(η^5 -cyclopentadienyl)- $\mu\text{-}\eta^{1:2}$ -vinyliden-dimolybdän (U. Kern, C.G. Kreiter, S. Müller-Becker and W. Frank), (444) C31

Dealkylation of thiols by a phosphido-bridged dimolybdenum complex. Crystal structure of $[\text{Mo}_2(\mu\text{-S})_2(\mu\text{-S'Pr})(\mu\text{-PPH}_2)(\eta^5\text{-C}_5\text{H}_5)_2]$ (H. Adams, N.A. Bailey, A.P. Bisson and M.J. Morris), (444) C34

Unexpected ring-expansion of "axially prosterogenic" biaryl lactones by methylenephosphoranyl-substituted imido complexes of molybdenum and tungsten (J. Sundermeyer, K. Weber, H. Werner, N. Mahr, G. Bringmann and O. Schupp), (444) C37

Darstellung und Charakterisierung von Indenyl- und Fluorenylfunktionalisierten Cyclopentadienylcarbonyl-Methylkomplexen des Molybdäns und Wolframs. Molekülstrukturen von $(\eta^5\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_9\text{H}_7)\text{M}(\text{CO})_3\text{Me}$ ($\text{M} = \text{Mo, W}$), $(\eta^5\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_{13}\text{H}_9)\text{M}(\text{CO})_3\text{Me}$ und $(\eta^5\text{:}\eta^2\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_9\text{H}_7)\text{Mo}(\text{CO})_2\text{Me}$ (H.G. Alt, J.S. Han and R.D. Rogers), (445) 115

Studies on molybdenocene derivatives: reactions of $[\text{Cp}_2\text{Mo}(\eta^2\text{-NCMe})]$ and preparation of alkyl hydride complexes. Crystal structure of $[\text{Cp}_2\text{Mo}(\text{PMMe}_3)]$ (C.G. De Azevedo, M.A.A.F. De C.T. Carrondo, A.R. Dias, A.M. Martins, M.F.M. Piedade and C.C. Romão), (445) 125

Electrospray mass spectrometric studies of some cationic and anionic Group 6 and Group 7 metal carbonyl complexes (I. Ahmed, A.M. Bond, R. Colton, M. Jurcevic, J.C. Traeger and J.N. Walter), (447) 59

Darstellung und Struktur von 1-Metalla-2,3,4-triazolen des Molybdäns und Wolframs (C.M. Stegmair, W. Schütt, W. Ullrich, P. Kiprof, J. Ostermeier und F.R. Kreißl), (447) 251

Zur Reaktion von metallkoordinierten 1-Molybda-2-phospha-1,3-dienen mit Alkinen (H. Lang, M. Leise und L. Zsolnai), (447) C1

Phenylbis(2-pyridyl)phosphine: *P*- vs. *N,N'*-coordination in carbonylmolybdenum-(0) and -(II) complexes (P. Espinet, P. Gómez-Elipe and F. Villafaña), (450) 145

Ferrocene derivatives. XXVIII. The metallation of phenylazoferrrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177

Intramolekulare Basenkoordination je nach Bedarf: Der 1-(Dimethylaminoethyl)-2,3,4,5-tetramethylcyclopentadienyl-Ligand in der Komplexchemie von s-, p-, d- und f-Block-Elementen (P. Jutzi, J. Dahlhaus und M.O. Kristen), (450) C1

Neodymium

Synthesis and crystal structure of the tris(cyclopentadienyl)phenyl complex of neodymium, $(\text{Li}\cdot 3\text{DME})(\eta^5\text{-C}_5\text{H}_5)_3\text{NdC}_6\text{H}_5]$ (H. Gao, Q. Shen, J. Hu, S. Jin and Y. Lin), (427) 141

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Metallogands containing samarium(III). Access to binuclear *f*-metal (samarium)-*d*-metal (rhodium) complexes (M. Visseaux, A. Dor-

mond, M.M. Kubicki, C. Moïse, D. Baudry and M. Ephritikhine), (433) 95

Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXXII. Absorptionsspektroskopische Charakterisierung von $\text{Cp}_3\text{Nd}(\text{NCCH}_3)_2$ (H. Schulz, H. Reddmann und H.-D. Amberger), (440) 317

Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXXIII. Esteraddukte des Grundkörpers $\text{Tris}(\eta^5\text{-cyclopentadienyl})\text{-lanthan(III)}$ —ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilian und F.T. Edelmann), (444) C15

Studies on organolanthanide complexes. XLVI. Synthesis and characterization of N-containing ring-linked biscyclopentadienyl lanthanide and yttrium chlorides (C. Qian and D. Zhu), (445) 79

Nickel

Neuartige Synthesen monosubstituierter Nickelocene als Zugang zu mehrkernigen Metallocensystemen (H. Schottenberger, G. Ingram and D. Obendorf), (426) 109

Struktur und Dynamik von $[(2,6\text{-Me}_2\text{Ph-dad})\text{Ni}_2]\sigma^2(1,4,\eta^4(1\text{-}4)\text{C}_4\text{H}_4)$ (S. Michaelis, K.R. Pörschke, R. Mynott, R. Goddard and C. Krüger), (426) 131

Metall- π -Komplexe von Benzolderivaten. XXXVIX. Di[1,4-bis(diphenylphosphano)- η^6 -benzol]chrom: ein Baustein für polynukleare Bimetallkomplexe? (Ch. Elschenbroich, J. Sebbach, B. Metz and G. Heikenfeld), (426) 173

Nickelalactone als Synthesebausteine: Sonochemische und Bimetallaktivierung der Kreuzkopplungsreaktion mit Alkyl-halogeniden (R. Fischer, D. Walther, G. Bräunlich, B. Undeutsch, W. Ludwig and H. Bandmann), (427) 395

Mechanism of C-C coupling reactions of aromatic halides, promoted by $\text{Ni}(\text{COD})_2$ in the presence of 2,2'-bipyridine and PPh_3 , to give biaryls (T. Yamamoto, S. Wakabayashi and K. Osakada), (428) 223

Synthesis and reactivities of novel $\eta^2\text{-(C}_6\text{O)}$ alkylphenylketene complexes of nickel. Coordination-mode switching reaction of the ketene ligand (A. Miyashita, R.-j. Sugai and J.-i. Yamamoto), (428) 239

Komplexbekatalyse. XXXVII. $\eta^3\text{,}\eta^2\text{,}\eta^2\text{-Dodeca-2(}E\text{,}6\text{(}E\text{,}10\text{(Z)-triene-1-yl-nickel(II)-tetrakis[3,5-bis(trifluormethyl)phenyl]borat}$ als Katalysator für die 1,4-*cis*-Polymerisation des Butadiens. Aktivität und Selektivität des $\text{C}_{12}\text{-Allylnickel(II)-Kations}$ (R. Taube and S. Wache), (428) 431

Reactions of pentafluorophenyl germanium hydrides with nickelocene (L.V. Pankratov, V.I. Nevodchikov, L.N. Zakharov, M.N. Bochkarev, I.V. Zdanovich, V.N. Latyaeva, A.N. Lineva, A.S. Batsanov and Y.T. Struchkov), (429) 13

Rupture of the triple $\text{C}\equiv\text{C}$ and $\text{C}\equiv\text{O}$ bonds in the mixed-metal clusters with Ni_2Mo_2 and Mo_4Co_3 cores. Synthesis and molecular structures of $\text{Cp}_4\text{Ni}_2\text{Mo}_2(\mu_3\text{-CPh})_2(\mu_3\text{-CO})_2$ and $\text{Cp}_4\text{Mo}_4(\mu_3\text{-OH})(\mu_3\text{-CPh})_2(\mu_6\text{-C})\text{Co}_3(\mu\text{-CO})_3(\text{CO})_3$ (A.D. Shaposhnikova, M.V. Drab, G.L. Kamalov, A.A. Pasynskii, I.L. Ereminko, S.E. Nefedov, Y.T. Struchkov and A.I. Yanovsky), (429) 109

Reactions of unstable compound $\{\text{CpNiCH}_3\}$ with acetylenes (S. Pasynkiewicz, M. Popławska and R. Mynott), (429) 135

Photolytisch induzierte Erzeugung und komplexchemische Stabilisierung eines Molybdän-Arsen-Dreifachbindungssystems; Röntgenstrukturanalysen der Cluster $\mu_3\text{-AsCp}_3\text{Mo}_2\text{Ni}(\text{CO})_4$, $\text{As}_2\text{Cp}_6\text{Mo}_4\text{Ni}_2(\text{CO})_7$ und $\mu_3\text{-AsCpMoPt}_2(\text{CO})_4(\text{PPh}_3)_2$ (M. Gorzellik, B. Nuber, T. Bohn and M.L. Ziegler), (429) 173

- Reactions of neutral and ionic square-planar organometallic nickel compounds with ethylene (R.M. Ceder, J. Cubillo, G. Muller, M. Rocamora and J. Sales), (429) 391
- Synthesis and properties of 2- or 2,5-substituted thiophene and 2- or 2,5'-substituted dithiophene derivatives of platinum (S. Kotani, K. Shiina and K. Sonogashira), (429) 403
- Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. IV. Synthese von $\text{Li}_2\text{Ni}-(\text{CH}_2\text{CH}_2-\text{CH}_2\text{CH}_2)_2(\text{Et}_2\text{O})$ und seine Umsetzung mit Kohlenmonoxid zu $\text{Li}_2\text{Ni}_6(\text{CO})_{12}(\text{thf})_6$ (H.-O. Fröhlich, B. Hippler and B. Hofmann), (430) 133
- The crystal structures of $(\text{R}_3\text{P})_2\text{Ni}$ -anthracene ($\text{R} = \text{Et}, \text{Bu}$) (A. Stanger and R. Boese), (430) 235
- Optically active complexes of transition metals (Rh^{I} , Ru^{II} , Co^{II} and Ni^{II}) with 2-aminocarbonylpyrrolidine ligands. Selective catalysts for hydrogenation of prochiral olefins (A. Corma, M. Iglesias, C. Del Pino and F. Sánchez), (431) 233
- Homoleptische metallorganische Innerkomplexe des 8-Dimethylaminonaphthalyl-1-Anions (H. Dreys), (433) C1
- Reactivity of the $[(\text{Ni}(\text{C}_6\text{F}_5)_2(\mu\text{-OH}))_2]^{2-}$ ion towards β -diketones, 8-hydroxiquinoline and heterocyclic thiones. Crystal structure of $[\text{NBu}_4]\text{Ni}(\text{C}_6\text{F}_5)_2(\text{C}_5\text{H}_4\text{NS-2})$ (G. López, G. Sánchez, G. García, J. García, A. Martínez, J.A. Hermoso and M. Martínez-Ripoll), (435) 193
- A kinetic study of the coordination template effect in the nickel-catalysed formation of arylphosphonium salts (D.W. Allen and P.E. Cropper), (435) 203
- Darstellung und Eigenschaften von und Reaktionen mit metallhaltigen Heterocyclen. LXXVIII. Das Verhalten von monomeren und dimeren Thiophosphinitonickel-Komplexen gegenüber Alkinen und Phosphaalkinen (E. Lindner, A. Nothdurft, R. Fawzi and C. Maichle), (435) 213
- Aktivierung von CO_2 an Übergangsmetallzentren: Zum Ablauf der homogen-katalytischen Bildung von 2-Pyron aus Kohlendioxid und Hex-3-in an Nickel(0)-Fragmenten (D. Walther, G. Bräunlich, R. Kempe und J. Sieler), (436) 109
- Organoerdmetall-Komplexe von *d*-Block-Elementen. V. Selektive Reduktion von Bis[carbonyl(η^5 -cyclopentadienyl)nickel] zum Carbonyl(η^5 -cyclopentadienyl)nickel-Anion durch C_8K . Synthese und Struktur von ($\eta^5\text{-C}_5\text{H}_5$) 2 (CO) Ni^- -Ga[$\text{CH}_2\text{C}(\text{CH}_3)_3\text{L}_2$] 2 (OC $_4\text{H}_8$) (R.A. Fischer, J. Behm, E. Herdtweck und C. Kroneder), (437) C29
- Nickel-catalyzed reactions of 3,4-benzo-1,1,2,2-tetraethyl-1,2-disilacyclobutene (M. Ishikawa, H. Sakamoto, S. Okazaki and A. Naka), (439) 19
- η^1 - und η^2 -Komplexe von Cp^* -substituierten Diphosphenen und Arsaphosphenen: Synthese, Struktur und Umlagerung unter Cp^* -Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammmer), (439) 279
- Reactions of biscyclopentadienylnickel with lithium and magnesium aryls (A. Pietrzykowski and S. Pasynkiewicz), (440) 401
- Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. V. Synthese von $\text{Li}_2\text{Ni}-(\text{CH}_2\text{CH}_2-\text{CH}_2\text{CH}_2)_2(\text{Solv})$, aus $\text{K}[\text{Ni}(\text{NPh}_2)_3](\text{THF})$ und $\text{LiCH}_2\text{CH}_2\text{CH}_2-\text{CH}_2\text{Li}$; Molekülstruktur von $(\text{Li}(\text{THF})_2)_2\text{Ni}(\text{CH}_2\text{CH}_2\text{CH}_2-\text{CH}_2)_2$ (H.-O. Fröhlich, R. Wyrwa und H. Görts), (441) 169
- Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241
- Synthesis, electrochemistry and complexation studies of new redox active bisferrrocene acyclic and macrocyclic thioethers (P.D. Beer, J.E. Nation, M.E. Harman and M.B. Hursthouse), (441) 465
- Reactions of the unstable compound $\{\text{CpNiCH}_3\}$ with 2-butyne. Formation of tetrameric complex $[\text{CpNi}-\text{C}(\text{CH}_3)=\text{C}(\text{CH}_3)_2]_4$ (S. Pasynkiewicz, A. Pietrzykowski and M. Popławska), (443) 131
- Dialkyl complexes of nickel(II) containing chelating diphosphines. The crystal structure of $[\text{Ni}(\text{CH}_2\text{SiMe}_2\text{Ph})_2(\text{iPr}_2\text{PCH}_2\text{CH}_2\text{P}^{\text{i}}\text{Pr}_2)]$ (O. Boutry, M.C. Nicasio, M. Paneque, E. Carmona, E. Gutiérrez and C. Ruiz), (444) 245
- Nickelocene-catalyzed polymerization of acetylenes (W.E. Douglas and A.S. Overend), (444) C62
- Coordination of a tripodal phosphine sulphide P_2S towards various transition metal ions (H.-E. Wang, M.-C. Cheng, G.-H. Lee, S.-M. Peng and S.-T. Liu), (445) 171
- Metallacyclische Acyl-carboxylate des Nickels: Reaktivität der Acylgruppe und Synthesepotential bei Kreuzkopplungsreaktionen mit Alkyhalogeniden (R. Fischer, D. Walther, R. Kempe, J. Sieler und B. Schönecker), (447) 131
- Stereospecific reactions of the " $\text{M}(\text{C}_6\text{F}_5)_2$ " ($\text{M} = \text{Pd}$ or Pt) synthons towards *cis*-dithiolato complexes (R. Usón, J. Forniés, M.A. Usón and S. Herrero), (447) 137
- Relative metal-ion affinities of alkylamines in the gas-phase (L.-Z. Chen and J.M. Miller), (448) 225
- Ferrocene derivatives. XXVIII. The metallation of phenylazoferrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177
- Niobium**
- Electrochemical studies on organometallic compounds. XL. Reactivity of organometallic niobium complexes towards diphenylacetaldehyde. Electrogeneration and ESR characterization of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2(\eta^2\text{-Ph}_2\text{CHCHO-C},\text{O})]$ (D. Lucas, H. Chollet, Y. Mugnier, A. Antíñolo, A. Otero and M. Fajardo), (426) C4
- Electrochemistry and bonding in biscyclopentadienyl complexes of molybdenum, tungsten, and niobium with *o*-phenanthroline and bipyridine. Crystal and molecular structures of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_5)_2(\text{bpy})]\text{PF}_6$ and $[\text{M}(\eta^5\text{-C}_5\text{H}_5)_2(\text{o-phen})]\text{PF}_6$ ($\text{M} = \text{Mo}$ or W) (M.J. Calhorda, M.A.A.F. de C.T. Carrondo, A. Bram, P.N. Olsen, A.R. Dias, A.M. Freitas, M.H. Garcia and M.F.M. Piedade), (426) 195
- Pentamethylcyclopentadienyl niobium(III) nitrene complexes containing phosphine, carbonyl, olefin and acetylene ligands (U. Siemeling and V.C. Gibson), (426) C25
- Agostic interaction in early transition metal alkyls and their role in catalytic activity for olefin polymerizations (K. Mashima and A. Nakamura), (428) 49
- Facile synthesis of α,β -unsaturated niobium acyls via γ -hydride abstraction (A.S. Hneihen, M.C. Fermin, J.J. Maas and J.W. Bruno), (429) C33
- Early-transition-metal ketene complexes: synthesis, reactivity and structure of ketene complexes of bis(trimethylsilyl)niobocene, X-ray structure of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Br}(\text{Ph}_2\text{C}=\text{C}-\text{O}-\text{C},\text{O})]$ (A. Antíñolo, A. Otero, M. Fajardo, C. Lopez-Mardomingo, D. Lucas, Y. Mugnier, M. Lanfranchi and M.A. Pellinghelli), (435) 55
- Electrochemical studies on organometallic compounds. XLI. Electrogeneration and ESR characterization of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Cl}(\text{POMe})_3]^+$ (D. Lucas, Y. Mugnier, A. Antíñolo, A. Otero and M. Fajardo), (435) C3
- Präparative, ^{11}B , ^{93}Nb -NMR-spektroskopische und strukturelle Untersuchungen an $\text{Cp}_2\text{NbBH}_4^-$ und $[\text{CpNb}(\text{B}_2\text{H}_6)_2]$ -Komplexen (H. Brunner, G. Gehart, W. Meier, J. Wachter, B. Wrackmeyer, B. Nuber und M.L. Ziegler), (436) 313
- Synthesis of η -cyclopentadienylidene-4-imidopropylniobium derivatives $[\text{Nb}(\eta : \sigma\text{-C}_5\text{H}_4(\text{CH}_2)_3\text{N})\text{Cl}_2]$ and $[\text{Nb}(\eta : \sigma\text{-C}_5\text{H}_4(\text{CH}_2)_3\text{N})(\text{PMMe}_3)\text{Cl}_2]$ (D.M. Antonelli, M.L.H. Green and P. Mountford), (438) C4
- Electrochemical studies on organometallic compounds. XLII. Electrogeneration and spectroscopic characterization of $[\text{Nb}(\eta^5$

$C_5H_4SiMe_3)_2(PhC \equiv CPh-C,C)$ (H. Chollet, D. Lucas, Y. Mugnier, A. Antiñolo, A. Otero and M. Fajardo), (441) 45

Osmium

Synthesis and spectroscopic characterization of the heteronuclear diphosphine linked cluster $Os_3(CO)_{11}(Ph_2PCH_2PPh_2)Rh_6(CO)_{15}$ (S.P. Tunik, M.V. Osipov and A.B. Nikolskyi), (426) 105

Synthesis and X-ray structure of the heteronuclear cluster, $(\mu-H)_2(\eta^5-C_5H_5)IrOs_3(CO)_{10}$ (L.-Y. Hsu, W.-L. Hsu, D.A. McCarthy, J.A. Krause, J.-H. Chung and S.G. Shore), (426) 121

Reactions of carbonyl clusters with potentially tridentate thioethers: crystal and molecular structures of $[Os_4(CO)_{13}([12]aneS_3)]$ and $[Ru_6(CO)_{15}(\mu_4-\eta^2-CO)([12]aneS_3)]$ ($[12]aneS_3 = [(CH_2)_3S_3]$) (A.J. Edwards, B.F.G. Johnson, F.K. Khan, J. Lewis and P.R. Raithby), (426) C44

Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CI. Stereoselektive Synthese gemischtalkylierter Osmiumoxide (W.A. Herrmann, S.J. Eder, P. Kiprof and P. Watzlowik), (428) 187

$[Rh(CO)_4]^-$, $[Rh_5(CO)_{15}]^-$, and bimetallic clusters as catalysts for the carbonylation of nitrobenzene to methyl phenylcarbamate (F. Ragaini, S. Cenini, A. Fumagalli and C. Crotti), (428) 401

Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CIII. *cis*-Dimethylbis(2,2-dimethylpropyl)oxoosmium(VI): Röntgenstrukturanalyse eines gemischtalkylierten Osmiumoxids (S.J. Eder, W.A. Herrmann and P. Kiprof), (428) 409

The synthesis and crystal structure of 1,2,3-trithiaf[3]osmocenophane (E.W. Abel, N.J. Long, A.G. Osborne, M.B. Hursthouse and M.A. Mazid), (430) 117

Synthesis of molybdenum-triosmium cluster complexes: reaction of $(\mu-H)_2Os_3(CO)_{10}$ with $(\eta^5-C_5H_5)CO_2Mo=CC_6H_4Me$ (J.T. Park, J.-J. Cho, K.-M. Chun and S.-S. Yun), (433) 295

Synthesis, characterization and hindered rotation of triosmium clusters $(\mu-H)Os_3(CO)_{10}L[ON(CF_3)_2]$ [$L = CO, CH_3CN$ and $PhCN$] (H.G. Ang, C.H. Koh and W.L. Kwik), (435) 149

Triosmium clusters containing bridging sulphur, hydrido, and diphosphine ligands: a study of hydride site preferences and mobility (K.A. Azam, S.E. Kabir, A. Miah, M.W. Day, K.I. Hardcastle, E. Rosenberg and A.J. Deeming), (435) 157

Synthesis and structural characterization of $[HOs_3(CO)_9(OEt)(R_2C_2)]$ ($R = Me$ or Ph) (D. Braga, P. Sabatino, B.F.G. Johnson, J. Lewis and A. Massey), (436) 73

Triosmium carbonyl cluster complexes of bridging phosphonate, arsonate, and phosphinate ligands (G.R. Frauenhoff, J.-C. Liu, S.R. Wilson and J.R. Shapley), (437) 347

Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXV. Dihalogenoxoosmium(VI)-Komplexe des Pinakols: Synthese, Spektroskopie und Alkylierung (W.A. Herrmann und P. Watzlowik), (437) 363

Novel synthesis and vibrational analysis of (arene)osmium(II) complexes of the type $[C_6H_3(CH_3)_3]OsH_2(L)$ (T. Polzer, A. Ellebracht, W. Kiefer, U. Wecker and H. Werner), (438) 319

Structural aspects of the osmium and ruthenium clusters $[M_3(CO)_{10}(\text{alkyne})]$ and related substituted compounds (A.J. Deeming and A.M. Senior), (439) 177

Synthesis of tetrานuclear heterometallic cluster complexes via condensation of triosmium alkyne complexes $Os_3(CO)_{10}(C_2R_2)$, $R = Tol$ and Me , and mononuclear tungsten acetylide complexes $LW(CO)_3C \equiv CR'$, $L = Cp$ and Cp^* , $R' = Ph$ and $'Bu$ (Y. Chi, R.-C. Lin, C.-C. Chen, S.-M. Peng and G.-H. Lee), (439) 347

The use of bis(diphenylphosphinoacetylene) and its digold derivative as linking groups in osmium cluster chemistry. Crystal structures

of $[(Os_3(CO)_{11})_2(dppa)]$, $[Os_3(CO)_{10}(dppa)]_2$ and $[Os_4H(CO)_{12}-Au(dppa)]_2$ ($dppa = Ph_2PC \equiv CPh_2$) (A.J. Amoroso, B.F.G. Johnson, J. Lewis, A.D. Massey, P.R. Raithby and W.T. Wong), (440) 219

Das Verhalten von stereoisomeren Bistriflaten des Tetrahydrofurans gegenüber ein- und zweiwertigen Carbonylmittallaten (E. Lindner, M. Pabel, R. Fawzi und M. Steimann), (441) 63

On the problem of stabilization of α -metallocenylcarbocations. Synthesis, 1H and ^{13}C NMR spectra of metallocenyldiphenylmethyl hexafluorophosphates and X-ray investigation of $(C_5H_5OsC_5H_4-CPh_2)^+PF_6^-$ (U. Turpeinen, A.Z. Kreindlin, P.V. Petrovskii and M.I. Rybinskaya), (441) 109

Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXI. Perfluorpinakolat-Komplexe von Rhenium(VII) und Osmium(VI) (W.A. Herrmann und P. Watzlowik), (441) 265

Reaction of $Ru_3(CO)_{12}$ with ferrocenylacetylene. Crystal and molecular structure of $Ru_2(CO)_6[C_4H_2(C_5H_4FeC_5H_5)_2]$ (A.A. Koridze, A.I. Yanovsky and Y.T. Struchkov), (441) 277

Synthesis of bridged and linked ruthenium and osmium carbonyl clusters containing a $[Au_2(Ph_2PCH_2CH_2PPh_2)]^{2+}$ unit. The crystal and molecular structures of $Ru_5C(CO)_{14}Au_2(Ph_2PCH_2CH_2PPh_2)$ and $\{Os_4H_3(CO)_{12}\}_2Au_2(Ph_2PCH_2CH_2PPh_2)$ (A.J. Amoroso, A.J. Edwards, B.F.G. Johnson, J. Lewis, M.R. Al-Mandhary, P.R. Raithby, V.P. Saharan and W.T. Wong), (443) C11

The controlled assembly of high nuclearity osmium clusters from smaller fragments (J. Lewis and J.R. Moss), (444) C51

Isolation and structural characterisation of $Os_5(CO)_{18}$, a postulated intermediate in the conversion of $Os_5(CO)_{19}$ into $Os_5(CO)_{16}$ (D. Coughlin, J. Lewis, J.R. Moss, A.J. Edwards and M. McPartlin), (444) C53

Synthesis and X-ray structure of the octaosmium carbido cluster $Os_8C(CO)_{22}$ (A.J. Amoroso, B.F.G. Johnson, J. Lewis, C.-K. Li, P.R. Raithby and W.-T. Wong), (444) C55

Synthesis of novel selenium-containing osmium carbonyl clusters: X-ray structures of $[Os_5(\mu-H)(CO)_{15}(\mu_3-SePh)]$ and $[Os_6(\mu-H)(CO)_{18}(\mu-SeH)]$ (T.M. Layer, J. Lewis, A. Martín, P.R. Raithby and W.-T. Wong), (444) C57

Stereoselektive Bildung von Organoosmium-Komplexen mit Aminocarboxylat-Liganden (H. Werner, T. Daniel, O. Nürnberg, W. Knaup und U. Meyer), (445) 229

Cluster chemistry. LXXXVII. Some homo- and hetero-nuclear complexes derived from $C_2(PPh_2)_2$: Crystal structures of $Re_3(\mu-H)_3(\mu-dppa)(CO)_{10}$ ($dppa = C_2(PPh_2)_2$), $Re_3(\mu-H)_3(CO)_{11}-(PPh_2\mu-C_2Ru_2(\mu-PPh_2)CO)_6$) and $Os_3Ru_2(\mu_5-C_2PPh_2\mu-PPh_2)CO)_{13}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (447) 91

Interaction between $[(\eta^5-C_5Me_5)_2OsH]^+PF_6^-$ and nitrogen or oxygen bases. Hydrogen bonds of the type $[OsH]^+ \cdots B$ (L.M. Epstein, E.S. Shubina, A.N. Krylov, A.Z. Kreindlin and M.I. Rybinskaya), (447) 277

Synthesis, and spectral and X-ray structural studies of a novel delocalised triphosphazene $(CF_3)_2P=N=P(CF_3)_2=N=P(CF_3)_2 = N=PPh_3$ (H.G. Ang, Y.M. Cai and W.L. Kwik), (448) 219

Cyclopentadienyl-ruthenium and -osmium chemistry. XXXIX. Syntheses of novel alkynyl-ruthenium complexes. X-Ray structures of two forms of $\{Ru(PPh_3)_2(\eta-C_5H_5)_2(\mu-C_4)\}$ and of $Ru(C \equiv CC(O)Me)(PPh_3)_2(\eta-C_5H_5)$ (M.I. Bruce, P. Hinterding, E.R.T. Tiekkink, B.W. Skelton and A.H. White), (450) 209

Reaction of $[(\mu-H)Os_3(CO)_{11}]^-$ anion with dioxygen (C.J. Cathey, B.F.G. Johnson, J. Lewis, P.R. Raithby and W.T. Wong), (450) C12

Palladium

Reactions of η^2 -13-acyltetracarbonylmanganese complexes derived from podocarpic acid with alkenes; cyclopentaannulation of ring C (R.C. Cambie, M.R. Metzler, C.E.F. Rickard, P.S. Rutledge and P.D. Woodgate), (426) 213

Synthesis and evaluation of the catalytic properties of homo- and hetero-bimetallic complexes containing bridging diphenylphosphido ligands (Z. He, N. Lugar, D. Neibecker, R. Mathieu and J.-J. Bonnet), (426) 247

Synthese und Struktur von (η^3 -Allyl)palladium(II)benzamidinaten (F.T. Edelmann, W. Ziegler and U. Behrens), (426) 261

Cyclometallated complexes of bis(*N*-benzylidene)-1,4-phenylenediamines. Synthesis and crystal structure of [1,4-(Pd[2,3,4-(MeO)₃C₆HC(H)=N](Br))₂C₆H₄]₂; a novel tetrานuclear cyclometallated palladium(II) complex (J.M. Vila, M. Gayoso, M^a.T. Pereira, M^a.C. Rodriguez, J.M. Ortigueira and M. Thornton-Pett), (426) 267

Synthesis and structure of binuclear thioazobenzene pallada-cycles and their reaction with *m*-chloroperbenzoic acid: formation and structure of dinuclear azophenolates (S. Chattopadhyay, C. Sinha, S.B. Choudhury and A. Chakravorty), (427) 111

Metal-ion dependent reactivity of 2-(2'-thienyl)pyridine (Hthpy) (E.C. Constable and L.R. Sousa), (427) 125

Cyclopalladated complexes. Synthesis and crystal structure of di- μ -chloro-bis([2,6-dimethyl-*N*-(benzylidene)phenylaminato-*C*^{2'},*N*]-palladium(II)) (A. Crispini, G. De Munno, M. Ghedini and F. Neve), (427) 409

Metallation of aliphatic carbon atoms. VI. Reactions of 2-pivaloylpyridine with palladium(II) acetate resulting in cyclopalladation and nucleophilic attacks on carbonyl carbon (K. Hiraki, M. Nakashima, T. Uchiyama and Y. Fuchita), (428) 249

Isomerization of alkynemono-ols catalyzed by palladium(0) complex and diols (X. Lu, J. Ji, C. Guo and W. Shen), (428) 259

Catalytic asymmetric arylation of *N*-substituted 2-pyrrolines with aryl triflates (F. Ozawa and T. Hayashi), (428) 267

Palladium- and platinum-catalyzed cyclocarbonylation reactions of substituted allyl acetates (Y. Ishii and M. Hidai), (428) 279

Recent developments in the chemistry of olefin complexes of palladium and platinum: some novel findings as well as elaborate extensions (H. Kurosawa and I. Ikeda), (428) 289

Synthesis and properties of 2- or 2,5-substituted thiophene and 2- or 2,5'-substituted dithiophene derivatives of platinum (S. Kotani, K. Shiina and K. Sonogashira), (429) 403

Palladium-catalyzed cycloaddition of carbon dioxide with methoxyalene (T. Tsuda, T. Yamamoto and T. Saegusa), (429) C46

Palladium(II) coordination and cyclometallated complexes derived from 3- and 5-aryl-substituted pyrazoles (M.T. Alonso, O. Juanes, J. de Mendoza and J.C. Rodríguez-Ubis), (430) 335

Palladium(II) coordination and cyclometallated complexes derived from 1-tert-butylpyrazole (M.T. Alonso, O. Juanes, J. de Mendoza and J.C. Rodríguez-Ubis), (430) 349

Influence of ligands and anions on the insertion of alkenes into palladium-acyl and palladium-carbomethoxy bonds in the neutral complex (dppp)Pd(C(O)CH₃)Cl and the ionic complexes [(P-P)PdR(L)]⁺SO₃CF₃⁻ (P-P = dppe, dppp, dppb; R = C(O)CH₃, L = CH₃CN, PPh₃; R = C(O)OCH₃, L = PPh₃) (G.P.C.M. Dekker, C.J. Elsevier, K. Vrieze, P.W.N.M. Van Leeuwen and C.F. Roobeek), (430) 357

Trimethylsilylcyclooctadiene-transition metal complexes: metal-catalysed protodesilylation of cyclic vinylsilanes, and transfer hydrogenation promoted by the displaced silyl group (B.S. Bandodkar and G. Nagendrappa), (430) 373

Reactions with nucleophiles and cathodic electrochemical behaviour of some neutral and cationic diamino-, dioxy- and aminoxycar-

bene complexes of palladium(II) and platinum(II) (R. Bertani, M. Mozzon, R.A. Michelin, T.J. Castilho and A.J.L. Pombeiro), (431) 117

Cyclometallation of phenylhydrazones of ketones. Crystal and molecular structure of [PdCl(C₆H₄C(C₆H₅)=NNH(C₆H₄-4-Cl))- (PPh₃)₂] (J. Granell, R. Moragas, J. Sales, M. Font-Bardía and X. Solans), (431) 359

Synthesis of halogeno, pseudohalogeno, and carboxylatopalladium(IV) complexes by halogen exchange. Crystal structure of azido(2,2'-bipyridyl)-benzylpalladium(II), formed on reductive elimination of ethane from Pd(N₃)Me₂(CH₂Ph)(bpy) (A.J. Canty, P.R. Traill, B.W. Skelton and A.H. White), (433) 213

Conformational studies in palladium(IV) and platinum(IV) chemistry. Crystal structure of the 1,1-bis(pyrazol-1-yl)-ethane complex fac-PtIMe₃((pz)₂CHMe-*N,N'*) (P.K. Byers, A.J. Canty, R.T. Honeyman, B.W. Skelton and A.H. White), (433) 223

The synthesis and structural characterization of a penta-palladium cluster, [Pd₅(μ -SO₂)₃(μ -CNXyl)₂(CNXyl)₅], with linear bridging isocyanide ligands (A.D. Burrows, H. Fleischer and D.M.P. Mingos), (433) 311

Palladium- and light-enhanced ring-opening of oxiranes by copper chloride (J. Muzart and A. Riahi), (433) 323

Pallada(IV)cyclopentane chemistry, including the synthesis of tris(pyrazol-1-yl)borate complexes (A.J. Canty and P.R. Traill), (435) C8

Komplexchemie perhalogenierter Cyclopentadiene und Alkine. IX. Darstellung einiger Phosphoniumacetylid-Komplexe und Struktur von (Ph₃P)Cl₂Pd(C≡CPPPh₃) (K. Sünkel), (436) 101

Synthesis of palladium(II) and platinum(II) *N,N*-dialkyldithiocarbamates starting from hydroxo-halophenyl complexes (G. López, J. Ruiz, G. García, C. Vicente, J.M. Martí and V. Rodríguez), (436) 121

Palladium-catalyzed rearrangement of silanes containing oxygen or halogen α to silicon (V. Gevorgyan, L. Borisova and E. Lukevics), (436) 277

Palladium-assisted formation of carbon-carbon bonds. Stoichiometric synthesis of indenols and indenones. Catalytic synthesis of an indenol (J. Vicente, J.-A. Abad and J. Gil-Rubio), (436) C9

A simple direct synthesis of an alkylaromatic palladacycle (M. Catellani and G.P. Chiusoli), (437) 369

Palladium-catalyzed norbornene–carbon monoxide co-oligomerization initiated by aryl groups and terminated by double bond formation (E. Dalcanale, Z. An, L.P. Battaglia, M. Catellani and G.P. Chiusoli), (437) 375

Anodic oxidation of the ferrocenyl-derived β -ketophosphine [Ph₂PCH₂C(O)(η^5 -C₅H₄)Fe(η^5 -C₅H₅)] and of two of its palladium complexes. Ferricinium-mediated oxidation of palladium (II)-phosphine complexes (A. Louati, M. Gross, L. Douce and D. Matt), (438) 167

Organometallics of diphosphazanes. VI. Reactions of bicyclo-1,3,2- λ^3 ,4- λ^3 -diazadiphosphetidines (bicyclic diphosphazanes) with Group 6 metal carbonyls and palladium and platinum(II) complexes (T.K. Prakash, S.S. Krishnamurthy, S.E. Davies, M.F. Meidine and J.F. Nixon), (438) 241

Organometallic nucleophiles. Mechanism of halide displacement at saturated carbon by 2-pyridyl and 4-pyridyl complexes [M(dmtc)(C₆H₄N-CⁿXL)] (M = Pd, Pt; dmtc = dimethyldithiocarbamate; n = 2,4; L = tertiary phosphine) (L. Canovese, P. Uguagliati, F. Di Bianca and B. Crociani), (438) 253

Palladium(II) complexes with the cyclometallated C–Y–N(*sp*²)-Pd–C(*sp*²) ring (Y = N,C) and (R)-(+)-hydroxy-methylene camphor (M. Ghedini, F. Neve, D. Pucci, E. Cesariotti and M. Grassi), (438) 343

Synthesis and structure of a thioazobenzene palladacycle: oxygen insertion into the Pd-C bond by *m*-chloroperbenzoic acid (C.K. Pal, S. Chattopadhyay, C. Sinha and A. Chakravorty), (439) 91
 The palladium-catalysed addition of hexamethylditin to 1,3-dienes (T.N. Mitchell, B. Kowall, H. Killing and C. Nettelbeck), (439) 101

Synthesis and characterization of dinuclear palladium(II) complexes containing both chloro and organochalcogenides as bridging ligands (V.K. Jain and S. Kannan), (439) 231

Bildung eines oxalatverbrückten Phenylpalladium(II)-Komplexes aus Phenylglyoxylsäure und Palladiumacetat: Synthese und Struktur von $[Bu_2S(Ph)Pd(\mu-C_2O_4)Pd(Ph)(SBu_2)]$ (R. Krämer, K. Polborn und W. Beck), (441) 333

Synthesis, electrochemistry and complexation studies of new redox active bisferrocene acyclic and macrocyclic thioethers (P.D. Beer, J.E. Nation, M.E. Harman and M.B. Hursthouse), (441) 465

Some attempts to prepare chelating nitrophenylpalladium(II) complexes. Synthesis of covalent arylpalladium(II) perchlorate complexes. Crystal and molecular structure of *catena-trans*- $\{[Pd(C_6H_2(NO_2)_3-2,4,6)(tht)_2(H_2O)]ClO_4\}_n$ (tht = tetrahydrothiophene) (J. Vicente, A. Arcas, M.V. Borrachero, E. Molins and C. Miravitles), (441) 487

Solvent, *N*-alkyl substituent and coordination effects on the electrochemistry of 2,11-diaza-5,8-dithia[12](1,1')ferrocenophanes (R.A. Holwerda, J.S. Kim, T.W. Robison, R.A. Bartsch and B.P. Czech), (443) 123

Sodium tetraphenylborate as a phenylating reagent in the palladium-catalyzed phenylation of alkenes and acid chlorides (C.S. Cho, K. Itoh and S. Uemura), (443) 253

A new type of stabilized carbonucleophiles with a carboranyl group (S.C. Shim, J.G. Shim, S.Y. Chae, S.Y. Lee and V.N. Kalinin), (443) C22

Trisubstituted alkenes from alkynes and aryl iodides by boron-to-zinc transmetalation and palladium(0) catalyzed cross coupling (K.A. Agrios and M. Srebnik), (444) 15

New aspects of the synthesis of dimethyl carbonate via carbonylation of methyl alcohol promoted by methoxycarbonyl complexes of palladium(II) (G. Cavinato and L. Toniolo), (444) C65

Coordination of a tripodal phosphine sulphide P_2S towards various transition metal ions (H.-E. Wang, M.-C. Cheng, G.-H. Lee, S.-M. Peng and S.-T. Liu), (445) 171

Cyclometallated complexes of Pd^{II} and Mn^I with *N,N*-terephthalylidenebis(cyclohexylamine) (J.M. Vila, M. Gayoso, M.T. Pereira, M. López, G. Alonso and J.J. Fernández), (445) 287

Convenient synthesis of $[\eta^3\text{-}1\text{-}(formyl)allyl]$ - and $[\eta^3\text{-}1\text{-}(dimethoxy-methyl)allyl]palladium$ chlorides (S. Ogoshi, K. Hirako, J. Nakaniishi, K. Ohe and S. Murai), (445) C13

Synthèse d'aryl- et hétéroarylsilanes par scission de l'hexaméthyl-disilane (P. Babin, B. Bennetau, M. Theurig et J. Dunoguès), (446) 135

Stereospecific reactions of the " $M(C_6F_5)_2$ " (M = Pd or Pt) synthons towards *cis*-dithiolato complexes (R. Usón, J. Forniés, M.A. Usón and S. Herrero), (447) 137

Synthesis and characterization of *cis*-bis[(*p*-tolylsulfonyl)methyl]palladium(II) complexes (P. Espinet, J.M. Martínez-llarduya and C. Pérez-Briso), (447) 145

Studies on Pd^{II} -catalyzed cyclization of 4'-hydroxy-2'-alkenyl 2-alkynoates (S. Ma and X. Lu), (447) 305

An easy approach to 1-silylated ketones and asymmetrical 1,6- and 1,8-dicarbonyl compounds (V. Fiandanese, A. Punzi and N. Ravasio), (447) 311

Cyclometallated complexes of palladium(II) with the diphosphines *trans*- $Ph_2PCH=CHPPh_2$, *cis*- $Ph_2PCH=CHPPh_2$, and $Ph_2P(CH_2)_4PPh_2$. The X-ray crystal structure of $[\{Pd(2,4-Me_2C_6H_2-$

$C(H)\equiv NCy\}]_2(\mu\text{-}Ph_2P(CH_2)_4PPh_2)(\mu\text{-}Cl)_2]$ (J.M. Vila, M. Gayoso, A. Fernandez, N.A. Bailey and H. Adams), (448) 233

Cyclopalladated compounds. Structural studies on dinuclear azobenzene complexes (A. Crispini, M. Ghedini and F. Neve), (448) 241

Synthesis and characterisation of new 2-bromovinyl selenides and their platinum group metal complexes (P.K. Khanna and C.P. Morley), (450) 109

Ferrocene derivatives. XXVIII. The metallation of phenylazoferrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177

Stereo and regioselectivity in the phenylation of cationic allylpalladium(II) α -diimine complexes by tetraphenylborate anion (B. Crociani, S. Antonaroli, F. Di Bianca and A. Fontana), (450) 218

Hydroformylation and hydrocarbethoxylation of 1,2-dicarbethoxy-1,2,3,6-tetrahydropyridazine and 1,2-dicarbethoxy-1,2,3,4-tetrahydropyridazine (G. Menchi, S. Paganelli, U. Matteoli, A. Scrivanti and C. Botteghi), (450) 229

Steric effects of the 2-(diphenylphosphino)pyridine bridging ligand in the synthesis of binuclear palladium(II) complexes (G. De Munno, G. Bruno, C.G. Arena, D. Drommi and F. Faraone), (450) 263

Cyclometallated derivatives of palladium(II) and platinum(II) derived from 6-t-butyl-2,2'-bipyridine (S. Stoccero, G. Chelucci, M.A. Cinelli, A. Zucca and G. Minghetti), (450) C15

Phosphorus

Phenylantimony(III) diorganophosphorodithioates: the crystal structure of diphenylantimony(III) di-isopropylphosphorodithioate, $Ph_2Sb_2P(O^1Pr)_2$; unusual polymerisation through semibonding (C. Silvestru, M. Curtui, I. Haiduc, M.J. Begley and D.B. Sowerby), (426) 49

Aminosubstituenten als Elektronendonatoren in Clustern (B. Eber, G. Huttner, W. Imhof, J.C. Daran and Y. Jeannin), (426) 87

Metall- π -Komplexe von Benzolderivaten. XXXIX. $Dil[1,4\text{-bis}(di-phenylphosphano)\text{-}\eta^6\text{-benzol}]chrom$: ein Baustein für polynukleare Bimetallkomplexe? (Ch. Elschenbroich, J. Sebbach, B. Metz and G. Heikenfeld), (426) 173

Metallorganische Lewis-Säuren. XLIV. Rheniooxy(organyl)carben-Komplexe von Chrom, Molybdän und Wolfram (P.M. Fritz, J. Breimair, B. Wagner and W. Beck), (426) 343

Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphan-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe $(CO)_3(R^1_3P)(R^1_3Si)Fe-ML_n$ (M = Cu, Ag, Au, Hg) (G. Reinhard, B. Hirle and U. Schubert), (427) 173

Syntheses of $W(PhC\equiv CPh)_3(\eta^1\text{-}PPh_2CH_2PPh_2)$ and $W(PhC\equiv CPh)_3(PPh_3)$ from $W(PhC\equiv CPh)_3(NCCH_3)$ (W.-Y. Yeh, C.-S. Ting and C.-f. Chih), (427) 257

Übergangsmetall-Heteroallen-Komplexe. XXV. Reaktionen von Ketenimin mit $Ru_3(CO)_12$ (W. Ziegler and U. Behrens), (427) 379

Isolation of cationic cyclopentadienyl iron complexes containing phosphinite and formation of metallaphosphineoxide complexes by the Arbuzov-like dealkylation reaction (H. Nakazawa, M. Yamaguchi, K. Kubo and K. Miyoshi), (428) 145

Asymmetric hydrogenation of prochiral carboxylic acids catalyzed by the five-coordinate ruthenium(II)-hydride complex $[RuH(binap)_2]PF_6$ (*binap* = (*R*)- or (*S*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) (M. Saburi, H. Takeuchi, M. Ogasawara, T. Tsukahara, Y. Ishii, T. Ikariya, T. Takahashi and Y. Uchida), (428) 155

Chemoselective asymmetric hydrogenation of α,β -unsaturated carbonyl compounds to allylic alcohols catalysed by $[Ir(binap)(cod)]BF_4$ -aminophosphine (K. Mashima, T. Akutagawa, X. Zhang, H. Takaya, T. Taketomi, H. Kumobayashi and S. Akutagawa), (428) 213

- Synthese, Struktur und Reaktivität der Ferriophosphonium-Salze $[CpFe(CO)(L)(PPh_2CH_2R')]BF_4$ ($L = CO, CH_3CN, P^nBu_3; R' = Ph, CN, COOEt, PPh_2, SiMe_3$); Kristallstrukturen von $[CpFe(CO)_2PPh_2CH_2SiMe_3]BF_4$ und $[CpFe(PPh_2CH_2Ph)CpFe(CO)](\mu-CO)_2$ (C. Klasen, I.-P. Lorenz, S. Schmid and G. Beuter), (428) 363
- Darstellung μ_3 -Chalkogenverbrückter dreikerniger Carbonyleisencluster durch die Reaktionsfolge XCN^- , Et_3O^+ . Aufbau und Dynamik von $(\mu_3-X)(\mu_3-PR)Fe_3(CO)_7L_2$ ($X = S, Se, Te; L = CO, EtNC, P(OMe)_3$) (W. Imhof, G. Huttner, B. Eber and D. Günauer), (428) 379
- Metallorganische Verbindungen der Lanthanoide. LXXI. Synthese und Röntgenstrukturanalyse von $CpLa(C_5H_4PPh_2)_2(THF)$, einem zweizähnigen Liganden zur Herstellung heterobimetallischer Lanthanoidkomplexe (H. Schumann, J.A. Meese-Markscheffel, B. Gorella and F.H. Görlich), (428) C27
- Chemie polyfunktioneller Moleküle. CIX. Synthese, Molekül- und Kristallstruktur des über $(NH \cdots O)$ - und $(OH \cdots NC)$ -Wasserstoffbrückenbindungen verknüpften tetraedrischen Chelatkomplexes $[Cu(CN)(Ph_2P-NH-PPh_2)PPh_3] \cdot CH_3OH$ (J. Ellermann, F.A. Knoch, K.J. Meier and M. Moll), (428) C44
- Organometallic complexes of the diphosphene $[2,4,6-(CF_3)_3C_6H_2]_2P_2$ (K.B. Dillon and H.P. Goodwin), (429) 169
- Photolytisch induzierte Erzeugung und komplexchemische Stabilisierung eines Molybdän–Arsen-Dreifachbindungssystems; Röntgenstrukturanalysen der Cluster μ_3 -AsCp₃Mo₂Ni(CO)₄, As₂Cp₆Mo₄Ni₂(CO)₇ und μ_3 -AsCpMoPt₂(CO)₄(PPh₃)₂ (M. Gorzellik, B. Nuber, T. Bohn and M.L. Ziegler), (429) 173
- Organoerdmetallkomplexe von d -Block-Elementen. IV. "Ein-Topf-Synthese" flüchtiger Organogalliocobalt-Komplexe, Molekülstruktur von $(\mu-[3-(Diethylamino)propyl]gallio)-[bis(tetracarbonylcobalt)]$ (R.A. Fischer, J. Behm and T. Priermeier), (429) 275
- Influence of ligands and anions on the insertion of alkenes into palladium–acyl and palladium–carbomethoxy bonds in the neutral complex $(dppp)Pd(C(O)CH_3)Cl$ and the ionic complexes $[(P-P)PdR(L)]^+SO_3CF_3^-$ ($P-P = dppe, dppp, dppb; R = C(O)CH_3, L = CH_3CN, PPh_3; R = C(O)OCH_3, L = PPh_3$) (G.P.C.M. Dekker, C.J. Elsevier, K. Vrieze, P.W.N.M. Van Leeuwen and C.F. Roobeek), (430) 357
- Penta- and hexaphosphapherrocenes as ligands: crystal and molecular structures of $[Fe(\eta^5-P_3C_2^1Bu_2)_2W(CO)_5]$, $[Fe(\eta^5-P_3C_2^1Bu_2)\chi(\eta^5-P_2C_3^1Bu_3)W(CO)_5]$ and the novel triruthenium carbonyl cluster complex $[Fe(\eta^5-P_3C_2^1Bu_2)_2Ru_3(CO)_{10}]$ containing two inter-linked $\eta^5-P_3C_2^1Bu_2$ ring systems (R. Bartsch, A. Gelessus, P.B. Hitchcock and J.F. Nixon), (430) C10
- Syntheses of low-valent nitrosyl complexes of rhenium and X-ray structure of $trans$ -[ReCl(NO)(Ph₂PCH₂CH₂PPh₂)₂][NO₃]₂ with nitrosyl derived nitrates (Y. Wang, J.J.R. Frausto Da Silva, A.J.L. Pombeiro, M.A. Pellinghelli and A. Tiripicchio), (430) C56
- Ein neues Verfahren zur Synthese von Metall-funktionalisierten Diphosphphenen, Arsaphosphphenen und Tetraphospha-bicyclobutanen (P. Jutzi and S. Opiela), (431) C29
- Mehrkernige Zirkonocen-Ethylkomplexe (H.G. Alt, C.E. Denner und R. Zenk), (433) 107
- Synthesis and characterisation of a bulky chelating bis(phosphine) ligand, 1,2-bis(diⁿbutylphosphino)ethane (DBPE), and its iron metal coordinated complexes, Fe(DBPE)₂Cl₂ and Fe(DBPE)₂($\text{C}\equiv\text{C}-\text{C}_6\text{H}_5$)₂ (J. Lewis, M.S. Khan, A.K. Kakkar, P.R. Raithby, K. Fuhrmann and R.H. Friend), (433) 135
- Synthesis and characterization of bis-(diphenylphosphinoethane)-substituted hexarhodium clusters: $Rh_6(CO)_{14}(\mu_2,\eta^2-Ph_2-P(CH_2)_2PPh_2)$, $(Rh_6(CO)_{15})_2(\mu_2,\eta^1:\eta^1-Ph_2P(CH_2)_2PPh_2)$ and $Rh_6(CO)_{15}(\eta^1-Ph_2P(CH_2)_2P(O)Ph_2)$. X-Ray crystal structures of $Rh_6(CO)_{15}PPh_3$ and $Rh_6(CO)_{14}(\mu_2,\eta^2-Ph_2P(CH_2)_2PPh_2)$ clusters (S.P. Tunik, A.V. Vlasov, N.I. Gorshkov, G.L. Starova, A.B. Nikol'skii, M.I. Rybinskaya, A.S. Batsanov and Y.T. Struchkov), (433) 189
- Sulphurisation reactions of $P_5C_5^1Bu_5$ and $P_5C_5^1Bu_5H_2$: crystal and molecular structure of the ethanol adduct of $P_5C_5^1Bu_5H_2S_3$ (R. Bartsch, P.B. Hitchcock and J.F. Nixon), (433) C11
- Die Kristallstruktur des Tris(dimethylthiophosphinyl)phosphans (M. Scheer und P.G. Jones), (434) 57
- Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XLI. Das dynamische Verhalten von 1,2-Bis(dimethylphosphino)ethan-dicarbonyl-(η^4 -dien)-wolfram(0) (S. Özkar, C. Kayran und C.G. Kreiter), (434) 79
- 3,5-Di-tert-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151
- Synthese und Charakterisierung von Hydridotrispyrazolylborat-Heterobimetall-Komplexen mit Titan und Molybdän (J. Ipaktschi und W. Sulzbach), (434) 287
- Neutrale und kationische Rhodiumkomplexe mit den potentiell zweizähnigen Phosphanliganden 1Pr_2PCH_2CH_2OMe , 1Pr_2PCH_2CH_2NMe_2 und 1Pr_2PCH_2CH_2CH_2NMe_2 (H. Werner, A. Hampp und B. Windmüller), (435) 169
- A kinetic study of the coordination template effect in the nickel-catalysed formation of arylphosphonium salts (D.W. Allen and P.E. Cropper), (435) 203
- Darstellung und Eigenschaften von und Reaktionen mit metallhalogenen Heterocylen. LXXVIII. Das Verhalten von monomeren und dimeren Thiophosphinitonickel-Komplexen gegenüber Alkinen und Phosphaalkinen (E. Lindner, A. Nothdurft, R. Fawzi und C. Maichle), (435) 213
- Approaches to new water-soluble phosphines (T.N. Mitchell and K. Heesche-Wagner), (436) 43
- Catalytic activity of *cis*- $[\mu-(bis(1,1-dimethylethyl)-phosphino)]dicarbonyl-\mu$ -chlorobis[tris(1,1-dimethylethyl)-phosphine]dirhodium and of its chiral analog (+)-*cis*- $[\mu-(bis(1,1-dimethylethyl)phosphino)]dicarbonyl-\mu$ -chlorobis[[β -methyl-2- α -(α -dimethylethyl)cyclohexyl]-diphenylphosphine]dirhodium (E. Berkovich, A. Jacobson, J. Blum, H. Schumann, M. Schäfers and H. Hemling), (436) 95
- Komplexchemie perhalogenierter Cyclopentadiene und Alkine. IX. Darstellung einiger Phosphoniumacetylid-Komplexe und Struktur von $(Ph_3P)Cl_2Pd(C\equiv CPPh_3)$ (K. Sünkel), (436) 101
- Reactivity of functionalized diphosphiranes and 1,3-diphosphapropenes towards Grignard reagents (R. El-Ouatib, C. Garot, G. Etemad-Moghadam and M. Koenig), (436) 169
- σ -Ferrocenyl-Komplexe des Rutheniums (M. Herberhold, W. Feger und U. Kölle), (436) 333
- Zur Funktionalisierung von Decaphenylcyclopentasilan (W. Uhlig und C. Tretner), (436) C1
- Zur Addition des Thiomethyl-Kations an die Wolfram–Kohlenstoff-Doppelbindung in $[Cp(CO)(PMo_3)W=C(Tol)PR_2][PF_6]$ (J. Ostermeier, W. Schütt und F.R. Kreißl), (436) C17
- Quasiphosphonium intermediates. VI. Crystal structure of neopentylloxy(phenacyl)diphenylphosphonium bromide: a stable Michaelis–Arbusov intermediate formed in the reaction of neopentyl diphenylphosphinite with phenacyl bromide (K. Henrick, H.R. Hudson, M. McPartlin, L. Powrozyk and L.S. Shaw), (437) 157
- Synthese, Charakterisierung und Reaktivität heterodinukleärer Komplexe: $[(CO)_4Fe(\mu-P^1Bu_2)Rh(CO)L](Fe-Rh)$ ($L = CO, HP^1Bu_2$), $[(CO)_3Co(\mu-H)(\mu-P^1Bu_2)Rh(CO)(HP^1Bu_2)](Co-Rh)$ (B. Walther, H.-C. Böttcher, M. Scheer, G. Fischer, D. Fenske und G. Süss-Fink), (437) 307
- Reaktionen von $[Cp^*Ru(OMe)]_2$. XIII. Pentaalkylcyclopentadienyl Ru und Rh Alkoxo-, Hydroxo- und Thiolato-Komplexe (U. Kölle, A. Hörnig und U. Englert), (438) 309

- Synthesis and characterization of dinuclear palladium(II) complexes containing both chloro and organochalcogenides as bridging ligands (V.K. Jain and S. Kannan), (439) 231
- η^1 - und η^2 -Komplexe von Cp*-substituierten Diphosphenen und Arsaphosphenen: Synthese, Struktur und Umlagerung unter Cp*-Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammmer), (439) 279
- Cycloadditionsreaktionen von Phosphoniumkomplexen. VIII. Regioselektive Cycloaddition chiraler Phosphonium-Komplexe $\text{Cp}(\text{OC})(\text{Ph}_2(\text{R})\text{P})\text{M}=\text{PPH}_2$ ($\text{M} = \text{Mo}, \text{W}; \text{R} = \text{H}, \text{Me}$) mit Isothiocyanaten (H. Pfister und W. Malisch), (439) C11
- Phosphonium-Übergangsmetallkomplexe. XX. Stabilisierung von Mesitylphosphiniden durch ein $\text{Cp}(\text{CO})_2\text{W}$ - und ein $\text{Cp}(\text{CO})_2\text{Fe}$ -Fragment (U.-A. Hirth und W. Malisch), (439) C16
- Phosphonium-Übergangsmetallkomplexe. XXI. Schwefel- und Selenaddition an die Metall-Phosphor-Doppelbindung der zweikernigen Phosphinidenkomplexe $\text{Cp}(\text{CO})_2\text{M}=\text{PMes}[\text{M}(\text{CO})_3\text{Cp}]$ ($\text{M} = \text{Mo}, \text{W}$) (U.-A. Hirth, W. Malisch und H. Käb), (439) C20
- A new reaction of $[\text{Cp}(\text{CO})_2\text{Fe}(\text{PPh}_2\text{H})]\text{PF}_6$ with NaBH_4 to give $\text{Cp}(\text{CO})(\text{H})\text{Fe}(\text{PPh}_2\text{H})$ via $\text{Cp}(\text{CO})_2\text{FeH}$ and PPh_2H (H. Nakazawa, K. Kubo, C. Kai and K. Miyoshi), (439) C42
- SO und NO als Konkurrenzliganden im neuen SO-Komplex $\text{RuCl}(\text{NO})(\text{SO})(\text{PPh}_3)_2$ (O. Heyke, G. Beuter und I.-P. Lorenz), (440) 197
- A remarkable accelerating effect of iodide ions in the photostimulated phosphorylation of bromoaromatic compounds (A. Boumekouez, E. About-Jaudet, N. Collignon and P. Savignac), (440) 297
- P_4 -Liganden mit maximaler Elektronendonorfähigkeit. III. Zum Reaktionsverhalten von Phosphankomplexen des Platins mit P_4 -Phosphor (M. Scheer, M. Dargatz und A. Rufińska), (440) 327
- Elektrophile Addition des Thiomethyl-Kations an η^5 -Cyclopentadienyl(ethoxymethylcarben)- (nitrosyl)(trimethylphosphin)molybdän (F.R. Kreißl und N. Ullrich), (440) 335
- Schwefel(IV)-Verbindungen als Liganden. XVII. Übergangsmetall-vermittelte Insertion von Schwefeldioxid in die C–O-Einfachbindung (W.A. Schenk und J. Pfeffermann), (440) 341
- Cycloadditionsreaktionen von Organometall-Komplexen. XV. [2 + 3]- und [2 + 1]-Cycloadditionsreaktionen von Carbonylrhodium- und Carbonylcobalt-Komplexen mit Nitriloxiden und Aziden. Die Kristall- und Molekülstruktur von $\text{C}_5\text{H}_5(\text{P}^{\text{t}}\text{Pr}_3)\text{RhC}(\text{O})\text{ONC}(\text{C}_6\text{H}_5)$ (H. Werner, U. Brekau, O. Nürnberg und B. Zeier), (440) 389
- Bis(diisopropylgalliumdiisopropylphosphid) und -arsenid. Kristallstruktur des Phosphidderivats — ein “angedeuteter” butterfly (G.G. Hoffmann, R. Fischer, U. Schubert und B. Hirle), (441) 7
- Zweikernige Rutheniumcluster mit $\mu_2\text{-}\eta^2$ -Sulfonatoliganden. Synthese und Molekülstruktur von $\text{Ru}_2(\text{CO})_4[\mu_2\text{-}\eta^2\text{-OOS(O)Tol}]_2(\text{PPh}_3)_2$ (G. Rheinwald, H. Stoeckli-Evans und G. Süss-Fink), (441) 295
- Synthesis and reactivity of dithiodiphenylphosphinato-derivatives of rhodium. Crystal structure of the square-pyramidal rhodium(III) complex $[\text{RhI}(\eta^2\text{-S}_2\text{Ph}_2)(\text{COMe})(\text{PPh}_3)]$ (J.A. Cabeza, V. Riéra, M.A. Villa-García, L. Ouahab and S. Triki), (441) 323
- Stereoselektive Additionsreaktionen von Komplex-gebundenen Phosphanen und Arsanen an Alkine (H. Lang und U. Lay), (441) 389
- Viergliedrige anorganische Ringsysteme des zweiwertigen Zinns und Bleis: Synthese und Struktur von chelatstabilisierten Stannylenen und Plumbylenen (U. Kilimann, M. Noltemeyer und F.T. Edelmann), (443) 35
- Übergangsmetall-substituierte Diphosphene. XXX. η^1 - und η^2 -Komplexe der Metalloidiphosphene ($\eta^5\text{-L}(\text{CO})_2\text{Fe}-\text{P}=\text{P}-\text{Mes}^*$ ($\text{L} = \text{C}_5\text{Me}_5$, $\text{C}_5\text{Me}_4\text{Et}$, $\text{C}_5\text{Me}_4^{\text{t}}\text{Bu}$, $1,3^{\text{t}}\text{Bu}_2\text{C}_5\text{H}_3$; $\text{Mes}^* = 2,4,6^{\text{t}}\text{Bu}_3\text{C}_6\text{H}_2$). Röntgenstrukturanalyse von $\text{Pt}\{\eta^2\text{-}[(\eta^5$ -C₅Me₄Et)(CO)₂Fe-P=Mes*](PPh₃)₂ (L. Weber, I. Schumann, H.-G. Stammmer und B. Neumann), (443) 175
- The structure of diphosphine radical cations (M.C.R. Symons, P. Tordo and J. Wyatt), (443) C29
- Darstellung und Röntgenstrukturuntersuchung von [1-Carbonyl-1-(η^5 -cyclopentadienyl)-2,3-dimethyl-1-trimethylphosphin-3-trimethylphosphoranylidien-1-wolframa-2-thiacyclopropan]-tetrafluoroborat (W. Schütt, E. Herdtweck, F. Hahn and F.R. Kreißl), (443) C33
- Additionsreaktionen von $(\text{CO})_5\text{CrEPhH}_2$ ($\text{E} = \text{P}, \text{As}$) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), (444) 53
- Wolfram(0)-Komplexe des π -aciden N -Methylpyrazinium-Kations und des N -Methylpyrazinium-Radikals. Kristall- und Molekülstruktur von $[(\text{C}_4\text{H}_4\text{N}_2\text{Me})\text{W}(\text{CO})_3(\text{PCy}_3)_2]\text{PF}_6$, Cy = Cyclohexyl (W. Bruns, H.-D. Hausen, W. Kaim und A. Schulz), (444) 121
- Reactions of coordinated phosphines and arsines: stereoselective reactions at the chlorophosphine- P stereocentre in the complex $[(\text{R}^*), (\text{R}^*, \text{R}^*)\text{-}(\pm)\text{-}[(\eta^5\text{-C}_5\text{H}_5)(1,2\text{-C}_6\text{H}_4(\text{PMePh})_2)\text{Fe}(\text{PClMePh})]\text{PF}_6 \cdot \text{CH}_2\text{Cl}_2$ (G. Salem, G.B. Shaw, A.C. Willis and S.B. Wild), (444) 147
- Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), (444) 155
- Ein- bis dreikernige Metall- π -Derivate von Hericen mit $(\text{Me}_3\text{P})_3\text{Fe}$ - und CpCo -Fragmenten (F.H. Köhler und A. Steck), (444) 165
- Chloro-, Methyl- und Hydrido-Iridiumkomplexe mit Ph_2CN_2 und anderen Diazoalkan-Liganden (L. Brandt, J. Wolf und H. Werner), (444) 235
- Übergangsmetall-Silyl-Komplexe. XLV. Erzeugung von Fluorsilyl-Liganden durch selektive Spaltung metallgebundener Si-Ph-Gruppen (S. Gilbert und U. Schubert), (444) C12
- Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), (444) C15
- First example of the cycloaddition of a terminal phosphinidene across the Rh=Rh bond of $[\text{Rh}_2(\eta^5\text{-C}_5\text{Me}_5)_2\{\mu\text{-}(\text{CO})_2\}]$. Crystal structure of $[\text{Rh}_2(\eta^5\text{-C}_5\text{Me}_5)_2(\text{CO})_2\{\text{MePW}(\text{CO})_5\}]$ (F.A. Ajulu, P.B. Hitchcock, F. Mathey and J.F. Nixon), (444) C60
- 1:1 Complexes of dimethyltin dihalides with tris(2,6-dimethoxyphenyl)-phosphine oxide, its sulphide, and related compounds (M. Wada, T. Fujii, S. Iijima, S. Hayase, T. Erabi and G.-e. Matsubayashi), (445) 65
- Atropisomerism in $(\pm)\text{-}7\text{-methyl- and -phenyl-substituted dinaphtho}[2,1-b;1',2'-d]\text{phospholes and dinaphth}[2,1-b;1',2'-d]\text{jarsoles}$ (A.A. Watson, A.C. Willis and S.B. Wild), (445) 71
- Metallorganische Lewissäuren. XLVII. Kationische Komplexe $[(\text{OC})_5\text{ReL}]^+$ und $[\text{Cp}(\text{OC})(\text{Ph}_3\text{P})\text{RuL}]^+$ mit O-Donoren und N-Donoren ($\text{L} = \text{Carbonsäureester, Ethanol, Lacton, Lactam, Diphenylketimin}$) (J. Milke, C. Missling, K. Sünkel und W. Beck), (445) 219
- Stereoselektive Bildung von Organoosmium-Komplexen mit Aminocarboxylat-Liganden (H. Werner, T. Daniel, O. Nürnberg, W. Knaup und U. Meyer), (445) 229
- Zur Reaktion von Vinyl- und Allylphosphonaten mit Dihalocarbenen — Dihalocyclopropyl- und Dihalocyclopropylmethylphosphonate (S. Bieler und K. Kellner), (447) 15
- Umsetzung von Alkinylphosphanen mit Hauptgruppenelement-verbrückten Eisencarbonylclustern (W. Imhof, B. Eber, G. Huttner und C. Emmerich), (447) 21
- Photochemisch induzierte Addition von Alkinen und Phosphaalkinen an die Cluster $\text{Fe}_3(\text{CO})_6(\mu_3\text{-PR})(\mu_3\text{-X})$ ($\text{X} = \text{Se, Te}; \text{R} = ^{\text{i}}\text{Pr}, ^{\text{t}}\text{Bu}$) (W. Imhof und G. Huttner), (447) 31

- Darstellung und Struktur von 1-Metalla-2,3,4-triazolen des Molybdäns und Wolframs (C.M. Stegmair, W. Schütt, W. Ullrich, P. Kiprof, J. Ostermeier und F.R. Kreißl), (447) 251
- P_4 -Liganden mit maximaler Elektronendonorfähigkeit. V. Zur Chlorigierung eines *cyclo-P₄*-Ligand-Komplexes (M. Scheer, M. Dargatz und P.G. Jones), (447) 259
- Interaction between $[(\eta^5-C_5Me_5)_2OsH]^+PF_6^-$ and nitrogen or oxygen bases. Hydrogen bonds of the type $[OsH]^+ \cdots B$ (L.M. Epstein, E.S. Shubina, A.N. Krylov, A.Z. Kreindlin and M.I. Rybinskaya), (447) 277
- Zur Reaktion von metallkoordinierten 1-Molybda-2-phospho-1,3-dienen mit Alkinen (H. Lang, M. Leise und L. Zsolnai), (447) C1
- Synthesis, and spectral and X-ray structural studies of a novel delocalised triphosphazene $(CF_3)_2P=N=P(CF_3)_2=N=PPPh_3$ and the triosmium cluster complex $Os_3(CO)_{11}[(CF_3)_2P=N=P(CF_3)_2=N=PPPh_3]$ (H.G. Ang, Y.M. Cai and W.L. Kwik), (448) 219
- Tertiary phosphine adducts of alane and gallane (F.M. Elms, M.G. Gardiner, G.A. Koutsantonis, C.L. Raston, J.L. Atwood and K.D. Robinson), (449) 45
- Synthesis and characterization of bridged dimeric and trimeric Group 13/15 complexes containing primary phosphido $[(^tBu)PH]^-$ and arsenido $[(^tBu)AsH]^-$ units (D.A. Atwood, A.H. Cowley, P.R. Harris, R.A. Jones, S.U. Koschmieder and C.M. Nunn), (449) 61
- Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of $[R(Me_3SiCH_2)InE(SiMe_3)_2]_2$ ($R = Ph, E = As; R = Me, E = As, P; R = Cl, E = P$) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), (449) 85
- The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), (449) 167
- Monomeric to polymeric gold(I) complexes from bifunctional ligands $R_2PC\equiv CH$ (G. Jia, R.J. Puddephatt and J.J. Vittal), (449) 211
- An indium-phosphorus heterocubane as a new type of indium phosphide precursor (D.A. Atwood, A.H. Cowley, R.A. Jones and M.A. Mardones), (449) C1
- Tetra-*ortho*-substituierte *cis*-Bis(phenyl)platin(II)-Verbindungen: Synthesen und photochemisches Verhalten (H.-A. Brune, J. Lerche, G. Schmidtberg und A. Baur), (450) 269
- ## Platinum
- Synthesis and evaluation of the catalytic properties of homo- and heterobiometallic complexes containing bridging diphenylphosphido ligands (Z. He, N. Lugan, D. Neibecker, R. Mathieu and J.-J. Bonnet), (426) 247
- Platinum complex-catalyzed dehydrogenative double silylation of olefins and dienes with *o*-bis(dimethylsilyl)benzene (M. Tanaka, Y. Uchimaru and H.-J. Lautenschlager), (428) 1
- Palladium- and platinum-catalyzed cyclocarbonylation reactions of substituted allyl acetates (Y. Ishii and M. Hidai), (428) 279
- Recent developments in the chemistry of olefin complexes of palladium and platinum: some novel findings as well as elaborate extensions (H. Kurosawa and I. Ikeda), (428) 289
- Organometallic complexes of the diphosphene $[2,4,6-(CF_3)_3C_6H_2]_2P_2$ (K.B. Dillon and H.P. Goodwin), (429) 169
- Photolytisch induzierte Erzeugung und komplexchemische Stabilisierung eines Molybdän-Arsen-Dreifachbindungssystems; Röntgenstrukturanalysen der Cluster $\mu_3\text{-AsCp}_3\text{Mo}_2\text{Ni}(\text{CO})_4$, $\text{As}_2\text{Cp}_6\text{Mo}_4\text{Ni}_2(\text{CO})_7$ und $\mu_3\text{-AsCpMoPt}_2(\text{CO})_4(PPh_3)_2$ (M. Gorzellik, B. Nuber, T. Bohn and M.L. Ziegler), (429) 173
- Synthesis and properties of 2- or 2,5-substituted thiophene and 2- or 2,5'-substituted dithiophene derivatives of platinum (S. Kotani, K. Shiina and K. Sonogashira), (429) 403
- Heteronuclear clusters containing the hexafluorobut-2-yne ligand. Crystal structures of $(C_5H_5)_2Rh_2Pt(\text{CO})(CF_3C_2CF_3)(\text{COD})$ (2) and $(C_5H_5)_4Rh_4Pt(\text{CO})_2(CF_3C_2CF_3)_2$ (3), and facile Rh–Pt bond making and breaking for complex 2 in solution (R.S. Dickson, G.D. Fallon, K.D. Heazle and M.J. Liddell), (430) 221
- Structural studies of complexes containing cycloplatinated tris(pyrazol-1-yl)methane (A.J. Canty, R.T. Honeyman, B.W. Skelton and A.H. White), (430) 245
- Trimethylsilylcyclooctadiene-transition metal complexes: metal-catalysed protodesilylation of cyclic vinylsilanes, and transfer hydrogenation promoted by the displaced silyl group (B.S. Bandodkar and G. Nagendrappa), (430) 373
- Reaction of (dichloro)(1,5-cyclooctadiene)platinum(II) with dimethyl diazomethylphosphonate. Crystal structure analysis of a bis-insertion product, $(R,R)\text{-bis}[\text{chloro}(\text{dimethoxyphosphonyl})\text{methyl}]\text{-}(1,5\text{-cyclooctadiene})\text{platinum(II)}$ (G. Ferguson, J.F. Gallagher, A.J. McAlees, R. McCrindle, J. Phillips and G.J.B. Williams), (430) C23
- Reactions with nucleophiles and cathodic electrochemical behaviour of some neutral and cationic diamino-, dioxy- and aminooxycarbene complexes of palladium(II) and platinum(II) (R. Bertani, M. Mozzon, R.A. Michelin, T.J. Castilho and A.J.L. Pombeiro), (431) 117
- Reactivity studies of square planar platinum, iridium, and rhodium triflate complexes with alkynols (P.J. Stang and Y.-H. Huang), (431) 247
- Conformational studies in palladium(IV) and platinum(IV) chemistry. Crystal structure of the 1,1-bis(pyrazol-1-yl)- ethane complex *fac*- $PtIMe_3(pz)_2CHMe-N,N'$ (P.K. Byers, A.J. Canty, R.T. Honeyman, B.W. Skelton and A.H. White), (433) 223
- Photochemical reactions of diphosphineplatinum(II) oxalate complexes (G.K. Anderson, G.J. Lumetta and J.W. Siria), (434) 253
- Electron transfer reactions involving *trans*- $[PtH_2(PCy_3)_2]$ and fluorinated benzonitriles (S. Hintermann, P.S. Pregosin, H. Rüegger and H.C. Clark), (435) 225
- Komplexchemie perhalogenierter Cyclopentadiene und Alkine. IX. Darstellung einiger Phosphoniumacetylid-Komplexe und Struktur von $(Ph_3P)Cl_2Pd(C=CPPPh_3)$ (K. Sünkel), (436) 101
- Synthesis of palladium(II) and platinum(II) *N,N*-dialkylthiocarbamates starting from hydroxo-halophenyl complexes (G. López, J. Ruiz, G. García, C. Vicente, J.M. Martí and V. Rodríguez), (436) 121
- Electrochemistry, electron spin resonance, optical spectroscopy, and reactivity of organometallic platinum(II) complexes containing strongly π -accepting aromatic α -diimine ligands (C. Vogler, B. Schwiderski, A. Klein and W. Kaim), (436) 367
- Photochemistry of mononuclear and binuclear tetramethylplatinum(IV) complexes: reactivity of an organometallic free radical (J.E. Hux and R.J. Puddephatt), (437) 251
- Steric and metal-centre control of two-, three- and four-orbital carborane–metal interaction in tin and platinum complexes of the $\{C_2B_8H_9X\}$ ligand, where $X = H$ and Ph (J.D. Kennedy, K. Nestor, B. Šíbr, M. Thornton-Pett and G.S.A. Zammit), (437) C1
- Organometallics of diphosphazanes. VI. Reactions of bicyclo-1,3,2- $\lambda^3,4\text{-}\lambda^3$ -diaza-diphosphetidines (bicyclic diphosphazanes) with Group 6 metal carbonyls and palladium and platinum(II) complexes (T.K. Prakasha, S.S. Krishnamurthy, S.E. Davies, M.F. Meidine and J.F. Nixon), (438) 241
- Organometallic nucleophiles. Mechanism of halide displacement at saturated carbon by 2-pyridyl and 4-pyridyl complexes $[M(\text{dmtc})(C_5H_4N-C^n)(L)]$ ($M = Pd, Pt$; dmtc = dimethylthiocarbamate; $n = 2,4$; L = tertiary phosphine) (L. Canovese, P. Uguagliati, F. Di Bianca and B. Crociani), (438) 253

- Five-coordinate hydrido/olefin platinum(II) complexes (V. De Felice, M.L. Ferrara, A. Panunzi and F. Ruffo), (439) C49
- Uptake of a Hg or Hg₂ unit into the center of cage-type platinum clusters supported by diphosphines having long methylene carbon chains: X-ray structure of [Hg₂Pt₆(dpphex)₃(RNC)₆]·2C₆H₆ (dpphex = 1,6-bis(diphenylphosphino)hexane, R = 2,6-dimethylphenyl) (T. Tanase, T. Horiuchi, Y. Yamamoto and K. Kobayashi), (440) 1
- Axiale Abschirmung des quadratisch-planar konfigurierten Platin(II) in strukturell charakterisiertem (bpy)PtMes₂: Lumineszenz und elektrochemisch reversible Bildung von Platin(III) (A. Klein, H.-D. Hausen und W. Kaim), (440) 207
- Synthesen und photochemische Untersuchungen von *ortho*- und *meta*-substituierten Bis(phenyl)[η^2 -1,2-bis(diphenylarsino)ethan bzw. -ethen]platin(II)-Verbindungen (H.-A. Brune, E. Hupfer, G. Schmidtberg und U. Ziegler), (440) 303
- Highly selective hydroformylation and dimerization reactions of 2-ferroenylpropene (L. Kollár and B. Floris), (441) 117
- Dimeric complexes of trimethylplatinum(IV) with ethylacetoacetate, methylacetoacetate, dimethylmalonate and diethylmalonate, and their pyridine and 2,2'-bipyridyl adducts (K. Kite and A.F. Psaila), (441) 159
- Synthesis and structural characterisation of some triangular and tetrahedral mixed metal platinum-gold clusters (C.M. Hill, D.M.P. Mingos, H. Powell and M.J. Watson), (441) 499
- Coordination of a tripodal phosphine sulphide P₂S towards various transition metal ions (H.-E. Wang, M.-C. Cheng, G.-H. Lee, S.-M. Peng and S.-T. Liu), (445) 171
- Highly stereoselective hydroformylation of a (2*R*)-2-tert-butyl-Δ⁴-1,3-oxazoline derivative (L. Kollár and P. Sándor), (445) 257
- Synthesis, molecular structure and NMR behaviour of [9,9-(PMe₂Ph)₂-*arachno*-9,6,8-PtS₂B₇H₇]: Direct evidence for the ten-vertex borane vertex-flip mechanism of isomerization (J.H. Jones, X.L.R. Fontaine, N.N. Greenwood, J.D. Kennedy, M. Thornton-Pett, B. Štíbr and H. Langhoff), (445) C15
- Cyclohydrosilylation dimer formation: evidence for Pt⁰ ↔ Pt^{II} ↔ Pt^{IV} catalysis (H.K. Chu and C.L. Frye), (446) 183
- Stereospecific reactions of the "M(C₆F₅)₂" (M = Pd or Pt) synths towards *cis*-dithiolato complexes (R. Usón, J. Forniés, M.A. Usón and S. Herrero), (447) 137
- Hydroformylation of norbornene and 2,5-norbornadiene catalysed by platinum(0)-alkene complexes in the presence of methanesulfonic acid: determination of the stereochemistry of the reaction (C. Botteghi, S. Paganelli, A. Perosa, R. Lazzaroni and G. Uccello-Barretta), (447) 153
- Binuclear pyrazolato-bridged organoplatinum(II) complexes: synthesis and characterization (A. Singhal, V.K. Jain and S. Kannan), (447) 317
- Synthesis of the heterodinuclear Fe-Pt complex [(OC)₃Fe{μ-CN(2,6-xyl)}₂{μ-dppm}Pt(PPh₃)] containing a bridging isonitrile ligand and electrophilic additions leading to μ-aminocarbyne complexes (M. Knorr, T. Faure and P. Braunstein), (447) C4
- Sterically-induced low temperature polyhedral rearrangements of carbaplatinaboranes: synthesis and crystal structures of 1-Ph-3,3-(PMe₂Ph)₂-3,1,2-PtC₂B₉H₁₀, 1-Ph-3,3-(PMe₂Ph)₂-3,1,11-PtC₂B₉H₁₀, 11-Ph-3,3-(PMe₂Ph)₂-3,1,11-PtC₂B₉H₁₀ and 1,11-Ph₂-3,3-(PMe₂Ph)₂-3,1,11-PtC₂B₉H₉ (D.R. Baghurst, R.C.B. Copley, H. Fleischer, D.M.P. Mingos, G.O. Kyd, L.J. Yellowlees, A.J. Welch, T.R. Spalding and D. O'Connell), (447) C14
- Triphenylphosphanchalkogenide als selektive Reagentien zum Aufbau von μ₃-Chalkogenverbrückten Eisencarbonylclustern (W. Imhof und G. Huttner), (448) 247
- Synthesis and characterization of sterically encumbered Li, Na, and K aryl tellurolates, and some Pt(II), Ir(I), and Cd(II) derivatives (Ph.J. Bonasia and J. Arnold), (449) 147

Synthesis and characterisation of new 2-bromovinyl selenides and their platinum group metal complexes (P.K. Khanna and C.P. Morley), (450) 109

Hydroformylation and hydrocarbethoxylation of 1,2-dicarbethoxy-1,2,3,6-tetrahydropyridazine and 1,2-dicarbethoxy-1,2,3,4-tetrahydropyridazine (G. Menchi, S. Paganelli, U. Matteoli, A. Scrivanti and C. Botteghi), (450) 229

Tetra-*ortho*-substituierte *cis*-Bis(phenyl)platin(II)-Verbindungen: Synthesen und photochemisches Verhalten (H.-A. Brune, J. Lerche, G. Schmidtberg und A. Baur), (450) 269

Cyclometallated derivatives of palladium(II) and platinum(II) derived from 6-t-butyl-2,2'-bipyridine (S. Stoccoro, G. Chelucci, M.A. Cinelli, A. Zucca and G. Minghetti), (450) C15

Potassium

The preference of 1-methylallyl polar organometallics and carbanions for *cis* rather than for *trans* geometries (P. von R. Schleyer, J. Kaneti, Y.-D. Wu and J. Chandrasekhar), (426) 143

Reactions of alkali metal macrocyclic complexes XIX. Reaction of potassium or sodium complexes with oxetane – novel route to organometallic compounds (Z. Jedliński, A. Misiołek, A. Jankowski and H. Janeczek), (433) 231

Cis/trans conversion of potassium derivatives of 2- and 4-nitrostilbenes (Z.V. Todres), (441) 349

Praseodymium

Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXIX. Optische und magnetochemische Untersuchungen an Mono- und Bis-Acetonitriladdukten des Grundkörpers Tris(η^5 -cyclopentadienyl)praseodym(III) (C. Apostolidis, B. Kanellakopoulos, R. Klenze, H. Reddmann, H. Schulz and H.-D. Amberger), (426) 307

Molekülstruktur und Ladungsverteilung in metallorganischen Verbindungen der 4-*f*- und 5-*f*-Elemente. II. Dielektrische Untersuchungen an metallorganischen Komplexen der 4-*f*- und Übergangselemente mit C_{3v}- und D_{3h}-Molekülsymmetrie (R. Maier, B. Kanellakopoulos and C. Apostolidis), (427) 33

Studies on organolanthanide complexes. XLII. Synthesis of the unsolvated monomeric complexes (MeOCH₂CH₂C₅H₄)₃Ln (Ln = La, Pr) and X-ray structure of (MeOCH₂CH₂C₅H₄)₃Pr (C. Qian, B. Wang, D. Deng, G. Wu and P. Zheng), (427) C29

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXXIII. Esteraddukte des Grundkörpers Tris(η^5 -cyclopentadienyl)-lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), (444) C15

Rhenium

Synthesis and spectroscopic characterization of the heteronuclear diphosphine linked cluster Os₃(CO)₁₁(Ph₂PCH₂PPh₂)Rh₆(CO)₁₅ (S.P. Tunik, M.V. Osipov and A.B. Nikolskyi), (426) 105

Metallorganische Lewis-Säuren. XLIV. Rhenooxy(organyl)carben-Komplexe von Chrom, Molybdän und Wolfram (P.M. Fritz, J. Breimair, B. Wagner and W. Beck), (426) 343

Photochemical reactions of [(Ph₃P)₂ReH₇] and [(Ph₂PCH₂CH₂PPh₂)ReH₇] (D. Baudry, J.-M. Cormier and M. Ephritikhine), (427) 349

The first examples of insertion of SnCl₂ into the Mn–Cl and Re–Cl bonds of octahedral complexes: X-ray structure of [Mn(CO)₃

- (SnCl₃)₂CPCy₃] · CH₂Cl₂ (B. Alvarez, D. Miguel, J.A. Pérez-Martínez, V. Riera and S. García-Granda), (427) C33
- Phosphine induced cyclopentadienyl ring slippage catalyzes CO insertion into a methyl rhenium compound to produce an acetyl rhenium compound (C.P. Casey, R.A. Widenhoefer and J.M. O'Connor), (428) 99
- A convenient lactonization of diols to γ - and δ -lactones catalyzed by transition metal polyhydrides (Y. Lin, X. Zhu and Y. Zhou), (429) 269
- Photochemical and thermal disproportionation of [(CO)₄CoM(CO)₃-(LL)] (M = Mn, Re; LL = 2,2'-bipyridine, 2-pyridinecarbaldehyde N-isopropylimine) complexes (H. Knoll, W.J.G. De Lange, H. Hennig, D.J. Stufkens and A. Oskam), (430) 123
- Syntheses of low-valent nitrosyl complexes of rhenium and X-ray structure of *trans*-[ReCl(NO)(Ph₂PCH₂CH₂PPh₂)₂]NO₃₂ with nitrosyl derived nitrates (Y. Wang, J.J.R. Frausto Da Silva, A.J.L. Pombeiro, M.A. Pellinghelli and A. Tiripicchio), (430) C56
- The reaction between unsaturated tetrarhenium cluster H₄Re₄(CO)₁₂ and tris(diphenylphosphino)methane (S.R. Wang, S.-L. Wang, C.P. Cheng and C.S. Yang), (431) 215
- The *cis*-{Re(Ph₂PCH₂CH₂PPh₂)₂}⁺ metal centre. Synthesis and molecular structure of the dinitrile complex *cis*-[Re(NCC₆H₄Me-4)₂(Ph₂PCH₂CH₂PPh₂)₂]BF₄ (M.F.C. Guedes da Silva, M.T. Duarte, A.M. Galvão, J.J.R. Frausto da Silva and A.J.L. Pombeiro), (433) C14
- 3,5-Di-tert-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151
- Hydrogen bonds and conformations of α -carbinol derivatives of cyclopentadienyltricarbonyls of manganese and rhenium (E.S. Shubina, A.N. Krylov, T.V. Timofeeva, Y.T. Struchkov, A.G. Ginzburg, N.M. Loim and L.M. Epstein), (434) 329
- Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109
- Novel concepts in directed biaryl synthesis. XV. Chiral rhenium complexes [(η^5 -C₅H₅)Re(NO)(PPh₃)*X*(R,R-lactone)]⁺BF₄⁻ of lactone-bridged biaryls as ligands: synthesis, structure, and stereochemical properties (G. Bringmann, O. Schupp, K. Peters, L. Walz and H.G. von Schnering), (438) 117
- Phosphinic complexes of rhenium and tungsten in reactions with stereoindicators (Z.V. Todres, E.A. Ionina and A.J.L. Pombeiro), (438) C23
- The synthesis and reactions of Re¹ carboxylic acids. The X-ray structure of *fac*-Re(CO)₃(dppp)CO₂D · 0.5C₆D₆ (S.K. Mandal, D.M. Ho and M. Orchin), (439) 53
- Synthesis and reactivity of bridging bis(alkyl) and bis(alkylidene) complexes derived from the chiral rhenium fragment [(η^5 -C₅H₅)Re(NO)(PPh₃)₂]⁺ (C. Roger, T.-S. Peng and J.A. Gladysz), (439) 163
- Wittig and Reformatsky chemistry of the formyl and acetyl derivatives of (η^5 -C₅H₅)Mn(CO)₃, (η^5 -C₅H₅)Re(CO)₃ and (η^6 -C₆H₆)Cr(CO)₃ (T.E. Bitterwolf and X. Dai), (440) 103
- The reactions of rhenium isocyanide complexes with acetonitrile. Formation of carbon–carbon and carbon–nitrogen bonds en route to rhenium containing heterocycles (L.L. Padolik, J.J. Alexander and D.M. Ho), (440) 153
- Structural characterization in solid state and solution of the hydrido-carbonyl cluster anion [Re₄(μ -H)₄(CO)₁₃]²⁻ (T. Beringhelli, G. D'Alfonso, A.P. Minoja, G. Ciani and A. Sironi), (440) 175
- Syntheses of gold–manganese and gold–rhenium clusters, [(Ph₃PAu)₃M(CO)₄] and [(Ph₃PAu)₄M(CO)₄]⁺ (M = Mn, Re), and the crystal structure of [(Ph₃PAu)₄Mn(CO)₄]BF₄ (B.K. Nicholson, M.I. Bruce, O. Bin Shawkataly and E.R.T. Tieckink), (440) 411
- Das Verhalten von stereoisomeren Bistriflaten des Tetrahydrofurans gegenüber ein- und zweiwertigen Carbonylmittallen (E. Lindner, M. Pabel, R. Fawzi und M. Steimann), (441) 63
- Metallorganische Lewis-Säuren. XLV. Pentacarbonylrhenium-Komplexe mit Oxoanionen (Trifluoromethoxyborat, Phosphat, Sulfat, Dichromat) sowie von Heteropolysäuren. Chemisorption von kationischen Carbonylrhenium-Komplexen auf γ -Al₂O₃, SiO₂, TiO₂ und in Zeolithen (E. Fritsch, J. Heidrich, K. Polborn und W. Beck), (441) 203
- Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXI. Perfluoropinakolat-Komplexe von Rhenium (VII) und Osmium(VI) (W.A. Herrmann und P. Watzlowik), (441) 265
- Umsetzungen der lithiierten Halbsandwich-Chalkogenole, (CO)₃MC₅H₄-ELi (M = Mn, Re; E = S, Se, Te) (M. Herberhold und M. Biersack), (443) 1
- Reaction of C₅Me₅(CO)₂Re(HC≡CCH₃) with Ph₃C⁺PF₆⁻ produces the η^3 -allyl complex C₅Me₅(CO)₂Re[η^3 -Ph₂CC(Ph)CHCH₃]⁺ PF₆⁻ (C.P. Casey, C.S. Yi and J.A. Gavney, Jr.), (443) 111
- Antiferromagnetic complexes with metal–metal bonds. XXVII. Synthesis, molecular structures and magnetic properties of Cr,Re-clusters Cp₂Cr₂(μ -SCMe₃)₂(μ ₃-S)Re₂(CO)₉ and Cp₂Cr₂(μ ₃-SCMe₃)₂(μ ₃-S)₂Re₂(CO)₆ (A.A. Pasynskii, I.L. Eremenko, S.E. Nefedov, B. Orazsakhatov, A.A. Zharkikh, O.G. Ellert, V.M. Novotortsev, A.I. Yanovsky and Y.T. Struchkov), (444) 101
- Metallorganische Lewissäuren. XLVII. Kationische Komplexe [(OC)₅ReL]⁺ und [Cp(OC)(Ph₃P)RuL]⁺ mit O-Donoren und N-Donoren (L = Carbonsäureester, Ethanol, Lacton, Lactam, Diphenylketimin) (J. Milke, C. Missling, K. Sünkel und W. Beck), (445) 219
- Electrospray mass spectrometric studies of some cationic and anionic Group 6 and Group 7 metal carbonyl complexes (I. Ahmed, A.M. Bond, R. Colton, M. Jurcevic, J.C. Traeger and J.N. Walter), (447) 59
- Cluster chemistry. LXXXVII. Some homo- and hetero-nuclear complexes derived from C₂(PPh₂)₂: Crystal structures of Re₃(μ -H)₃(μ -dppa)(CO)₁₀ (dppa = C₂(PPh₂)₂), Re₃(μ -H)₃(CO)₁₁(PPh₃)₂C₂Ru₂(μ -PPh₂)(CO)₆] and Os₃Ru₂(μ ₅-C₂PPh₂)(μ -PPh₂)(CO)₁₃ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (447) 91
- Carbon is not the initial site of attack in the protonation of an allene ligand to give an η^2 -vinyl species: kinetic studies on *trans*-[ReCl(CH₂CCPh)(Ph₂PCH₂CH₂PPh₂)₂] (R.A. Henderson, A.J.L. Pombeiro, R.L. Richards and Y. Wang), (447) C11
- Metallorganische Chemie des Technetiums. VIII. Technetium(I)-carbonyl-Komplexe mit Polypyrazol-1-yl-borato-Liganden im Vergleich mit seinen Mn- und Re-Homologen (J.E. Joachim, C. Apostolidis, B. Kanellakopulos, R. Maier, N. Marques, D. Meyer, J. Müller, A. Pires de Matos, B. Nuber, J. Rebizant und M.L. Ziegler), (448) 119
- Substituted metal carbonyls. XX. Unexpected formation of two diphenylphosphine complexes from the oxidation of pentacarbonylrhenate by chlorodiphenylphosphine. Crystal and molecular structures of *eq*-Re₂(CO)₉(PPh₂H) and *cis*-ReCl(CO)₄(PPh₂H) (T.S. Andy Hor, P.M.N. Low, Y.K. Yan, L.-K. Liu and Y.-S. Wen), (448) 131
- A facile synthetic route to (η^5 -benzoylcyclopentadienyl)metal compounds (S.S. Jones, M.D. Rausch and T.E. Bitterwolf), (450) 27
- Structures of propionaldehyde, butyraldehyde, and pivalaldehyde complexes of the chiral rhenium fragment [η^5 -C₅H₅)Re(NO)(PPh₃)₂]⁺ (D.P. Klein, N. Quirós Méndez, J.W. Seyler, A.M. Arif and J.A. Gladysz), (450) 157
- Electrochemical and chemical generation of a mixed-valent organorhenium oxide and its subsequent aggregation (R.W. Al-

- bach, U. Küsthadt, J. Behm, B. Ebert, M.-H. Delville and D. Astruc), (450) 165
- Carbyne-fluoro complexes of rhenium. Single-pot synthesis of *trans*-[ReF₃=CCH₂R)(Ph₂PCH₂CH₂PPh₂)₂]BF₄] (S.S.P.R. Almeida, J.J.R. Frausto Da Silva and A.J.L. Pombeiro), (450) C7
- Rhodium**
- Novel active dirhodium complexes for the low pressure hydroformylation reaction. Unexpected inhibiting effect of high partial pressures of hydrogen (Ph. Kalck, Y. Peres, R. Queau, J. Molinier, P. Escaffre, E. Leandro de Oliveira and B. Peyrille), (426) C16
- Crystal structure of [Rh(SnCl₃)₂(norbornadiene)(dppp)]. A ¹H NMR study of [Rh(SnCl₃)₂(norbornadiene)(diphosphine)] complexes (M.A. Garralda, E. Pinilla and M.A. Monge), (427) 193
- Rhodium(I) complex-catalyzed hydrosilylation of dimethyl muconates (K. Yamamoto and T. Tabei), (428) C1
- Rhodium hydride (HRh(CO)(PPh₃)₃) and rhodium carbonyl (Rh₄(CO)₈L₄) complexes obtained by reaction of Rh(acac)(CO)-L type complexes with methanol and formaldehyde (A.M. Trzezak and J.J. Ziolkowski), (429) 239
- Substituted cyclopentadienyl compounds. I. Formylation of methylcyclopentadienide and the synthesis and NMR study of some thallium(I) and rhodium(I) derivatives of cyano-, methyl-, methanoyl- and dimethylamidocyclopentadienes (M. Arthurs, J.C. Bickerton, M. Kirkley, J. Palin and C. Piper), (429) 245
- Substituted cyclopentadienyl compounds. II. Molecular orbital study of ring slippage in some rhodium(I) derivatives and the X-ray structure of η^5 -phenylcyclopentadienylbis(η^2 -ethene)rhodium(I) (M. Arthurs, C. Piper, D.A. Morton-Blake and M.G.B. Drew), (429) 257
- Heteronuclear clusters containing the hexafluorobut-2-yne ligand. Crystal structures of (C₅H₅)₂Rh₂Pt(CO)(CF₃C₂CF₃)(COD) (2) and (C₅H₅)₂Rh₄Pt(CO)₂(CF₃C₂CF₃)₂ (3), and facile Rh-Pt bond making and breaking for complex 2 in solution (R.S. Dickson, G.D. Fallon, K.D. Heazle and M.J. Liddell), (430) 221
- Trimethylsilylcyclooctadiene-transition metal complexes: metal-catalysed protodesilylation of cyclic vinylsilanes, and transfer hydrogenation promoted by the displaced silyl group (B.S. Bandodkar and G. Nagendrappa), (430) 373
- Sterically induced opening of a *clos*o carbometallaborane: synthesis and characterisation of 1,2-Ph₂-3-(η -C₅Me₅)-3,1,2-pseudocloso-RhC₂B₉H₉ (Z.G. Lewis and A.J. Welch), (430) C45
- Binuclear hydroxo-, methoxo-, and pyrazolate-bridged complexes of ruthenium(II). Molecular structure of [(*p*-cymene)Ru]₂(μ -pz)(μ -OH)₂]BF₄ (*p*-cymene = *p*-MeC₆H₄CHMe₂, pz = pyrazolate) (D. Carmona, A. Mendoza, J. Ferrer, F.J. Lahoz and L.A. Oro), (431) 87
- Rhodium pentafluorophenylthiolate complexes derived from [Rh₂(μ -SC₆F₅)₂(PPh₃)₄] (L. Carlton), (431) 103
- Synthesis and chemistry of 1-acetyl, 2-methylsubstituted cyclopentadienyl thallium and its chiral manganese and rhodium carbonyl derivatives. The molecular structure of [1-CH₃CO, 2-CH₃C₅H₃]Rh(CO)[P(C₆H₅)₃] (T.E. Bitterwolf, T.L. Hubler and A.L. Rheingold), (431) 199
- Methylation of bis(oxazolinyl)pyridinerhodium(III) chlorides with trimethylaluminum (H. Nishiyama, T. Hirai and K. Itoh), (431) 227
- Optically active complexes of transition metals (Rh^I, Ru^{II}, Co^{II} and Ni^{II}) with 2-aminocarbonylpiperidine ligands. Selective catalysts for hydrogenation of prochiral olefins (A. Corma, M. Iglesias, C. Del Pino and F. Sánchez), (431) 233
- Reactivity studies of square planar platinum, iridium, and rhodium trflate complexes with alkynols (P.J. Stang and Y.-H. Huang), (431) 247
- Metallocigands containing samarium(III). Access to binuclear *f*-metal (samarium)-*d*-metal (rhodium) complexes (M. Visseaux, A. Dormond, M.M. Kubicki, C. Moise, D. Baudry and M. Ephritikhine), (433) 95
- μ -Methylene rhodium complexes with SH ligand: synthesis and structures of [Rh₂Cp^{*}₂(μ -CH₂)₂(μ -SH)]BPh₄ and *trans*-[Rh₂Cp^{*}₂(μ -CH₂)₂(SH)₂] (Cp^{*} = η^5 -C₅Me₅) (Y. Ozawa, A.V. de Miguel and K. Isobe), (433) 183
- Synthesis and characterization of bis-(diphenylphosphinoethane)-substituted hexarhodium clusters: Rh₆(CO)₁₄(μ , η^2 -Ph₂P(CH₂)₂PPh₂), (Rh₆(CO)₁₅)₂(μ , η^1 : η^1 -Ph₂P(CH₂)₂PPh₂) and Rh₆(CO)₁₅(η^1 -Ph₂P(CH₂)₂P(O)Ph₂). X-Ray crystal structures of Rh₆(CO)₁₅PPh₃ and Rh₆(CO)₁₄(μ , η^2 -Ph₂P(CH₂)₂PPh₂) clusters (S.P. Tunik, A.V. Vlasov, N.I. Gorshkov, G.L. Starova, A.B. Nikol'skii, M.I. Rybinskaya, A.S. Batsanov and Y.T. Struchkov), (433) 189
- Homoleptische metallorganische Innerkomplexe des 8-Dimethylaminonaphthalyl-1-Anions (H. Dreys), (433) C1
- Binuclear rhodium complexes immobilized on functionalized diphenylphosphino-(styrene-divinylbenzene) resins giving high selectivities for linear aldehydes in hydroformylation reactions (P. Kalck, E. Leandro de Oliveira, R. Queau, B. Peyrille and J. Molinier), (433) C4
- Coordination chemistry of [CH(P(S)Ph₂)₂]⁻: X-ray diffraction studies of S,S-chelate complexes of iridium and rhodium (J. Brownning, G.W. Bushnell, K.R. Dixon and R.W. Hiltz), (434) 241
- Neutrale und kationische Rhodiumkomplexe mit den potentiell zweizähnigen Phosphanliganden ¹Pr₂PCH₂CH₂OMe, ¹Pr₂PCH₂CH₂NMe₂ und ¹Pr₂PCH₂CH₂CH₂NMe₂ (H. Werner, A. Hampp und B. Windmüller), (435) 169
- Ligand redistribution reactions of some organometallic rhodium and iridium complexes (P.J. Stang and Y.-H. Huang), (435) 185
- Catalytic activity of *cis*-[μ -[bis(1,1-dimethylethyl)-phosphino]]dicarbonyl- μ -chlorobis[tris(1,1-dimethylethyl)-phosphine]dirhodium and of its chiral analog (+)-*cis*-[μ -[bis(1,1-dimethylethyl)phosphino]]dicarbonyl- μ -chlorobis[[5 β -methyl-2 α -(1 α -methylethyl)cyclohexyl]-diphenylphosphine]dirhodium (E. Berkovich, A. Jacobson, J. Blum, H. Schumann, M. Schäfers and H. Hemling), (436) 95
- Komplexchemie perhalogenierter Cyclopentadiene und Alkine. IX. Darstellung einiger Phosphoniumacylid-Komplexe und Struktur von (Ph₃P)Cl₂Pd(C=CPh₃) (K. Sünkel), (436) 101
- Synthese, Charakterisierung und Reaktivität heterodinukleärer Komplexe: [(CO)₄Fe(μ -P^tBu₂)₂Rh(CO)L](Fe-Rh) (L = CO, HP^tBu₂), [(CO)₃Co(μ -H)(μ -P^tBu₂)₂Rh(CO)(HP^tBu₂)](Co-Rh) (B. Walther, H.-C. Böttcher, M. Scheer, G. Fischer, D. Fenske und G. Süss-Fink), (437) 307
- Chiral sulfonated phosphines. VIII. Hydrogenation of dehydropeptides in a two-phase system (M. Laghmari, D. Sinou, A. Masdeu and C. Claver), (438) 213
- Synthesis and characterization of rhodium complexes [Rh(Az)(L)Cl] (Az = 2,2-dimethyl-3-phenyl-3-allylaziridine; L = C₂H₄, CO, or PPh₃). Crystal and molecular structure of [Rh(Az)(CO)Cl] (R.B. Cheikh, M.C. Bonnet, R. Chaabouni and F. Dahan), (438) 217
- Pentafluorophenyl compounds of rhodium(III) with bidentate phosphorus or nitrogen ligands (M.P. García, M.V. Jiménez and L.A. Oro), (438) 229
- Fulvenylsubstituierte Cp-Liganden. III. Komplexe des Anions [C₃H₄CH=C₉H₆]⁻ mit Molybdän, Mangan und Rhodium (P. Härtter, J. Behm und K.J. Burkert), (438) 297
- Reaktionen von [Cp^{*}Ru(OMe)]₂. XIII. Pentaalkylcyclopentadienyl Ru und Rh Alkoxo-, Hydroxo- und Thiolato-Komplexe (U. Kölle, A. Hörrig und U. Englert), (438) 309
- Heterobimetallic Group 6-rhodium complexes. II. Reactions of [Rh(S₂CNEt₂)(CO)₂] with the complexes [M(CO)₃(η^2 -phen)(η^1 -

- dppm)] ($M = Mo$ or W) and $[Mo(CO)_4(\eta^2\text{-dppm})]$ as P-donor systems (M. Cano, P. Ovejero and J.V. Heras), (438) 329
- Indenyl carbametallaboranes. III. The synthesis and characterization of novel rhodium carbaborane compounds with $3-(C_2B_9)\text{-}8\text{-}\{(C_9H_6)\text{Rh}(C_9H_7)\}\text{-}3,1,2\text{-clos}\text{o}\text{-}RhC_2B_9$, $10\text{-}\{(C_9H_6)\text{Rh}(C_9H_7)\}\text{-}7,8\text{-nido}\text{-}C_2B_9$, $9\text{-}\{(C_9H_6)\text{Rh}(C_9H_7)\}\text{-}7,8\text{-nido}\text{-}C_2B_9$ and $3\text{-}(\eta\text{-}C_9H_7)\text{-}3,1,2\text{-clos}\text{o}\text{-}RhC_2B_9$ structures (Z.G. Lewis and A.J. Welch), (438) 353
- ^{13}C NMR studies of silyl and germinal complexes of rhodium and iridium containing carbonyl ligands (E.A.V. Ebsworth and S. Moreton), (439) 213
- Cycloadditionsreaktionen von Organometall-Komplexen. XV. $[2 + 3]$ -und $[2 + 1]$ -Cycloadditionsreaktionen von Carbonylrhodium- und Carbonylcobalt-Komplexen mit Nitriloxiden und Aziden. Die Kristall- und Molekülstruktur von $C_5H_5(P^iPr_3)RhC(O)ONC(C_6H_5)$ (H. Werner, U. Brekau, O. Nürnberg und B. Zeier), (440) 389
- Highly selective hydroformylation and dimerization reactions of 2-ferrocenylpropene (L. Kollár and B. Floris), (441) 117
- Synthesis and molecular structure of new families of iridium(III)- Cp^* and rhodium(III)- Cp^* complexes derived from 1,2-di-cyanoethene-1,2-dithiolate, 2,2'-biimidazole or 2,2'-bithiazole. Single crystal structures of $[(\eta^5\text{-Me}_5C_5)\text{Ir}(\text{biimH}_2)\text{Cl}]Cl$ and $[(\eta^5\text{-Me}_5C_5)\text{Rh}(\text{dcdt})]$ (R. Ziessel, M.-T. Youinou, F. Balegroune and D. Grandjean), (441) 143
- Synthesis and reactivity of dithiodiphenylphosphinato-derivatives of rhodium. Crystal structure of the square-pyramidal rhodium(III) complex $[\text{Rh}(\eta^2\text{-S}_2\text{PPh}_2)(\text{COMe})(\text{PPh}_3)]$ (J.A. Cabeza, V. Riera, M.A. Villa-García, L. Ouahab and S. Triki), (441) 323
- Synthesis, electrochemistry and complexation studies of new redox active bisferrocene acyclic and macrocyclic thioethers (P.D. Beer, J.E. Nation, M.E. Harman and M.B. Hursthouse), (441) 465
- Rhodium(I) complexes of 4-t-butylcyclohexyliaziridine and adamantlylaziridine: synthesis, structure and catalytic activity (A. Adedapo, S.A. Benyunes, P.A. Chaloner, C. Claver, P.B. Hitchcock, A. Ruiz and N. Ruiz), (443) 241
- Selective *ortho*-metallation reaction of the phosphine $P(o\text{-ClC}_6H_4\text{Ph}_2)$. Synthesis and crystal structure of $[\text{Rh}_2(\mu\text{-O}_2\text{CCH}_3)_3(\mu\text{-}(\text{ClC}_6\text{H}_3)\text{PPh}_2)(\text{OH}_2)_2]\cdot\text{CHCl}_3$ (P. Lahuerta, J. Payá, S. García-Granda, F. Gómez-Beltrán and A. Anillo), (443) C14
- Synthèse de phosphinites dérivés de sucres: applications en cyclodimérisation du penta 2,4-diénoate de méthyle, en hydrogénéation et en hydrosilylation de la pentolactone (A. Bendayan, H. Masotti, G. Peiffer, C. Siv et A. Archavlis), (444) 41
- Synthesis and reactivity of substituted cyclopentadienyl diphenylphosphido complexes of zirconium(IV). X-Ray crystal structure of $[(C_5H_4\text{SiMe}_3)_2\text{Zr}(\text{PPh}_2)_2]$ (A.-M. Larsonneur, R. Choukroun, J.-C. Daran, T. Cuena, J.C. Flores and P. Royo), (444) 83
- Umsetzungen von Metallorganylen mit dem capto-dativ substituierten Ethylen $H_2C=C(\text{CN}^+\text{Butyl})$ (J. Herbig, J. Köhler, B. Nuber, H.G. Stammle and M.L. Ziegler), (444) 107
- Photochemistry of tris(dimethylpyrazolyl)-boratorrhodium dicarbonyl and bis(dimethylpyrazolyl)-boratorrhodium dicarbonyl complexes in low temperature media at 12–298 K: some insights into C–H activation processes (P.E. Bloyce, J. Mascetti and A.J. Rest), (444) 223
- First example of the cycloaddition of a terminal phosphinidene across the Rh=Rh bond of $[\text{Rh}_2(\eta^5\text{-C}_5\text{Me}_5)_2(\mu\text{-}(\text{CO})_2)]$. Crystal structure of $[\text{Rh}_2(\eta^5\text{-C}_5\text{Me}_5)_2(\text{CO})_2(\text{MePW}(\text{CO})_5)]$ (F.A. Ajulu, P.B. Hitchcock, F. Mathey and J.F. Nixon), (444) C60
- Coordination of a tripodal phosphine sulphide P_2S towards various transition metal ions (H.-E. Wang, M.-C. Cheng, G.-H. Lee, S.-M. Peng and S.-T. Liu), (445) 171
- Synthesis, structure and hydrogenation of $\eta^3\text{-benzyl diphosphine}$ complexes of rhodium and iridium (M.D. Fryzuk, D.H. McConville and S.J. Rettig), (445) 245
- Highly stereoselective hydroformylation of a $(2R)\text{-}2\text{-tert-butyl-}\Delta^4\text{-}1,3\text{-oxazoline derivative}$ (L. Kollár and P. Sándor), (445) 257
- Simultaneous substitution of bridging acetate groups and reversible Rh–C bond cleavage in $[\text{Rh}_2(\text{O}_2\text{CCH}_3)_3(\text{C}_6\text{H}_4)\text{PPh}_2](\text{HO}_2\text{CCH}_3)_2$ in the presence of $\text{CF}_3\text{CO}_2\text{H}$. Crystal structure of $[\text{Rh}_2(\text{O}_2\text{CCF}_3)_3(\text{C}_6\text{H}_4)\text{PPh}_2](\text{HO}_2\text{CCF}_3)_2$ (P. Lahuerta, J. Latorre, E. Peris, M. Sanaú, M.A. Ubeda and S. García-Granda), (445) C10
- Dehydrogenative silation, isomerization and the control of *syn*- vs. *anti*-addition in the hydrosilation of alkynes (C.-H. Jun and R.H. Crabtree), (447) 177
- Electronic and steric effects in the rhodium-complex catalysed cyclization of alkynes and nitriles to pyridine derivatives (P. Diversi, L. Ermini, G. Ingrosso and A. Lucherini), (447) 291
- Rhodium(I) complexes with aromatic diamines (L. Fidalgo, M.A. Garralda, R. Hernández and L. Ibarlucea), (447) 299
- Studies of the thermal and photochemical lability of olefin complexes of rhodium(I) to form Rh and Rh_2O_3 (Z. Duan, M.J. Hampden-Smith and A.P. Sylwester), (449) 173
- A facile synthetic route to $(\eta^5\text{-benzoylcyclopentadienyl})\text{metal}$ compounds (S.S. Jones, M.D. Rausch and T.E. Bitterwolf), (450) 27
- Ferrocene derivatives. XXVIII. The metallation of phenylazoferrrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177
- Hydroformylation and hydrocarbethoxylation of 1,2-dicarbethoxy-1,2,3,6-tetrahydropyridazine and 1,2-dicarbethoxy-1,2,3,4-tetrahydropyridazine (G. Menchi, S. Paganelli, U. Matteoli, A. Scrivanti and C. Botteghi), (450) 229
- The crystal and molecular structure of bis(indazol-1-yl)pyridin-2'-ylmethane (BIPM) and $[\text{Rh}(\text{BIPM})(\text{NBD})]\text{PF}_6$ (M.C. López Gallego-Preciado, P. Ballesteros, R.M. Claramunt, M. Cano, J.V. Heras, E. Pinilla and A. Monge), (450) 237
- ^{31}P and ^{13}C NMR investigation of the system tetracarbonyldi- μ -chlorodirhodium(I)-tertiary phosphine (E. Rotondo, G. Battaglia, G. Giordano and F.P. Cusmano), (450) 245
- ## Rubidium
- The preference of 1-methylallyl polar organometallics and carbanions for *cis* rather than for *trans* geometries (P. von R. Schleyer, J. Kaneti, Y.-D. Wu and J. Chandrasekhar), (426) 143
- ## Ruthenium
- Chemistry of $[\text{Cp}^*\text{RuOMe}]_2$. XI. Reactions with $\text{P}(\text{OMe})_3$ (U. Koelle, Th. Ruether and W. Kläui), (426) 99
- Synthesis and evaluation of the catalytic properties of homo- and hetero-bimetallic complexes containing bridging diphenylphosphido ligands (Z. He, N. Lukan, D. Neibecker, R. Mathieu and J.-J. Bonnet), (426) 247
- Reactions of alkenyl and alkynyl ruthenium(II) complexes with isocyanides: synthesis of α,β -unsaturated $\eta^1\text{-acylruthenium(II)}$ complexes and X-ray structure of $[\text{Ru}(\text{C}\equiv\text{CPh})(\text{CN}^+\text{Bu})_3(\text{PPh}_3)_2]\text{PF}_6$ (J. Montoya, A. Santos, J. López, A.M. Echavarren, J. Ros and A. Romero), (426) 383
- Synthesis, X-ray structure, and catalytic properties of a ruthenium cluster with a carbohydrate ligand (S. Bhaduri, N. Sapre, H. Khwaja and P.G. Jones), (426) C12
- Carbon–nitrogen bond cleavage of β -nitrostyrene promoted by acetoxyhydridercarbonylbis(triphenylphosphine)ruthenium(II) (A. Yousfi and R. Mathieu), (426) C33

A diruthenium complex containing an *ortho*-metallated bipyridyl ligand formed directly from $[Ru_3(CO)_{12}]$ and pyridine (B.R. Cockerton and A.J. Deeming), (426) C36

Organometallic macrocycle chemistry. I. The reactions of $[RuCl_2(PPPh_3)_3]$ with 1,4,7-trithiacyclononane and 1,4,8,11-tetra-thiacyclopentadecane (A.F. Hill, N.W. Alcock, J.C. Cannadine and G.R. Clark), (426) C40

Reactions of carbonyl clusters with potentially tridentate thioethers: crystal and molecular structures of $[Os_4(CO)_{13}([12]aneS_3)]$ and $[Ru_6(CO)_{15}(\mu_4-\eta^2-CO)([12]aneS_3)]$ ($[12]aneS_3 = \{[(CH_2)_3S]_3$) (A.J. Edwards, B.F.G. Johnson, F.K. Khan, J. Lewis and P.R. Raithby), (426) C44

Metal-ion dependent reactivity of 2-(2'-thienyl)pyridine (Hthpy) (E.C. Constable and L.R. Sousa), (427) 125

Synthesis and reactivity of triruthenium carbonyl clusters containing bis(diphenylphosphino)methane and the face-bridging ligand 2-amido-6-methylpyridine (P.L. Andreu, J.A. Cabeza, J.L. Cuyás and V. Riera), (427) 363

Synthesis, crystal structure and spectroelectrochemistry of $[(C_5Me_5Ru)_2\cdot\eta^6,\eta^6\text{-chrysene}]^{2+}(O_3SCF_3^-)_2$. Formation of $[(C_5Me_5Ru^I)(C_5Me_5Ru^{II})\cdot\eta^6,\eta^6\text{-chrysene}]O_3SCF_3$ (I. Chavez, M. Otero, E. Román and U. Müller), (427) 369

Übergangsmetall-Heteroallen-Komplexe. XXV. Reaktionen von Ketenimin mit $Ru_3(CO)_{12}$ (W. Ziegler and U. Behrens), (427) 379

Novel organometallic charge transfer salts derived from electron-rich diruthenium species and the electron-acceptor ligands 7,7,8,8-te-tracyano-*p*-quinodimethane and tetracyanoethylene and containing the radical anions of these ligands in both the inner and the outer coordination spheres (S.E. Bell, J.S. Field, R.J. Haines and J. Sundermeyer), (427) C1

Effect of ion-pairs on the stereochemistry of decarbonylation of acetyl iodide complexes of ruthenium: evidence of anion catalysis in the CO insertion (G. Bellachoma, G. Cardaci, A. Macchioni and G. Reichenbach), (427) C37

Asymmetric hydrogenation of prochiral carboxylic acids catalyzed by the five-coordinate ruthenium(II)-hydride complex $[RuH(binap)_2]PF_6$ (binap = (*R*)- or (*S*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) (M. Saburi, H. Takeuchi, M. Ogasawara, T. Tsukahara, Y. Ishii, T. Ikariya, T. Takahashi and Y. Uchida), (428) 155

Chemistry of hexaruthenium carbidocarbonyl cluster complexes. Syntheses and X-ray crystal structures of $Ru_6C(CO)_{15}(\mu-SePh)(\mu-AuPPh_3)$, $Ru_6C(CO)_{14}(\mu-SePh)_2$, and $Ru_6C(CO)_{14}(\mu-SePh)(\mu,\eta^3-C_3H_5)$ (T. Chihara and H. Yamazaki), (428) 169

$[Rh(CO)_4]^-$, $[Rh_5(CO)_{15}]^-$, and bimetallic clusters as catalysts for the polymerization of nitrobenzene to methyl phenylcarbamate (F. Ragagni, S. Cenini, A. Fumagalli and C. Crotti), (428) 401

Thermolysis of $Cp(CO)_nM-CH(CH_3)C_6H_5$ ($n = 2$, M = Fe, Ru; $n = 3$, M = Mo, W). Some vertical trends in transition metal-alkyl bond dissociation energies (C. Mancuso and J. Halpern), (428) C8

Single and unsupported Ru-H-Ru bond: preparation and crystal structure of $\{Ru_2H(\mu-H)Cl(syn-Me_4[14]aneS_4)_2\}Cl$ ($Me_4[14]aneS_4 = 1,4,8,11\text{-tetra-thiacyclopentadecane}$) (T. Yoshida, T. Adachi, T. Tanaka and F. Goto), (428) C12

Synthesis and protonation of new 1,3-diene complexes of zerovalent ruthenium (M.A. Bennett and X.-q. Wang), (428) C17

Oligomerization of butadiene with ($\eta^5-C_5R_5$)Ru^{II} complexes; stoichiometric and catalytic chemistry (K. Masuda, K. Nakano, T. Fukahori, H. Nagashima and K. Itoh), (428) C21

Cluster chemistry. LXXXI. Reactions of $Ru_3(\mu-dppm)(CO)_{10}$ with $C_2(CO_2Me)_2$. X-Ray structures of $Ru_3(\mu_3-C_4(CO_2Me)_4)(\mu-dppm)(CO)_6$, $Ru_3(\mu-H)_2(\mu_3-C_2(CO_2Me)_2)(\mu-dppm)(CO)_7$ and $Ru_4(\mu_3-H)(\mu_4-C_2(CO_2Me)_2)(\mu-PPh_2CH_2PPh(C_6H_4))(CO)_9$ (M.I. Bruce, P.A. Humphrey, H. Miyamae, B.W. Skelton and A.H. White), (429) 187

Cluster chemistry. LXXXII. Reactions between $Ru_3(\mu-dppm)(CO)_{10}$ and 1-alkynes and related chemistry. Crystal structures of $Ru_3(\mu-H)(\mu_3-C_2C_6F_5)(\mu-dppm)(CO)_7$, $Ru_3(\mu_3-PPhCH_2PPh(C_6H_4))(\mu-C_4H_2Ph_2)(CO)_6$, $AuRu_3(\mu_3-C_2Ph)(\mu-dppm)(CO)_7\cdot\{P(C_6H_4Me-p)_3\}$ and $Ru_2(\mu-I)(\mu-dppm)(\mu-C_2Ph)(CO)_4$ (M.I. Bruce, P.A. Humphrey, E. Horn, E.R.T. Tiekkink, B.W. Skelton and A.H. White), (429) 207

Polyazolyl chelate chemistry. IV. Dihydrobis(benzotriazolyl)borato complexes of ruthenium(II) (J. Cartwright and A.F. Hill), (429) 229

A convenient lactonization of diols to γ - and δ -lactones catalyzed by transition metal polyhydrides (Y. Lin, X. Zhu and Y. Zhou), (429) 269

Facile conversion of ($\eta^5-C_5R_5$)M(CO)₂-halide complexes to halomethyl, alkoxymethyl, and cyanomethyl derivatives (R = H, CH₃; M = Fe, Ru; halide = Cl, Br, I) (J.L. Hubbard and W.K. McVicar), (429) 369

Synthesis and characterisation of novel selenium-containing clusters; crystal structures of $[Ru_4(\mu_4-Se)_2(CO)_8(\mu-CO)_3]$ and $[Ru_3(\mu_3-Se)_2(CO)_7(Ph_2P(CH_3)_3PPh_2)]$ (B.F.G. Johnson, T.M. Layer, J. Lewis, A. Martín and P.R. Raithby), (429) C41

Carbonylation of methanol to acetic acid using homogeneous Ru complex catalyst (A.A. Kelkar, D.S. Kolhe and R.V. Chaudhari), (430) 111

Cluster chemistry. LXXX. Reactions of $Ru_5(\mu_5-C_2PPh_2)(\mu-PPh_2)(CO)_{13}$ with 1,3-butadiene: addition to C_α and formation of a μ_4 -cyclohexenyne complex. X-Ray structures of $Ru_5(\mu_4-CC(PPh_2)_2CH_2\cdot\eta^3-CHCH_2)(\mu-PPh_2)(\mu-CO)(CO)_{11}$, $Ru_5(\mu_4-CC(PPh_2)_2CHCH=CHMe)(\mu-PPh_2)(\mu-CO)_2(CO)_9$ and $Ru_5(\mu_4-C_6H_6)(\mu-PPh_2)_2(CO)_{11}$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (430) 181

Stereochemical aspects of the synthesis and reactivity of (diphosphine)(carbonyl)(cyclopentadienyl)ruthenium complexes (A. Mezzetti, G. Consiglio and F. Morandini), (430) C15

Half-way coordination state of a butadienyl group on ruthenium. η^3 -Allylic bonding with η^1 -character or *vice versa* (Y. Wakatsuki, H. Yamazaki, Y. Maruyama and I. Shimizu), (430) C60

Binuclear hydroxo-, methoxo-, and pyrazolate-bridged complexes of ruthenium(II). Molecular structure of $[(p\text{-cymene})Ru]_2(\mu\text{-pz})(\mu-OH)_2BF_4$ (*p*-cymene = *p*-Me₆H₄CHMe₂, pz = pyrazolate) (D. Carmona, A. Mendoza, J. Ferrer, F.J. Lahoz and L.A. Oro), (431) 87

Optically active complexes of transition metals (Rh^I, Ru^{II}, Co^{II} and Ni^{II}) with 2-aminocarbonylpyrrolidine ligands. Selective catalysts for hydrogenation of prochiral olefins (A. Corma, M. Iglesias, C. Del Pino and F. Sánchez), (431) 233

Struktur von $[Cp^*Ru(CO)_2]_2$ ($Cp^* = \eta^5\text{-Pentamethylcyclopentadienyl}$) (A. Steiner, H. Gornitzka, D. Stalke and F.T. Edelmann), (431) C21

Ein neues Verfahren zur Synthese von Metall-funktionalisierten Diphosphenen, Arsaphosphenen und Tetraphospha-bicyclobutanen (P. Jutzi and S. Opiela), (431) C29

Fixierung von Harnstoffderivaten an einem dreikernigen Rutheniumverband: Synthese und Molekülstruktur von $(\mu_2-H)Ru_3(CO)_9(\mu_3-\eta^2-HNCONHCH=CHMe)$ (U. Bodensieck, J. Santiago, H. Stoeckli-Evans und G. Süss-Fink), (433) 141

Rutheniuminduzierter Abbau von Tetramethylthioharnstoff zu Carbénliganden. Synthese und Molekülstruktur der Cluster $Ru_4(\mu_2-CO)_9\cdot_n(\mu_4-S_2)[CNMe_2]_2$ ($n = 1, 2$) und $(\mu_2-H)Ru_3(CO)_9(\mu_3-\eta^2-HNRu_3(CO)_9(\mu_3-S)RuC_6H_4NMeCNMe_2)$ (U. Bodensieck, H. Stoeckli-Evans und G. Süss-Fink), (433) 149

Rutheniuminduzierter Abbau von Tetramethylthioharnstoff zu Carbén- und Dicarbénliganden. Synthese und Molekülstruktur der Cluster $Ru_4(CO)_{10}(\mu_2-CNMe_2)(\mu_3-S)(\mu_3-SRu(CO)_3(\eta^2-CH_2NMeCNMe_2))$, $Ru_5(CO)_{11}(\mu_2-CNMe_2)_2(\mu_4-S)_2$ und

- $\text{Ru}_4(\mu_2\text{-CO})_3(\text{CO})_6(\mu_4\text{-S})_2(\eta^2\text{-Me}_2\text{NCNMeCH}_2\text{CNMe}_2)$ (U. Bodensieck, H. Stoeckli-Evans, G. Rheinwald und G. Süss-Fink), (433) 167
- Selective activation of carbon-chlorine bonds by the electrophilic fragment “ Cp^*Ru^+ ” (D. Rondon, X.-D. He and B. Chaudret), (433) C18
- Mercury-bridged transition-metal clusters. Synthesis of pentanuclear Ru_3HgM ($M = \text{Mo, W, or Co}$) clusters and X-ray structure of $[(\text{Ru}_3(\mu_3,\eta^2\text{-ampy})(\text{CO})_9\text{-}(\mu_3\text{-Hg})\text{Co}(\text{CO})_4)]$ (Hampy = 2-amino-6-methylpyridine) (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, S. García-Granda and J.F. Van der Maelen), (434) 123
- [Tris(1-pyrazolyl)hydroborato]ruthenium chemistry: preparation and reactions of $[\text{Ru}(\eta^3\text{-HB[1-pyrazolyl]})_3\text{-}(\text{PPh}_3)(\text{CO})(X)]$ ($X = \text{H, Cl}$). The X-ray crystal structure of $[\text{Ru}(\eta^3\text{-HB[1-pyrazolyl]})_3\text{-}(\text{PPh}_3)(\text{CO})(\text{PMe}_3)]\text{PF}_6$ (N.-Y. Sun and S.J. Simpson), (434) 341
- The synthesis and characterisation of an ortho-metallated carbeneruthenium(II) complex, containing a metallated *N*-benzyl substituent with a 6-membered metallate nucleus, crystal structure of $[\text{RuCl}(\text{CN}(\text{CH}_2\text{C}_6\text{H}_5)\text{CH}_2\text{CH}_2\text{NCH}_2\text{C}_6\text{H}_4)\text{(PPh}_3)_2]$ (M.A. Owen, P.L. Pye, B. Piggott and M.V. Capparelli), (434) 351
- Synthesis, crystal structures, and solution dynamics of some mono(2,4-dimethylpentadienyl)ruthenium(II) complexes (T. Luminii, D.N. Cox, R. Roulet and K. Schenk), (434) 363
- Synthesis, characterisation and molecular structure of $[\text{Ru}_5\text{C}(\text{CO})_{15}\text{-}(\text{HgCF}_3\text{-}(\text{CF}_3\text{CO}_2))]$ (B.F.G. Johnson, J. Lewis, P.R. Raithby, V.P. Saharan and W.T. Wong), (434) C10
- Synthesis and solution properties of the boron-containing clusters $\text{HM}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf})$ ($M = \text{Fe or Ru}$) and the competitive formation of $\{\text{HM}_4(\text{CO})_{12}\text{BAu}_2\}_2(\text{dppf})$ for $M = \text{Ru}$ ($\text{dppf} = 1,1'\text{-bis(diphenylphosphino)ferrocene}$). Molecular structure of $\text{HRu}_4(\text{CO})_{12}\text{BAu}_2(\text{dppf})\cdot 2\text{CHCl}_3$ (S.M. Draper, C.E. Housecroft and A.L. Rheingold), (435) 9
- Asymmetric hydrogenation of prochiral carboxylic acids and functionalized carbonyl compounds catalysed by ruthenium(II)-binap complexes with aryl nitriles (binap = (*R*)- or (*S*)-2,2'-bis(diphenylphosphino)-1,1'-binaphthyl) (L. Shao, K. Takeuchi, M. Ikemoto, T. Kawai, M. Ogasawara, H. Takeuchi, H. Kawano and M. Saburi), (435) 133
- Reactions of 1,3,5-trithiacyclohexane with tetrahedral ruthenium-cobalt clusters (S. Rossi, J. Pursiainen and T.A. Pakkanen), (436) 55
- o*-Ferrocenyl-Komplexe des Rutheniums (M. Herberhold, W. Feger und U. Kölle), (436) 333
- Hydrogen migration to acyclic pentadienyl ligands: the reactions of (η^5 -2,4-dimethylpentadienyl) (η^4 -2,4-dimethylpenta-1,3-diene)(L)ruthenium(II) tetrafluoroborates ($L = {}^t\text{BuNC, CO, P(OMe)}_3$) with dienes (D.N. Cox, T. Luminii and R. Roulet), (438) 195
- Synthesis of 1,3-dienyl complexes of ruthenium(II) by facile hydro-ruthenation/dehydration of propargylic alcohols (M.C.J. Harris and A.F. Hill), (438) 209
- Reaktionen von $[\text{Cp}^*\text{Ru}(\text{OMe})]_2$. XIII. Pentaalkylcyclopentadienyl Ru und Rh Alkoxo-, Hydroxo- und Thiolato-Komplexe (U. Kölle, A. Hörnig und U. Englert), (438) 309
- Structural aspects of the osmium and ruthenium clusters $[\text{M}_3(\text{CO})_{10}(\text{alkyne})]$ and related substituted compounds (A.J. Deeming and A.M. Senior), (439) 177
- Co-ordinatively unsaturated metal cluster compounds: facile reactivity of the phosphinidene-capped phosphido-bridged triruthenium cluster $[\text{Ru}_3(\mu_3\text{-PPh})(\mu_2\text{-PPh}_2)_2(\text{CO})_7]$ (J.S. Field, R.J. Haines and F. Mulla), (439) C56
- The mechanism of the addition of haloalkanes to alkenes in the presence of $[\text{RuH}_3(\text{SiMe}_2\text{Ph})(\text{PPh}_3)_3]$ and $[\text{RuH}_2(\text{PPh}_3)_4]$ (R. Davis, J.D. Furze, D.J. Cole-Hamilton and P. Pogorzelec), (440) 191
- SO und NO als Konkurrenzliganden im neuen SO-Komplex $\text{RuCl}(\text{NO})(\text{SO})(\text{PPh}_3)_2$ (O. Heyke, G. Beuter und I.-P. Lorenz), (440) 197
- Reactions of cyclic thioethers with $\text{Ru}_3(\text{CO})_{12}$. Synthesis, reactivity and crystal structures of $\text{HRu}_3(\text{CO})_9(\mu_3\text{-}\eta^3\text{-C}_5\text{S}_2\text{-}1,3,5\text{-dithiacyclohexane})$, $\text{Ru}_3(\text{CO})_9(\mu_3\text{-}\eta^3\text{-}1,3,5\text{-trithiacyclohexane})$ and $\text{Ru}_3(\text{CO})_9(\mu_2\text{-}\eta^3\text{-}1,4,7\text{-trithiaheptane})$ (S. Rossi, K. Kallinen, J. Pursiainen, T.T. Pakkanen and T.A. Pakkanen), (440) 367
- On the problem of stabilization of α -metallocenylcarbocations. Synthesis, ^1H and ^{13}C NMR spectra of metallocenyldiphenylmethyl hexafluorophosphates and X-ray investigation of $(\text{C}_5\text{H}_5\text{OsC}_5\text{H}_4\text{-CPh}_2)^+\text{PF}_6^-$ (U. Turpeinen, A.Z. Kreindlin, P.V. Petrovskii and M.I. Rybinskaya), (441) 109
- Reaction of $\text{Ru}_3(\text{CO})_{12}$ with ferrocenylacetylene. Crystal and molecular structure of $\text{Ru}_2(\text{CO})_6[\text{C}_4\text{H}_2(\text{C}_5\text{H}_4\text{FeC}_5\text{H}_5)_2]$ (A.A. Koridze, A.I. Yanovsky and Y.T. Struchkov), (441) 277
- Zweikernige Rutheniumcluster mit $\mu_2\text{-}\eta^2\text{-Sulfonatoliganden}$. Synthese und Molekülstruktur von $\text{Ru}_2(\text{CO})_4[\mu_2\text{-}\eta^2\text{-OOS(O)Tol}]_2\text{-}(\text{PPh}_3)_2$ (G. Rheinwald, H. Stoeckli-Evans und G. Süss-Fink), (441) 295
- Mixed-valence state of 1',1'''-dialkylbiruthenocenium(II,IV) salts (M. Watanabe, T. Iwamoto, H. Sano, A. Kubo and I. Motoyama), (441) 309
- Neue Derivierungswege ethinylsubstituierter Metallocene (M. Buchmeiser und H. Schottenberger), (441) 457
- Synthesis of Ru^{II} hydride and alkenyl amidine complexes. The crystal structure of $[\text{Ru}(\text{CO})(\text{CH}=\text{CHCMe}_3)(\text{NH}=\text{C}(\text{Me})\text{-}(\text{Me}_2\text{pz}))\text{(PPh}_3)_2]\text{PF}_6$ (J. López, A. Santos, A. Romero and A.M. Echavarren), (443) 221
- Synthesis of bridged and linked ruthenium and osmium carbonyl clusters containing a $[\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)]^{2+}$ unit. The crystal and molecular structures of $\text{Ru}_5\text{C}(\text{CO})_{14}\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{-CH}_2\text{PPh}_2)$ and $\{\text{Os}_4\text{H}_3(\text{CO})_{12}\}_2\text{Au}_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)$ (A.J. Amoroso, A.J. Edwards, B.F.G. Johnson, J. Lewis, M.R. Al-Mandhary, P.R. Raithby, V.P. Saharan and W.T. Wong), (443) C11
- Übergangsmetallkomplexe von Diazenen. XXXI. Orthometallierte Diazenkomplexe von Molybdän, Eisen und Ruthenium: Struktur und Umsetzung zu 2,3-Diphenylindol (D. Garn, F. Knoch und H. Kisch), (444) 155
- Synthesis and properties of the ruthenium vinylidene and acetylide complexes containing 1,1'-bis(diphenylphosphino)ferrocene (M. Sato and M. Sekino), (444) 185
- Photochemical reactivity of $[\text{Ru}^{II}(\text{L})(\text{CO})_2\text{Cl}_2]$ and $[\text{Me}_4\text{N}^+ \text{Ru}^{II}(\text{L})(\text{CO})\text{Cl}_3]$ ($\text{L} = 2,2'\text{-bipyridine or }4,4'\text{-di(isopropoxy-carbonyl)-2,2'-bipyridine}$) in CH_3CN and the redox properties of the resulting new complexes (M.-N. Collomb-Dunand-Sauthier, A. Derouzier and R. Ziessel), (444) 191
- Formal insertion of carbon into a $\text{C}=\text{C}$ double bond on an Ru_5 cluster. X-Ray structure of $\text{Ru}_5(\mu_4\text{-CC}_11\text{H}_{12})\text{-}(\mu_3\text{-SMe})_2(\mu\text{-PPh}_2)_2(\text{CO})_9$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (444) C44
- Cyclohexenyl [2.2]paracyclophane complexes of ruthenium(II): highly fluxional agostics from the sequential reduction of arenes (J.W. Steed and D.A. Tocher), (444) C47
- Synthesis of 1-alkynyl(phthalocyaninato)iron(II) and -ruthenium(II) complexes (M. Hanack, S. Knecht and H.-J. Schulze), (445) 157
- Cluster chemistry. LXXXIII. Reactions of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)(\mu\text{-PPh}_2)_2(\text{CO})_{13}$ with olefins: Products formed by addition of an alkene to the C_2 fragment. X-ray structures of $\text{Ru}_5(\mu_4\text{-PPh})(\mu_4\text{-C}_2\text{CHCHR})\text{CHCH}_2\text{R}(\mu\text{-PPh}_2)_2(\text{CO})_{12}$ ($\text{R} = \text{H, Me}$) and $\text{Ru}_5(\mu_4\text{-PPh})(\mu\text{-PPh}_2)(\mu\text{-CO})(\text{CO})_{10}(\eta^5\text{-C}_5\text{H}_3\text{Me-1-(CH=CH}_2\text{-)3})$ (C.J. Adams, M.I. Bruce, M.J. Liddell, E.R.T. Tiekkink, B.W. Skelton and A.H. White), (445) 187

- Cluster chemistry. LXXXIV. Reactions of $\text{Ru}_5(\mu_5\text{-C}_2\text{PPh}_2)(\mu\text{-PPh}_2)\text{(CO)}_{13}$ with alkenes: attack at the C_2PPh_2 ligand without cleavage of the C-P bond. X-Ray structures of $\text{Ru}_5(\mu_5\text{-C})(\mu\text{-PPh}_2)(\mu\text{-RC}_3\text{H}_3\text{PPh}_2)(\text{CO})_{11}$ ($\text{R} = \text{Me, Ph}$) and $\text{Ru}_5(\mu_5\text{-C})(\mu\text{-PPh}_2)(\mu\text{-PhC}_3\text{H}_3\text{PPh}_2)(\text{CO})_{11-n}\text{P(OMe)}_n$ ($n = 0, 1$) (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (445) 199
- Cluster chemistry. LXXXVI. Metallation of pyridine in reactions with an Ru_5 cluster. X-Ray structures of $\text{Ru}_4(\mu_4\text{-C}_2\text{H}(\text{PPh}_2))(\mu\text{-PPh}_2)(\mu\text{-NC}_5\text{H}_4)(\text{CO})_{10} \cdot \text{CH}_2\text{Cl}_2$ and $\text{Ru}_6(\mu_6\text{-C}_2\text{H})(\mu_3\text{-NC}_5\text{H}_4)(\mu\text{-PPh}_2)_2(\mu\text{-CO})(\text{CO})_{12} \cdot 1.25\text{CHCl}_3$ (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (445) 211
- Metallorganische Lewissäuren. XLVII. Kationische Komplexe $[(\text{OC})_5\text{ReL}]^+$ und $[\text{Cp}(\text{OC})(\text{Ph}_3\text{P})\text{RuL}]^+$ mit O-Donoren und N-Donoren ($\text{L} = \text{Carbonsäureester, Ethanol, Lacton, Lactam, Diphenylketimin}$) (J. Milke, C. Missling, K. Sünkel und W. Beck), (445) 219
- Stereoselektive Bildung von Organoosmium-Komplexen mit Aminocarboxylat-Liganden (H. Werner, T. Daniel, O. Nürnberg, W. Knaup und U. Meyer), (445) 229
- Reactions of Ru_3 clusters with functional phosphines: formation of $\mu_3\text{-acyl}$ and $\mu_3\text{-imidoyl}$ complexes (C.J. Adams, M.I. Bruce, O. Kühl, B.W. Skelton and A.H. White), (445) C6
- ^1H , ^{13}C NMR spectroscopic studies of iodoferrocenylruthenocenium $^+$ salts (M. Watanabe, T. Iwamoto, H. Sano and I. Motoyama), (446) 177
- Ruthenium complexes with bridging functionalized alkylidene ligands. Synthesis of $[\text{Ru}_2\text{Cp}_2(\text{CO})_2(\mu\text{-CO})(\mu\text{-CX})\text{N}(\text{Me})\text{CH}_2\text{Ph}]$ ($\text{X} = \text{H, CN}$) and molecular structure of the CN derivative (L. Busetto, L. Carlucci, V. Zanotti, V.G. Albano and M. Monari), (447) 271
- Hydride-phosphoniodithiocarboxylate/phosphonium-betaine isomerism in Cy_3PCS_2 complexes of ruthenium (A.L. Hector and A.F. Hill), (447) C7
- Heterometallic clusters in homogeneous catalysis. Hydrogenation of diphenylacetylene and isomerization of *cis*-stilbene in the presence of $[\text{Fe}_2\text{Ru}(\text{CO})_{12}]$, $[\text{FeRu}_2(\text{CO})_{12}]$, $[\text{H}_2\text{FeRu}_3(\text{CO})_{13}]$, $[\text{H}_2\text{Ru}_4(\text{CO})_{13}]$, and related compounds. Identification of organometallic complexes and a discussion of their role (R. Giordano and E. Sappa), (448) 157
- ^1H , ^{13}C -NMR and ^{57}Fe -Mössbauer spectroscopic studies of α,α -ferrocenylruthenocenylmethylium $^+$ ion (M. Watanabe, T. Iwamoto, S. Nakashima, H. Sakai, I. Motoyama and H. Sano), (448) 167
- Addition of carboxylic acids to alkynes catalysed by ruthenium complexes (M. Rotem and Y. Shvo), (448) 189
- Reactions of dicarboxylic acids with $[\text{Ru}(\text{CO})_3(\text{PPh}_3)_2]$ (M. Pizzotti, S. Cenini and F. Porta), (448) 205
- Facile synthesis of $\text{Ru}_4(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and its reactions with phosphine ligands (P. Mathur, M.M. Hossain and R.S. Rashid), (448) 211
- Transition metal polyhydrides-catalyzed addition of activated nitriles to aldehydes and ketones via Knoevenagel condensation (Y. Lin, X. Zhu and M. Xiang), (448) 215
- Stability of the silicon-ruthenium bond in the presence of a strong nucleophile. Phase sensitive ^1H NMR COSY and crystal structure of $\text{Ru}(\text{CO})_2(\text{NH}_2\text{CH}_2\text{C}_6\text{H}_5)_2(\text{Si}(\text{C}_6\text{H}_5)_2\text{CH}_3)_2\text{I}$ (B.J. Rapoport, K.J. McGrath, C.F. George and J.C. Cooper), (450) 85
- Ferrocene derivatives. XXVIII. The metallation of phenylazoferrrocene and the synthesis of phenylazocyclopentadienyl derivatives of other metals (G.R. Knox, P.L. Pauson and D. Willison), (450) 177
- Ruthenium complex-catalyzed allylic alkylation of carbonucleophiles with allylic carbonates (S.-W. Zhang, T.-a. Mitsudo, T. Kondo and Y. Watanabe), (450) 197
- Cyclopentadienyl-ruthenium and -osmium chemistry. XXXIX. Syntheses of novel alkynyl-ruthenium complexes. X-Ray structures of two forms of $\{\text{Ru}(\text{PPh}_3)_2(\eta\text{-C}_5\text{H}_5)\}_2(\mu\text{-C}_4)$ and of $\text{Ru}[\text{C}\equiv\text{CC}(\text{O})\text{-Me}](\text{PPh}_3)_2(\eta\text{-C}_5\text{H}_5)$ (M.I. Bruce, P. Hinterding, E.R.T. Tiekkink, B.W. Skelton and A.H. White), (450) 209
- A new route to complexes containing the tetracarbon (C_4) ligand (C.J. Adams, M.I. Bruce, B.W. Skelton and A.H. White), (450) C9
- Samarium**
- Dimorphe von kristallinem Acetonitril-tris(η^5 -cyclopentadienyl)samarium(III) (M. Adam, H. Schultze and R.D. Fischer), (429) C1
- Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79
- Metalloligands containing samarium(III). Access to binuclear f -metal (samarium)- d -metal (rhodium) complexes (M. Visseaux, A. Dormond, M.M. Kubicki, C. Moise, D. Baudry and M. Ephritikhine), (433) 95
- Ferrocenhaltige Liganden in der Lanthanid-Chemie: Alkoxide und Thiolate (H. Gornitzka, F.T. Edelmann und K. Jacob), (436) 325
- Kinetically stable adduct of samarocene with aluminium deuteride ($\eta\text{-C}_5\text{H}_3^+\text{Bu}_2^2\text{Sm}(\mu_2\text{-D})_2\text{AlD} \cdot \text{Me}_2\text{NC}_2\text{H}_4\text{NMe}_2$) (Y.K. Gun'ko, B.M. Bulychev, V.K. Belsky and G.L. Soloveichik), (440) 47
- Zur Elektronenstruktur metallorganischer Komplexe der f -Elemente. XXXIII. Esteraddukte des Grundkörpers $\text{Tris}(\eta^5\text{-cyclopentadienyl})$ -lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71
- Studies on organolanthanide complexes. XLIV. Synthesis of bis(2-methoxyethylcyclopentadienyl) divalent organolanthanides ($\text{Ln} = \text{Sm, Yb}$) and molecular structure of solvated $(\text{MeOCH}_2\text{CH}_2\text{C}_5\text{H}_4)_2\text{Yb} \cdot \text{OC}_4\text{H}_8$ (D. Deng, C. Qian, F. Song, Z. Wang, G. Wu and P. Zheng), (443) 79
- Novel polynuclear organotin complexes of samarium and ytterbium (L.N. Bochkarev, O.V. Grachev, N.E. Molosnova, S.F. Zhiltsov, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), (443) C26
- Synthesis and structure of mono-THF solvates of bis(cyclopentadienyl)samarium(II) complexes: $(\text{C}_5\text{Me}_5)_2\text{Sm}(\text{THF})$ and $[\text{C}_5\text{H}_2(\text{SiMe}_3)_3]\text{C}_5\text{H}_3(\text{SiMe}_3)_2\text{Sm}(\text{THF})$ (W.J. Evans, G. Kociok-Köhne, S.E. Foster, J.W. Ziller and R.J. Doedens), (444) 61
- Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), (444) C15
- Studies on organolanthanide complexes. XLVI. Synthesis and characterization of N-containing ring-linked biscyclopentadienyl lanthanide and yttrium chlorides (C. Qian and D. Zhu), (445) 79
- Synthesis and characterization of novel covalent biphenyldiyli Sm^{11} and Yb^{11} complexes (Y. Chauvin, N. Marchal, H. Olivier and L. Saussine), (445) 93
- Metallorganische Verbindungen der Lanthanoide. LXXIV. Tris(trimethylcyclopentadienyl)komplexe der Lanthanoide (H. Schumann, M. Glanz und H. Hemling), (445) C1
- ansa*-Metallocenderivate des Samariums und Ytterbiums mit "weichen" Donorliganden (F.T. Edelmann, M. Rieckhoff, I. Haiduc und I. Silaghi-Dumitrescu), (447) 203
- Tetrahedral versus square planar arrangement of cyclopentadienyl ligands in bimetallic organosamarium complexes. X-ray crystal structure of $[(\text{C}_5\text{H}_4\text{Me})_2(\text{THF})\text{Sm}(\mu\text{-Cl})]_2$ (W.J. Evans, R.A. Keyer and J.W. Ziller), (450) 115
- Intramolekulare Basenkoordination je nach Bedarf: Der 1-(Dimethylaminoethyl)-2,3,4,5-tetramethylcyclopentadienyl-Ligand in der

Komplexchemie von s-, p-, d- und f-Block-Elementen (P. Jutzi, J. Dahlhaus und M.O. Kristen), (450) C1

Scandium

Metallorganische Verbindungen der Lanthanoide. LXX. Bis(cyclopentadienyl)seltenerdkomplexe: Synthese und Röntgenstrukturanalyse von dem intramolekular stabilisierten Lutetiumpentamethyl $\text{Cp}_2\text{Lu}(\text{CH}_2)_3\text{NMe}_2$ und dem monomeren basenfreien Yttriumcarboxylat $\text{Cp}_2\text{Y}[\eta^5\text{-O}_2\text{C}(\text{CH}_2)_3\text{NMe}_2]$ (H. Schumann, J.A. Meese-Märktscheffel, A. Dietrich and F.H. Görlitz), (430) 299

Selenium

Zur Addition von Schwefel und Selen an Dicarbonyl(η^5 -cyclopentadienyl)[η^2 -(methylthio)methylcarben]wolfram-tetrafluoroborat (F.R. Kreißl and N. Ullrich), (426) C8

Struktur und Reaktivität von Ytterbiumbenzamidinaten (M. Wedler, A. Recknagel, J.W. Gilje, M. Nottemeyer and F.T. Edelmann), (426) 295

Neue selenreiche 1,3-Dichalkogenol-2-selone via 4,5-Dilithio-1,3-dichalkogenol-2-selone (H. Poleschner, R. Radeglia and J. Fuchs), (427) 213

Chemistry of hexaruthenium carbido carbonyl cluster complexes. Syntheses and X-ray crystal structures of $\text{Ru}_6(\text{CO})_{15}(\mu\text{-SePh})(\mu\text{-AuPPh}_3)$, $\text{Ru}_6(\text{CO})_{14}(\mu\text{-SePh})_2$, and $\text{Ru}_6(\text{CO})_{14}(\mu\text{-SePh})(\mu,\eta^3\text{-C}_3\text{H}_5)$ (T. Chihara and H. Yamazaki), (428) 169

Structure and dynamics of $[\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{E}(\text{CH}_2)_2)]\text{BF}_4$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$) complexes studied by ^{13}C solid state NMR spectroscopy (A.P.M. Kentgens, H. Karrenbeld, E. De Boer and H. Schumann), (429) 99

3,5-Di-tert-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151

Ferrocenhaltige Liganden in der Lanthanid-Chemie: Alkoxide und Thiolate (H. Gornitzka, F.T. Edelmann and K. Jacob), (436) 325

An *ab initio* study of some free radical homolytic substitution reactions at selenium (J.E. Lyons and C.H. Schiesser), (437) 165

Diferrocenyl dichalkogenide: Synthese und Struktur von Bis[2-(dimethylaminomethyl)ferrocenyl]diselenid (H. Gornitzka, S. Besser, R. Herbst-Irmer, U. Kilimann, F.T. Edelmann and K. Jacob), (437) 299

New cyclic selenide. Synthesis, characterization and reactions of 2-methylselenolan (A.Z. Al-Rubaie), (438) 77

Synthesis and characterization of dinuclear palladium(II) complexes containing both chloro and organochalcogenides as bridging ligands (V.K. Jain and S. Kannan), (439) 231

Trigonal-planar koordiniertes Selen bzw. Tellur in $[\text{Cp}'(\text{CO})_2\text{Mn}]_2\text{EMes}^+$ ($\text{E} = \text{Se}, \text{Te}$): Verbindungen vom "Iniden"-Typ (P. Lau, G. Huttner and L. Zsolnai), (440) 41

Cyclopentadienyl chromium chalcogenide complexes. Synthesis of $[\text{CpCr}(\text{EPH})_2]_2\text{E}$ ($\text{E} = \text{S}, \text{Se}$) and their $\text{Cr}(\text{CO})_5$ adducts. Crystal structures of $[\text{CpCr}(\text{SePh})_2]_2\text{Se} \cdot 1/2\text{C}_6\text{H}_6$ and $[\text{CpCr}(\text{SePh})_2]\text{Se} [\text{Cr}(\text{CO})_5]$ (L.Y. Goh, M.S. Tay, Y.Y. Lim, W. Chen, Z.-Y. Zhou and T.C.W. Mak), (441) 51

Umsetzungen der lithierten Halbsandwich-Chalkogenole, $(\text{CO})_3\text{MC}_5\text{H}_4\text{-ELi}$ ($\text{M} = \text{Mn}, \text{Re}; \text{E} = \text{S}, \text{Se}, \text{Te}$) (M. Herberhold and M. Biersack), (443) 1

The redox behaviour of ferrocene derivatives. I. 1,2,3-Tritellura[3]ferrocenophane and 1,3-ditellura[3]ferrocenophanes (P. Zanello, G. Oppomolla, M. Casarin, M. Herberhold and P. Leitner), (443) 199

Ligand exchange in cyclopentadienylcobalt diselenolenes. Isolation and characterization of the Se_3 -chelate, $\text{C}_5\text{H}_5\text{Co}[(\text{SeC}_8\text{H}_{12})_2\text{Se}]$ (C.P. Morley and R.R. Vaughan), (444) 219

Synthesis of novel selenium-containing osmium carbonyl clusters: X-ray structures of $[\text{Os}_5(\mu\text{-H})(\text{CO})_{15}(\mu_3\text{-SePh})]$ and $[\text{Os}_6(\mu\text{-H})(\text{CO})_{18}(\mu\text{-SeH})]$ (T.M. Layer, J. Lewis, A. Martín, P.R. Raithby and W.-T. Wong), (444) C57

Stereochemistry of the compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$. Crystal and molecular structures of the mixed triiron carbonyl compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$ ($\text{X} = \text{S}, \text{Se}, \text{or Te}$) (G. Gervasio), (445) 147

Un nouveau type de ferrocénophane, premier exemple de structure dimère (R. Broussier, Y. Gobet, R. Amardeil, A. Da Rold, M.M. Kubicki et B. Gautheron), (445) C4

Photochemisch induzierte Addition von Alkinen und Phosphaalkinen an die Cluster $\text{Fe}_3(\text{CO})_9(\mu_3\text{-PR})(\mu_3\text{-X})$ ($\text{X} = \text{Se}, \text{Te}; \text{R} = ^1\text{Pr}, ^1\text{Bu}$) (W. Imhof and G. Huttner), (447) 31

Substitution, nukleophile und oxidative Addition an Arsinidenkomplexe. Zur Reaktion von $[\text{Cp}'(\text{CO})_2\text{Mn}]_2\text{AsCl}$ mit PhSeH und verwandten Chalkogenverbindungen (C. Emmerich und G. Huttner), (447) 81

ansa-Metallocenderivate des Samariums und Ytterbiums mit "weichen" Donorliganden (F.T. Edelmann, M. Rieckhoff, I. Haiduc und I. Silaghi-Dumitrescu), (447) 203

Facile synthesis of $\text{Ru}_4(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and its reactions with phosphine ligands (P. Mathur, M.M. Hossain and R.S. Rashid), (448) 211

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), (449) 167

Bis(2-bromoethyl)selenium dibromide as the selenium-introducing reagent: One-pot preparation of 2,5-bis(alkoxymethyl)tetrahydro-selenophenes by the cyclization of 1,5-hexadiene (Y. Takanohashi, N. Tabata, T. Tanase and S. Akabori), (450) 103

Synthesis and characterisation of new 2-bromovinyl selenides and their platinum group metal complexes (P.K. Khanna and C.P. Morley), (450) 109

Silicon

Silaheterocyclen. XI. Erzeugung und Cycloadditionsverhalten des Diphenyleneopentylsilaethens, $\text{Ph}_2\text{Si}=\text{CHCH}_2^1\text{Bu}$ (N. Auner, W. Ziche and E. Herdtweck), (426) 1

Laser-powered homogeneous pyrolysis of 1,1-dimethyl-1-silacyclobutane in the presence of some common monomers (J. Pola, D. Čukanová, M. Minářík, A. Lyčka and J. Tláskal), (426) 23

Siliciumverbindungen mit starken intramolekularen sterischen Wechselwirkungen. XLVI. Neue 2,4-Disilacyclobutan-1,3-diimine (M. Weidenbruch, J. Hamann, H. Piel, D. Lentz, K. Peters and H.G. von Schnering), (426) 35

Fast atom bombardment mass spectra of silatrane and silocanes (I. Mažeika, S. Grinberga, A.P. Gaukhman, G.I. Zelchan and E. Lukevics), (426) 41

Darstellung und Reaktionen von Alkyl-Hydridotrispyrazolylborat-Komplexen des Titans (J. Ipaktschi and W. Sulzbach), (426) 59

Le 3,4-diméthylsilole (J.-P. Béteilie, A. Laporterie and J. Dubac), (426) C1

Lithio-di-t-butylsilyldiamid, ein cyclisches Anion (B. Tecklenburg, U. Klingebiel and D. Schmidt-Bäse), (426) 287

Struktur und Reaktivität von Ytterbiumbenzamidinaten (M. Wedler, A. Recknagel, J.W. Gilje, M. Nottemeyer and F.T. Edelmann), (426) 295

Alkinkomplexe des Titanocens und Permethyltitanoçens ohne zusätzliche Liganden — Erste Strukturvergleiche (U. Rosenthal, H. Görls, V.V. Burlakov, V.B. Shur and M.E. Vol'pin), (426) C53

Structure and dynamics of hindered organosilicon compounds. The conformations of symmetrical $(\text{Me}_3\text{Si})_3\text{C}$ and $(\text{PhMe}_2\text{Si})_3\text{C}$ derivatives (A.G. Avent, S.G. Bott, J.A. Ladd, P.D. Lickiss and A. Pidcock), (427) 9

- Tetra-*p*-tolyl-Verbindungen *p*-Tol₄Si und *p*-Tol₄Ge: ein Beitrag zur Konfiguration der Tetraaryl-Methan-Analoga Ar₄M (M = C, Si, Ge, Sn, Pb) (M. Charissé, S. Roller and M. Dräger), (427) 23
- Molekülstruktur und Ladungsverteilung in metallorganischen Verbindungen der 4f- und 5f-Elemente. II. Dielektrische Untersuchungen an metallorganischen Komplexen der 4f- und Übergangselemente mit C_{3v}- und D_{3h}-Molekülsymmetrie (R. Maier, B. Kanellakopoulos and C. Apostolidis), (427) 33
- The behaviour of medium-sized permethylated cyclosilanes towards SOCl₂ and SOCl₂–HC(OCH₃)₃ (P.K. Jenkner, A. Spielberger and E. Hengge), (427) 161
- Platinum catalyzed hydrosilylation of alkynes: comparison of rates of addition of terminal olefins to internal alkynes (L.N. Lewis, K.G. Sy and P.E. Donahue), (427) 165
- Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphanyl-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe (CO)₃(R'₃P)(R₃Si)Fe–ML_n (M = Cu, Ag, Au, Hg) (G. Reinhard, B. Hirle and U. Schubert), (427) 173
- Substitution of the acetoxy groups of dialkoxyacetates by organometallic reagents: a route to allyl-, propargyl-, homoallyl-, homopropargyl- and α-stannylacetals (I. Beaudet, A. Duchêne, J.-L. Parrain and J.-P. Quintard), (427) 201
- 2,2'-Bipyridyl-bis[bis(trimethylsilyl)methyl]zink — Synthese, spektroskopische Charakterisierung und Struktur (M. Westerhausen, B. Rademacher and W. Schwarz), (427) 275
- Regioselectivity of inter- and intramolecular hydrosilylation of the vinyloxy group (M.G. Voronkov, S.V. Kirpichenko, V.V. Keiko and A.I. Albanov), (427) 289
- The synthesis and properties of (CH₂F)SiH₃ and related monofluoromethylsilanes (H. Bürger and P. Moritz), (427) 293
- Silicon–carbon unsaturated compounds. XXXIX. Photolysis of 3,4-benzo-1,1,2,2-tetraethyl-1,2-disilacyclobutene (H. Sakamoto and M. Ishikawa), (427) C26
- Platinum complex-catalyzed dehydrogenative double silylation of olefins and dienes with *o*-bis(dimethylsilyl)benzene (M. Tanaka, Y. Uchimaru and H.-J. Lautenschlager), (428) 1
- Photoreactions of silyliron(II) complexes Cp*Fe(CO)₂SiMe₃ (Cp* = η⁵-C₅H₅, η⁵-C₅Me₅) in the presence of trihydrosilanes (Y. Kawano, H. Tobita and H. Ogino), (428) 125
- Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CI. Stereoselektive Synthese gemischalkylierter Osmiumoxide (W.A. Herrmann, S.J. Eder, P. Kiprof and P. Watzlowik), (428) 187
- Conversion of hydrosilanes to alkoxy silanes catalyzed by Cp₂TiCl₂/ⁿBuLi (T.C. Bedard and J.Y. Corey), (428) 315
- Synthese und Charakterisierung von 1,3-disubstituierten Decamethylcyclohexasilanen (E. Hengge and M. Eibl), (428) 335
- Synthese, Struktur und Reaktivität der Ferriophosphonium-Salze [CpFe(CO)(L)(PPh₂CH₂R')]**BF**₄ (L = CO, CH₃CN, PⁿBu₃; R' = Ph, CN, COOEt, PPh₂, SiMe₃); Kristallstrukturen von [CpFe(CO)₂PPh₂CH₂SiMe₃]**BF**₄ und [CpFe(PPh₂CH₂Ph)CpFe(CO)(μ-CO)₂ (C. Klasen, I.-P. Lorenz, S. Schmid and G. Beuter), (428) 363
- Rhodium(I) complex-catalyzed hydrosilylation of dimethyl muconates (K. Yamamoto and T. Tabei), (428) C1
- Photoelektronen-Spektren und Moleküleigenschaften. CXXXI. C(SiH₃)₄, HC(SiH₃)₃ und (H₃O)HC(SiH₃)₂ (H. Bock, M. Kremer and H. Schmidbaur), (429) 1
- Convenient route to di- and triorganosilyl ethyl ethers and the corresponding di- and triorganosilanes (J.M. Tour, J.A. John and E.B. Stephens), (429) 301
- Crystallographic evidence for intermolecular hydrogen bonding between aryl groups and OH groups in silanols (S.S. Al-Juaid, A.K.A. Al-Nasr, C. Eaborn and P.B. Hitchcock), (429) C9
- Cyclopentadienyl-Liganden mit hydrophilen Tentakeln (U. Sieling), (429) C14
- Silicon–silicon interaction in bis(silylene)iron, disilanyliiron, and bis(silyl)iron complexes (K. Ueno, H. Tobita and H. Ogino), (430) 93
- Darstellung und Eigenschaften neuer Monosilylcyclopentasilan-Derivate (P.K. Jenkner and E. Hengge), (430) 259
- Synthesis of silatranyl- and 3,7,10-trimethylsilatranyl-cyclopropanes (M. Nasim, V.S. Petrosyan, G.S. Zaitseva and J. Lorberth), (430) 269
- Organometalloidal derivatives of the transition metals. XXX. Mass spectrometry of transition metal substituted disilanes (A. Guerrero, J. Gómez-Lara, J. Cervantes, L. Velasco, H. Sharma and K.H. Pannell), (430) 273
- Synthese und ungewöhnliche Reaktivität eines Cyclotrisilans (J. Belzner), (430) C51
- Permethylheteroatomcyclopolsilane mit Heteroatomen der VI. Hauptgruppe. Darstellung und spektroskopische Eigenschaften (H. Stüger, M. Eibl, E. Hengge and I. Kovacs), (431) 1
- Si-, Ge- und Sn-diine derivatives in 1,3-dipolar cycloaddition reactions (A.D. Jurayev, A.G. Makhsumov, K. Yuldasheva and M.A. Tolipova), (431) 129
- Dimere Alkalimetallbenzamidinate: Einfluß des Metallions auf die Struktur (D. Stalke, M. Wedler and F.T. Edelmann), (431) C1
- Cleavage of siloxanes with organyltrifluoro- and diorganyldifluorosilanes (M.G. Voronkov, S.V. Basenko, I.A. Gebel, V.Y. Vitkovskii and R.G. Mirskov), (433) 1
- Synthese 1,4-disubstituierter Tetrasilane durch selektive Spaltung von Si–Phenyl-Bindungen (H. Stüger), (433) 11
- Dehydrierende Polymerisation von Methyldisilanen mittels Metallocen-Katalyse zu den Oligomeren Me₃Si[Me₂Si]_nSiMe₂H, HMe₂Si[Me₂Si]_nSiMe₂H mit n = 3–9, H[MeSiH]_nH mit n = 3–14 und hochpolymerem Poly(methylsilan) (MeSiH_x)_n (E. Hengge und M. Weinberger), (433) 21
- Ab initio* study of the molecule C₆H₅SiH₃ and the molecular anion C₆H₅SiH₂⁻: geometries and rotational barriers (F. Ramondo), (434) 19
- Reaktionen subvalenter Verbindungen des Siliciums mit Benzol und mit Biphenyl (F. Huppmann, W. Maringgele, T. Kottke und A. Meller), (434) 35
- Multinuclear NMR spectroscopic studies of some organometallic N-substituted anilines (B.K. Barr, A.J. Herman, L.K. Myers, P.I. Young, C.D. Schaeffer, Jr., H.J. Eppley, J.C. Otter and C.H. Yoder), (434) 45
- 3,5-Di-tert-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151
- Polycyclopentadienyls: synthesis of arylsilylcyclopentadienyl compounds and their η⁵-complexes with tungsten (H. Plenio), (435) 21
- Silicon–carbon unsaturated compounds. XL. Photolysis of 1,4-bis(2-phenyltetramethylsilyl)benzene (M. Ishikawa, K. Watanabe, H. Sakamoto and A. Kunai), (435) 249
- Amination of vinylsilanes and vinylgermanes containing an activated double bond (V. Dirnens, D.I. Barabanov, E. Liepiņš, L.M. Ignatovich and E. Lukevics), (435) 257
- Synthese und Kristallstruktur von Cp'₂Zr{P(SiMe₃)₂}₂ (Cp' = η⁵-C₅H₄Me) — Erste röntgenstrukturanalytische Untersuchung eines Zirkonocen–Bispophidido-Komplexes (F. Lindenberg und E. Hey-Hawkins), (435) 291
- Spectroscopic and AM1 semiempirical studies of vibrations of tris(trimethylsilyl)silane derivatives, (Me₃Si)₃SiX (X = H, Cl, OH, OD, Me, OMe, SiMe₃) (B.H. Boo, H.K. Kang, S.K. Kang, S.S. Lee, D. Kim and M.H. Lee), (436) 1

- Synthesis and structural characterization of cycloborasiloxanes: the X-ray crystal structures of *cyclo*-1,3,3,5,5-pentaphenyl-1-bora-3,5-disiloxane and *cyclo*-1,3,3,5,7,7-hexaphenyl-1,5-dibora-3,7-disiloxane (B.J. Brisdon, M.F. Mahon, K.C. Molloy and P.J. Schofield), (436) 11
- Molecular geometry and ring deformation of trimethylsilylbenzenes: gas-phase electron-diffraction study of 1,3-bis(trimethylsilyl)benzene and 1,3,5-tris(trimethylsilyl)benzene (B. Rozsondai and I. Hargittai), (436) 127
- Titanium-catalyzed cycloaddition reactions of phenyl(trimethylsilyl)acetylene to conjugated dienes and 1,3,5-cycloheptatriene. 1-Phenyl-2-(trimethylsilyl)-cyclohexa-1,4-dienes and their aromatization (R. Klein, P. Sedmera, J. Čejka and K. Mach), (436) 143
- Ethynylferrocene durch direkte Einführung von Acetylenen (M. Buchmeiser und H. Schottenberger), (436) 223
- A study by electron spin resonance spectroscopy of the di-t-butyl-silanone radical anion (A.G. Davies and A.G. Neville), (436) 255
- Cobaltcarbonyl-Komplexe von Alkinyl-Silanen (H. Lang, U. Lay und M. Weinmann), (436) 265
- Palladium-catalyzed rearrangement of silanes containing oxygen or halogen α to silicon (V. Gevorgyan, L. Borisova and E. Lukevics), (436) 277
- Zur Funktionalisierung von Decaphenylcyclopentasilan (W. Uhlig und C. Tretner), (436) C1
- Silicon knocked into a cocked hat (I.M.T. Davidson), (437) 1
- Reactions of sterically hindered organozinc and organocadmium compounds containing functional silicon centres. Crystal structures of $Zn[C(SiMe_3)_2(SiMe_2OCOCF_3)]_2$ and $Cd[C(SiMe_3)_2(SiMe_2OMe)]_2$ (S.S. Al-Juaid, C. Eaborn, A. Habtemariam, P.B. Hitchcock and J.D. Smith), (437) 41
- Carbanion mechanisms. XVIII. Generation of silyl anions by nucleophilic cleavage of disilanes (E. Buncel, T.K. Venkatachalam and U. Edlund), (437) 85
- Regioselectivity of dechlorination: reductive dechlorination of polychlorobiphenyls by polymethylhydrosiloxane-alkali metal (J. Hawari), (437) 91
- Reactions of trimethylaluminium with 2-[methyl-bis(trimethylsiloxy)silyl]but-2-ene-1,4-diol: synthesis and structure of $[Al(CH_3)_3][OCH_2(SiMe_3)_2C=C(H)CH_2O_2][Al(CH_3)_2]$ (S. Pasynkiewicz and W. Ziemkowska), (437) 99
- Addition of sulphenyl halides to alkenyl-metal compounds. III. Arenesulphenyl chloride additions to triethoxy- and triphenylvinylsilanes and to triphenylvinylgermanium. Crystal structure of (4-Me-2-O₂NC₆H₃S)CH(SiPh₃)CH₂Cl (R.A. Howie, G.M. Spencer, J.L. Wardell and J.N. Low), (437) 111
- A convenient synthesis of 1,3-disilacyclobutanes (J.-H. Hong, J.S. Han, G.-H. Lee and I.N. Jung), (437) 265
- Chemical vapour deposition of metals and metal silicides on the internal surfaces of porous silicon (D.G. Anderson, N. Anwar, B.J. Aylett, L.G. Earwaker, M.I. Nasir, J.P.G. Farr, K. Stiebahl and J.M. Keen), (437) C7
- Organoerdmetall-Komplexe von *d*-Block-Elementen. V. Selektive Reduktion von Bis[carbonyl(η^5 -cyclopentadienyl)nickel] zum Carbonyl(η^5 -cyclopentadienyl)nickel-Anion durch C₈K. Synthese und Struktur von (η^5 -C₅H₅)(CO)Ni-Ga[CH₂C(CH₃)₃]₂(OC₄H₈) (R.A. Fischer, J. Behm, E. Herdtweck und C. Kronseder), (437) C29
- Solid-state and solution structures of three lithiumsulfonimidamides: identification of two distinct structural types (F.T. Edelmann, F. Knösel, F. Pauer, D. Stalke and W. Bauer), (438) 1
- Condensation of primary silanes in the presence of Cp₂MCl₂/ⁿBuLi (M = Ti, Zr, Hf) (J.Y. Corey and X.-H. Zhu), (439) 1
- Nickel-catalyzed reactions of 3,4-benzo-1,1,2,2-tetraethyl-1,2-disilacyclobutene (M. Ishikawa, H. Sakamoto, S. Okazaki and A. Naka), (439) 19
- Generation and reactivities of methoxyphenylsilylene (Y.-W. Kwak, I.-H. Jeong, J.-Y. Ko and B.H. Boo), (439) 107
- Synthesis of polymetallacycloalkanes (P. Mazerolles and C. Laurent), (439) 115
- ¹³C NMR studies of silyl and germyl complexes of rhodium and iridium containing carbonyl ligands (E.A.V. Ebsworth and S. Moreton), (439) 213
- Reaktionen von Komplexliganden. LII. Sechsgliedrige Aminocarben-Chelatkomplexe: Synthese, Alkin/Carben-Verknüpfung und Carbenanellierung (K.H. Dötz, A. Rau und K. Harms), (439) 263
- Hydride transfer reactions between penta- and tetra-coordinate silicon derivatives (R.J.P. Corriu, C. Guerin, B.J.L. Henner und Q. Wang), (439) C1
- Die Bildung von [Cp₂Ti(C≡CSiMe₃)₂] aus Titanocen und 1,4-Bis(trimethylsilyl)-1,3-Butadien (U. Rosenthal und H. Görts), (439) C36
- Intramolecular Peterson olefination of *ortho*-trimethylsilylmethyl-N-acyl-N-alkylbenzamides. A new route to 2-alkyl-1(2*H*)isoquinolones (A. Couture, H. Cornet und P. Grandclaudon), (440) 7
- An investigation of steric effects along a cyclopentadienyl substituent: a study by ¹H, ¹³C and ²⁹Si NMR spectroscopy of bis(trimethylsilyl)cyclopentadienyl iron complexes (D. White, L. Carlton und N.J. Coville), (440) 15
- The mechanism of the addition of haloalkanes to alkenes in the presence of [RuH₃(SiMe₂Ph)(PPh₃)₃] and [RuH₂(PPh₃)₄] (R. Davis, J.D. Furze, D.J. Cole-Hamilton und P. Pogorzelec), (440) 191
- Le (chlorodiisopropyl)silyl fluorenyllithium: un equivalent synthétique du diisopropyl(fluorenylidène)silene (C. Couret, J. Escudé, G. Delpon-Lacaze et J. Satgé), (440) 233
- Allénylsilanes fonctionnels. II. Synthèse par voie organoaluminique du 2-triméthylsilylbuta-2,3-diénal et application à la synthèse d'alcôols secondaires α -alléniques triméthylsilylés (D. Mesnard et L. Miginiac), (440) 277
- Heterozyklen als Liganden. XVI. (2,5-C₄-¹Bu₂RHN)MCl₃ (M = Ti, Zr, Hf; R = H, SiMe₃)—Azacyclopentadienyl-Komplexe der Gruppe 4-Metalle (N. Kuhn, S. Stubenrauch, R. Boese und D. Bläser), (440) 289
- On the synthesis of siloxanes. XXI. Verification of inductive and steric substituent constants for siloxy groups by reaction of triorganylchlorosilanes and chlorosiloxanes with lithium silanolates and lithium isopropylate (K. Käppler, U. Scheim, K. Rühlmann und A. Porzel), (441) 15
- Rearrangement of silatranyl- and 3,7,10-trimethylsila-tranyloxiranes to silatranylaldehydes and the crystal structure of silatranylacetraldehyde (M. Nasim, V.S. Petrosyan, G.S. Zaitseva, J. Lorberth, S. Wocadlo und W. Massa), (441) 27
- Siliciumverbindungen mit starken intramolekularen sterischen Wechselwirkungen. XLVIII. Trisilacyclobutanimine: Molekülstrukturen und lichtinduzierte Reaktionen (M. Weidenbruch, J. Hamann, K. Peters, H.G. Von Schnering und H. Marsmann), (441) 185
- Metallation reactions. XVIII. Regioselective metallation of (alkylthio)benzenes by superbases (S. Cabiddu, C. Fattuoni, C. Floris, G. Gelli und S. Melis), (441) 197
- Metallorganische Lewis-Säuren. XLV. Pentacarbonylrhenium-Komplexe mit Oxoanionen (Trifluoromethoxyborat, Phosphat, Sulfat, Dichromat) sowie von Heteropolysäuren. Chemisorption von kationischen Carbonylrhenium-Komplexen auf γ -Al₂O₃, SiO₂, TiO₂ und in Zeolithen (E. Fritsch, J. Heidrich, K. Polborn und W. Beck), (441) 203

- Dimethylvinylsilylation of $\text{Si}_8\text{O}_{20}^{8-}$ silicate anion in methanol solutions of tetramethylammonium silicate (I. Hasegawa and S. Motojima), (441) 373
- Cleavage of Si-C and Ge-C bonds in heterysilanes and -germanes by organolithium reagents (V. Gevorgyan, L. Borisova and E. Lukevics), (441) 381
- Stereoselective Additionsreaktionen von Komplex-gebundenen Phosphanen und Arsanen an Alkine (H. Lang und U. Lay), (441) 389
- Dehydrierende Copolymerisation zu den neuen Polysilan-Copolymeren Poly(methyl-co-phenylsilan) $\text{H}[(\text{MeSiH})_x(\text{PhSiH})_y]_n\text{H}$ und Poly(methyl-co-dimethylsilan) $\text{H}[(\text{MeSiH})_x(\text{Me}_2\text{Si})_y]_n\text{H}$ (E. Hengge und M. Weinberger), (441) 397
- Die Reaktion des Tetrakis[bis(trimethylsilyl)methyl]dialans(4) mit Schwefelkohlenstoff: Molekülstrukturen mit Al-S-Al- und Al- CS_2 -Al-Fragment (W. Uhl, A. Vester und W. Hiller), (443) 9
- First observation of stable primary radical cations ${}^1\text{Bu}_3\text{SiH}^+$ formed by directed radiolysis of a pure compound: ${}^1\text{Bu}_3\text{SiH}$ (C.J. Rhodes), (443) 19
- Bis[2,2,4,4,6,6-hexamethyl-2,4,6-trisila-cyclo-hexyl]zink und sein 2,2'-Bipyridyl-Addukt – spektroskopische Charakterisierung und Struktur (M. Westerhausen und B. Rademacher), (443) 25
- Viergliedrige anorganische Ringsysteme des zweiwertigen Zinns und Bleis: Synthese und Struktur von chelatstabilisierten Stannylenen und Plumbylenen (U. Kilimann, M. Noltmeyer und F.T. Edelmann), (443) 35
- Sila-Riechstoffe und Riechstoffisostere. XII. Geruchsvergleiche homologer Organoelementverbindungen der vierten Hauptgruppe (C, Si, Ge, Sn) (U. Wannagat, V. Damrath, V. Huch, M. Veith und U. Harder), (443) 153
- Mechanistische Aspekte zur Übergangsmetall-katalysierten dehydrierenden Polymerisation von Disilanen und höheren Silanen (E. Hengge und M. Weinberger), (443) 167
- Preparation of new C-alkyl-C-stannyliamines (B. Jousseaume, M. Pereyre, N. Petit, J.-B. Verlhac and A. Ricci), (443) C1
- Silicon-carbon unsaturated compounds. XLII. Novel dimerization of a silene generated from 1-pentamethyldisilanyl-4-(trimethylsilyl)benzene (M. Ishikawa, M. Kikuchi, K. Watanabe, H. Sakamoto and A. Kunai), (443) C3
- Exclusive β -hydrosilylation of acrylates catalyzed by copper-tetramethylethylenediamine (P. Boudjouk, S. Kloos and A.B. Rajkumar), (443) C41
- Ultrasound in organometallic chemistry. The effects of temperature, metal purity and power source on the ultrasound-promoted reaction between trimethylchlorosilane and lithium (P.D. Lickiss and R. Lucas), (444) 25
- Synthese und Charakterisierung von 1,3,5-trisubstituierten Nonamethylcyclohexasilanen (M. Eibl, U. Katzenbeißer und E. Hengge), (444) 29
- Regioselectivity in the trichlorosilylation of allylic halides. Synthesis of 1,1-dichloro-3-(methyldichlorosilyl)-1-silacyclopent-3-ene (R. Boukherroub, G. Manuel and W.P. Weber), (444) 37
- Synthèse de phosphinites dérivés de sucres: applications en cyclodimérisation du penta 2,4-diénoate de méthyle, en hydrogénéation et en hydrosilylation de la pentolactone (A. Bendayan, H. Masotti, G. Peiffer, C. Siv and A. Archavlis), (444) 41
- Synthesis of the square-bipyramidal cluster $[\text{Co}_4(\mu_4\text{-SiMe})_2(\text{CO})_{11}]$ by two routes and its reaction with GeMe_2H_2 . The crystal structures of $[\text{Co}_4(\mu_4\text{-SiMe})_2(\text{CO})_{11}]$ and $[\text{Co}_4(\mu_4\text{-SiMe})_2(\mu\text{-GeMe}_2)_2(\text{CO})_{10}]$ (S.G. Anema, S. Kim Lee, K.M. Mackay and B.K. Nicholson), (444) 211
- Dérivés silylés issus d'aldimines α,β -insaturées: préparations et applications en synthèse (M. Bellassoued et A. Majidi), (444) C7
- Comb-like polycarbosilanes-routes to a new group of liquid crystal polymers (E. Bialecka-Florjanczyk, T. Ganicz, M. Pluta and W. Stańczyk), (444) C9
- Übergangsmetall-Silyl-Komplexe. XLV. Erzeugung von Fluorsilyl-Liganden durch selektive Spaltung metallgebundener Si-Ph-Gruppen (S. Gilbert und U. Schubert), (444) C12
- Neue Cyclooctatetraenyl-lanthanid(III)-Komplexe (U. Kilimann und F.T. Edelmann), (444) C15
- Synthesis of $\text{Me}_2\text{Si}-\eta^5\text{C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{Fe}(\text{CO})-\eta^5\text{C}_5\text{H}_4\text{SiMe}_2$ and a novel reaction between its Si-Si and Fe-Fe bonds (H. Sun, S. Xu, X. Zhou, H. Wang, R. Wang and X. Yao), (444) C41
- Sterically overcrowded or charge perturbed molecules. XXXIII. Hexakis(trimethylsilyl)disilane: structure and photoelectron spectrum of a sterically overcrowded molecule (H. Bock, J. Meuret and K. Ruppert), (445) 19
- Photoelectron spectrum and reactivity of silylalkyl sulphides. Stabilization of radical cations by the β -silyl effect (L. Nyulászi, T. Vesprémi and J. Réffy), (445) 29
- Synthesis and characterization of bis(trichlorosilylmethyl)benzenes (M. Mottevalli, D.L. Ou, A.C. Sullivan and S.W. Carr), (445) 35
- Über gemischte Gruppe 14-Gruppe 14-Bindungen. V. Sechs Perphenylierte Propan-Analoga $\text{Ph}_3\text{M}-\text{M}'\text{Ph}_2-\text{M}''\text{Ph}_3$ ($\text{M}, \text{M}', \text{M}'' = \text{Si}/\text{Ge}$): ein Beitrag zum Elektronegativitätsunterschied zwischen Silicium und Germanium (M. Charissé, M. Mathes, D. Simon und M. Dräger), (445) 39
- Synthesis, reactivity, and molecular structure of $[(\text{N}(\text{CH}_2\text{CH}_2\text{O})_3\text{Si-C}_6\text{H}_5)\text{W}(\text{CO})_3]$ (J.-S. Lee, Y.K. Chung, D. Whang and K. Kim), (445) 49
- Molecular λ^5 -silicates: synthesis, structure and properties of zwitterionic λ^5 -organospirosilicates and λ^5 -organofluorosilicates (R. Tacke, J. Becht, A. Lopez-Mras and J. Sperlich), (446) 1
- Organosiloxanes with functional groups – a short review (G. Schmaucks, R. Wagner and R. Wersig), (446) 9
- Recent advances in catalytic hydrosilylation (B. Marciniec and J. Guliński), (446) 15
- Dimethylsiloxy-aluminiumdihalide dimers. Molecular structures of $[\text{Me}_2\text{HSiOAlX}_2]_2$ ($\text{X} = \text{Cl}, \text{I}$) and $[\text{Me}_3\text{SiOAl}(\text{BH}_4)_2]_2$ (P. Bissinger, P. Mikulcik, J. Riede, A. Schier and H. Schmidbaur), (446) 37
- A stable aquo-complex of lithiated di-tert-butylfluorosilanol. Synthesis and crystal structure (S. Schütte, U. Pieper, U. Klingebiel and D. Stalke), (446) 45
- Beiträge zur Chemie der Halogensilan-Addukte. XXIII. Über die Ionisierung der SiCl -Bindung. Temperaturabhängige Übergänge zwischen penta- und tetrakoordinierten SiCl -Verbindungen. Die Kristall- und Molekülstruktur von Chlоро-dimethyl-(3,4,7,8-tetrahydro-2H,6H-pyrimido[1,2- α]pyrimidin-1-ylmethyl- C^1,N^9)-silicium (D. Kummer, S.H.A. Halim, W. Kuhs und G. Mattern), (446) 51
- Linear 2,2-diaryl-substituted trisilanes: structure and photolysis to disilenes (R.S. Archibald, Y. Van den Winkel, D.R. Powell and R. West), (446) 67
- Synthesis of siloxanes. XX. Stereochemical investigations of substitution reactions at cyclotrisiloxanes with new types of model compounds (R. Gewald, U. Scheim, R. Lang, K. Rühlmann and R. Lehnert), (446) 79
- Synthesis, properties and characterization of octadecamethylbicyclo[4.4.0]decasilane (P.K. Jenkner, E. Hengge, R. Czaputa and C. Kratky), (446) 83
- Disproportionation of oligodimethylsiloxanols in the presence of a protic acid in dioxane (M. Cypryk, S. Rubinsztajn and J. Chojnowski), (446) 91

- MNDO, AM1 and PM3 semiempirical molecular orbital study of 1-fluorosilatrane (G.I. Csonka and P. Hencsei), (446) 99
- Synthesis and characterization of (4S)-silatrane-4-carboxylic acids; molecular structure of (3R,4S)-1-vinyl-3-methylsilatrane-4-carboxylic acid (R.-X. Zhuo, Z.-R. Lu, J. Liao and L.-F. Shen), (446) 107
- Photoelectron spectra and molecular properties. CXXXVIII. 1,4-Di[tris(trimethylsilyl)silyl]benzene: synthesis, structural analogy, photoelectron spectrum and ESR/ENDOR characterization of its radical anion (H. Bock, J. Meuret, R. Baur and K. Ruppert), (446) 113
- A novel route to the trisilacyclobutane moiety. A possible silene–disilene reaction (D. Bravo-Zhivotovskii, Y. Apeloig, Y. Ovchinnikov, V. Igonin and Y.T. Struchkov), (446) 123
- Excimer laser photolysis of hexamethyldisilazane (J. Pola and R. Taylor), (446) 131
- Synthèse d'aryl- et hétéroarylsilanes par scission de l'hexaméthyl-disilane (P. Babin, B. Bennetau, M. Theurig et J. Dunoguès), (446) 135
- Chemistry of decamethylsilicocene: oxidative addition of compounds with X–H bonds (X = F, Cl, Br, O, S) (P. Jutzi, E.-A. Bunte, U. Holtmann, B. Neumann and H.-G. Stamm), (446) 139
- Cyclohydrosilylation dimer formation: evidence for $\text{Pt}^0 \leftrightarrow \text{Pt}^{\text{II}} \leftrightarrow \text{Pt}^{\text{IV}}$ catalysis (H.K. Chu and C.L. Frye), (446) 183
- Direct and sequential synthesis of polysilazanes using lithiated (n-butylamino)silanes (B.J. Aylett and C.-F. Liaw), (446) C1
- The effect of pentacoordination on silicon-29 NMR chemical shifts and silicon–hydrogen coupling constants (A.R. Bassindale and J. Jiang), (446) C3
- Unexpected behaviour of a hexacoordinate silicon compound (F. Carré, C. Chuit, R.J.P. Corriu, A. Mehdi and C. Reyé), (446) C6
- Insertion of dimethylsilylene into the sulphur–hydrogen bond of trialkoxysilanethiols (S. Konieczny, K. Wrzesień and W. Wojnowski), (446) 73
- Metathesis of silicon-containing olefins. VIII. Synthesis of alkenyl (phenyl, methyl) substituted silanes by olefin metathesis (B. Marciniec and C. Pietraszuk), (447) 163
- Sila et germacyclopentanes C-bifonctionnels. I. 1-Sila et 1-germacyclopentan-2,3 et 3,4-diols (G. Manuel et R. Boukherroub), (447) 167
- Dehydrogenative silylation, isomerization and the control of *syn*- vs. *anti*-addition in the hydrosilation of alkynes (C.-H. Jun and R.H. Crabtree), (447) 177
- Organometalloid derivatives of the transition metals. XXXIII. Reactions of ferrocenylcarboxaldehyde with base and trimethylsilylcyanide. A new high yield reaction for the formation of cyanoamines or silylethers (E. Peña, M. Rivera-Claudio, R.N. Kapoor and K.H. Pannell), (447) 265
- An easy approach to 1-silylated ketones and asymmetrical 1,6- and 1,8-dicarbonyl compounds (V. Fiandanese, A. Punzi and N. Ravasio), (447) 311
- Synthesis of the heterodinuclear Fe-Pt complex $[(\text{OC})_3\text{Fe}(\mu\text{-CN}(2,6\text{-xylyl}))(\mu\text{-dppm})\text{Pt}(\text{PPh}_3)]$ containing a bridging isonitrile ligand and electrophilic additions leading to μ -aminocarbyne complexes (M. Knorr, T. Faure and P. Braunstein), (447) C4
- 4-Phenylbenzylidene benzylamine: a new and convenient reagent for the titration of solutions of lithium alkyls and metal amides (L. Duhamel and J.-C. Plaquevent), (448) 1
- The chemistry of the Si–H bond with acid treated clays (L.N. Lewis, D.S. Johnson and J.D. Rich), (448) 15
- Characterization of new cyclosiloxazanes using both GC/MS and GC/FT-IR (A. Marchand, G. Bourgeois, P. Gerval, E. Duguet and A. Soum), (448) 19

Über gemischte Gruppe 14–Gruppe 14-Bindungen. VII. Ein Inkrementensystem für ^{13}C -NMR chemische Verschiebungen bei arylierten Gruppe 14-Verbindungen (C. Schneider-Koglin, B. Mathiasch und M. Dräger), (448) 39

Über Tetraaryl-Methan-Analoga in der Gruppe 14. II. $\text{Ph}_{4-n}\text{Si}-p\text{-Tol}_n$ und $\text{Ph}_{4-n}\text{Sn}-p\text{-Tol}_n$ ($n = 0$ bis 4): Einfluß der *p*-Tolyl-Substituenten auf Struktur und NMR-chemische Verschiebungen (M. Charissé, V. Gauthey und M. Dräger), (448) 47

Synthesis, isolation, and characterization of dimeric compounds containing two different exocyclic ligands on the Group 13 center. X-Ray crystal structures of $[\text{R}(\text{Me}_3\text{SiCH}_2)\text{InE}(\text{SiMe}_3)_2]_2$ ($\text{R} = \text{Ph}, \text{E} = \text{As}; \text{R} = \text{Me}, \text{E} = \text{As}, \text{P}; \text{R} = \text{Cl}, \text{E} = \text{P}$) (R.L. Wells, A.T. McPhail, L.J. Jones III and M.F. Self), (449) 85

Measurement of surface tension for polysilanes (T. Fujisaka, R. West and C. Murray), (449) 105

Pyrolysis of poly(silylene)diacetylenes: direct evidence between their morphology and thermal behavior (R. Corriu, Ph. Gerbier, C. Guérin, B. Henner and R. Fourcade), (449) 111

Siliciumverbindungen mit starken intramolekularen sterischen Wechselwirkungen. XLIX. Disilane und ein Disilen mit unsymmetrischer Substitution an den Siliciumatomen (M. Weidenbruch, A. Pellmann, Y. Pan, S. Pohl, W. Saak und H. Marsmann), (450) 67

Synthesis of siloxanes. XXII. Synthesis and crystal structure of 2-t-butoxy-2,4,6-trimethyl-4,6-diphenyl-(2 α ,4 α ,6 α)-cyclotrisiloxane (R. Gewald, U. Scheim, K. Rühlmann, H. Goesman and D. Fenske), (450) 73

Darstellung und spektroskopische Charakterisierung verzweigter funktioneller Hexasilangerüste (H. Stüger und P. Lassacher), (450) 79

Stability of the silicon–ruthenium bond in the presence of a strong nucleophile. Phase sensitive ^1H NMR COSY and crystal structure of $\text{Ru}(\text{CO})_2(\text{NH}_2\text{CH}_2\text{C}_6\text{H}_5)_2(\text{Si}(\text{C}_6\text{H}_5)(\text{CH}_3)_2)\text{I}$ (B.J. Rappoli, K.J. McGrath, C.F. George and J.C. Cooper), (450) 85

Rearrangements from the reactions of silenes with trimethylsilylketene (A.G. Brook and A. Baumegger), (446) C9

Silver

Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphanyl-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe $(\text{CO})_3(\text{R}'_3\text{P})(\text{R}_3\text{Si})\text{Fe}-\text{ML}_n$ ($\text{M} = \text{Cu}, \text{Ag}, \text{Au}, \text{Hg}$) (G. Reinhard, B. Hirle and U. Schubert), (427) 173

Fast-atom bombardment (FAB) mass spectra of coordination complexes: aggregation processes in the mass spectra of Group 11 alkynyl complexes (M.I. Bruce and M.J. Liddell), (427) 263

Trimethylsilylcyclooctadiene-transition metal complexes: metal-catalysed protodesilylation of cyclic vinylsilanes, and transfer hydrogenation promoted by the displaced silyl group (B.S. Bandodkar and G. Nagendrappa), (430) 373

A novel high nuclearity bimetallic cluster: the crystal structure of $[\text{N}(\text{PPh}_3)_2]\text{Ag}_6\text{Cu}_7(\text{C}_2\text{Ph})_{14}$ (M.S. Hussain and O.M. Abu-Salah), (445) 295

Crystal structure of $[\text{CF}_3\text{CF}=\text{C}(\text{CF}_3)\text{Ag}]_4$ and its use as a chemical vapor deposition precursor for silver films (P.M. Jeffries, S.R. Wilson and G.S. Girolami), (449) 203

Sodium

The preference of 1-methylallyl polar organometallics and carbanions for *cis* rather than for *trans* geometries (P. von R. Schleyer, J. Kaneti, Y.-D. Wu and J. Chandrasekhar), (426) 143

Reactions of alkali metal macrocyclic complexes XIX. Reaction of potassium or sodium complexes with oxetane – novel route to

- organometallic compounds (Z. Jedliński, A. Misiołek, A. Jankowski and H. Janeczek), (433) 231
- Synthesis of polymetallacycloalkanes (P. Mazerolles and C. Laurent), (439) 115
- Reactions of chelating alkali metal amides with the Grignard reagent $^n\text{BuMgCl}$: crystal structures and solution ^1H NMR spectroscopic studies of the expected mono(amido)-product [$\text{PhCH}_2(\text{Me}_2\text{NCH}_2\text{CH}_2)\text{NMg}^n\text{Bu}_2$] and the unexpected bis(amido)-product [$([\text{Ph}(2-\text{Pyr})\text{N}]_2\text{Mg} \cdot (\text{THF})_2)$] (K.W. Henderson, R.E. Mulvey, W. Clegg and P.A. O'Neil), (439) 237
- A facile synthetic route to (η^5 -benzoylcyclopentadienyl)metal compounds (S.S. Jones, M.D. Rausch and T.E. Bitterwolf), (450) 27
- Sulphur**
- Zur Addition von Schwefel und Selen an Dicarbonyl(η^5 -cyclopentadienyl) \cdot η^2 -(methylthio)methylcarben[wolfram-tetrafluoroborat] (F.R. Kreißl and N. Ullrich), (426) C8
- Organometallic macrocycle chemistry. I. The reactions of $[\text{RuCl}_2(\text{PPh}_3)_3]$ with 1,4,7-trithiacyclononane and 1,4,8,11-tetra-thiacyclotetradecane (A.F. Hill, N.W. Alcock, J.C. Cannadine and G.R. Clark), (426) C40
- Structure and dynamics of $[\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{E}(\text{CH}_3)_2)]\text{BF}_4$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$) complexes studied by ^{13}C solid state NMR spectroscopy (A.P.M. Kentgens, H. Karrenbeld, E. De Boer and H. Schumann), (429) 99
- Übergangsmetall-Heteroallen-Komplexe. XXVI. Synthese und Struktur von zweikernigen Molybdän-Thioketen-Komplexen (W. Ziegler, D. Wormsbächer and U. Behrens), (431) C11
- Thiran-Komplexe aus Thioketonen und $(\text{CO})_5\text{W}=\text{C}(\text{Ph})\text{H}$ bzw. aus Diphenyldiazomethan und $(\text{CO})_5\text{W}[\text{S}=\text{C}(\text{Ph})\text{H}]$ (H. Fischer, K. Treier and U. Gerbing), (433) 127
- 3,5-Di-tert-butylthiobenzoyltrifluorosilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151
- The redox behaviour of ferrocene derivatives. I. 1,2,3-Tritellura[3]ferrocenophane and 1,3-ditellura[3]ferrocenophanes (P. Zanello, G. Opronolla, M. Casarin, M. Herberhold and P. Leitner), (433) 199
- Stereochemistry of the compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$. Crystal and molecular structures of the mixed triiron carbonyl compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$ ($\text{X}, \text{Y} = \text{S}, \text{Se}, \text{or Te}$) (G. Gervasio), (445) 147
- Indium tert-butylthiolates as single source precursors for indium sulfide thin films: is molecular design enough? (A.N. MacInnes, M.B. Power, A.F. Hepp and A.R. Barron), (449) 95
- The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), (449) 167
- Tantalum**
- Aspects of carbonyltantalum chemistry (M. Fornalczyk, F. Süssmilch, W. Priebisch and D. Rehder), (426) 159
- Convenient synthesis of pentamethylcyclopentadienyl-tantalum-diene complexes via the reaction of Cp^*TaCl_4 with methylated-allyl anions (K. Mashima, Y. Yamanaka, S. Fujikawa, H. Yasuda and A. Nakamura), (428) C5
- Reactions of tetrachlorocyclopentadienyltantalum(V) derivatives with hexamethylaluminium: Crystal and molecular structure of dichlorodimethylpentamethylcyclopentadienyltantalum(V) (M. Gómez, G. Jimenez, P. Royo, J.M. Selas and P.R. Raithby), (439) 147
- Methylation of (pentamethylcyclopentadienyl)trichloro(diphenyldimethylenephosphoranyl-*C,C*)tantalum(V). Crystal structures of $[\text{TaCp}^*\text{Cl}_3((\text{CH}_2)_2\text{PPh}_2)]$ and $[\text{TaCp}^*\text{Me}_2((\text{CH}(\text{CH}_2))\text{PPh}_2)]$ (M. Gómez, G. Jimenez, P. Royo, M.A. Pellinghelli and A. Tiripicchio), (439) 309
- Alkoxide, phenoxide and hydrido-phenoxide compounds of tantalum bearing the pentamethylcyclopentadienyl ligand (V.C. Gibson and T.P. Kee), (444) 91
- Rigidity of the unsymmetric coordination geometry of the two dithiolates in $(\text{C}_5\text{Me}_5)\text{Ta}(\text{norbornane}-\text{exo}-2,3\text{-dithiolate})_2$ (K. Tatsumi, Y. Inoue and A. Nakamura), (444) C25
- Technetium**
- Metallorganische Chemie des Technetiums. VIII. Technetium(I)-carbonyl-Komplexe mit Polypyrazol-1-yl-borato-Liganden im Vergleich mit seinen Mn- und Re-Homologen (J.E. Joachim, C. Apostolidis, B. Kanellakopulos, R. Maier, N. Marques, D. Meyer, J. Müller, A. Pires de Matos, B. Nuber, J. Rebizant und M.L. Ziegler), (448) 119
- Tellurium**
- Struktur und Reaktivität von Ytterbiumbenzamidinaten (M. Wedler, A. Recknagel, J.W. Gilje, M. Nottemeyer and F.T. Edelmann), (426) 295
- Structure and dynamics of $[\text{C}_5\text{H}_5\text{Fe}(\text{CO})_2(\text{E}(\text{CH}_3)_2)]\text{BF}_4$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$) complexes studied by ^{13}C solid state NMR spectroscopy (A.P.M. Kentgens, H. Karrenbeld, E. De Boer and H. Schumann), (429) 99
- FT Raman spectroscopy—the preferred technique for routine vibrational spectroscopic studies of organotellurium compounds (M.R. Greaves, W.R. McWhinnie, S.L.W. McWhinnie and D.N. Waters), (430) 37
- Homolytic reactions of diorganotellurium and diorganoditellurium compounds in solution; an EPR study (W. Bell, D.J. Cole-Hamilton, P.N. Culshaw, A.E.D. McQueen, D.V. Shenai-Khatkhate, J.C. Walton and J.E. Hails), (430) 43
- Ferrocenhaltige Liganden in der Lanthanid-Chemie: Alkoxide und Thiolate (H. Gornitzka, F.T. Edelmann und K. Jacob), (436) 325
- Diferrocenyldichalkogenide: Synthese und Struktur von Bis[2-(dimethylaminomethyl)ferrocenyl]diselenid (H. Gornitzka, S. Besser, R. Herbst-Irmer, U. Kilimann, F.T. Edelmann und K. Jacob), (437) 299
- Synthesis and characterization of dinuclear palladium(II) complexes containing both chloro and organochalcogenides as bridging ligands (V.K. Jain and S. Kannan), (439) 231
- Trigonal-planar koordiniertes Selen bzw. Tellur in $[\text{Cp}'(\text{CO})_2\text{Mn}]_2\text{EMes}^+$ ($\text{E} = \text{Se}, \text{Te}$): Verbindungen vom "Iniden"-Typ (P. Lau, G. Huttner und L. Zsolnai), (440) 41
- Crystal and molecular structure of bis(μ_3 -tellurido)-decacarbonyltri-iron, $[\text{Fe}_3(\text{CO})_{10}\text{Te}_2]$ (G. Gervasio), (441) 271
- Umsetzungen der lithiierten Halbsandwich-Chalkogenole, $(\text{CO})_3\text{MC}_5\text{H}_4\text{--ELi}$ ($\text{M} = \text{Mn, Re}; \text{E} = \text{S}, \text{Se}, \text{Te}$) (M. Herberhold und M. Biersack), (443) 1
- Synthesis and characterization of new organotellurium compounds containing an *ortho*-amino group (A.Z. Al-Rubaie, N.I. Al-Salim and S.A.N. Al-Jadaan), (443) 67
- The redox behaviour of ferrocene derivatives. I. 1,2,3-Tritellura[3]ferrocenophane and 1,3-ditellura[3]ferrocenophanes (P. Zanello, G. Opronolla, M. Casarin, M. Herberhold and P. Leitner), (443) 199
- Stereochemistry of the compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$. Crystal and molecular structures of the mixed triiron carbonyl compounds $[\text{Fe}_3(\text{CO})_9\text{XY}]$ ($\text{X}, \text{Y} = \text{S}, \text{Se}, \text{or Te}$) (G. Gervasio), (445) 147
- Photochemisch induzierte Addition von Alkinen und Phosphaalkinen an die Cluster $\text{Fe}_3(\text{CO})_9(\mu_3\text{-PR})(\mu_3\text{-X})$ ($\text{X} = \text{Se}, \text{Te}; \text{R} = ^i\text{Pr}, ^t\text{Bu}$) (W. Imhof und G. Huttner), (447) 31
- Evidence for a free radical mechanism in the decomposition of bis(but-2-enyl)tellurium (J. Stevenson, W. Bell, J. Ferry, D.J. Cole-Hamilton and J.E. Hails), (449) 141

The use of phosphine chalcogenides in the preparation of cobalt chalcogenides (S.M. Stuczynski, Y.-U. Kwon and M.L. Steigerwald), (449) 167

Terbium

Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXXIII. Esteraddukte des Grundkörpers Tris(η^5 -cyclopentadienyl)-lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Metallorganische Verbindungen der Lanthanoide. LXXIV. Tris(tetramethylcyclopentadienyl)komplexe der Lanthanoide (H. Schumann, M. Glanz und H. Hemling), (445) C1

Thallium

Metallorganische Verbindungen der Lanthanoide. LXXI. Synthese und Röntgenstrukturanalyse von CpLa(C₅H₄PPPh₂)₂(THF), einem zweizähnigen Liganden zur Herstellung heterobimetallischer Lanthanoidkomplexe (H. Schumann, J.A. Meese-Märtscheffel, B. Gorella and F.H. Görslitz), (428) C27

Substituted cyclopentadienyl compounds. I. Formylation of methylcyclopentadienide and the synthesis and NMR study of some thallium(I) and rhodium(I) derivatives of cyano-, methyl-, methanoyl- and dimethylamido cyclopentadienes (M. Arthurs, J.C. Bickerton, M. Kirkley, J. Palin and C. Piper), (429) 245

Dimethyl(diorganophosphinodithioformato-*S,S'*)-thallium(III) compounds. X-Ray structure of [Ti(CH₃)₂S₂CP(C₆H₅)₂](THF)] (E.M. Vázquez-López, A. Sánchez, J.S. Casas, J. Sordo and E.E. Castellano), (438) 29

A facile synthetic route to (η^5 -benzoylcyclopentadienyl)metal compounds (S.S. Jones, M.D. Rausch and T.E. Bitterwolf), (450) 27

Molecular structure of *meso*-tetraphenylporphyrinatothallium(III) cyanide: Ti(tpy)(CN) (W.-B. Lee, S.-C. Suen, T.-T. Jong, F.-E. Hong, J.-H. Chen, H.-J. Lin and L.-P. Hwang), (450) 63

Thorium

Organo-*f*-element thermochemistry. Thorium-ligand bond disruption enthalpies in [(CH₃)₃SiC₆H₆)₂]ThX (X = H or D) and in [(CH₃)₃SiC₅H₄)₂]ThH complexes (X. Jemine, J. Goffart, M. Ephritikhine and J. Fuger), (448) 95

Thulium

Zur Elektronenstruktur metallorganischer Komplexe der *f*-Elemente. XXXIII. Esteraddukte des Grundkörpers Tris(η^5 -cyclopentadienyl)-lanthan(III)—ideale Wirtsgitter für einkristalloptische Untersuchungen von Lanthanoid-Ionen in metallorganischer Umgebung (H. Schulz und H.-D. Amberger), (443) 71

Tin

Alkinkomplexe des Titanocens und Permethyltitanocens ohne zusätzliche Liganden — Erste Strukturvergleiche (U. Rosenthal, H. Görls, V.V. Burlakov, V.B. Shur and M.E. Vol'pin), (426) C53

Tetra-*p*-tolyl-Verbindungen *p*-Tol₄Si und *p*-Tol₄Ge: ein Beitrag zur Konfiguration der Tetraaryl-Methan-Analoga Ar₄M (M = C, Si, Ge, Sn, Pb) (M. Charissé, S. Roller and M. Dräger), (427) 23

Übergangsmetall-Silyl-Komplexe. XLII. Einfluß des Phosphan-Liganden auf Bildung, Struktur und Stabilität der Hetero-Zweikernkomplexe (CO)₃(R'₃P)(R₃Si)Fe—ML_n (M = Cu, Ag, Au, Hg) (G. Reinhard, B. Hirle and U. Schubert), (427) 173

Crystal structure of [Rh(SnCl₃)₂(norbornadiene)(dppp)]. A ¹H NMR study of [Rh(SnCl₃)₂(norbornadiene)(diphosphine)] complexes (M.A. Garralda, E. Pinilla and M.A. Monge), (427) 193

Substitution of the acetoxy groups of dialkoxyacetates by organometallic reagents: a route to allyl-, propargyl-, homoallyl-, homopropargyl- and α -stannylacetals (I. Beaudet, A. Duchêne, J.-L. Parrain and J.-P. Quintard), (427) 201

The structure of ('Bu₂Sn)₂CO₃ and (Me₃Sn)₂CO₃ in solution and in the solid state studied by ¹³C/¹¹⁹Sn NMR spectroscopy and X-ray diffraction (J. Kümmelen, A. Sebald and H. Reuter), (427) 309

Synthesis and spectral characterization of medium ring distannacycloalkanes and their Lewis acid derivatives (D. Farah, K. Swami and H.G. Kuivila), (429) 311

Structural chemistry of organotin carboxylates. XVII. Diorganotin (IV) derivatives of *N*-phthaloyl-DL-valine (G.K. Sandhu, R. Hundal and E.R.T. Tiekkink), (430) 15

Organotin mediated cycloaddition reactions: a re-investigation of the reaction between organotin azides and isothiocyanates (R.J. Deeth, K.C. Molloy, M.F. Mahon and S. Whittaker), (430) 25

Dicyclohexylammonium (dicarboxylato)triorganostannates. Crystal structure of bis(dicyclohexylammonium) tris(malonato)tetrakis (tributylstannate) (S.W. Ng, V.G. Kumar Das, B.W. Skelton and A.H. White), (430) 139

Synthesis and spectroscopic studies of thienyl triorganotin(IV) compounds. Crystal structures of [2-(4,4-dimethyl-2-oxazolinyl)-3-thienyl]dimethyltin *N,N*-dimethylthiocarbamate and [2-(4,4-dimethyl-2-oxazolinyl)-3-thienyl]diphenyltin chloride (K.M. Lo, S. Selvaratnam, S.W. Ng, C. Wei and V.G. Kumar Das), (430) 149

Molecular structure of 1,2;3,4,di- μ -nitrobenzoato-OO'-1,3-bis(*o,m*, or *p*-nitrobenzoato)-1,2,4;2,3,4-di- μ_3 -oxotetrakis [di-*n*-butyltin(IV)] compounds (S.P. Narula, S.K. Bharadwaj, Y. Sharda, D.C. Povey and G.W. Smith), (430) 167

The effect of ligands, solvent and temperature on the reactions of allyltin(IV) compounds with singlet oxygen (H.-S. Dang and A.G. Davies), (430) 287

Addition of sulphenyl halides and thiocyanates to alkenyl-metal compounds. Structures of [1-chloro-2-(4-methyl-2-nitrophenyl)ethyl]-triphenylstannane and [2-(2-nitrophenyl)-1-thiocyanato-ethyl]triphenylstannane (R.A. Howie, J.L. Wardell, E. Zanetti, P.J. Cox and S.M.S.V. Dodge-Harrison), (431) 27

Si-, Ge- and Sn-diine derivatives in 1,3-dipolar cycloaddition reactions (A.D. Jurayev, A.G. Makhsumov, K. Yuldasheva and M.A. Tolipova), (431) 129

Structural chemistry of organotin carboxylates. XVIII. Synthesis, characterization and X-ray crystal structure of bis(μ -hydroxo)-bis[di-*t*-butyltin(IV)acetate]: [¹Bu₂Sn(O₂CCH₃)₂(μ -OH)]₂ (V.B. Mokal, V.K. Jain and E.R.T. Tiekkink), (431) 283

Darstellung und spektroskopische Eigenschaften der Halogen-(trifluormethyl)stannane, (CF₃)_nSnX_{4-n} (R. Eujen und U. Thurmann), (433) 63

Multinuclear NMR spectroscopic studies of some organometallic N-substituted anilines (B.K. Barr, A.J. Herman, L.K. Myers, P.I. Young, C.D. Schaeffer, Jr., H.J. Eppley, J.C. Otter and C.H. Yoder), (434) 45

Reactivity of main-group-transition-metal bonds. X. The kinetics of iodination of compounds containing one, two, three or four tin–cobalt bonds (J.R. Chipperfield, D.E. Webster and H. Yusof), (434) 53

3,5-Di-*tert*-butylthiobenzoyltriphenylsilane: a versatile spin trapping agent (A. Alberti and M. Benaglia), (434) 151

Darstellung und Eigenschaften trifluormethylsubstituierter Stannane, (CF₃)_nSnH_{4-n} (*n* = 1–3) (R. Eujen, N. Jahn und U. Thurmann), (434) 159

Übergangsmetall-Stannyly-Komplexe. V. Darstellung der anionischen Stannyly-Komplexe K[(π -Aromat)(CO)₂CrSnPh₃] und Reaktion mit Organozinnhalogeniden zu Bis(stannyly)-Komplexen (π -

- Aromat)(CO)₂Cr(SnPh₃)(SnR₃) (π -Aromat = C₆H_{6-n}Me_n) (U. Schubert und J. Schubert), (434) 169
- Organotin compounds with 2,2'-biimidazole derivatives. The crystal structure of dibromo(*N,N'*-dimethyl-2,2'-biimidazole)dimeethyltin(IV) (C. López, A. Sánchez González, M.E. García, J.S. Casas, J. Sordo, R. Graziani and U. Casellato), (434) 261
- Stereoselective hydrostannation: synthesis and absolute configuration of (-)-methyl 2,3-diphenyl-3-(trimethylstannylyl) propanoates and derivatives (J.C. Podestá, A.B. Chopra, L.C. Koll and S.D. Mandolesi), (434) 269
- Complexes of dibutyltin(IV) with imidazole and pyrazole. The crystal structure of dibutylchlorobis(pyrazole)tin(IV) (A. Sánchez González, J.S. Casas, J. Sordo and G. Valle), (435) 29
- Sterically hindered organotin compounds. II. Synthesis of the organotin halides (mesityl)₂SnX_nY_{2-n}, (mesityl)SnX_nY_{3-n} and R₄Sn₂X₂ (R = 2,4,6-ⁱPr₃C₆H₂, X = Cl, Y = Br). Crystal structure of the ditin species (R₂BrSn)₂ (P. Brown, M.F. Mahon and K.C. Molloy), (435) 265
- Main group metal 1,3-dithiole-2-thione-4,5-dithiolate (DMIT) compounds. II. Synthesis and structures of tetrabutylammonium butyl- and phenyl-bis(1,3-dithiole-2-thione-4,5-dithiolato)stannates, [Bu₄N]RSn(DMIT)₂] (R = Bu or Ph) (S.M.S.V. Dodge-Harrison, R.A. Howie, J.T.S. Irvine, G.M. Spencer and J.L. Wardell), (436) 23
- A Mössbauer study of tin(IV) and organotin(IV) adducts of bis(pyrazolyl)methane-type ligands (G. Gioia Lobbia, A. Cingolani, P. Cecchi, S. Calogero and F.E. Wagner), (436) 35
- Some triphenyltin(IV) complexes containing potentially bidentate, biologically active anionic groups (B.D. James, S. Gioskos, S. Chandra, R.J. Magee and J.D. Cashion), (436) 155
- Preparation, IR, ¹³C and ¹¹⁹Sn Mössbauer spectral studies of organotin(IV) derivatives of ethylenediaminetetraacetic acid (G.K. Sandhu and R. Hundal), (436) 287
- Synthesis and structure of organotin complexes of ytterbium (L.N. Bochkarev, O.V. Grachev, S.F. Ziltsov, L.N. Zakharov and Y.T. Struchkov), (436) 299
- Zur Funktionalisierung von Decaphenylcyclopentasilan (W. Uhlig und C. Tretner), (436) C1
- Anchimerically assisted cleavage of a one-electron bond in the [Me₂N(CH₂)₃SnⁿBu₃]⁺ radical cation (E. Butcher, C.J. Rhodes, R.S. Davidson and R. Bowser), (436) C5
- Use of organotin halides as catalytic precursors in dehydration processes (G. Tagliavini), (437) 15
- Tin for organic synthesis. VI. The new role of organotin reagents in organic synthesis Part 1: Stannyl groups as leaving groups in electrophilic substitutions Part 2: Polymer-supported organotin reagents for organic synthesis (W.P. Neumann), (437) 23
- The first synthesis of 1,1,2-tristannylalkenes (T.N. Mitchell and B. Kowall), (437) 127
- Preparation, spectroscopic studies and structure of bis(triorganostannyl) esters of substituted aliphatic dicarboxylic acids (A. Samuel-Lewis, P.J. Smith, J.H. Aupers, D. Hampson and D.C. Povey), (437) 131
- FT-IR study of the gas phase thermal decomposition of dimethyldivinyltin in the presence of oxygen (P.G. Harrison and E.N. Clark), (437) 145
- Polysulfonylamine. XXXV. Synthese, Röntgenstrukturanalysen und hochauflöste Festkörper-NMR-Spektren der ionischen Organozinn(IV)-dimesylamide [Me₂Sn(NH₃)₂]IN(SO₂Me)₂] und [Me₂Sn(DMSO)₄]IN(SO₂Me)₂ (A. Blaschette, I. Hippel, J. Krahl, E. Wieland, P.G. Jones und A. Sebald), (437) 279
- Steric and metal-centre control of two-, three- and four-orbital carborane–metal interaction in tin and platinum complexes of the (C₂B₈H₉X) ligand, where X = H and Ph (J.D. Kennedy, K. Nestor, B. Štíbr, M. Thornton-Pett and G.S.A. Zammit), (437) C1
- Synthesis of acylsilanes via transmetalation of 1-triorganosilyloxyvinyltin derivatives (J.-B. Verlhac, H.A. Kwon and M. Pereyre), (437) C13
- Stereoselective opening of chiral α -stannylacetals with organometallic reagents (J.-L. Parrain, J.-C. Cintrat and J.-P. Quintard), (437) C19
- Organoerdmetall-Komplexe von *d*-Block-Elementen. V. Selektive Reduktion von Bis[carbonyl(η^5 -cyclopentadienyl)nickel] zum Carbonyl(η^5 -cyclopentadienyl)nickel-Anion durch C₈K. Synthese und Struktur von (η^5 -C₅H₅)(CO)Ni-Ga[CH₂C(CH₃)₃]₂(OC₄H₈) (R.A. Fischer, J. Behm, E. Herdtweck und C. Kronseder), (437) C29
- Synthesis and coordination chemistry of itaconic esteritin(IV) complexes (O.-S. Jung, J.H. Jeong and Y.S. Sohn), (439) 23
- The palladium-catalysed addition of hexamethylditin to 1,3-dienes (T.N. Mitchell, B. Kowall, H. Killing and C. Nettelbeck), (439) 101
- Some further studies on acyltrimethylstannanes (T.N. Mitchell and K. Kwetkat), (439) 127
- Complex salts derived from the reactions of Me₂SnCl₂ with some aromatic Schiff bases. Crystal structure of bis[N-(2-methoxy-1-naphthalidene)-3-methoxyanilinium]hexachlorotetramethyldistannate(IV)dihydrate: [CH₃OC₁₀H₆CH:NHC₆H₄OCH₃]₂[(CH₃)₂Sn₂Cl₆]₂H₂O]₂ (S.-G. Teoh, S.-B. Teo, G.-Y. Yeap and H.-K. Fun), (439) 139
- Mössbauer and NMR study of tin(IV) and organotin(IV) hydridotris(3-Me-1*H*-pyrazol-1-yl)borates. The X-ray crystal structure of phenyldichlorotin(IV) derivative (G. Gioia Lobbia, S. Calogero, B. Bovio and P. Cecchi), (440) 27
- Steric effects on the formation of isolable products in the reactions of dibutyltin oxides with carboxylic acids (V.B. Mokal and V.K. Jain), (441) 215
- Synthesis of stanna- and germa-cycloheptatrienes (Y. Nakadaira, R. Sato and H. Sakurai), (441) 411
- Insertions- und Immintransfer-Reaktionen mit Schwefel und N-stannylierten Schwefelimidinen (D. Hänsgen, H. Salz, S. Rheindorf und C. Schrage), (443) 61
- Sila-Riechstoffe und Riechstoffisostere. XII. Geruchsvergleiche homologer Organoelementverbindungen der vierten Hauptgruppe (C, Si, Ge, Sn) (U. Wannagat, V. Damrath, V. Huch, M. Veith und U. Harder), (443) 153
- Selectivity of mercury(II) salts in reactions with α,β -unsaturated stannyl esters (C. Deb and B. Basu), (443) C24
- Novel polynuclear organotin complexes of samarium and ytterbium (L.N. Bochkarev, O.V. Grachev, N.E. Molosnova, S.F. Ziltsov, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), (443) C26
- A tetranuclear β -diketonato complex of organotin(IV): synthesis and crystal structure of (benzoyltrifluoroacetonato)-(trifluoroacetato)tetramethyldistannoxane dimer [(C₆H₅C(O)CHCO(CF₃)Sn(CH₃)₂O(Sn(CH₃)₂O₂CCF₃)₂] (B.K. Agarwal, Y.P. Singh, R. Bohra, G. Srivastava and A.K. Rai), (444) 47
- Additionsreaktionen von (CO)₅CrEPhH₂ (E = P, As) an Di(ethinyl)-Phosphane und Stannane; Stufenweise Darstellung von Komplex-gebundenen 1,4-Dihetero-cyclohexa-2,5-dienen (H. Lang, U. Lay und W. Imhof), (444) 53
- Subvalent Group 14 metal compounds. XV. Germanium and tin amides M(NR₂)₂ as ligands in carbonylchromium(0) chemistry. The crystal structure of the compounds *cis*-[Cr(CO)₄{Sn(NR₂)₂}₂] [M = Ge or Sn, R' = SiMe₃ or NR₂ = NCMe₂(CH₂)₃CMe₂] (S.L. Ellis, P.B. Hitchcock, S.A. Holmes, M.F. Lappert and M.J. Slade), (444) 95
- Un nouvel hydrure organostannique greffé sur support insoluble (G. Ruel, N.K. The, G. Dumartin, B. Delmond et M. Pereyre), (444) C18

- Polysulfonylamine.** XXXVIII. Darstellung und Festkörperstruktur von Bis(trimethylzinn)-dimesylamid-hydroxid $[Me_3Sn(\mu-(MeSO_2)_2N)Me_3Sn(\mu-OH)]_n$ (A. Blaschette, E. Wieland, P.G. Jones und I. Hippel), (445) 55
- Zur Struktur sterisch gehinderter Zinn-Elementverbindungen: Kristallstruktur der Zinn-Schwefel-Stickstoff-Ringe $[(^tBu_2Sn)_2N_2S_2]$ und $[(^tBu_2Sn)N_2S_2]_2$ (D. Hänssgen, M. Jansen, W. Assenmacher und H. Salz), (445) 61
- 1:1 Complexes of dimethyltin dihalides with tris(2,6-dimethoxyphenyl)-phosphine oxide, its sulphide, and related compounds (M. Wada, T. Fujii, S. Iijima, S. Hayase, T. Erabi and G.-e. Matsabayashi), (445) 65
- Preparation and crystal structures of the crowded triorganotin isothiocyanate $(PhMe_2Si)_3CSnMe_2NCS$ and nitrate $(PhMe_2Si)_3CSnMe_2ONO_2$ (S.S. Al-Juaid, M. Al-Rawi, C. Eaborn, P.B. Hitchcock and J.D. Smith), (446) 161
- The first authenticated example of triorganotin(IV) perchlorate complexes of some heterocyclic bases. I. Synthesis, characterization and assessment of bonding and structure (T.S. Basu Baul, D. Dey, D.D. Mishra, W.L. Basaiawmoit and E. Rivarola), (447) 9
- Synthesis and molecular structure of $^nBu_2(MeO)_3C_6H_2C(O)N_2CHC_6H_4O]Sn$ formed in the reaction of di-n-butyltin(IV) oxide with 3,4,5-trimethoxy-benzoyl salicylhydrazone (Z.-K. Yu, S.-H. Wang, Z.-Y. Yang, X.-M. Liu and N.-H. Hu), (447) 189
- Über gemischte Gruppe 14-Gruppe 14-Bindungen. VI. Hexa-*o*-tolylethan-Analoga *o*-Tol₆Sn₂, *o*-Tol₆PbSn und *o*-Tol₆Pb₂: ein Vergleich von Bindungsstärke und Polarität in der Reihung Sn-Sn, Pb-Sn, Pb-Pb (C. Schneider-Koglin, K. Behrends und M. Dräger), (448) 29
- Über gemischte Gruppe 14-Gruppe 14-Bindungen. VII. Ein Inkrementensystem für ¹³C-NMR chemische Verschiebungen bei arylierten Gruppe 14-Verbindungen (C. Schneider-Koglin, B. Mathiasch und M. Dräger), (448) 39
- Über Tetraaryl-Methan-Analoga in der Gruppe 14. II. Ph_{4-n}Si-*p*-Tol_n und Ph_{4-n}Sn-*p*-Tol_n (*n* = 0 bis 4): Einfluß der *p*-Tolyl-Substituenten auf Struktur und NMR-chemische Verschiebungen (M. Charissé, V. Gauthey und M. Dräger), (448) 47
- The structure and Lewis acidity of some triorganotin carboxylates (C.H. Yoder, R.A. Morreall, C.I. Butoi, W.J. Kowalski and J.N. Spencer), (448) 59
- Polysulfonylamine.** XXXIX. Charakterisierung des Kations $[Me_2Sn(H_2O)_4]^{2+}$ im kristallinen Zustand: Bildung und Röntgenstrukturanalyse von Tetraaquadimethylzinn(IV)-bis(1,1,3,3-tetraoxo-1,3,2-benzodithiazolid) (I. Hippel, P.G. Jones und A. Blaschette), (448) 63
- Synthesis of α -stannyli α , β -unsaturated carboxylic esters (A.J. Zapata, C. Fortoul R. and C. Acuña A.), (448) 69
- Organotin diphenylmonothiophosphinates. Crystal structure of bis(diphenylmonothiophosphinato)dimethyltin(IV), $Me_2Sn(OS-PPh_2)_2$ (C. Silvestru, I. Haiduc, F. Caruso, M. Rossi, B. Mahieu and M. Gielen), (448) 75
- Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors $M^{II}M^{IV}P_2$ where M^{II} = Zn and Cd, and M^{IV} = Ge and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conrad), (449) 9
- Synthesis and thermal decomposition of substituted [2-(acyloxy)alkyl]diorganotins compounds, $Bu_2Sn(X)CH_2CHR^1OCOR^2$ (X = halogen, 2,4-pentanedionate, or OCOR), potential sources of organotin catalysts (B. Joussemae, V. Guillou, N. Noiret, M. Pereyre and J.-M. Francès), (450) 97
- Titanium**
- Darstellung und Reaktionen von Alkyl-Hydridotrispyrazolylborat-Komplexen des Titans (J. Ipaktschi and W. Sulzbach), (426) 59
- Direct proof of the molecular structure of dimeric titanocene; the X-ray structure of $\mu-(\eta^5:\eta^5\text{-fulvalene})\text{-di-(}\mu\text{-hydrido)\text{-bis}(\eta^5\text{-cyclopentadienyltitanium)} \cdot 1.5\text{benzene}$ (S.I. Troyanov, H. Antropiusová and K. Mach), (427) 49
- Conversion of hydrosilanes to alkoxy silanes catalyzed by $Cp_2TiCl_2/nBuLi$ (T.C. Bedard and J.Y. Corey), (428) 315
- The crystal structure of $(\eta^6-C_6Me_6)Ti[(\mu-Cl)_2(AlClEt)_2]$ and the catalytic activity of the $(C_6Me_6)TiAl_2Cl_{8-x}Et_x$ (x = 0–4) complexes towards butadiene (S.I. Troyanov, J. Poláček, H. Antropiusová and K. Mach), (430) 317
- Die Struktur von $[Cp_2Ti^{III}(CN)]_4$ (P. Schinnerling and U. Thewalt), (431) 41
- Dehydrierende Polymerisation von Methyldisilanen mittels Metallocen-Katalyse zu den Oligomeren $Me_3Si[Me_2Si]_nSiMe_2H$, $HMe_2Si[Me_2Si]_nSiMe_2H$ mit n = 3–9, $H[MeSiH]_nH$ mit n = 3–14 und hochpolymerem Poly(methylsilan) ($MeSiH_x$)_n (E. Hengge und M. Weinberger), (433) 21
- Controlled synthesis of $(1,3-^tBu_2-C_5H_3)_2TiCl_2$ and $(1,3-^tBu_2-C_5H_3)_2TiCl_3$. Crystal structure and reactions of $(1,3-^tBu_2-C_5H_3)_2TiCl_3$ (I. Jibril, S. Abu-Orabi, S.A. Klaib, W. Imhof and G. Hüttnér), (433) 253
- Base-free cationic 14-electron alkyls of Ti, Zr and Hf as polymerisation catalysts: a comparison (M. Bochmann and S.J. Lancaster), (434) C1
- ansa*-Metallocene derivatives of Ti^{IV} and Zr^{IV} with the shortest $-C(CH_3)_2-$ bridge (I.E. Nifant'ev, A.V. Churakov, I.F. Urazowski, S.G. Mkoyan and L.O. Atovmyan), (435) 37
- Synthesis and characterization of Group IV metal–iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $Cp_2Ti[(\mu-O_2CC_5H_4)Fe(CO)_2(CH_2C_6H_5)]_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43
- Titanium-catalyzed cycloaddition reactions of phenyl(trimethylsilyl)acetylene to conjugated dienes and 1,3,5-cycloheptatriene. 1-Phenyl-2-(trimethylsilyl)-cyclohexa-1,4-dienes and their aromatization (R. Klein, P. Sedmera, J. Čejka and K. Mach), (436) 143
- Zum Koordinationsverhalten der Lewis-Basen HCN und DCN gegenüber $Cp_2Ti(AsF_6)_2$ (A. Schulz und T.M. Klapötke), (436) 179
- Condensation of primary silanes in the presence of $Cp_2MCl_2/nBuLi$ (M = Ti, Zr, Hf) (J.Y. Corey and X.-H. Zhu), (439) 1
- Catalytic activity of the titanocene complex with tolane in homogeneous hydrogenation of unsaturated hydrocarbons (V.B. Shur, V.V. Burlakov and M.E. Vol'pin), (439) 303
- Die Bildung von $[Cp_2Ti(C\equiv CSiMe_3)]_2$ aus Titanocen und 1,4-Bis(trimethylsilyl)-1,3-Butadiin (U. Rosenthal und H. Görls), (439) C36
- Heterozyklen als Liganden. XVI. $(2,5-C_4-^tBu_2RHN)MCl_3$ (M = Ti, Zr, Hf; R = H, SiMe₃)—Azacyclopentadienyl-Komplexe der Gruppe 4-Metalle (N. Kuhn, S. Stubenrauch, R. Boese und D. Bläser), (440) 289
- Dehydrierende Copolymerisation zu den neuen Polysilan-Copolymeren Poly(methyl-co-phenylsilan) $H[(MeSiH)_x(PhSiH)_y]_nH$ und Poly(methyl-co-dimethylsilan) $H[(MeSiH)_x(Me_2Si)_y]_nH$ (E. Hengge und M. Weinberger), (441) 397
- Reactions of $TiCl_4$ with diesters. Crystal structures of $[CH_2(CO_2Et)_2Cl_4Ti]$ and $[C_2H_4(CO_2CH_2CH_2OPh)_2Cl_4Ti]$ (P. Sobota, S. Szafert and T. Lis), (443) 85
- Benzylverbindungen elektronenärmer Übergangsmetalle. Molekülstrukturen von $Cp_2Ti(CH_2Ph)_2$ und $CpZr(CH_2Ph)_3$ (J. Scholz, F. Rehbaum, K.-H. Thiele, R. Goddard, P. Betz und C. Krüger), (443) 93
- Mechanistische Aspekte zur übergangsmetall-katalysierten dehydrierenden Polymerisation von Disilanen und höheren Silanen (E. Hengge und M. Weinberger), (443) 167
- Carbonyl adducts of open titanocenes – syntheses and structures (Y. Wang, Y.-Q. Liang, Y.-J. Liu and J.-Z. Liu), (443) 185

Photochemistry of $\text{Cp}_2\text{Ti}(\text{CO})_2$ and $\text{Cp}_2\text{Ti}(\text{N}_3)_2$ in solid matrices and in solution at low temperatures (M. Tacke, C. Klein, D.J. Stufkens and A. Oskam), (444) 75

Cyclische Metallamide. III. Flüchtige Tetrahydroborat-Komplexe von Titan und Zirkonium (W.A. Herrmann, M. Denk, W. Scherer und F.-R. Klingan), (444) C21

Ketene acetal metallocene complexes. Crystal structure of $(\text{C}_5\text{H}_5)_2\text{Ti}(\text{Cl})\text{OC}(\text{OCH}_3)=\text{C}(\text{CH}_3)_2$ (K. Hortmann, J. Diebold and H.-H. Brintzinger), (445) 107

Ti-Verbindungen mit Fulvalenbrücken. Die Struktur von $[\text{CpTiBr}]_2$ ($\text{C}_{10}\text{H}_8\text{O}$) Toluol (T. Wöhrel und U. Thewalt), (447) 45

Fulvalene cyclopentadienyl titanium and zirconium(III) and -(IV) complexes. X-Ray crystal structure of $[(\text{Ti}(\eta^5\text{-C}_5\text{H}_5)\text{Cl})_2(\mu\text{-O})(\mu\text{-}\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)]$ (A. Cano, T. Cuenca, G. Rodriguez, P. Royo, C. Cardin and D.J. Wilcock), (447) 51

Synthesis and structure of polymeric $[1,3,5\text{-C}_6\text{H}_3(\text{CO}_2\text{Me})_3 \cdot 3\text{TiCl}_4]$ and dimeric $[(\mu\text{-1,3,5-C}_6\text{H}_3(\text{CO}_2\text{Me})_3)_2\text{Cl}_8\text{Ti}_2] \cdot 2\text{CH}_2\text{Cl}_2$ compounds (P. Sobota, J. Utko and T. Lis), (447) 213

Crystal and molecular structure of bis(tetramethylcyclopentadienyl)titanium halides, $(\text{C}_5\text{HMe}_4)_2\text{TiCl}$, $(\text{C}_5\text{HMe}_4)_2\text{TiI}$ and $(\text{C}_5\text{HMe}_4)_2\text{TiCl}_2$ (S.I. Troyanov, V.B. Rybakov, U. Thewalt, V. Varga and K. Mach), (447) 221

Isodicyclopentadienes and related molecules. LVII. Solid state structural studies of the diastereomeric *exo*- and *endo*- $(\eta^5\text{-cyclopentadienyl})\chi\eta^5\text{-}(1S,8R)\text{-9,9-dimethyltricyclo[6.1.1.0}^{2,6}\text{]deca-2,5-dienyl})$ dichlorotitaniums and a stereopure *exo*- $(\eta^5\text{-penta-methyl-cyclopentadienyl})$ congener (R.D. Rogers, M.R. Sivik and L.A. Paquette), (450) 125

Tungsten

Zur Addition von Schwefel und Selen an Dicarbonyl($\eta^5\text{-cyclopentadienyl})\chi\eta^2\text{-}(methylthio)methylcarben]wolfram-tetrafluoroborat (F.R. Kreißl and N. Ullrich), (426) C8$

Electrochemistry and bonding in biscyclopentadienyl complexes of molybdenum, tungsten, and niobium with *o*-phenanthroline and bipyridine. Crystal and molecular structures of $[\text{Nb}(\eta^5\text{-C}_5\text{H}_5)_2(\text{bpy})]\text{PF}_6$ and $[\text{M}(\eta^5\text{-C}_5\text{H}_5)_2(\text{o-phen})]\text{PF}_6$ ($\text{M} = \text{Mo}$ or W) (M.J. Calhorda, M.A.A.F. de C.T. Carrondo, A. Bram, P.N. Olsen, A.R. Dias, A.M. Freitas, M.H. Garcia and M.F.M. Piedade), (426) 195

Synthesis and evaluation of the catalytic properties of homo- and hetero-bimetallic complexes containing bridging diphenylphosphido ligands (Z. He, N. Lusan, D. Neibecker, R. Mathieu and J.-J. Bonnet), (426) 247

A study of polarities, polarisabilities and infrared spectra of $\text{LM}(\text{CO})_5$ complexes ($\text{M} = \text{Cr, Mo or W; L} = \text{trimethylamine, quinuclidine, pyridine, } p\text{-methyl- or } p\text{-t-butylpyridine}$) and of the X-ray crystal structure of quinuclidine-Cr(CO)₅: evidence for π -acceptor behaviour of coordinated pyridine (M.J. Aroney, R.M. Clarkson, T.W. Hambley and R.K. Pierens), (426) 331

Metallorganische Lewis-Säuren. XLIV. Rhenoxy(organyl)carben-Komplexe von Chrom, Molybdän und Wolfram (P.M. Fritz, J. Breimair, B. Wagner and W. Beck), (426) 343

Metathesis of 2-pentene with the $\text{W}(\text{CO})_6/\text{CCl}_4/\text{hv}$ -system. Role and physical properties of the photogenerated precipitate (P.G.M. Schilder, D.J. Stufkens, A. Oskam and J.C. Mol), (426) 351

Reaktionen von Komplexliganden. XLVIII. Carben-Liganden in intramolekularen Pauson-Khand-Reaktionen (K.H. Dötz and J. Christoffers), (426) C58

Kinetische und mechanistische Untersuchungen von Übergangsmetall-Komplex-Reaktionen. XXVI. Kinetische Untersuchungen zur Cyclopropanierung von Olefinen mit Benzyliden(pentacarbonyl)wolfram-Komplexen (H. Fischer, E. Mauz, M. Jaeger and R. Fischer), (427) 63

Syntheses of $\text{W}(\text{PhC}\equiv\text{CPh})_3(\eta^1\text{-PPPh}_2\text{CH}_2\text{PPh}_2)$ and $\text{W}(\text{PhC}\equiv\text{CPh})_3(\text{PPh}_3)$ from $\text{W}(\text{PhC}\equiv\text{CPh})_3(\text{NCCH}_3)$ (W.-Y. Yeh, C.-S. Ting and C.-f. Chih), (427) 257

The synthesis and substitution reactions of $\text{CpM}(\text{CO})_2(\mu\text{-RC}\equiv\text{CR})\text{Co}(\text{CO})_3$ ($\text{M} = \text{Mo, W; R} = \text{COOEt, COOMe}$) (C.-A. Dickson and N.J. Coville), (427) 335

Synthese und Kristallstrukturanalyse von Alkinylcarbenkomplexverbrückten Dicobalthexacarbonylkomplexen (A. Wienand, H.-U. Reissig, H. Fischer, D. Pflumm and C. Troll), (427) C9

Reaction of bis(cyclopentadienyl)hydridophenyltungsten(IV) with carboxylic acids, carbon dioxide, and related compounds (T. Ito, S. Sugimoto, T. Ohki, T. Nakano and K. Osakada), (428) 69

Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XXXIX. Synthese und dynamisches Verhalten von Tetracarbonyl- $\eta^4\text{-tricyclo[6.3.0.0}^{2,7}\text{]undeca-3,5-dien-metall(0)}$ - und Dicarbonyl- $\eta^4\text{-tricyclo[6.3.0.0}^{2,7}\text{]undeca-3,5-dien-metall(0)}$ -Komplexen der Elemente Chrom, Molybdän und Wolfram (S. Özkar, N.K. Tunali and C.G. Kreiter), (428) 345

Thermolysis of $\text{Cp}(\text{CO})_n\text{M}-\text{CH}(\text{CH}_3)\text{C}_6\text{H}_5$ ($n = 2, \text{M} = \text{Fe, Ru}; n = 3, \text{M} = \text{Mo, W}$). Some vertical trends in transition metal-alkyl bond dissociation energies (C. Mancuso and J. Halpern), (428) C8

Organometallic complexes of the diphosphene $[2,4,6\text{-}(\text{CF}_3)_3\text{C}_6\text{H}_2]_2\text{P}_2$ (K.B. Dillon and H.P. Goodwin), (429) 169

Tris($\eta^4\text{-1-oxa-1,3-diene})$ molybdenum and -tungsten: Preparation and properties of new homoleptic enone complexes (T. Schmidt and S. Neis), (430) C5

Reactivity of dinuclear tungsten carbonyls bridged both by hydride and by bidentate phosphine ligands, $(\mu\text{-H})(\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2)\text{W}_2(\text{CO})_7(\text{NO})$ ($n = 1\text{-}3$) (J.T. Lin, K.-Y. Chang, M.L. Gong, C.-C. Chang, H.-M. Gau, L.-H. Tsen and Y.S. Wen), (431) 65

Synthese und strukturelle Charakterisierung von bcpm-Metallcarbonyl-Dimeren (bcpm = Biscyclopentadienylmethan) mit Metallen der 6. Nebengruppe (H. Brunner, R. Graßl, J. Wachter, B. Nuber and M.L. Ziegler), (431) 143

Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XL. Synthese und dynamisches Verhalten von Tricarbonyl- $\eta^4\text{-tricyclo[6.3.0.0}^{2,7}\text{]undeca-3,5-dien-trimethylphosphit-metall(0)}$ - und Dicarbonyl- $\eta^4\text{-tricyclo[6.3.0.0}^{2,7}\text{]undeca-3,5-dien-bis(trimethylphosphit)metall(0)}$ -Komplexen der Elemente Chrom, Molybdän und Wolfram (C.G. Kreiter, M. Kotzian, S. Özkar and I. Abu-Mour), (431) 159

Dihydropyridine-induced reductions of methyl(ethoxy)carbene complexes of chromium and tungsten: formation and X-ray crystal structure of the adduct between pyridine and the corresponding ethylidene carbene complexes (F. Cohen, R. Goumont, H. Rudler, J.C. Daran and R.A. Toscano), (431) C6

Thiiran-Komplexe aus Thioketonen und $(\text{CO})_5\text{W}=\text{C}(\text{Ph})\text{H}$ bzw. aus Diphenyldiazomethan und $(\text{CO})_5\text{W}[\text{S}=\text{C}(\text{Ph})\text{H}]$ (H. Fischer, K. Treier und U. Gerbing), (433) 127

Reactions of α -lithio (aminocarbene)tungsten anions with diiodoalkanes: synthesis of $(\mu\text{-bis(aminocarbene)})\text{ditungsten}$ complexes (D.W. Macomber and P. Madhukar), (433) 279

Synthesis of molybdenum-triosmium cluster complexes: reaction of $(\mu\text{-H})_2\text{Os}_3(\text{CO})_{10}$ with $(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\text{Mo}\equiv\text{CC}_6\text{H}_4\text{Me}$ (J.T. Park, J.-J. Cho, K.-M. Chun and S.-S. Yun), (433) 295

Einfluß verschiedenartig substituierter Cyclopentadienylliganden auf die Chemie schwefelreicher Organometallkomplexe mit Metallen der 6. Nebengruppe (H. Brunner, R. Graßl, W. Meier, J. Wachter, B. Nuber und M.L. Ziegler), (434) 63

Gehinderte Ligandbewegungen in Übergangsmetallkomplexen. XLI. Das dynamische Verhalten von 1,2-Bis(dimethylphosphino)ethan-dicarbonyl-($\eta^4\text{-dien})$ wolfram(0) (S. Özkar, C. Kayran und C.G. Kreiter), (434) 79

Mercury-bridged transition-metal clusters. Synthesis of pentanuclear Ru_3HgM ($\text{M} = \text{Mo, W, or Co}$) clusters and X-ray structure of

- [{Ru₃(μ₃,η²-ampy)(CO)₉}- (μ₃-Hg)Co(CO)₄] (Hampy = 2-amino-6-methylpyridine) (P.L. Andreu, J.A. Cabeza, A. Llamazares, V. Riera, S. García-Granda and J.F. Van der Maelen), (434) 123
- Carbyne–carbyne coupling at a single metal center. Synthesis and structure of bis(diethylamino)acetylene complexes of molybdenum(II) and tungsten(II) (A.C. Filippou, C. Völkl, W. Grünleitner and P. Kiprof), (434) 201
- Reactions of [WI₂(CO)(NCMe)(η²-RC₂R)] (R = Me or Ph) with t-butylisonitrile (E.M. Armstrong, P.K. Baker, T. Callow, K.R. Flower, P.D. Jackson and L.M. Severs), (434) 321
- Polyclopentadienyls: synthesis of arylsilylcyclopentadienyl compounds and their η⁵-complexes with tungsten (H. Plenio), (435) 21
- Preparation and properties of molybdenum and tungsten dinitrogen complexes. XXXV. Synthesis of diazoalkane and hydrazido(2-) complexes of tungsten containing a π-acceptor ligand (T. Aoshima, T. Tamura, Y. Mizobe and M. Hidai), (435) 85
- Synthese und Reaktivität kohlenwasserstoffüberbrückter Komplexe von Mangan, Rhenium, Molybdän und Wolfram (E. Lindner, M. Pabel, R. Fawzi, H.A. Mayer und K. Wurst), (435) 109
- Structural properties of cationic molybdenum and tungsten allyl derivatives (B.J. Briddon, M. Cartwright, A.G.W. Hodson, M.F. Mahon and K.C. Molloy), (435) 319
- EPR studies of photochemical reactions of hexacarbonylbis(η⁵-2,4-cyclopentadien-1-yl)ditungsten, [CpW(CO)₃]₂, with quinones (M. Hanaya and M. Iwaizumi), (435) 337
- Reactions of silyl enol ethers with [ethoxy(phenylethynyl)carbene] chromium and -tungsten complexes (F. Camps, L. Jordi, J.M. Moretó, S. Ricart, A.M. Castaño and A.M. Echavarren), (436) 189
- Oxidative addition reactions of [Os₃(CO)₁₀(MeCN)₂] with 2-imine-substituted derivatives of furan, thiophene, selenophene and pyrrole (A.J. Arce, Y. De Sanctis, L. Hernandez, M. Marquez and A.J. Deeming), (436) 351
- Zur Addition des Thiomethyl-Kations an die Wolfram–Kohlenstoff-Doppelbindung in [Cp(CO)(PM₃)₃W=C(Tol)PR₂]PF₆] (J. Ostermeier, W. Schütt und F.R. Kreißl), (436) C17
- Static and dynamic structures of pentacarbonyl-chromium(0) and -tungsten(0) complexes of dithioether ligands. I. Symmetrical dithioether complexes and X-ray crystal structure of [W(CO)₅(^tBuSCH₂S^tBu)] (E.W. Abel, K.G. Orrell, H. Rahoo, V. Šík, M.A. Mazid and M.B. Hursthouse), (437) 191
- Organometallics of diphosphazanes. V. The different coordination behaviour of *cis*- and *trans*-cyclodiphosphazanes towards Group 6 metal carbonyl moieties (V.S. Reddy, S.S. Krishnamurthy and M. Nethaji), (438) 99
- Heterobimetallic Group 6-rhodium complexes. II. Reactions of [Rh(S₂CNEt₂)(CO)₂] with the complexes [M(CO)₃(η²-phen)(η¹-dppm)] (M = Mo or W) and [Mo(CO)₄(η²-dppm)] as P-donor systems (M. Cano, P. Ovejero and J.V. Heras), (438) 329
- Übergangsmetall-Carbin-Komplexe. CIII. Aktivierung von Ethylisocyanid an elektronenreichen Metallzentren; Synthese von einkernigen Wolfram-Aminocarbon-Komplexen des Typs Tp'(CO)₂W≡CN(R)Et (R = Me, Et) (A.C. Filippou, C. Wagner, E.O. Fischer und C. Völkl), (438) C15
- Phosphinic complexes of rhenium and tungsten in reactions with stereoindicators (Z.V. Todres, E.A. Ionina and A.J.L. Pombeiro), (438) C23
- η¹ und η²-Komplexe von Cp*-substituierten Diphosphenen und Arsaphosphenen: Synthese, Struktur und Umlagerung unter Cp*-Wanderung (P. Jutzi, U. Meyer, S. Opiela, B. Neumann und H.-G. Stammmer), (439) 279
- The synthesis of dinuclear molybdenum and tungsten complexes with the bridging heterodifunctional ligand C₅H₄PPh₂: X-ray crystal structure of Mo₂(μ-C₅H₄PPh₂)(η⁵-C₅H₅)(CO)₅ (T.J. Duckworth, M.J. Mays, G. Conole and M. McPartlin), (439) 327
- Synthesis of tetranuclear heterometallic cluster complexes via condensation of triosmium alkyne complexes Os₃(CO)₁₀(C₂R₂), R = Tol and Me, and mononuclear tungsten acetylidyne complexes LW(CO)₃C≡CR', L = Cp and Cp*, R' = Ph and 'Bu (Y. Chi, R.-C. Lin, C.-C. Chen, S.-M. Peng and G.-H. Lee), (439) 347
- Cycloadditionsreaktionen von Phospheniumkomplexen. VIII. Regioselektive Cycloaddition chiraler Phosphenium-Komplexe Cp(CO)[Ph₂(R)P]M=PPh₂ (M = Mo, W; R = H, Me) mit Isothiocyanaten (H. Pfister und W. Malisch), (439) C11
- Phosphenium-Übergangsmetallkomplexe. XX. Stabilisierung von Mesitylphosphiniden durch ein Cp(CO)₂W- und ein Cp(CO)₂Fe-Fragment (U.-A. Hirth und W. Malisch), (439) C16
- Phosphenium-Übergangsmetallkomplexe. XXI. Schwefel- und Sele-naddition an die Metall–Phosphor-Doppelbindung der zweikernigen Phosphinidenkomplexe Cp(CO)₂M=PMes[M(CO)₃Cp] (M = Mo, W) (U.-A. Hirth, W. Malisch und H. Käb), (439) C20
- Studies on olefin-coordinating transition metal carbene complexes. XIX. Synthesis and structural characterization of the cycloalkene ligated carbonyl-chromium and -tungsten carbene complexes (J. Chen and B. Wang), (440) 67
- Conjugate additions to alkynylalkoxycarbenemetal (Cr or W) complexes (A. Llebaria, J.M. Moretó, S. Ricart, J. Ros, J.M. Viñas and R. Yáñez), (440) 79
- Methylisocyanide derivatives of molybdenocene and tungstenocene: preparation, reactivity and electronic structure: crystal structures of [(η⁵-C₅H₅)₂WBr(CNMe)]Br and [(η⁵-C₅H₅)₂MoC(N(H)CH₃)-N(HC(N(H)CH₃)₂BF₄)₂]CH₃CN (M.J. Calhorda, A.R. Dias, M.T. Duarte, A.M. Martins, P.M. Matias and C.C. Romão), (440) 119
- Electron transfer catalyzed substitution in carbonyl complexes. VI. Highly variable rates of substitution by tetracyanoethylene (B. Schwederski, W. Kaim, B. Olbrich-Deussner and T. Roth), (440) 145
- Schwefel(IV)-Verbindungen als Liganden. XVII. Übergangsmetall-vermittelte Insertion von Schwefeldioxid in die C–O-Einfachbindung (W.A. Schenk und J. Pfeffermann), (440) 341
- Das Verhalten von stereoisomeren Bistriflaten des Tetrahydrofurans gegenüber ein- und zweiwertigen Carbonylmetallaten (E. Lindner, M. Pabel, R. Fawzi und M. Steimann), (441) 63
- Reversible Acyl-Keten-Umwandlung (F.R. Kreißl, H. Keller und W. Schütt), (441) 75
- Static and dynamic structures of pentacarbonyl-chromium(0) and -tungsten(0) complexes of dithioether ligands. II. Unsymmetrical dithioether ligand complexes of general type [M(CO)₄(MeSCH₂-SR)] (E.W. Abel, K.G. Orrell, H. Rahoo and V. Šík), (441) 255
- Kohlenwasserstoffverbrückte Komplexe. XXVI. Reaktionen von anionischen Aromatenkomplexen [(OC)₃Cr(Ar)]⁻ mit metallorganischen Elektrophilen. Darstellung von (OC)₃Cr(μ-η⁶:η⁶-C₆H₅-C₇H₄), M(CO)₃ (M = Cr, Mo), (OC)₃Cr(μ-η⁶:η⁵-C₆H₅-C(O)C₅H₄), Fe(η⁵-C₅H₅), (OC)₃Cr(μ-η⁶:η⁶-C₈H₅S-C₇H₇), Mo(CO)₃, (OC)₃Cr(μ-η⁶:η⁶-C₉H₇-C₇H₇), Mo(CO)₃, (OC)₃Cr(μ-η⁶:η⁶-C₁₃H₉-C₇H₇), M(CO)₃ und (OC)₃Cr(μ-η⁶:η⁶-C₁₃H₁₁-C₇H₇), M(CO)₃ (M = Cr, Mo, W) (J. Breimair, M. Wieser und W. Beck), (441) 429
- Static and dynamic structures of pentacarbonyl-chromium(0) and tungsten(0) complexes of dithioether ligands. III. 1,3-Metallocropic shifts of sulphur coordination sites (E.W. Abel, K.G. Orrell, H. Rahoo and V. Šík), (441) 441
- Synthesis, structure, and chemistry of heterobimetallic phosphido-bridged complexes CpFe(CO)₂(μ-PPh₂)M(CO)₅ (M = Cr, Mo, W) (S.-G. Shyu, P.-J. Lin and Y.-S. Wen), (443) 115

Darstellung und Röntgenstrukturuntersuchung von [1-Carbonyl-1-(η^5 -cyclopentadienyl)-2,3-dimethyl-1-trimethylphosphin-3-trimethylphosphoranylidene-1-wolframa-2-thiacyclopropan]-tetrafluoroborat (W. Schütt, E. Herdtweck, F. Hahn and F.R. Kreißl), (443) C33

Wolfram(0)-Komplexe des π -aciden N-Methylpyrazinium-Kations und des N-Methylpyrazinium-Radikals. Kristall- und Molekülstruktur von $[(C_4H_4N_2\text{-Me})W(\text{CO})_3(\text{PCy}_3)_2](\text{PF}_6)$, Cy = Cyclohexyl (W. Bruns, H.-D. Hauser, W. Kaim und A. Schulz), (444) 121

Unexpected ring-expansion of "axially prosterogenic" biaryl lactones by methylenephosphoranyl-substituted imido complexes of molybdenum and tungsten (J. Sundermeyer, K. Weber, H. Werner, N. Mahr, G. Bringmann and O. Schupp), (444) C37

Synthesis, reactivity, and molecular structure of $[(N(CH_2CH_2O)_3Si-C_6H_5)W(\text{CO})_3]$ (J.-S. Lee, Y.K. Chung, D. Whang and K. Kim), (445) 49

Darstellung und Charakterisierung von Indenyl- und Fluorenylfunktionalisierten Cyclopentadienylcarbonyl-Methylkomplexen des Molybdäns und Wolframs. Molekülstrukturen von $(\eta^5-C_5H_4CMe_2C_9H_7)M(\text{CO})_3\text{Me}$ ($M = \text{Mo}, \text{W}$), $(\eta^5-C_5H_4CMe_2C_{13}H_9)M(\text{CO})_3\text{Me}$ und $(\eta^5:\eta^2-C_5H_4CMe_2C_9H_7)Mo(\text{CO})_2\text{Me}$ (H.G. Alt, J.S. Han und R.D. Rogers), (445) 115

Electrospray mass spectrometric studies of some cationic and anionic Group 6 and Group 7 metal carbonyl complexes (I. Ahmed, A.M. Bond, R. Colton, M. Jurcevic, J.C. Traeger and J.N. Walter), (447) 59

Reactions of the cationic bis(but-2-yne) complex $[W(\text{CO})(\text{NCMe})_2(\text{S}_2\text{CNC}_4\text{H}_8)(\eta^2\text{-MeC}_2\text{Me})_2]\text{[BF}_4]$ with bidentate anionic oxygen and sulphur donor ligands (P.K. Baker and K.R. Flower), (447) 67

Darstellung und Struktur von 1-Metalla-2,3,4-triazolen des Molybdäns und Wolframs (C.M. Stegmair, W. Schütt, W. Ullrich, P. Kiprof, J. Ostermeier und F.R. Kreißl), (447) 251

P_4 -Liganden mit maximaler Elektronendonorfähigkeit. V. Zur Chlorierung eines cyclo-P₄-Ligand-Komplexes (M. Scheer, M. Dargatz und P.G. Jones), (447) 259

Uranium

Anionic triscyclopentadienyluranium(III) hydrides (J.-C. Berthet, C. Villiers, J.-F. Le Maréchal, B. Delavaux-Nicot, M. Lance, M. Nierlich, J. Vigner and M. Ephritikhine), (440) 53

Synthesis, structure and oxidative addition reactions of triscyclopentadienyluranium(III) nitrile complexes (R. Adam, C. Villiers, M. Ephritikhine, M. Lance, M. Nierlich and J. Vigner), (445) 99

Vanadium

ansa-Metallocene derivatives. XXV. Synthesis, crystal structure and reactions of a tetramethylethano-bridged vanadocene dichloride, $(\text{CH}_3)_4\text{C}_2(\text{C}_5\text{H}_4)_2\text{VCl}_2$ (B. Dorer, J. Diebold, O. Weyand and H.-H. Bräntzinger), (427) 245

Cyclopentadienyl- and methylcyclopentadienyl-vanadium complexes containing chalcogen bridges: dechalcogenation to pseudocubane clusters (M. Herberhold, M. Schrepfermann and J. Darkwa), (430) 61

Organometallic Chemie hochfluorierter Ligand-Systeme. XIV. Vanadocenehexafluoroarsenat-Komplexe: Struktur von $\text{Cp}_2\text{V}(\text{AsF}_6)_2$ und $^{51}\text{V-NMR}$ von $[\text{Cp}_2\text{VCl}_2]^+[\text{AsF}_6]^-$ (P. Gowik, T.M. Klapötke, K. Siems and U. Thewalt), (431) 47

Preparation and structure of half-open vanadocene, $\text{V}[2,4-(\text{CH}_3)_2\text{C}_5\text{H}_5](\text{C}_5\text{H}_5)\text{CO}$ (Y. Wang, Y. Liang, J. Liu, S. Jin and Y. Lin), (435) 311

Reactions of metallocenes with organocadmium and organomercury compounds (V.P. Maryin), (441) 241

Ytterbium

Struktur und Reaktivität von Ytterbiumbenzamidinaten (M. Wedler, A. Recknagel, J.W. Gilje, M. Nottemeyer and F.T. Edelmann), (426) 295

Synthesis and characterization of pentaphenylytterbium $\text{Ph}_2\text{Yb}(\text{THF})(\mu\text{-Ph})_3\text{Yb}(\text{THF})_3$ (M.N. Bochkarev, V.V. Khramenkov, Y.F. Rad'kov, L.N. Zakharov and Y.T. Struchkov), (429) 27

Bis(cyclopentadienyl)ytterbium(III)-triflat: Ein Zweikernkomplex mit Trifluormethylsulfonat-Brücken (J. Stehr and R.D. Fischer), (430) C1

Synthesis, thermal stability and X-ray crystal structure of the bis(cyclopentadienyl)ytterbium naphthoxide derivative $(C_5H_5)_2\text{Yb}(\text{OC}_{10}\text{H}_7)(\text{THF})$ (X.-G. Zhou, Z.-Z. Wu and Z.-S. Jin), (431) 289

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Synthesis and structure of organotin complexes of ytterbium (L.N. Bochkarev, O.V. Grachev, S.F. Ziltsov, L.N. Zakharov and Y.T. Struchkov), (436) 299

The first examples of intermolecular weak (agostic) γ -methyl–metal interactions in organolanthanide complexes: the synthesis and X-ray structures of $[(\text{Yb}(\eta\text{-Cp}'')_2)_2]$ and $[(\text{Eu}(\eta\text{-Cp}'')_2)_2]$ [$\text{Cp}'' = C_5H_3(\text{SiMe}_3)_2\text{-}1,3$] (P.B. Hitchcock, J.A.K. Howard, M.F. LapPERT and S. Prashar), (437) 177

Alpha-metalated diazoalkanes: synthesis of yttrium, lutetium and ytterbium trimethylsilyldiazomethyl complexes (H. Siebald, M. Dartiguenave and Y. Dartiguenave), (438) 83

Ferrocenliganden in der Lanthanid-Chemie: Chelatstabilisierung des ersten Diorganyllanthanid(III)-halogenids (H. Gornitzka, A. Steiner, D. Stalke, U. Kilmann, F.T. Edelmann, K. Jacob und K.-H. Thiele), (439) C6

Studies on organolanthanide complexes. XLIV. Synthesis of bis(2-methoxyethylcyclopentadienyl) divalent organolanthanides ($\text{Ln} = \text{Sm}, \text{Yb}$) and molecular structure of solvated $(\text{MeOCH}_2\text{CH}_2\text{C}_5\text{H}_4)_2\text{Yb} \cdot \text{OC}_4\text{H}_8$ (D. Deng, C. Qian, F. Song, Z. Wang, G. Wu and P. Zheng), (443) 79

Novel polynuclear organotin complexes of samarium and ytterbium (L.N. Bochkarev, O.V. Grachev, N.E. Molosnova, S.F. Ziltsov, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), (443) C26

Umlagerung von Tetrahydrofuran im Lanthanoid-Organyl: Die Kristall- und Molekülstrukturen von Bis(cyclopentadienyl)ytterbium(III) (μ -n-butene-2-olat) und Bis(methylcyclopentadienyl)ytterbium(III) (μ - η^2 : η^1 -tetrahydrofuran-2-olat) (G. Massarweh und R.D. Fischer), (444) 67

Studies on organolanthanide complexes. XLVI. Synthesis and characterization of N-containing ring-linked biscyclopentadienyl lanthanide and yttrium chlorides (C. Qian and D. Zhu), (445) 79

Synthesis and characterization of novel covalent biphenyldiyl Sm^{II} and Yb^{II} complexes (Y. Chauvin, N. Marchal, H. Olivier and L. Saussine), (445) 93

ansa-Metallocenderivate des Samariums und Ytterbiums mit "weichen" Donorliganden (F.T. Edelmann, M. Rieckhoff, I. Haiduc und I. Silaghi-Dumitrescu), (447) 203

Structural investigation of the organolanthanide di- η^5 -cyclopentadienyl 1,1,1-trifluoroacetylacetato ytterbium (Z. Ye, Y. Yu, S. Wang and X. Jin), (448) 91

Synthesis and molecular structure of $(\text{CH}_3\text{Cp})_2\text{Yb} \cdot \text{DME}$ and its catalytic activity for the polymerization of methyl methacrylate (T. Jiang, Q. Shen, Y. Lin and S. Jin), (450) 121

Yttrium

Studies on organolanthanide complexes. XLI. Catalytic isomerization of olefins by organolanthanide complex/sodium hydride systems (C. Qian, D. Zhu and D. Li), (430) 175

Metallorganische Verbindungen der Lanthanoide. LXX. Bis(cyclopentadienyl)seltenerdkomplexe: Synthese und Röntgenstrukturanalyse von dem intramolekular stabilisierten Lutetiumalkyl $Cp_2Lu(CH_2)_3NMe_2$ und dem monomeren basenfreien Yttriumcarboxylat $Cp_2Y[\eta^2-O_2C(CH_2)_3NMe_2]$ (H. Schumann, J.A. Meese-Markscheffel, A. Dietrich and F.H. Görilitz), (430) 299

Structural trends in bis(pentamethylcyclopentadienyl)lanthanide and yttrium complexes (W.J. Evans and S.E. Foster), (433) 79

Alpha-metatalated diazoalkanes: synthesis of yttrium, lutetium and ytterbium trimethylsilyldiazomethyl complexes (H. Siebold, M. Dartiguenave and Y. Dartiguenave), (438) 83

Studies on organolanthanide complexes. XLVI. Synthesis and characterization of N-containing ring-linked biscyclopentadienyl lanthanide and yttrium chlorides (C. Qian and D. Zhu), (445) 79

Zinc

Substitution of the acetoxy groups of dialkoxy methylacetates by organometallic reagents: a route to allyl-, propargyl-, homoallyl-, homopropargyl- and α -stannylacetals (I. Beaudet, A. Duchêne, J.-L. Parrain and J.-P. Quintard), (427) 201

$2,2'$ -Bipyridyl-bis[bis(trimethylsilyl)methyl]zink — Synthese, spektroskopische Charakterisierung und Struktur (M. Westerhausen, B. Rademacher and W. Schwarz), (427) 275

Complexation of diphenylzinc with simple ethers. Crystal structures of the complexes $Ph_2Zn \cdot$ glyme and $Ph_2Zn \cdot$ diglyme (P.R. Markies, G. Schat, O.S. Akkerman, F. Bickelhaupt and A.L. Spek), (430) 1

Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CX. Methylierte Nitridomolybdän(VI)-Komplexe (W.A. Herrmann, S. Bogdanović, J. Behm and M. Denk), (430) C33

Reactions of sterically hindered organozinc and organocadmium compounds containing functional silicon centres. Crystal structures of $Zn[C(SiMe_3)_2(SiMe_2OCOCF_3)]_2$ and $Cd[C(SiMe_3)_2(SiMe_2OMe)]_2$ (S.S. Al-Juaid, C. Eaborn, A. Habtemariam, P.B. Hitchcock and J.D. Smith), (437) 41

Mehrachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXV. Dihalogenoxoosmium(VI)-Komplexe des Pinakols: Synthese, Spektroskopie und Alkylierung (W.A. Herrmann and P. Watzlowik), (437) 363

Preparation of polyfunctional diorganomercurials and their transmetalation to diorganozincs. Applications to the preparation of optically active secondary alcohols (M.J. Rozema, D. Rajagopal, C.E. Tucker and P. Knochel), (438) 11

Synthesis of polymetallacycloalkanes (P. Mazerolles and C. Laurent), (439) 115

Etude de la fixation du groupe tertiaire *gem*-diméthylpropynyle $HC\equiv CC(CH_3)_2$ par voie organométallique; applications (F. Barbot et P. Miginiac), (440) 249

Allénysilanes fonctionnels. II. Synthèse par voie organoaluminique du 2-triméthylsilylbuta-2,3-diénal et application à la synthèse d'alcools secondaires α -alléniques triméthylsilylés (D. Mesnard et L. Miginiac), (440) 277

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindingen. VI. Synthese und Eigenschaften von Zincacyclohexankomplexen des Typs $\{Li(D)_2\}_2Zn\{CH_2CH_2CH_2-CH_2CH_2\}_2$ ($D = (C_2H_5)_2O$; $2D = (CH_3)_2NC_2H_4N(CH_3)_2$); Molekülstruktur von $\{Li(CH_3)_2NC_2H_4N(CH_3)_2\}_2Zn\{CH_2CH_2-CH_2CH_2CH_2\}_2$ (H.-O. Fröhlich, B. Kosan, B. Müller und W. Hiller), (441) 177

Bis[2,2,4,4,6,6-hexamethyl-2,4,6-trisila-cyclo-hexyl]zink und sein 2,2'-Bipyridyl-Addukt — spektroskopische Charakterisierung und Struktur (M. Westerhausen und B. Rademacher), (443) 25

New method of zinc activation by electrochemistry: synthetic applications to the Blaise reaction (N. Zylber, J. Zylber, Y. Rollin, E. Duñach and J. Perichon), (444) 1

Trisubstituted alkenes from alkynes and aryl iodides by boron-to-zinc transmetalation and palladium(0) catalyzed cross coupling (K.A. Agrios and M. Srebnik), (444) 15

Structural studies of some Group 12 metal alkyl adducts: the X-ray crystal structures of $Me_2Zn[Me_2N(CH_2)_2NMe_2]$, $Me_2Cd[Me_2N(CH_2)_2NMe_2]$, $(Me_3CCH_2)_2Zn[Me_2N(CH_2)_2NMe_2]$ and $(Me_3CCH_2)_2Cd[Me_2N(CH_2)_2NMe_2]$ (P. O'Brien, M.B. Hursthause, M. Mottevalli, J.R. Walsh and A.C. Jones), (449) 1

Low-temperature organometallic synthesis of crystalline and glassy ternary semiconductors $M^{II}M^{IV}P_2$ where $M^{II} = Zn$ and Cd , and $M^{IV} = Ge$ and Sn (S.C. Goel, W.E. Buhro, N.L. Adolphi and M.S. Conradi), (449) 9

Formation of a high surface area, regular porous solid from the cluster of clusters, $Zn_4^{II}O[(CO)_9Co_2CCO_2]_6$ (W. Cen, B. Ladna, T.P. Fehlner, A.E. Miller and D. Yue), (449) 19

Etude de la régiosélectivité de l'action des organozinciques sur les α -iminoesters issus de 2-amino ou 3-aminoalcools. I. Synthèse d' α -aminoesters polyfonctionnels (G. Courtois et L. Miginiac), (450) 33

Zirconium

Di(tertiobutylcyclopentadienyl)zircona[1]ferrocenophane précurseur de zirconocenes disubstitués dissymétriquement (R. Broussier, A. Da Rold and B. Gautheron), (427) 231

Insertion of alkynes into zirconium–nitrogen bonds leading to 2,3-diaza-metallocyclopentenes. Generation and reactivity of $Cp_2Zr(N_2Ph_2)$ (P.J. Walsh, F.J. Hollander and R.G. Bergman), (428) 13

Agostic interaction in early transition metal alkyls and their role in catalytic activity for olefin polymerizations (K. Mashima and A. Nakamura), (428) 49

Cyclohexyl[*trans*-1,2-bis(1-indenyl)]zirconium(IV)dichlorid: Ein chiraler Polymerisationskatalysator mit stereochemisch starrer Brücke (B. Rieger), (428) C33

Hydrozirconation of oleyl sulphides and oleyl phenylsulphone (S. Karlsson, A. Hallberg and S. Gronowitz), (430) 53

Structure of an oxazirconacyclohexadiene synthesized by thermolysis of difurylzirconocene (G. Erker, R. Petrenz, C. Krüger and M. Nolte), (431) 297

Dehydrierende Polymerisation von Methyldisilanen mittels Metallocen-Katalyse zu den Oligomeren $Me_3Si[Me_2Si]_nSiMe_2H$, $HMe_2Si[Me_2Si]_nSiMe_2H$ mit $n = 3-9$, $H[MeSiH]_nH$ mit $n = 3-14$ und hochpolymerem Poly(methylsilan) ($MeSiH_x$) $_n$ (E. Hengge und M. Weinberger), (433) 21

Mehrketige Zirkonocen-Ethylkomplexe (H.G. Alt, C.E. Denner und R. Zenk), (433) 107

Base-free cationic 14-electron alkyls of Ti, Zr and Hf as polymerisation catalysts: a comparison (M. Bochmann and S.J. Lancaster), (434) C1

ansa-Metallocene derivatives of Ti^{IV} and Zr^{IV} with the shortest $-C(CH_3)_2-$ bridge (I.E. Nifant'ev, A.V. Churakov, I.F. Urazowski, S.G. Mkoyan and L.O. Atovmyan), (435) 37

Synthesis and characterization of Group IV metal–iron complexes bridged by carboxylate substituted cyclopentadienyl group. The structure of $Cp_2Ti(\mu-O_2CC_5H_4)Fe(CO)_2(CH_2C_6H_5)_2$ (H.-M. Gau, C.-C. Schei, L.-K. Liu and L.-H. Luh), (435) 43

Synthese und Kristallstruktur von $Cp'_2Zr[P(SiMe_3)_2]_2$ ($Cp' = \eta^5-C_5H_4Me$) — Erste röntgenstrukturanalytische Untersuchung

- eines Zirkonocen-Bisphosphido-Komplexes (F. Lindenberg und E. Hey-Hawkins), (435) 291
- Preparation and crystal structures of the complexes ($\eta^5\text{-C}_5\text{H}_4\text{CMe}_2$ - $\eta^5\text{-C}_{13}\text{H}_8\text{MCl}_2$ (M = Zr, Hf) and their role in the catalytic formation of syndiotactic polypropylene (A. Razavi and J. Ferrara), (435) 299
- Ferrocenhaltige Liganden in der Lanthanid-Chemie: Alkoxide und Thiolate (H. Gornitzka, F.T. Edelmann und K. Jacob), (436) 325
- Condensation of primary silanes in the presence of Cp_2MCl_2 / $^\text{n}\text{BuLi}$ (M = Ti, Zr, Hf) (J.Y. Corey and X.-H. Zhu), (439) 1
- Characterization of 1:1 heterobimetallic complexes of ($\eta^5\text{-C}_5\text{Me}_5\text{Si}$) $\text{Ir}(\text{CO})_2$ with zinc, cadmium and mercury(II) chloride and the X-ray structure determination of the 1:2 mercury complex (F.W.B. Einstein, X. Yan, X. Zhang and D. Sutton), (439) 221
- Heterozyklen als Liganden. XVI. (2,5-C₄¹Bu₂RHN)MCl₃ (M = Ti, Zr, Hf; R = H, SiMe₃)—Azacyclopentadienyl-Komplexe der Gruppe 4-Metalle (N. Kuhn, S. Stubenrauch, R. Boese und D. Bläser), (440) 289
- Dehydrierende Copolymerisation zu den neuen Polysilan-Copolymeren Poly(methyl-co-phenylsilan) H[(MeSiH)_x(PhSiH)_y]_nH und Poly(methyl-co-dimethylsilan) H[(MeSiH)_x(Me₂Si)_y]_nH (E. Hengge und M. Weinberger), (441) 397
- Rearrangement processes of bis-(η^5 -alkylcyclopentadienyl)-zirconium dichlorides under electron impact (Y.A. Andrianov and V.P. Maryin), (441) 419
- Benzylverbindungen elektronenärmer Übergangsmetalle. Molekülsstrukturen von $\text{Cp}_2\text{Ti}(\text{CH}_2\text{Ph})_2$ und $\text{CpZr}(\text{CH}_2\text{Ph})_3$ (J. Scholz, F. Rehbaum, K.-H. Thiele, R. Goddard, P. Betz und C. Krüger), (443) 93
- Mechanistische Aspekte zur übergangsmetall-katalysierten dehydrierenden Polymerisation von Disilanen und höheren Silanen (E. Hengge und M. Weinberger), (443) 167
- Synthesis and reactivity of substituted cyclopentadienyl diphenylphosphido complexes of zirconium(IV). X-Ray crystal structure of [$(\text{C}_5\text{H}_4\text{SiMe}_3)_2\text{Zr}(\text{PPh}_2)_2$] (A.-M. Larsonneur, R. Choukroun, J.-C. Daran, T. Cuenca, J.C. Flores and P. Royo), (444) 83
- Cyclische Metallamide. III. Flüchtige Tetrahydroborat-Komplexe von Titan und Zirkonium (W.A. Herrmann, M. Denk, W. Scherer und F.-R. Klingan), (444) C21
- Ketene acetal metallocene complexes. Crystal structure of $(\text{C}_5\text{H}_5)_2\text{Ti}(\text{Cl})\text{OC}(\text{OCH}_3)=\text{C}(\text{CH}_3)_2$ (K. Hortmann, J. Diebold and H.-H. Brintzinger), (445) 107
- Stabilization of a catalytically syndiotactic-specific metallocene cation by trimethylphosphine. The crystal structure of [$(\eta^5\text{-C}_5\text{H}_4)\text{CMe}_2$ - $\eta^5\text{-C}_{13}\text{H}_8\text{ZrMePME}_3]^+ [\text{B}(\text{C}_6\text{F}_5)_4]^-$ (A. Razavi and U. Thewalt), (445) 111
- Un nouveau type de ferrocénophane, premier exemple de structure dimère (R. Broussier, Y. Gobet, R. Amardeal, A. Da Rold, M.M. Kubicki et B. Gautheron), (445) C4
- Fulvalene cyclopentadienyl titanium and zirconium(III) and -(IV) complexes. X-Ray crystal structure of [$\{\text{Ti}(\eta^5\text{-C}_5\text{H}_5\text{Cl})_2\}(\mu\text{-O})(\mu\text{-}\eta^5\text{-}\eta^5\text{-C}_{10}\text{H}_8)$] (A. Cano, T. Cuenca, G. Rodríguez, P. Royo, C. Cardin and D.J. Wilcock), (447) 51
- Studies on the reactions of transition metal hydrides with heterocumulenes. VI. Reaction of $\text{Cp}_2\text{Zr}(\text{H})\text{Cl}$ with RNCS (R = C₆H₅, 2-C₁₀H₇, c-C₆H₁₁ and n-C₄H₉) (M. Wang, S. Lu, M. Bei, H. Guo and Z. Jin), (447) 227
- Group IV metal-chromium complexes bridged by a benzoate group (H.-M. Gau, C.-T. Chen, T.-T. Jong and M.-Y. Chien), (448) 99
- A comparison of heterometallic alkoxide molecules containing copper(I) or copper(II) and zirconium (B.A. Vaartstra, J.A. Samuels, E.H. Barash, J.D. Martin, W.E. Streib, C. Gasser and K.G. Caulton), (449) 191
- Stereoselective propene polymerization at a metallocene/alumoxane catalyst derived from the chirally-substituted “meso-like” (*p*-R,*p*-S)bis[1-(neoisopinocampyl)-4,5,6,7-tetrahydroindenyl]zirconium dichloride (G. Erker, M. Aulbach, C. Krüger and S. Werner), (450) 1
- Carbonylation of 2-bornenyl(methyl)zirconocene (G. Erker, R. Noe, M. Albrecht and A.L. Rheingold), (450) 137
- ### Book reviews
- Gmelin handbook of inorganic and organometallic chemistry, (C. Eaborn), (426) C21
- Advances in free radical chemistry, vol. 1, D.D. Tanner (ed.) (C. Eaborn), (426) C21
- Reductions by the aluminato- and borohydrides in organic synthesis, J. Seyden-Penne (J.R. Hanson), (426) C22
- Organic synthesis, the roles of boron and silicon, S.E. Thomas (J.R. Hanson), (426) C23
- Gmelin handbook of inorganic and organometallic chemistry, 8th edition, Mo, part 7, organomolybdenum compounds, Vol. XI (G.J. Leigh), (426) C50
- Gmelin handbook of inorganic and organometallic chemistry, 8th edition, Fe-iron supplement volume B1, compounds with noble gases and hydrogen, Vol. XX (G.J. Leigh), (426) C50
- Gmelin handbook of inorganic and organometallic chemistry, 8th edition, Fe supplement B18, organo-iron compounds, Vol. XI (G.J. Leigh), (426) C51
- Kinetics and mechanism of reactions of transition metal complexes, R.G. Wilkins (R.A. Henderson), (426) C62
- Gmelin handbook of inorganic and organometallic chemistry, 8th edition, Br-bromine, supplement Vol. B1, compounds with rare gases and hydrogen, Vol. XVIII (G.J. Leigh), (426) C63
- Silicon-Based Polymer Science. A Comprehensive Resource, J.M. Zeigler and F.W.G. Fearon (Eds.) (C. Eaborn), (427) C6
- Inorganic Polymers, J.E. Mark, H.R. Allcock and R. West (C. Eaborn), (427) C19
- The Silicon-Heteroatom Bond, D.A. Armitage, R.J.P. Corru, T.C. Kendrick, B. Parbhov, T.D. Tilley, J.W. White and J.C. Young (C. Eaborn), (427) C41
- Frontiers of Organosilicon Chemistry, A.R. Bassindale and P.P. Gaspar (Eds.) (C. Eaborn), (427) C42
- Preparative Polar Organometallic Chemistry, Vol. 2, L. Brandsma (B.J. Wakefield), (427) C42
- Aspects of homogeneous catalysis, Vol. 7, Renato Ugo (ed.) P.A. Chaloner), (429) C4
- Advances in metal-organic chemistry, Vol. 2, Lanny S. Liebeskind (Ed.) (P.A. Chaloner), (429) C5
- Inorganic biochemistry of iron metabolism, R.R. Crichton (G.J. Leigh), (429) C6
- Dioxygen activation and homogeneous catalytic oxidation, L.I. Simándi (G.J. Leigh), (429) C6
- Nucleoside synthesis — organosilicon methods, E. Lukevics (D.W. Young), (429) C49
- Electron deficient boron and carbon clusters, G.A. Olah (C.E. Housecroft), (429) C49
- Chemistry of the Platinum Group Metals: Recent Developments, F.R. Hartley (Ed.) (D.M.P. Mingos), (434) C15
- Reaction Mechanisms of Inorganic and Organometallic Systems, R.B. Jordan (D.A. Brown), (435) C1
- Inorganic Reactions and Methods. Vol. 9. Formation of Bonds to C, Si, Ge, Sn, Pb (Part I), A.P. Hagen (Ed.) (C. Eaborn), (436) C13
- Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Edition. Supplement Volume B5c. Silicon Nitride in Electronics (C. Eaborn), (436) C14

- Inorganic Reactions and Methods. Volume 4. Formations of Bonds to Halogens (Part 2), A.P. Hagen (Ed.) (C. Eaborn), (436) C21
- Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Ed. Sn. Organotin Compounds. Part 19. Organotin-nitrogen Compounds (concluded), Organotin-Phosphorus, -Arsenic, -Antimony, and -Bismuth Compounds, (C. Eaborn), (436) C22
- Advances in Sonochemistry, Volume 2, T.J. Mason (Ed.) (P.D. Lickiss), (436) C25
- Inorganic Reactions and Methods, Vol. 13, The Formation of Bonds to Group-I, -II, and -IIIB Elements, A.P. Hagen (Ed.) (C. Eaborn), (436) C130
- Free Radical Chain Reactions in Organic Synthesis, W.B. Motherwell and D. Crich (C. Eaborn), (436) C210
- Gmelin Handbook of Inorganic and Organometallic Chemistry. 8th Edn. F. Perfluorohalogenoorgano Compounds of Main Group Elements. Supplement Vol. 6. Aliphatic and Aromatic Compounds of Nitrogen (continued), (C. Eaborn), (436) C250
- Hazardous Metals in the Environment, M. Stoeppeler (Ed.) (G.J. Leigh), (437) C35
- Environmental analysis using chromatography interfaced with atomic spectroscopy, R.M. Harrison and S. Rapsomanikis (Eds.) (G.J. Leigh), (437) C35
- Introduction to cluster chemistry, D.M.P. Mingos and D.J. Wales (G. Cloke), (437) C36
- The Organometallic Chemistry of the Transition Elements, by R.H. Crabtree (G.J. Leigh), (438) C13
- A Guide for the Perplexed Organic Experimentalist, H.J.E. Loewenthal (C. Eaborn), (445) C17
- Advances in Physical Organic Chemistry, Vol. 27, D. Bethell (ed.), (C. Eaborn), (445) C17
- Directory of Graduate Research 1991 (C. Eaborn), (445) C18
- Organometallics: A Concise Introduction by Ch. Elschenbroich and A. Salzer (C. Eaborn), (445) C18
- Salt Effects in Organic and Organometallic Chemistry by A. Loupy and B. Tchoubar (C. Eaborn), (447) C10
- Inorganic reactions and methods, Vol. 5 The Formation of Bonds to Group VIB (O, S, Se, Te, Po) Elements (Part 1), A.P. Hagen (ed.) (C. Eaborn), (447) C10
- Gmelin Handbook of Inorganic and Organometallic Chemistry, 8th edition Iron, Organoiron Compounds, Part A, Ferrocene 10 (G.J. Leigh), (447) C18
- Gmelin Handbook of Inorganic and Organometallic Chemistry, 8th edition Iron, Organoiron Compounds, Part C 6a, Trinuclear Compounds (G.J. Leigh), (447) C18
- The Chemistry of Organophosphorus Compounds, Volume 2: Phosphine Oxides, Sulphides, Selenides, and Tellurides. Edited by Frank R. Hartley, in the series *The Chemistry of Functional Groups*, edited by Saul Patai, John Wiley and Sons, Chichester, 1992, pp. 647. (J.F. Nixon), (447) C18