

Subject Index of Volume 458

Antimony

1,3-Dioxa-[3]ferrocenophane (M. Herberhold und H.-D. Brendel), 205

Arsenic

1,3-Dioxa-[3]ferrocenophane (M. Herberhold und H.-D. Brendel), 205

Boron

Neue Synthesen und Reaktionen ungesättigter Heterotitanacyclen (S. Dürr, U. Höhlein und R. Schobert), 89

1,3-Dioxa-[3]ferrocenophane (M. Herberhold und H.-D. Brendel), 205

Chromium

Déprotonation des allylbenzenes chrometricarbonyle en milieu basique. Etude RMN ^1H et ^{13}C et réactivité des anions allyliques: influence électronique et stérique du groupement chrometricarbonyle (M.C. Sénéchal-Tocquer, D. Senechal, J.Y. Le Bihan, D. Gentric, B. Caro, M. Gruselle, C. Cordier et G. Jaouen), 105

Application of scalar ^{13}C , ^{19}F couplings between carbonyl carbons and aromatic fluorine to investigation of conformation of chelate phosphitodicarbonylchromium and dicarbonyl(phosphine)chromium complexes of fluorobenzenes (P. Szczeciński), 119

Heterozyklen als Liganden. XVIII. Synthese und Struktur von Tricarbonyl(1,2,3,4,5-pentamethylpyrrol)chrom(0)-der Einfluß der Methylsubstituenten auf die Koordinationseigenschaften des Pyrrol-Liganden (N. Kuhn, J. Kreutzberg, E.-M. Lampe, D. Bläser und R. Boese), 125

The synthesis of ring-substituted cycloheptatrienyl complexes $[\text{M}(\text{CO})_3(\eta^7\text{-C}_7\text{H}_6\text{R})]^+$ ($\text{M} = \text{Cr, Mo or W; R} = \text{Me, } ^{1}\text{Bu, } \text{C}_6\text{H}_4\text{-F-p or C}\equiv\text{CPh}$) (A. Roberts and M.W. Whiteley), 131

Metal-ligand bonding in $\text{LM}(\text{CO})_5$ complexes ($\text{L} = \text{NMe}_3, \text{ PMe}_3, \text{ PCl}_3, \text{ PBr}_3$ or $\text{AsMe}_3; \text{M} = \text{Cr, Mo or W}$) from dielectric, electro-optic and spectroscopic evidence (M.S. Davies, R.K. Pierens and M.J. Aroney), 141

Iron

Poly-linked cyclopentadienide ions. II. Synthesis and properties of the tetracyclopentadienylmethane tetraanion (T. Kinoshita, S. Tsuji, T. Otake and K. Takeuchi), 187

New synthesis of acylferrocene by hydroiminoacetylation of the terminal olefin with ferrocenecarboxaldimine and application to polymer-supported acylferrocene (C.-H. Jun, J.-B. Kang and J.-Y. Kim), 193

Synthesis, some reactions, and molecular structure of the $\text{Pd}(\text{BF}_4)_2$ complex of 1,1'-bis(diphenylphosphino)ferrocene (M. Sato, H. Shigeta, M. Sekino and S. Akabori), 199

1,3-Dioxa-[3]ferrocenophane (M. Herberhold und H.-D. Brendel), 205

Lithium

Poly-linked cyclopentadienide ions. II. Synthesis and properties of the tetracyclopentadienylmethane tetraanion (T. Kinoshita, S. Tsuji, T. Otake and K. Takeuchi), 187

Manganese

Metallorganische Schutzgruppen für den Aufbau ungewöhnlicher Zinn-Chalkogen-Käfigverbindungen: Thiotannat(II) $\text{Sn}_3\text{S}_4^{2-}$ und "Zinnoxidhydrat" $\text{Sn}_6(\mu_3\text{-OH})_4(\mu_3\text{-O})_4$ als Liganden in $^{\text{R}}\text{Cp}(\text{CO})_2\text{Mn}$ -Komplexen (B. Schiemenz, F. Ettel, G. Huttner und L. Zsolnai), 159

Activation of the carbon-carbon bond in the Mn-mediated cycloaddition reaction between disilacyclobutene and cyclic conjugated dienes (H.-J. Chiang and C.-S. Liu), 167

Kinetics and mechanism of the reaction of bis- μ -diethylphosphido-bis(tetracarbonylmanganese) with tri-*n*-butylphosphine (M. Basato), 173

Komplexechemie perhalogenierter Cyclopentadiene und Alkine. XIII. Darstellung, Reaktionen und Strukturen von $[\text{C}_5\text{Cl}_4\text{M}]\text{Mn}(\text{CO})_3$ ($\text{M} = \text{Mn}(\text{CO})_5, \text{Au}(\text{PPh}_3)$) (K. Sünkel und U. Birk), 181

Photochemistry of methanesulphinatopentacarbonylmanganese, $\text{Mn}(\text{CO})_5(\text{SO}_2\text{CH}_3)$, in frozen gas matrices at ca. 12 K (X. Ren and A.J. Rest), C5

Metallocenes

Synthesis, some reactions, and molecular structure of the $\text{Pd}(\text{BF}_4)_2$ complex of 1,1'-bis(diphenylphosphino)ferrocene (M. Sato, H. Shigeta, M. Sekino and S. Akabori), 199

1,3-Dioxa-[3]ferrocenophane (M. Herberhold und H.-D. Brendel), 205

Molybdenum

Chirale Tripodliganden: Die Eintopfreaktion $\text{MeC}(\text{CH}_2\text{PPh}_2)_3 \rightarrow \text{MeC}(\text{CH}_2\text{P}(\text{Ph})(\text{R}))_3$; Zwischenstufen, diastereoselektive Kontrolle und Komplexechemie (O. Walter, T. Klein, G. Huttner und L. Zsolnai), 63

The synthesis of ring-substituted cycloheptatrienyl complexes $[\text{M}(\text{CO})_3(\eta^7\text{-C}_7\text{H}_6\text{R})]^+$ ($\text{M} = \text{Cr, Mo or W; R} = \text{Me, } ^{1}\text{Bu, } \text{C}_6\text{H}_4\text{-F-p or C}\equiv\text{CPh}$) (A. Roberts and M.W. Whiteley), 131

Metal-ligand bonding in $\text{LM}(\text{CO})_5$ complexes ($\text{L} = \text{NMe}_3, \text{ PMe}_3, \text{ PCl}_3, \text{ PBr}_3$ or $\text{AsMe}_3; \text{M} = \text{Cr, Mo or W}$) from dielectric, electro-optic and spectroscopic evidence (M.S. Davies, R.K. Pierens and M.J. Aroney), 141

Synthesis, structure, and properties of molybdenum octahedral d^6 bis(alkyne) complexes (C.H. Lai, C.H. Cheng, C.Y. Cheng and S.L. Wang), 147

Nickel

A one-pot method for the preparation of $(\text{R}_3\text{P})_2\text{Ni}^0\text{L}$ complexes (A. Stanger and A. Shazar), 233

Synthesis and structure of the nickel A-frame complex $\text{Ni}_2(\mu-\text{CNC}_6\text{H}_5)(\text{dppm})_2\text{Cl}_2$ (S. Hinze, J.K. Gong, P.E. Fanwick and C.P. Kubiak), C10

Palladium

Synthesis, some reactions, and molecular structure of the $\text{Pd}(\text{BF}_4)_2$ complex of 1,1'-bis(diphenylphosphino)ferrocene (M. Sato, H. Shigeta, M. Sekino and S. Akabori), 199

1,8-Bis(diphenylphosphino)naphthalene: a rigid chelating, diphosphine analogue of proton sponge (R.D. Jackson, S. James, A.G. Orpen and P.G. Pringle), C3

Regioselective ring opening of a palladium(IV) alkylaromatic metalacycle by benzyl group migration from palladium to the aromatic carbon and X-ray structure of the resulting palladium(II) complex (G. Bögelj, M. Catellani and S. Ghelli), C12

Phosphorus

η^7 -Cycloheptatrienyl-carbonylvanadium-Komplexe mit mono- und bifunktionellen Phosphan-Liganden. Eine ^{51}V -NMR-spektroskopische Untersuchung (M. Herberhold und M. Schrepfermann), 19

Chirale Tripodliganden: Die Eintopfreaktion $\text{MeC}(\text{CH}_2\text{PPh}_2)_3 \rightarrow \text{MeC}(\text{CH}_2\text{P}(\text{Ph})(\text{R}))_3$; Zwischenstufen, diastereoselektive Kontrolle und Komplexchemie (O. Walter, T. Klein, G. Huttner und L. Zsolnai), 63

Kinetics and mechanism of the reaction of bis- μ -diethylphosphido-bis(tetracarbonylmanganese) with tri-*n*-butylphosphine (M. Basato), 173

Komplexchemie perhalogenierter Cyclopentadiene und Alkine. XIII. Darstellung, Reaktionen und Strukturen von $[\text{C}_5\text{Cl}_4\text{M}]\text{Mn}(\text{CO})_3$ ($\text{M} = \text{Mn}(\text{CO})_5, \text{Au}(\text{PPh}_3)$) (K. Sünkel und U. Birk), 181

1,3-Dioxa-[3]ferrocenophane (M. Herberhold und H.-D. Brendel), 205

Ligandeneffekte bei der Synthese der Komplexe $\text{ClRh}(\text{P} \sim \text{O})\text{P} \sim \text{O}$, $[\text{Rh}(\text{P} \sim \text{O})_2\text{BPh}_4]$ und $[\text{Rh}(\text{COD})(\text{P} \sim \text{O})]\text{BPh}_4$ (E. Lindner, Q. Wang, H.A. Mayer and A. Bader), 229

1,8-Bis(diphenylphosphino)naphthalene: a rigid chelating, diphosphine analogue of proton sponge (R.D. Jackson, S. James, A.G. Orpen and P.G. Pringle), C3

Platinum

1,8-Bis(diphenylphosphino)naphthalene: a rigid chelating, diphosphine analogue of proton sponge (R.D. Jackson, S. James, A.G. Orpen and P.G. Pringle), C3

Rhodium

New synthesis of acylferrocene by hydroiminoacetylation of the terminal olefin with ferrocenecarboxaldimine and application to polymer-supported acylferrocene (C.-H. Jun, J.-B. Kang and J.-Y. Kim), 193

Wolff rearrangement in a transition-metal diazoalkane complex. Synthesis of the first η^1 -ketenyl rhodium complex $[\text{Rh}(\text{C}(\text{CO})\text{SiMe}_3)(\text{CO})(\text{PET}_3)_2]$ (E. Deydier, M.-J. Menu, M. Dartiguenave and Y. Dartiguenave), 225

Ligandeneffekte bei der Synthese der Komplexe $\text{ClRh}(\text{P} \sim \text{O})\text{P} \sim \text{O}$, $[\text{Rh}(\text{P} \sim \text{O})_2\text{BPh}_4]$ und $[\text{Rh}(\text{COD})(\text{P} \sim \text{O})]\text{BPh}_4$ (E. Lindner, Q. Wang, H.A. Mayer and A. Bader), 229

Ruthenium

Ruthenium carbonyl cluster complexes with nitrogen ligands. II. Syntheses, X-ray structures and interconversion of $\text{Ru}_3(\mu-\text{H})(\mu-\eta^2-\text{NC}_5\text{H}_8)(\text{CO})_{10}$ and $\text{Ru}_3(\mu-\text{H})(\mu_3-\eta^2-\text{NC}_5\text{H}_8)(\text{CO})_9$ (M.P. Ci-fuentes, M.G. Humphrey, B.W. Skelton and A.H. White), 211

(η^6 -Aren)(η^4 -buta-1,3-dien)ruthenium(0)-Komplexe: Synthese und Eigenschaften sowie Kristallstruktur von (Benzol)(butadien)ruthenium (J. Müller, K. Qiao, M. Siewing und B. Westphal), 219
Insertion of diphenylacetylene into the carbon-sulfur bond of phenylisothiocyanate at a diruthenium centre: synthesis and X-ray structure of $[\text{Cp}_2\text{Ru}_2(\text{CO})(\mu-\text{CO})(\mu-\text{SC}(\text{Ph})=\text{C}(\text{Ph})\text{C}(=\text{NPh}))]$ (G. Hogarth and E. Skordalakis), C8

Samarium

Studies on organolanthanoid complexes. LI. Syntheses of bis- and tris(2-methoxyethylcyclopentadienyl) lanthanoid complexes and crystal structures of bis(2-methoxyethylcyclopentadienyl)ytterbium iodide and tris(2-methoxyethylcyclopentadienyl)samarium (D. Deng, C. Qian, F. Song, Z. Wang, G. Wu, P. Zheng, S. Jin and Y. Lin), 83

Silicon

Lineare Tetrasilane mit mittelständigen Substituenten—Oligosilane mit optischer Aktivität (H. Stüger), 1

New cyclodisilazane monomers (E. Duguet, M. Schappacher and A. Soum), 9

Synthesis of acetyltrimethyl(phenyl)silane and its enantioselective conversion into (*R*)-(1-hydroxyethyl)dimethyl(phenyl)silane by plant cell suspension cultures of *Sympyrum officinale* L. and *Ruta graveolens* L. (R. Tacke, S.A. Wagner, S. Brakmann, F. Wuttke, U. Eilert, L. Fischer and C. Syldatk), 13

η^7 -Cycloheptatrienyl-carbonylvanadium-Komplexe mit mono- und bifunktionellen Phosphan-Liganden. Eine ^{51}V -NMR-spektroskopische Untersuchung (M. Herberhold und M. Schrepfermann), 19
Activation of the carbon–carbon bond in the Mn-mediated cycloaddition reaction between disilacyclobutene and cyclic conjugated dienes (H.-J. Chiang and C.-S. Liu), 167

N-alkyldiene[bis(trimethylsilyl)methyl]amines: a new series of stable imines (A. Capperucci, A. Ricci, G. Seconi, J. Dunogues, S. Grelier, J.-P. Picard, C. Palomo and J.-M. Aizpurua), C1

Tin

Utilization of hypervalently activated organotin compounds in synthesis. Preparation and reactions of $\text{Me}_2\text{N}(\text{CH}_2)_3\text{SnPh}_3$ (D. Dakternieks, G. Dyson, K. Jurkschat, R. Tozer and E.R.T. Tiekkink), 29

Tin(IV) and organotin(IV) complexes containing the anion of some substituted-3-methyl-4-acyl-5-pyrazolones. Crystal structure of dimethylbis(1-phenyl-3-methyl-4-benzoyl pyrazolone-5-ato)tin(IV) (B. Bovio, A. Cingolani, F. Marchetti and C. Pettinari), 39

A compound with a tin–carbon double bond, bis(2,4,6-trisopropylphenyl)(fluorenylidene)stannane: aspects of its reactivity and X-ray structure of its dimer (G. Anselme, J.-P. Declercq, A. Dubourg, H. Ranaivojatovo, J. Escudé and C. Couret), 49

Polysulfonylamine. XLVII. Darstellung und Kristallstrukturanalyse der ionischen Verbindung catena-Poly(μ -4,4'-bipyridin)trimesitylzinno-dimesylamid-Acetonitril (1/1) (I. Lange, E. Wieland, P.G. Jones und A. Blaschette), 57

Titanium

Neue Synthesen und Reaktionen ungesättigter Heterotitanacyclen (S. Dürr, U. Höhlein und R. Schobert), 89

Tungsten

The synthesis of ring-substituted cycloheptatrienyl complexes $[\text{M}(\text{CO})_3(\eta^7\text{-C}_7\text{H}_6\text{R})]^+$ ($\text{M} = \text{Cr}, \text{Mo}$ or W ; $\text{R} = \text{Me}, ^1\text{Bu}, \text{C}_6\text{H}_4\text{-F}-\text{p}$ or $\text{C}\equiv\text{CPh}$) (A. Roberts and M.W. Whitley), 131

Metal-ligand bonding in $\text{LM}(\text{CO})_5$ complexes ($\text{L} = \text{NMe}_3$, PMe_3 , PCl_3 , PBr_3 or AsMe_3 ; $\text{M} = \text{Cr}$, Mo or W) from dielectric, electro-optic and spectroscopic evidence (M.S. Davies, R.K. Pierens and M.J. Aroney), 141

Vanadium

η^7 -Cycloheptatrienyl-carbonylvanadium-Komplexe mit mono- und bifunktionellen Phosphan-Liganden. Eine ^{51}V -NMR-spektroskopische Untersuchung (M. Herberhold und M. Schrepfermann), 19
Structure of a metallacyclic metaloxycarbene vanadium complex pre-

pared from $\text{CpV}(\text{CO})_4$ (M. Berlekamp, G. Erker and J.L. Petersen), 97

Ytterbium

Studies on organolanthanoid complexes. LI. Syntheses of bis- and tris(2-methoxyethylcyclopentadienyl) lanthanoid complexes and crystal structures of bis(2-methoxyethylcyclopentadienyl)ytterbium iodide and tris(2-methoxyethylcyclopentadienyl)samarium (D. Deng, C. Qian, F. Song, Z. Wang, G. Wu, P. Zheng, S. Jin and Y. Lin), 83