

Subject Index of Volume 454

Aluminium

Synthesis and molecular structure of $\text{AlMe}[\text{O}=\text{C}(\text{OMe})\text{C}_6\text{H}_4-\text{o}-\text{O}]_2$. The first structurally characterized five-coordinated simple alkyl-aluminium chelate complex (J. Lewiński, J. Zachara, B. Mańk and S. Pasynkiewicz), 5
 Dialan-Radikal-anionen $[\text{R}_2\text{Al}-\text{AlR}_2]^-$ (W. Uhl, A. Vester, W. Kaim und J. Poppe), 9

Antimony

On tertiary stibine. I. Reaction of ω -bromoacetophenone with aldehydes mediated by diphenylantimonyorganometallic reagents (L.-J. Zhang and Y.-Z. Huang), 101

Chromium

α -Addition von Aminen, Iminen und Hydrazinen an Allenyliden-Komplexe — Bildung von Carben-, Azetidinylen- und Nitril-Komplexen (H. Fischer, G. Roth, D. Reindl und C. Troll), 133
 Dimethylenyclobuten als Komplexligand. Synthese und Struktur von Tricarbonyl(η^6 -diisopropylidencyclobuten)chrom und Bis(η^2 -dimethylenyclobuten)(η^5 -indenyl)rhodium (D. Rau und U. Behrens), 151

Cobalt

Bidentate ligand substitution in $\text{PhCCo}_3(\text{CO})_9$. Synthesis, molecular structure, and redox reactivity of $\text{PhCCo}_3(\text{CO})_7(\text{cis}-\text{Ph}_2\text{PCH}=\text{CHPPh}_2)$ (K. Yang, S.G. Bott and M.G. Richmond), 273

Copper

Studies on binuclear complexes. Synthesis and catalytic behaviour of Fe-M complexes (M = Mo, Ni, Rh, Cu, Hg) using *trans*-Fe(Ph_2Ppy)₂(CO)₃ [Ph_2Ppy = 2-(diphenylphosphino)pyridine] as a neutral tridentate ligand. Molecular structure of $\text{FeHg}(\mu\text{-Ph}_2\text{Ppy})_2(\text{CO})_3(\text{SCN})_2$ (Z.-Z. Zhang, H.-P. Xi, W.-J. Zhao, K.-Y. Jiang, R.-J. Wang, H.-G. Wang and Y. Wu), 221

Germanium

An easy route to 1-germaindenes via a transmetalation zirconium-germanium reaction (A. Kanj, P. Meunier, B. Gautheron, J. Dubac and J.-C. Daran), 51

Gold

Substituted metal carbonyls. XXII. Synthesis of 1,1'-bis(diphenylphosphino)ferrocene-bridged complexes with two heterometallic Au-M (M = Mn and Re) bonds (P.M.N. Low, Y.K. Yan, H.S.O. Chan and T.S.A. Hor), 205

Iron

Übergangsmetall-Silyl-Komplexe. XLVII. Deprotonierung der elektronenreichen Hydrido-Silyl-Komplexe $\text{Fe}(\text{CO})_2(\text{dppe})(\text{H})(\text{SiR}_3)$ und $\text{Fe}(\text{CO})(\text{dppe})\text{H}_3(\text{SiR}_3)$; Darstellung der Komplexe $\text{Fe}(\text{CO})_2(\text{dppe})(\text{EMe}_3)(\text{SiR}_3)$ und $\text{Fe}(\text{CO})(\text{dppe})\text{H}_2(\text{EMe}_3)(\text{SiR}_3)$ (E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79

($\text{CO})_2(\text{dppe})(\text{EMe}_3)(\text{SiR}_3)$ und $\text{Fe}(\text{CO})(\text{dppe})\text{H}_2(\text{EMe}_3)(\text{SiR}_3)$ (E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79
 Substituted metal carbonyls. XXII. Synthesis of 1,1'-bis(diphenylphosphino)ferrocene-bridged complexes with two heterometallic Au-M (M = Mn and Re) bonds (P.M.N. Low, Y.K. Yan, H.S.O. Chan and T.S.A. Hor), 205
 The use of high reaction pressures in the synthesis of acetylide complexes of iron (L.D. Field and A.V. George), 217
 Studies on binuclear complexes. Synthesis and catalytic behaviour of Fe-M complexes (M = Mo, Ni, Rh, Cu, Hg) using *trans*-Fe(Ph_2Ppy)₂(CO)₃ [Ph_2Ppy = 2-(diphenylphosphino)pyridine] as a neutral tridentate ligand. Molecular structure of $\text{FeHg}(\mu\text{-Ph}_2\text{Ppy})_2(\text{CO})_3(\text{SCN})_2$ (Z.-Z. Zhang, H.-P. Xi, W.-J. Zhao, K.-Y. Jiang, R.-J. Wang, H.-G. Wang and Y. Wu), 221
 Synthesis and molecular structures of $\text{S}_2\text{Fe}_2(\text{CO})_6$ heterometallic derivatives, containing (Ph_3P)₂Pt or CpRe(CO) fragments (A.A. Pasynskii, B.I. Kolobkov, S.E. Nefedov, I.L. Eremenko, E.S. Koltun, A.I. Yanovsky and Y.T. Struchkov), 229
 The use of $[(\text{C}_6\text{H}_7)\text{Fe}(\text{CO})_3]^+$ as a recoverable protecting group in peptide synthesis (S. Fu, J.A. Carver and L.A.P. Kane-Maguire), C11

Lead

Übergangsmetall-Silyl-Komplexe. XLVII. Deprotonierung der elektronenreichen Hydrido-Silyl-Komplexe $\text{Fe}(\text{CO})_2(\text{dppe})(\text{H})(\text{SiR}_3)$ und $\text{Fe}(\text{CO})(\text{dppe})\text{H}_3(\text{SiR}_3)$; Darstellung der Komplexe $\text{Fe}(\text{CO})_2(\text{dppe})(\text{EMe}_3)(\text{SiR}_3)$ und $\text{Fe}(\text{CO})(\text{dppe})\text{H}_2(\text{EMe}_3)(\text{SiR}_3)$ (E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79

Lutetium

Metalcarbonylates of lanthanides. Unexpected formation of the tri-nuclear cluster $[\text{Na}(\text{DME})_3]_2[\text{W}_3(\text{CO})_{14}]$ (I.P. Beletskaya, A.Z. Voskoboynikov, E.B. Chuklanova, A.I. Gusev and A.V. Kisim), 1

Magnesium

A titanium-magnesium complex containing perpendicularly bridging bis(trimethylsilyl)acetylene ligands: $[(\eta^5\text{C}_5\text{H}_5)\text{Ti}][(\eta^5\text{C}_5\text{H}_5)\text{Mg}][\mu\text{-}\eta^2\text{:}\eta^2\text{-C}_2(\text{SiMe}_3)_2]_2$ (V. Varga, K. Mach, G. Schmid and U. Thewalt), C1

Manganese

Synthese und Struktur zweikerniger Cyclopentadienyltricarbonyl-mangan-Komplexe vom Typ $(\text{CO})_3\text{Mn}[(\eta^5\text{C}_5\text{H}_4)-\text{X}-(\eta^5\text{C}_5\text{H}_4)]\text{Mn}(\text{CO})_3$ (S. Pitter, G. Huttner, O. Walter und L. Zsolnai), 183

Photochemische Reaktionen von Übergangsmetall-Organyl-Komplexen mit Olefinen. VII. Photoreaktionen von Tricarbonyl- η^5 -dienyl-mangan-Komplexen mit 6,6-Dimethylfulven (C.G. Kreiter und K. Lehr), 199

Substituted metal carbonyls. XXII. Synthesis of 1,1'-bis(diphenylphosphino)ferrocene-bridged complexes with two heterometallic

Au–M (M = Mn and Re) bonds (P.M.N. Low, Y.K. Yan, H.S.O. Chan and T.S.A. Hor), 205

Mercury

Studies on binuclear complexes. Synthesis and catalytic behaviour of Fe–M complexes (M = Mo, Ni, Rh, Cu, Hg) using *trans*-Fe(Ph₂Ppy)₂(CO)₃ [Ph₂Ppy = 2-(diphenylphosphino)pyridine] as a neutral tridentate ligand. Molecular structure of FeHg(μ-Ph₂Ppy)₂(CO)₃(SCN)₂ (Z.-Z. Zhang, H.-P. Xi, W.-J. Zhao, K.-Y. Jiang, R.-J. Wang, H.-G. Wang and Y. Wu), 221

Metallocenes

Preparation and structure of bis(η^5 -methylcyclopentadienyl)titanaindane (G. Erker, U. Korek and A.L. Rheingold), 113
 New metallocene complexes of Nb and Ta with functionalized diazoalkanes (D.A. Lemenovskii, M. Putala, G.I. Nikonorov, N.B. Kazennova, D.S. Yufit and Y.T. Struchkov), 123
 Formation of [cp₂Zr(C≡C-SiMe₃)₂] from the reaction of zirconocene with trimethylsilylacetylene (N. Metzler and H. Nöth), C5

Molybdenum

Organometallic chemistry of diphosphazanes. VIII. Synthesis, spectroscopic studies and crystal structure of *fac*-[Mo(CO)₃-(MeCN)(Ph₂PN('Pr)P(Ph)DMMP)], and its reactions with tertiary phosphines (R.P.K. Babu, S.S. Krishnamurthy and M. Nethaji), 157

Photoinduzierte Desalkylierungsreaktionen an Indenyl- und Fluorenyl-funktionalisierten Cyclopentadienyltricarbonyl-Methylkomplexen des Molybdäns und Wolframs. Molekulstrukturen von (η^5 : η^1 -C₅H₄CMe₂C₉H₆)Mo(CO)₃, (η^5 : η^1 -C₅H₄CMe₂C₁₃-H₈)W(CO)₃ und [η^5 -C₅H₄(CH₂)₃C₁₃H₉]W(CO)₃₂ (H.G. Alt, J.S. Han und R.D. Rogers), 165

Studies on binuclear complexes. Synthesis and catalytic behaviour of Fe–M complexes (M = Mo, Ni, Rh, Cu, Hg) using *trans*-Fe(Ph₂Ppy)₂(CO)₃ [Ph₂Ppy = 2-(diphenylphosphino)pyridine] as a neutral tridentate ligand. Molecular structure of FeHg(μ-Ph₂Ppy)₂(CO)₃(SCN)₂ (Z.-Z. Zhang, H.-P. Xi, W.-J. Zhao, K.-Y. Jiang, R.-J. Wang, H.-G. Wang and Y. Wu), 221

Nickel

Catalysis of hydrosilylation. XXIII. Effect of substituents at silicon on unusual hydrosilylation of vinylsilanes catalysed by nickel acetylacetone (B. Marciniec and H. Maciejewski), 45

Studies on binuclear complexes. Synthesis and catalytic behaviour of Fe–M complexes (M = Mo, Ni, Rh, Cu, Hg) using *trans*-Fe(Ph₂Ppy)₂(CO)₃ [Ph₂Ppy = 2-(diphenylphosphino)pyridine] as a neutral tridentate ligand. Molecular structure of FeHg(μ-Ph₂Ppy)₂(CO)₃(SCN)₂ (Z.-Z. Zhang, H.-P. Xi, W.-J. Zhao, K.-Y. Jiang, R.-J. Wang, H.-G. Wang and Y. Wu), 221

Übergangsmetallallyle. VII. Synthese von η^3 -Allylnickel- und η^3 -Allylpalladiumkomplexen mit Trifluoracetatgruppen als Brückenliganden (R. Goddard, C. Krüger, R. Mynott, M. Neumann und G. Wilke), C20

Niobium

Von Cp⁺₂Nb(η^2 -S₂)H zu Cp⁺₃Nb₃S₁₂: Reaktivitätsstudien an Niobocensulfiden (H. Brunner, G. Gehart, W. Meier, J. Wachter und B. Nuber), 117

New metallocene complexes of Nb and Ta with functionalized diazoalkanes (D.A. Lemenovskii, M. Putala, G.I. Nikonorov, N.B. Kazennova, D.S. Yufit and Y.T. Struchkov), 123

Osmium

Stabile Allyl(hydrido)osmium-Komplexe ohne Olefin- und Arenliganden (C. Schlünken und H. Werner), 243

Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXVI. Tetrakis(2-methylpropyl)oxoosmium(VI), ein thermisch stabiles, sterisch überladenes d²-Alkylmetalloid mit β -ständigem Wasserstoff (W.A. Herrmann, S.J. Eder und W. Scherer), 257

Preparation of [Os₃(CO)₁₁]²⁻ and its reactions with Os₃(CO)₁₂: structures of [Et₄N][HOS₃(CO)₁₁] and H₂Os₃(CO)₁₃ (J.A. Krause, U. Siriwardane, T.A. Salupo, J.R. Werner, D.W. Knoepfel and S.G. Shore), 263

Palladium

Übergangsmetallallyle. VII. Synthese von η^3 -Allylnickel- und η^3 -Allylpalladiumkomplexen mit Trifluoracetatgruppen als Brückenliganden (R. Goddard, C. Krüger, R. Mynott, M. Neumann und G. Wilke), C20

Phosphorus

Übergangsmetall-Silyl-Komplexe. XLVII. Deprotonierung der elektronenreichen Hydrido-Silyl-Komplexe Fe(CO)₂(dppe)H(SiR₃) und Fe(CO)₂(dppe)H₃(SiR₃): Darstellung der Komplexe Fe(CO)₂(dppe)(EMe₃)(SiR₃) und Fe(CO)₂(dppe)H₂(EMe₃)(SiR₃) (E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79

Enantioselektive Katalyse. LXXXI. Optisch aktive Zweischalenphosphine (H. Brunner, J. Fürst und J. Ziegler), 87

Asymmetrische Katalyse. III. Synthese, spektroskopische Untersuchungen und katalytische Wirkung eines neuen chiralen Phosphanliganden (A. Tillack, M. Michalik, D. Fenske und H. Goessmann), 95

Stabile Allyl(hydrido)osmium-Komplexe ohne Olefin- und Arenliganden (C. Schlünken und H. Werner), 243

Platinum

Synthesis and molecular structures of S₂Fe₂(CO)₆ heterometallic derivatives, containing (Ph₃P)₂Pt or CpRe(CO) fragments (A.A. Pasynskii, B.I. Kolobkov, S.E. Nefedov, I.L. Eremenko, E.S. Koltun, A.I. Yanovsky and Y.T. Struchkov), 229

Rhenium

Syntheses and reactions of metal carbonyl complexes containing the trimethylsilylmethyl ligand: CO insertion reactions assisted by the trimethylsilylmethyl groups (J.-D. Chen, C.-K. Wu, I.-Y. Wu, B.-C. Huang, Y.-C. Lin and Y. Wang), 173

Substituted metal carbonyls. XXII. Synthesis of 1,1'-bis(diphenylphosphino)ferrocene-bridged complexes with two heterometallic Au–M (M = Mn and Re) bonds (P.M.N. Low, Y.K. Yan, H.S.O. Chan and T.S.A. Hor), 205

Synthesis of the pseudohalide-dinitrogen complexes *trans*-[ReX-(N₂)(Ph₂PCH₂CH₂PPh₂)₂] (X = NCS, NCO or N₃) and crystal structures of the isothiocyanate and azide complexes (Y. Wang, J.J.R. Frausto Da Silva, A.J.L. Pombeiro, M.A. Pellinghelli and A. Tiripicchio), 211

Synthesis and molecular structures of S₂Fe₂(CO)₆ heterometallic derivatives, containing (Ph₃P)₂Pt or CpRe(CO) fragments (A.A. Pasynskii, B.I. Kolobkov, S.E. Nefedov, I.L. Eremenko, E.S. Koltun, A.I. Yanovsky and Y.T. Struchkov), 229

Aminocarbyne coupling reactions leading to the bis(amino)acetylene complexes *trans*-[MF(η^2 -MeHNC≡CNHMe)(Ph₃PCH₂CH₂PPh₂)₂][BF₄]⁻ (M = Mo or W) and cleavage of the acetylenic-triple bond. A reformulation, and X-ray structure of the Mo complex

ple bond. A reformulation, and X-ray structure of the Mo compound (J.J.R. Fraústo da Silva, M.A. Pellinghelli, A.J.L. Pombeiro, R.L. Richards, A. Tiripicchio and Y. Wang), C8

Rhodium

Asymmetrische Katalyse. III. Synthese, spektroskopische Untersuchungen und katalytische Wirkung eines neuen chiralen Phosphanliganden (A. Tillack, M. Michalik, D. Fenske und H. Goessmann), 95

Dimethylenecyclobuten als Komplexligand. Synthese und Struktur von Tricarbonyl(η^6 -diisopropylidencyclobuten)chrom und Bis(η^2 -dimethylenecyclobuten)(η^5 -indenyl)rhodium (D. Rau und U. Behrens), 151

Studies on binuclear complexes. Synthesis and catalytic behaviour of Fe-M complexes (M = Mo, Ni, Rh, Cu, Hg) using trans-Fe(Ph₂Ppy)₂(CO)₃ [Ph₂Ppy = 2-(diphenylphosphino)pyridine] as a neutral tridentate ligand. Molecular structure of FeHg(μ -Ph₂Ppy)₂(CO)₃(SCN)₂ (Z.-Z. Zhang, H.-P. Xi, W.-J. Zhao, K.-Y. Jiang, R.-J. Wang, H.-G. Wang and Y. Wu), 221

Syntheses of nitrogen heterocycles by means of amine-directed carbonylation and hydrocarbonylation (Z. Zhang and I. Ojima), 281

Ruthenium

Syntheses and reactions of metal carbonyl complexes containing the trimethylsilylmethyl ligand; CO insertion reactions assisted by the trimethylsilylmethyl groups (J.-D. Chen, C.-K. Wu, I.-Y. Wu, B.-C. Huang, Y.-C. Lin and Y. Wang), 173

Paramagnetic organometallics formed by *o*-metallation of phenolic Schiff bases by ruthenium(III) (P. Ghosh, A. Pramanik, N. Bag, G.K. Lahiri and A. Chakravorty), 237

Reaktionen der Aren(carbonyl)ruthenium-Komplexe [C₆Me₆Ru(CO)Cl₂] und [(Mes)Ru(CO)Cl₂] mit Alkyl- und Allyl-Grignardreagenzien. Die Kristall- und Molekülstruktur von {[(η^5 -C₆Me₆(CH₂CH=CH₂)₂)Ru(exo- η^3 -C₃H₅)(CO)]} (H. Werner, G. Brauers und O. Nürnberg), 247

Regiospecific Cp^{*}Ru⁺ mediated codimerization of conjugated dienes, with non-conjugated 1,5-dienes and its application to catalytic cooligomerization (K. Masuda, H. Ohkita, S. Kurumata and K. Itoh), C13

A ruthenium dihydrogen complex as intermediate in the synthesis of [RuH(CO)₂(MeCO₂)(PⁿBu₃)₂] from [Ru(H)2(CO)₂(PⁿBu₃)₂] (P. Frediani, A. Salvini, M. Bianchi and F. Piacenti), C17

Silicon

Dialan-Radikalationen [R₂Al-AlR₂]⁺ (W. Uhl, A. Vester, W. Kaim und J. Poppe), 9

AM1 and PM3 semiempirical molecular orbital study of silatranes. III. 1-chlorosilatrane (G.I. Csonka and P. Hencsei), 15

Synthesis and crystal structure of *para*-substituted diphenylsilanes of interest for nonlinear optics: sulfonyl acceptor groups (D. Hissink, P.F. van Hutten, G. Hadzioannou and F. van Bolhuis), 25

Siliciumverbindungen mit starken intramolekularen sterischen Wechselwirkungen. L. Reaktionen von Silylenen und Disilenen mit 2,2'-Bipyridyl, Pyridin-2-aldiminen und α -Ketoiminen: Cycloadditionen *versus* C-H-Insertion (M. Weidenbruch, H. Piel, A. Lesch, K. Peters und H.G. von Schnerring), 35

Catalysis of hydrosilylation. XXIII. Effect of substituents at silicon on unusual hydrosilylation of vinylsilanes catalysed by nickel acetylacetone (B. Marciniec and H. Maciejewski), 45

Übergangsmetall-Silyl-Komplexe. XLVII. Deprotonierung der elektronenreichen Hydrido-Silyl-Komplexe Fe(CO)₂(dppe)(H)(SiR₃) und Fe(CO)(dppe)H₃(SiR₃); Darstellung der Komplexe Fe(CO)₂(dppe)(EMe₃)(SiR₃) und Fe(CO)(dppe)H₂(EMe₃)(SiR₃) (E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79

(E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79

Synthesis and characterization of ansa-dimethylsilylbiscyclopentadienyl titanium(II) complexes. Crystal structure of [Ti{Me₂Si(C₅H₄)₂}CN(2,6-Me₂C₆H₃)₂] (T. Cuenca, R. Gómez, P. Gómez-Sal and P. Royo), 105

Formation of [cp₂Zr(C≡C-SiMe₃)₂] from the reaction of zirconocene with trimethylsilylacetylene (N. Metzler and H. Nöth), C5

Sodium

Metalcarbylates of lanthanides. Unexpected formation of the tri-nuclear cluster [Na(DME)₃]₂[W₃(CO)₁₄] (I.P. Beletskaya, A.Z. Voskoboinikov, E.B. Chuklanova, A.I. Gusev and A.V. Kisim), 1

Tantalum

New metallocene complexes of Nb and Ta with functionalized diazoalkanes (D.A. Lemenovskii, M. Putala, G.I. Nikonor, N.B. Kazennova, D.S. Yufit and Y.T. Struchkov), 123

Tin

Tin(IV) and organotin(IV) complexes with heterocyclic β -diketonates. I. Bis-[4(1-phenyl-3-methylpyrazol-5-one)]-dioxaalkane derivatives (C. Pettinari, F. Marchetti, A. Cingolani, D. Leonesi, G. Gioia Lobbia and A. Lorenzotti), 59

A complex salt formed in reactions of ⁷BuSnCl₃ with 1-(2-methyl-2,3-dihydrobenzothiazol-2-yl)-2-ketones (S.-B. Teo, S.-G. Teoh, R.C. Okechukwu and H.-K. Fun), 67

The structures of the adducts formed from organotin moieties and 1-[(4-methoxyphenylimino)methyl]-2-naphthol. Crystal structure of chloro-trimethyl[1-[(4-methoxyphenylimino)methyl]-2-naphthol]tin(IV): Me₃SnCl · [HOC₁₀H₆CH : NC₆H₄OCH₃] (S.-G. Teoh, S.-B. Teo, G.-Y. Yeap, H.-K. Fun and P.J. Smith), 73
Übergangsmetall-Silyl-Komplexe. XLVII. Deprotonierung der elektronenreichen Hydrido-Silyl-Komplexe Fe(CO)₂(dppe)(H)(SiR₃) und Fe(CO)(dppe)H₃(SiR₃); Darstellung der Komplexe Fe(CO)₂(dppe)(EMe₃)(SiR₃) und Fe(CO)(dppe)H₂(EMe₃)(SiR₃) (E = Sn, Pb) aus den korrespondierenden anionischen Komplexen (U. Schubert, S. Gilbert und M. Knorr), 79

Titanium

Synthesis and characterization of ansa-dimethylsilylbiscyclopentadienyl titanium(II) complexes. Crystal structure of [Ti{Me₂Si(C₅H₄)₂}CN(2,6-Me₂C₆H₃)₂] (T. Cuenca, R. Gómez, P. Gómez-Sal and P. Royo), 105

Preparation and structure of bis(η^5 -methylcyclopentadienyl)titanaindane (G. Erker, U. Korek and A.L. Rheingold), 113

A titanium-magnesium complex containing perpendicularly bridging bis(trimethylsilyl)acetylene ligands: [(η^5 -C₅H₅)Ti][(η^5 -C₅H₅)-Mg] $[\mu\text{-}\eta^2:\eta^2\text{-C}_2(\text{SiMe}_3)_2]$ (V. Varga, K. Mach, G. Schmid and U. Thewalt), C1

Tungsten

Metalcarbylates of lanthanides. Unexpected formation of the tri-nuclear cluster [Na(DME)₃]₂[W₃(CO)₁₄] (I.P. Beletskaya, A.Z. Voskoboinikov, E.B. Chuklanova, A.I. Gusev and A.V. Kisim), 1

α -Addition von Aminen, Iminen und Hydrazinen an Allenyliden-Komplexe — Bildung von Carben-, Azetidinylen- und Nitril-Komplexen (H. Fischer, G. Roth, D. Reindl und C. Troll), 133

Photoinduzierte Desalkylierungsreaktionen an Indenyl- und Fluoren-funktionalisierten Cyclopentadienyltricarbonyl-Methylkomplexen des Molybdäns und Wolframs. Molekülstrukturen von

$(\eta^5:\eta^1\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_9\text{H}_6)\text{Mo}(\text{CO})_3$, $(\eta^5:\eta^1\text{-C}_5\text{H}_4\text{CMe}_2\text{C}_{13}\text{H}_8)\text{W}(\text{CO})_3$ und $[(\eta^5\text{-C}_5\text{H}_4\text{CH}_2)_3\text{C}_{13}\text{H}_9]\text{W}(\text{CO})_3$ (H.G. Alt, J.S. Han und R.D. Rogers), 165

Syntheses and reactions of metal carbonyl complexes containing the trimethylsilylmethyl ligand; CO insertion reactions assisted by the trimethylsilylmethyl groups (J.-D. Chen, C.-K. Wu, L.-Y. Wu, B.-C. Huang, Y.-C. Lin and Y. Wang), 173

Zirconium

An easy route to 1-germaindenes via a transmetallation zirconium–germanium reaction (A. Kanj, P. Meunier, B. Gautheron, J. Dubac and J.-C. Daran), 51

Formation of $[\text{cp}_2\text{Zr}(\text{C}\equiv\text{C-SiMe}_3)]_2$ from the reaction of zirconocene with trimethylsilylacetylene (N. Metzler and H. Nöth), C5

Book review

Comprehensive Handbook on Hydrosilylation. B. Marciniec (ed.) (C. Eaborn), C26