

Subject Index of Volume 470

Acetyl

Preparation and Mössbauer spectroscopic studies of 1',1''-diacetyl- and 1'-acetyl-1''-(1-naphthylimino-ethyl)biferrocenes and their partially oxidized biferrocenium salts (H.-H. Wei, I.-S. Hwang, M.-C. Cheng and Y. Wang), 161

Alkene

Elimination of β -palladium hydroxide in the Pd^{II}-catalyzed reaction of methyl (α -hydroxymethyl)acrylate with alcohols (T. Hosokawa, T. Sugafuji, T. Yamanaka and S.-I. Murahashi), 253

Alkoxide

Synthesis of bimetallic[Sn^{IV},Al^{III}]- μ -oxoisopropoxyacetate and μ -oxoisopropoxide and their acetylacetone and benzoylacetone derivatives (Sonika, A.K. Narula, O.P. Vermani, H.K. Sharma and A.S. Sarpal), 67

Alkoxyacetylene

A convenient synthesis of hexacarbonyldicobalt complexes of chiral (non-racemic) terminal alkoxyacetylenes (V. Bernardes, X. Verdager, A. Moyano, M.A. Pericàs, A. Riera and A.E. Greene), C12

Alkylidene

The reaction of (μ -H)₂Os₃(CO)₉(PPh₃) with ethylene affords the ethylidene complex (μ -H)₂Os₃(CO)₉(PPh₃)(μ -CHCH₃) (M. Koike and J.R. Shapley), 199

Alkyne

Ein Beitrag zur Darstellung und Reaktivität der η^3 -Allyliridium-Komplexe [Ir(η^3 -2-RC₃H₄)(PⁱPr₃)₂] (M. Schulz und H. Werner), 243

Reaction of sodium amalgam with (μ -enynne)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum(I)) and (μ - η^2 , η^3 -allenyl)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum) tetrafluoroborate complexes. Crystal structure of [(Mo(η^5 -C₅H₅)(CO)₂)(μ -HC≡CCH(CH₂CH₃))]₂ (J.F. Capon, S. Cornen, N. Le Berre-Cosquer, R. Pichon, R. Kergoat and P. L'Haridon), 137

Synthesis and characterisation of monomeric, dimeric and polymeric ferrocenylacetylides. Crystal structure of 1,1'-bis(phenylethynyl)ferrocene (S.L. Ingham, M.S. Khan, J. Lewis, N.J. Long and P.R. Raithby), 153

Alkynyl

Unusual stabilization of cationic "M(η^3 -allyl)⁺ⁿ (M = Pt, Pd) units by a dianionic *cis*-{Pt(C₆F₅)₂(C≡CSiMe₃)₂}²⁻ fragment (J.R. Berenguer, J. Fornies, E. Lalinde and F. Martínez), C15

Allyl

Ein Beitrag zur Darstellung und Reaktivität der η^3 -Allyliridium-Komplexe [Ir(η^3 -2-RC₃H₄)(PⁱPr₃)₂] (M. Schulz und H. Werner), 243

Aluminium

Synthesis of bimetallic[Sn^{IV},Al^{III}]- μ -oxoisopropoxyacetate and μ -oxoisopropoxide and their acetylacetone and benzoylacetone derivatives (Sonika, A.K. Narula, O.P. Vermani, H.K. Sharma and A.S. Sarpal), 67

Amine

Reactivity of mono- and di-methoxycarbonyl complexes of Pd^{II} towards amines and copper amine complexes: role of copper

in the catalyzed palladium-copper oxidative carbonylation of amines (P. Giannoccaro), 249

Amino acid

η^5 -Pentamethylcyclopentadienylruthenium(II) complexes containing η^6 -coordinated α -amino acids (W.S. Sheldrick and A. Gleichmann), 183

Aniline

Etude conformationnelle du complexe η^6 -(4-(trifluorométhoxy)-aniline)tricarbonylchrome (F. Rose-Munch, R. Khourzom, J.-P. Djukic, E. Rose, B. Langlois et J. Vaisserman), 131

Annulene

Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17,18:21,22-hexabenzocycloheptatriene, 1,2:5,6:9,10:13,14-tetrabenzocycloheptatriene (HBC), 1,2:5,6:9,10:13,14-tetrabenzocycloheptatriene (QBC) and a tetracobalt complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231

Antimony

Strukturen und Reaktionen von Methylantimonidihalogeniden und Versuche zur Darstellung von Methylantimon (H.J. Breunig, M. Denker und K.H. Ebert), 87

Arene

Direct synthesis of (η^6 -arene)- tricarbonylchromium complexes by arene displacement from tricarbonyl(η^5 -1-methylpyrrole)chromium(0) (A. Goti and M.F. Semmelhack), C4

Etude conformationnelle du complexe η^6 -(4-(trifluorométhoxy)-aniline)tricarbonylchrome (F. Rose-Munch, R. Khourzom, J.-P. Djukic, E. Rose, B. Langlois et J. Vaisserman), 131

Synthesis and derivative chemistry of [Ru₂(μ -PPh₂)(μ -OH)₂(η^6 -*p*-cymene)]⁺ (J.A. Cabeza, F. Mulla and V. Riera), 173

Asymmetric synthesis

A convenient synthesis of hexacarbonyldicobalt complexes of chiral (non-racemic) terminal alkoxyacetylenes (V. Bernardes, X. Verdager, A. Moyano, M.A. Pericàs, A. Riera and A.E. Greene), C12

Bismuth

Syntheses and structures of dimesitylbismuth(III) bromide, Mes₂BiBr, and bis(diphenyldithiophosphinato)mesitylbismuth(III), MesBi(S₂PPh₂)₂ (K.H. Ebert, R.E. Schulz, H.J. Breunig, C. Silvestru and I. Haiduc), 93

Bond activation

Thermally and photochemically induced vinyl-hydrogen activation of [η^4 -1,2:3,4-(*trans*-1,3,5-hexatriene)](η^5 -cyclopentadienyl)cobalt: regio- and stereospecific hydrogen migrations (J.A. King Jr. and K.P.C. Vollhardt), 207

π -Bonding

η^5 -Pentamethylcyclopentadienylruthenium(II) complexes containing η^6 -coordinated α -amino acids (W.S. Sheldrick and A. Gleichmann), 183

Bond nature

Relative stability of η^5 -cyclopentadienyl(olefin)ruthenium cations: the first estimation of the nature of the Ru^{II}-olefin bond (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 189

Boracycloalkene

Reaktion von konjugierten Doppelbindungssystemen mit Dihalo-(diorganyl-amino)boranen und Na/K-Legierung (W. Maringgele, S. Dielkus, R. Herbst-Irmer, H. Michaelsen und A. Meller), 23

Borane

Reaktion von konjugierten Doppelbindungssystemen mit Dihalo-(diorganyl-amino)boranen und Na/K-Legierung (W. Maringgele, S. Dielkus, R. Herbst-Irmer, H. Michaelsen und A. Meller), 23

Boron

Reaktion von konjugierten Doppelbindungssystemen mit Dihalo-(diorganyl-amino)boranen und Na/K-Legierung (W. Maringgele, S. Dielkus, R. Herbst-Irmer, H. Michaelsen und A. Meller), 23

Boron-nitrogen compounds

Reaktion von konjugierten Doppelbindungssystemen mit Dihalo-(diorganyl-amino)boranen und Na/K-Legierung (W. Maringgele, S. Dielkus, R. Herbst-Irmer, H. Michaelsen und A. Meller), 23

Boroxin

Can triorganoboroxins exist in a "monomeric" R-B=O form? MNDO calculations and ebulliometric molecular weight determination (A. Sporzyski and H. Szatyłowicz), 31

Bridging ligand

Synthesis and derivative chemistry of $[\text{Ru}_2(\mu\text{-PPh}_2)(\mu\text{-OH})_2(\eta^6\text{-}p\text{-cymene})_2]^+$ (J.A. Cabeza, F. Mulla and V. Riera), 173

Bromide

Syntheses and structures of dimesitylbismuth(III) bromide, Mes_2BiBr , and bis(diphenyldithiophosphinato)mesitylbismuth(III), $\text{MesBi}(\text{S}_2\text{PPh}_2)_2$ (K.H. Ebert, R.E. Schulz, H.J. Breunig, C. Silvestru and I. Haiduc), 93

Cage compound

Formation and structure of $[\text{Co}(\text{Ph}_2\text{PN}(\text{i-Bu})\text{PPh}_2\text{-}P, P')_2(\text{CO})][\text{Co}(\text{CO})_4]$ – a cage molecule-pair or an ion-pair complex? (Z.-Z. Zhang, A. Yu, H.-P. Xi, R.-J. Wang and H.-G. Wang), 223

Carbamate

Reactivity of mono- and di-methoxycarbonyl complexes of Pd^{II} towards amines and copper amine complexes: role of copper in the catalyzed palladium-copper oxidative carbonylation of amines (P. Giannoccaro), 249

Carbene

Heterozyklen als Liganden. XX. Spektroskopische und strukturelle Charakterisierung von Pentacarbonylmetall-Komplexen der Imidazol-2-ylidene (N. Kuhn, T. Kratz, R. Boese und D. Bläser), C8

Carbon-13

The reaction of $(\mu\text{-H})_2\text{Os}_3(\text{CO})_9(\text{PPh}_3)$ with ethylene affords the ethylidene complex $(\mu\text{-H})_2\text{Os}_3(\text{CO})_9(\text{PPh}_3)(\mu\text{-CHCH}_3)$ (M. Koike and J.R. Shapley), 199

Carbonyl

A convenient synthesis of hexacarbonyldicobalt complexes of chiral (non-racemic) terminal alkoxyacetylenes (V. Bernardes, X. Verdaguer, A. Moyano, M.A. Pericàs, A. Riera and A.E. Greene), C12

Etude conformationnelle du complexe $\eta^6\text{-}(4\text{-}(\text{trifluorométhoxy})\text{-aniline})\text{tricarbonylchrome}$ (F. Rose-Munch, R. Khourzom, J.-P. Djukic, E. Rose, B. Langlois et J. Vaisserman), 131

Formation and structure of $[\text{Co}(\text{Ph}_2\text{PN}(\text{i-Bu})\text{PPh}_2\text{-}P, P')_2(\text{CO})][\text{Co}(\text{CO})_4]$ – a cage molecule-pair or an ion-pair complex? (Z.-Z. Zhang, A. Yu, H.-P. Xi, R.-J. Wang and H.-G. Wang), 223

Heterozyklen als Liganden. XX. Spektroskopische und strukturelle Charakterisierung von Pentacarbonylmetall-Komplexen der Imidazol-2-ylidene (N. Kuhn, T. Kratz, R. Boese und D. Bläser), C8

Molecular orbital calculations on $\text{Fe}_4(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ (S.N. Datta, R.-K. Kondru and P. Mathur), 169

Carbonylation

Reactivity of mono- and di-methoxycarbonyl complexes of Pd^{II} towards amines and copper amine complexes: role of copper in the catalyzed palladium-copper oxidative carbonylation of amines (P. Giannoccaro), 249

Synthesis of aromatic acids via catalyzed methyl formate-chloroarene reactions (G. Jenner and A. Ben Taleb), 257

Catalysis

Elimination of β -palladium hydroxide in the Pd^{II} -catalyzed reaction of methyl (α -hydroxymethyl)acrylate with alcohols (T. Hosokawa, T. Sugafuji, T. Yamanaka and S.-I. Murahashi), 253

Synthesis of aromatic acids via catalyzed methyl formate-chloroarene reactions (G. Jenner and A. Ben Taleb), 257

Chloroarene

Synthesis of aromatic acids via catalyzed methyl formate-chloroarene reactions (G. Jenner and A. Ben Taleb), 257

Chromium

Direct synthesis of (η^6 -arene)-tricarbonylchromium complexes by arene displacement from tricarbonyl(η^5 -1-methylpyrrole)-chromium(0) (A. Goti and M.F. Semmelhack), C4

Etude conformationnelle du complexe $\eta^6\text{-}(4\text{-}(\text{trifluorométhoxy})\text{-aniline})\text{tricarbonylchrome}$ (F. Rose-Munch, R. Khourzom, J.-P. Djukic, E. Rose, B. Langlois et J. Vaisserman), 131

Strukturen und Reaktionen von Methylantimonodihalogeniden und Versuche zur Darstellung von Methylantimon (H.J. Breunig, M. Denker und K.H. Ebert), 87

Cluster

Molecular orbital calculations on $\text{Fe}_4(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ (S.N. Datta, R.-K. Kondru and P. Mathur), 169

The reaction of $(\mu\text{-H})_2\text{Os}_3(\text{CO})_9(\text{PPh}_3)$ with ethylene affords the ethylidene complex $(\mu\text{-H})_2\text{Os}_3(\text{CO})_9(\text{PPh}_3)(\mu\text{-CHCH}_3)$ (M. Koike and J.R. Shapley), 199

Cobalt

A convenient synthesis of hexacarbonyldicobalt complexes of chiral (non-racemic) terminal alkoxyacetylenes (V. Bernardes, X. Verdaguer, A. Moyano, M.A. Pericàs, A. Riera and A.E. Greene), C12

Formation and structure of $[\text{Co}(\text{Ph}_2\text{PN}(\text{i-Bu})\text{PPh}_2\text{-}P, P')_2(\text{CO})][\text{Co}(\text{CO})_4]$ – a cage molecule-pair or an ion-pair complex? (Z.-Z. Zhang, A. Yu, H.-P. Xi, R.-J. Wang and H.-G. Wang), 223

Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17,18:21,22-hexabenzocycloheptatriene-3,7,11,15,19,23-hexadehydro[24]annulene (HBC), 1,2:5,6:9,10:13,14-tetrabenzocycloheptatriene-3,7,11,15-tetrahydro[16]annulene (QBC) and a tetracobalt complex of QBC. The first example of a transition metal complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231

Thermally and photochemically induced vinyl-hydrogen activation of $[\eta^4\text{-}1,2:3,4\text{-}(\text{trans-}1,3,5\text{-hexatriene})(\eta^5\text{-cyclopentadienyl})\text{cobalt}]$: regio- and stereospecific hydrogen migrations (J.A. King Jr. and K.P.C. Vollhardt), 207

Conformation

NMR conformational studies of (1,2-bis(diarylphosphino)ethane)(diorgano sulfide)ruthenium complexes (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 179

- Zur Struktur des Diethylgermanium-diphenylsiliciumoxids und zur Konformation von Achtringen mit Silicium und Germanium (M. Akkurt, T.R. K ok, P. Faleschini, L. Randaccio, H. Puff und W. Schuh), 59
- Copper**
 Reactivity of mono- and di-methoxycarbonyl complexes of Pd^{II} towards amines and copper amine complexes: role of copper in the catalyzed palladium-copper oxidative carbonylation of amines (P. Giannoccaro), 249
- Crystal structure**
 A ¹⁹⁷Au M ssbauer study of reaction products of trimeric 1-benzyl-2-gold(I)-imidazole leading to Au^I carbene or Au^I imidazolone complexes and trinuclear Au^{III} imidazolyl derivatives. X-Ray crystal structure of [(μ-1-benzylimidazolato-N³,C²)Au]₃I₂] (B. Bovio, S. Calogero, F.E. Wagner, A. Burini and B.R. Pietroni), 275
 Etude conformationnelle du complexe η⁶-(4-(trifluorom thoxy)-aniline)tricarbonylchrome (F. Rose-Munch, R. Khouzom, J.-P. Djukic, E. Rose, B. Langlois et J. Vaisserman), 131
 Heterozyklen als Liganden. XX. Spektroskopische und strukturelle Charakterisierung von Pentacarbonylmetall-Komplexen der Imidazol-2-ylidene (N. Kuhn, T. Kratz, R. Boese und D. Bl aser), C8
 Pentacoordinate silicon compounds: stereochemical non-rigidity of chelates formed by intramolecular ring-closure. Crystal structure of 8-dimethylamino-1-trifluorosilylnaphthalene (F. Carr , R.J.P. Corriu, A. Kpoton, M. Poirier, G. Royo, J.C. Young and C. Belin), 43
 Reaction of sodium amalgam with (μ-enyne)bis(dicarbonyl-η⁵-cyclopentadienylmolybdenum(I)) and (μ-η²,η³-allenyl)bis(dicarbonyl-η⁵-cyclopentadienylmolybdenum) tetrafluoroborate complexes. Crystal structure of [(Mo(η⁵-C₅H₅)(CO)₂]₂(μ-HC≡CCH(CH₂CH₃))₂] (J.F. Capon, S. Cornen, N. Le Berre-Cosquer, R. Pichon, R. Kergoat and P. L'Haridon), 137
 Strukturen und Reaktionen von Methylantimonidihalogeniden und Versuche zur Darstellung von Methylantimon (H.J. Breunig, M. Denker und K.H. Ebert), 87
 Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17,18:21,22-hexabenzo-3,7,11,15,19,23-hexadehydro[24]annulene (HBC), 1,2:5,6:9,10:13,14-tetrabenzo-3,7,11,15-tetradehydro[16]annulene (QBC) and a tetracobalt complex of QBC. The first example of a transition metal complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231
 Syntheses and structures of dimesitylbismuth(III) bromide, Mes₂BiBr, and bis(diphenyldithiophosphinato)mesitylbismuth(III), MesBi(S₂PPh₂)₂ (K.H. Ebert, R.E. Schulz, H.J. Breunig, C. Silvestru and I. Haiduc), 93
 Synthesis and characterisation of monomeric, dimeric and polymeric ferrocenylacetylides. Crystal structure of 1,1'-bis(phenylethynyl)ferrocene (S.L. Ingham, M.S. Khan, J. Lewis, N.J. Long and P.R. Raithby), 153
 The reaction of dihydridotetrakis(triphenylphosphane)ruthenium(II) with methyl acrylate. Crystal structure of bis(methylacrylate)bis(triphenylphosphane) aqua ruthenium(0) (R. Sustmann, B. Patzke and R. Boese), 191
 Zur Struktur des Diethylgermanium-diphenylsiliciumoxids und zur Konformation von Achtringen mit Silicium und Germanium (M. Akkurt, T.R. K ok, P. Faleschini, L. Randaccio, H. Puff und W. Schuh), 59
- Cumulene**
 Delocalization of the unpaired spin density in some niobocene complexes with σ-donor, π-acceptors (A. Anti olo, M. Fajardo, E. De Jes s, Y. Mugnier and A. Otero), 127
- Cycloheptatrienyl**
 Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organo-holmium hydroxide complexes (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99
- Cyclopentadienyl**
 η⁵-Pentamethylcyclopentadienylruthenium(II) complexes containing η⁶-coordinated α-amino acids (W.S. Sheldrick and A. Gleichmann), 183
 Relative stability of η⁵-cyclopentadienyl(olefin)ruthenium cations: the first estimation of the nature of the Ru^{II}-olefin bond (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 189
 The electrochemical reduction of Cp₂TiCl₂ in the presence of trimethylphosphine (I.M.M. Fussing, D. Pletcher and R.J. Whitby), 119
 The synthesis and electrochemistry of CpTiCl₂(OR) (R = alkyl, aryl) complexes (I.M.M. Fussing, D. Pletcher and R.J. Whitby), 109
- Cyclosilazane**
 Cyclotetrasilazane als Precursor f r Cyclotrisilazane und Bis(cyclodisilazan-1-yl)silane (E. Werner und U. Klingebiel), 35
- Cyclcyne**
 Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17,18:21,22-hexabenzo-3,7,11,15,19,23-hexadehydro[24]annulene (HBC), 1,2:5,6:9,10:13,14-tetrabenzo-3,7,11,15-tetradehydro[16]annulene (QBC) and a tetracobalt complex of QBC. The first example of a transition metal complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231
- Dialkylamide**
 Reduction with lithium dialkylamides (M. Majewski and D.M. Gleave), 1
- Diastereoselectivity**
 Direct synthesis of (η⁶-arene)-tricarbonylchromium complexes by arene displacement from tricarbonyl(η⁵-1-methylpyrrole)chromium(0) (A. Goti and M.F. Semmelhack), C4
- Dinuclear**
 Unusual stabilization of cationic "M(η³-allyl)⁺" (M = Pt, Pd) units by a dianionic cis-(Pt(C₆F₅)₂(C≡CSiMe₃)₂)²⁻ fragment (J.R. Berenguer, J. Forni s, E. Lalinde and F. Mart nez), C15
- Diphosphine**
 NMR conformational studies of (1,2-bis(diarylphosphino)ethane)(diorgano sulfide)ruthenium complexes (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 179
- Dithioether**
 Nuclear magnetic resonance studies of sulfur inversion in bis(cyclopentadienyl)-molybdenum and -tungsten complexes with dithioethers (J.R. Ascenso, M. De Deus Carvalho, A.R. Dias, C.C. Rom o, M.J. Calhorda and L.F. Veiros), 147
- Dynamic NMR**
 Pentacoordinate silicon compounds: stereochemical non-rigidity of chelates formed by intramolecular ring-closure. Crystal structure of 8-dimethylamino-1-trifluorosilylnaphthalene (F. Carr , R.J.P. Corriu, A. Kpoton, M. Poirier, G. Royo, J.C. Young and C. Belin), 43
- Electrochemistry**
 Molecular structure and electrochemical properties of the Pt^{II} complex of 1,1'-bis(methylthio)ferrocene and the Pd^{II} complexes of 1,4,7-trithia[7]- and 1,5,9-trithia[9] (1,1')ferrocenophanes (M. Sato, K. Suzuki, H. Asano, M. Sekino, Y. Kawata, Y. Habata and S. Akabori), 263

- The electrochemical reduction of Cp_2TiCl_2 in the presence of trimethylphosphine (I.M.M. Fusing, D. Pletcher and R.J. Whitby), 119
- The synthesis and electrochemistry of $\text{CpTiCl}_2(\text{OR})$ (R = alkyl, aryl) complexes (I.M.M. Fusing, D. Pletcher and R.J. Whitby), 109
- Electron spin resonance**
Delocalization of the unpaired spin density in some niobocene complexes with σ -donor, π -acceptors (A. Antiñolo, M. Fajardo, E. De Jesús, Y. Mugnier and A. Otero), 127
- Enyne**
Reaction of sodium amalgam with (μ -enyne)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum(I)) and (μ - η^2 , η^3 -allenyl)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum) tetrafluoroborate complexes. Crystal structure of $[\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-HC}\equiv\text{CCH}(\text{CH}_2\text{CH}_3))_2]$ (J.F. Capon, S. Cornen, N. Le Berre-Cosquer, R. Pichon, R. Kergoat and P. L'Haridon), 137
- Extended Hückel calculations**
Delocalization of the unpaired spin density in some niobocene complexes with σ -donor, π -acceptors (A. Antiñolo, M. Fajardo, E. De Jesús, Y. Mugnier and A. Otero), 127
Nuclear magnetic resonance studies of sulfur inversion in bis(cyclopentadienyl)-molybdenum and -tungsten complexes with dithioethers (J.R. Ascenso, M. De Deus Carvalho, A.R. Dias, C.C. Romão, M.J. Calhorda and L.F. Veiros), 147
- Ferrocene**
Molecular structure and electrochemical properties of the Pt^{II} complex of 1,1'-bis(methylthio)ferrocene and the Pd^{II} complexes of 1,4,7-trithia[7]- and 1,5,9-trithia[9] (1,1')ferrocenophanes (M. Sato, K. Suzuki, H. Asano, M. Sekino, Y. Kawata, Y. Habata and S. Akabori), 263
ortho-Mercuration of ferrocenylimines (S.Q. Huo, Y.J. Wu, Y. Zhu and L. Yang), 17
Preparation and Mössbauer spectroscopic studies of 1',1''-diacetyl- and 1'-acetyl-1''-(1-naphthylimino-ethyl)biferrocenes and their partially oxidized biferrocenium salts (H.-H. Wei, I.-S. Hwang, M.-C. Cheng and Y. Wang), 161
Synthesis and characterisation of monomeric, dimeric and polymeric ferrocenylacetylides. Crystal structure of 1,1'-bis(phenylethynyl)ferrocene (S.L. Ingham, M.S. Khan, J. Lewis, N.J. Long and P.R. Raithby), 153
- Germanium**
Zur Struktur des Diethylgermanium-diphenylsiliciumoxids und zur Konformation von Achtringen mit Silicium und Germanium (M. Akkurt, T.R. Kök, P. Faleschini, L. Randaccio, H. Puff and W. Schuh), 59
- Gold**
A ^{197}Au Mössbauer study of reaction products of trimeric 1-benzyl-2-gold(I)-imidazole leading to Au^{I} carbene or Au^{I} imidazoline complexes and trinuclear Au^{III} imidazolyl derivatives. X-Ray crystal structure of $[\{(\mu\text{-1-benzylimidazolato-}N^3,C^2)\text{Au}\}_3\text{I}_2]$ (B. Bovio, S. Calogero, F.E. Wagner, A. Burini and B.R. Pietroni), 275
- Group 6**
Heterozyklen als Liganden. XX. Spektroskopische und strukturelle Charakterisierung von Pentacarbonylmetall-Komplexen der Imidazol-2-ylidene (N. Kuhn, T. Kratz, R. Boese und D. Bläser), C8
- Hexabenzocyclohexyne**
Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17, 18:21,22-hexabenzocyclohexyne-3,7,11,15,19,23-hexadehydro[24]annulene (HBC), 1,2:5,6:9,10:13,14-tetrabenzocyclohexyne-3,7,11,15-tetradehydro[16]annulene (QBC) and a tetracobalt complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231
- Holmium**
Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organoholmium hydroxide complexes (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99
- Hydride**
Ein Beitrag zur Darstellung und Reaktivität der η^3 -Allyliridium-Komplexe $[\text{Ir}(\eta^3\text{-2-RC}_3\text{H}_4)(\text{P}^i\text{Pr}_3)_2]$ (M. Schulz und H. Werner), 243
Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organoholmium hydroxide complexes (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99
The reaction of (μ -H) $_2\text{Os}_3(\text{CO})_9(\text{PPh}_3)_3$ with ethylene affords the ethylidene complex (μ -H) $_2\text{Os}_3(\text{CO})_9(\text{PPh}_3)(\mu\text{-CHCH}_3)$ (M. Koike and J.R. Shapley), 199
- Hydrogen migration**
Thermally and photochemically induced vinyl-hydrogen activation of $[\eta^4\text{-1,2:3,4-(trans-1,3,5-hexatriene)](\eta^5\text{-cyclopentadienyl})\text{cobalt}$: regio- and stereospecific hydrogen migrations (J.A. King Jr. and K.P.C. Vollhardt), 207
- Hydroxide**
Elimination of β -palladium hydroxide in the Pd^{II} -catalyzed reaction of methyl (α -hydroxymethyl)acrylate with alcohols (T. Hosokawa, T. Sugafuji, T. Yamanaka and S.-I. Murahashi), 253
- Imidazole**
A ^{197}Au Mössbauer study of reaction products of trimeric 1-benzyl-2-gold(I)-imidazole leading to Au^{I} carbene or Au^{I} imidazoline complexes and trinuclear Au^{III} imidazolyl derivatives. X-Ray crystal structure of $[\{(\mu\text{-1-benzylimidazolato-}N^3,C^2)\text{Au}\}_3\text{I}_2]$ (B. Bovio, S. Calogero, F.E. Wagner, A. Burini and B.R. Pietroni), 275
Heterozyklen als Liganden. XX. Spektroskopische und strukturelle Charakterisierung von Pentacarbonylmetall-Komplexen der Imidazol-2-ylidene (N. Kuhn, T. Kratz, R. Boese und D. Bläser), C8
- Imine**
ortho-Mercuration of ferrocenylimines (S.Q. Huo, Y.J. Wu, Y. Zhu and L. Yang), 17
- Indole**
Direct synthesis of (η^6 -arene)-tricarbonylchromium complexes by arene displacement from tricarbonyl(η^5 -1-methylpyrrole)chromium(0) (A. Goti and M.F. Semmelhack), C4
- Intramolecular coordination**
Pentacoordinate silicon compounds: stereochemical non-rigidity of chelates formed by intramolecular ring-closure. Crystal structure of 8-dimethylamino-1-trifluorosilylnaphthalene (F. Carré, R.J.P. Corriu, A. Kpoton, M. Poirier, G. Royo, J.C. Young and C. Belin), 43
- Iridium**
Ein Beitrag zur Darstellung und Reaktivität der η^3 -Allyliridium-Komplexe $[\text{Ir}(\eta^3\text{-2-RC}_3\text{H}_4)(\text{P}^i\text{Pr}_3)_2]$ (M. Schulz und H. Werner), 243

- Iron**
 Molecular orbital calculations on $\text{Fe}_2(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ (S.N. Datta, R.-K. Kondru and P. Mathur), 169
 Übergangsmetall-substituierte Acylphosphane und Phosphaalkene. XX. Dipolare [3 + 2]-Cycloadditionen eines Acetylendicarbonsäureesters an die Metallophosphaalkene ($\eta^5\text{-C}_5\text{Me}_5$)(CO) $_2$ Fe-P=C(R)SiMe $_3$ (R = Ph, SiMe $_3$) (L. Weber und A. Rühlicke), C1
- Isomerization**
 Cyclotetrasilazane als Precursor für Cyclotrisilazane und Bis(cyclodisilazan-1-yl)silane (E. Werner und U. Klingebiel), 35
- Lanthanum**
 Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organoholmium hydroxide complexes (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99
- Large ring**
 Zur Struktur des Diethylgermanium-diphenylsiliciumoxids und zur Konformation von Achtringen mit Silicium und Germanium (M. Akkurt, T.R. Kök, P. Faleschini, L. Randaccio, H. Puff und W. Schuh), 59
- Lithium**
 Reduction with lithium dialkylamides (M. Majewski and D.M. Gleave), 1
- Mechanism**
 Elimination of β -palladium hydroxide in the Pd^{II}-catalyzed reaction of methyl (α -hydroxymethyl)acrylate with alcohols (T. Hosokawa, T. Sugafuji, T. Yamanaka and S.-I. Murahashi), 253
- Mercury**
ortho-Mercuration of ferrocenylimines (S.Q. Huo, Y.J. Wu, Y. Zhu and L. Yang), 17
- Mesityl**
 Syntheses and structures of dimesitylbismuth(III) bromide, Mes $_2$ BiBr, and bis(diphenyldithiophosphinato)mesitylbismuth(III), MesBi(S $_2$ PPh $_2$) $_2$ (K.H. Ebert, R.E. Schulz, H.J. Breunig, C. Silvestru and I. Haiduc), 93
- Metallocarbene**
 Thermally and photochemically induced vinyl-hydrogen activation of [$\eta^4\text{-1,2 : 3,4-(trans-1,3,5\text{-hexatriene})$]($\eta^5\text{-cyclopentadienyl}$)cobalt: regio- and stereospecific hydrogen migrations (J.A. King Jr. and K.P.C. Vollhardt), 207
- Metallocenes**
 Delocalization of the unpaired spin density in some niobocene complexes with σ -donor, π -acceptors (A. Antiñolo, M. Fajardo, E. De Jesús, Y. Mugnier and A. Otero), 127
- Methyl acrylate**
 The reaction of dihydridotetrakis(triphenylphosphane)ruthenium(II) with methyl acrylate. Crystal structure of bis(methylacrylate)bis(triphenylphosphane)(aqua)ruthenium(0) (R. Sustmann, B. Patzke and R. Boese), 191
- MNDO calculations**
 Can triorganoboroxins exist in a "monomeric" R-B=O form? MNDO calculations and ebulliometric molecular weight determination (A. Sporyński and H. Szatyłowicz), 31
- Molecular mechanics**
 Molecular mechanics (MM2) calculations and cone angles of phosphine ligands (M. Chin, G.L. Durst, S.R. Head, P.L. Bock and J.A. Mosbo), 73
- Molecular orbital calculations**
 Molecular orbital calculations on $\text{Fe}_2(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ (S.N. Datta, R.-K. Kondru and P. Mathur), 169
- Molybdenum**
 Nuclear magnetic resonance studies of sulfur inversion in bis(cyclopentadienyl)-molybdenum and -tungsten complexes with dithioethers (J.R. Ascenso, M. De Deus Carvalho, A.R. Dias, C.C. Romão, M.J. Calhorda and L.F. Veiros), 147
 Reaction of sodium amalgam with (μ -enyne)bis(dicarbonyl- η^5 -cyclopentadienyl)molybdenum(I) and (μ - η^2, η^3 -allenyl)bis(dicarbonyl- η^5 -cyclopentadienyl)molybdenum) tetrafluoroborate complexes. Crystal structure of [[Mo($\eta^5\text{-C}_5\text{H}_5$)(CO) $_2$] $_2$ (μ -HC \equiv CCH(CH $_2$ CH $_3$))] $_2$] (J.F. Capon, S. Cornen, N. Le Berre-Cosquer, R. Pichon, R. Kergoat and P. L'Haridon), 137
- Mössbauer spectroscopy**
 A ^{197}Au Mössbauer study of reaction products of trimeric 1-benzyl-2-gold(I)-imidazole leading to Au^I carbene or Au^I imidazole complexes and trinuclear Au^{III} imidazolyl derivatives. X-Ray crystal structure of [(μ -1-benzylimidazolato- N^3, C^2)Au] $_3$ I $_2$] (B. Bovio, S. Calogero, F.E. Wagner, A. Burini and B.R. Pietroni), 275
 Preparation and Mössbauer spectroscopic studies of 1',1''-diacetyl- and 1'-acetyl-1''-(1-naphthylimino-ethyl)biferrocenes and their partially oxidized biferrocenium salts (H.-H. Wei, I.-S. Hwang, M.-C. Cheng and Y. Wang), 161
- Niobium**
 Delocalization of the unpaired spin density in some niobocene complexes with σ -donor, π -acceptors (A. Antiñolo, M. Fajardo, E. De Jesús, Y. Mugnier and A. Otero), 127
- Nuclear magnetic resonance**
 Analogues of DIOP. Synthesis, reactivity, and NMR behaviour (A. Börner, R. Kadyrov, M. Michalik and D. Heller), 237
 Determination of the free activation energy for the breaking of the $^{31}\text{P}-^{107(109)}\text{Ag}$ bond in [($\eta^6\text{-p-cymene}$)Ru(μ -pz) $_3$ Ag(PPh $_3$)] (D. Carmona, J. Ferrer, M.P. Lamata, L.A. Oro, H.-H. Limbach, G. Scherer, J. Elguero and M.L. Jimeno), 271
 NMR conformational studies of (1,2-bis(diarylphosphino)ethane)(diorgano sulfide)ruthenium complexes (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 179
 Nuclear magnetic resonance studies of sulfur inversion in bis(cyclopentadienyl)-molybdenum and -tungsten complexes with dithioethers (J.R. Ascenso, M. De Deus Carvalho, A.R. Dias, C.C. Romão, M.J. Calhorda and L.F. Veiros), 147
 The reaction of (μ -H) $_2$ Os $_3$ (CO) $_9$ (PPh $_3$) with ethylene affords the ethylidene complex (μ -H) $_2$ Os $_3$ (CO) $_9$ (PPh $_3$)(μ -CHCH $_3$) (M. Koike and J.R. Shapley), 199
- Olefin**
 Relative stability of η^5 -cyclopentadienyl(olefin)ruthenium cations: the first estimation of the nature of the Ru^{II}-olefin bond (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 189
- Osmium**
 The reaction of (μ -H) $_2$ Os $_3$ (CO) $_9$ (PPh $_3$) with ethylene affords the ethylidene complex (μ -H) $_2$ Os $_3$ (CO) $_9$ (PPh $_3$)(μ -CHCH $_3$) (M. Koike and J.R. Shapley), 199
- Oxide**
 Zur Struktur des Diethylgermanium-diphenylsiliciumoxids und zur Konformation von Achtringen mit Silicium und Germanium (M. Akkurt, T.R. Kök, P. Faleschini, L. Randaccio, H. Puff und W. Schuh), 59

Palladium

Elimination of β -palladium hydroxide in the Pd^{II}-catalyzed reaction of methyl (α -hydroxymethyl)acrylate with alcohols (T. Hosokawa, T. Sugafuji, T. Yamanaka and S.-I. Murahashi), 253

Molecular structure and electrochemical properties of the Pt^{II} complex of 1,1'-bis(methylthio)ferrocene and the Pd^{II} complexes of 1,4,7-trithia[7]- and 1,5,9-trithia[9] (1,1')ferrocenophanes (M. Sato, K. Suzuki, H. Asano, M. Sekino, Y. Kawata, Y. Habata and S. Akabori), 263

Reactivity of mono- and di-methoxycarbonyl complexes of Pd^{II} towards amines and copper amine complexes: role of copper in the catalyzed palladium-copper oxidative carbonylation of amines (P. Giannoccaro), 249

Synthesis of aromatic acids via catalyzed methyl formate-chloroarene reactions (G. Jenner and A. Ben Taleb), 257

Unusual stabilization of cationic "M(η^3 -allyl)⁺ⁿ (M = Pt, Pd) units by a dianionic *cis*-{Pt(C₆F₅)₂(C≡CSiMe₃)₂}²⁻ fragment (J.R. Berenguer, J. Forniés, E. Lalinde and F. Martínez), C15

Paramagnetism

Delocalization of the unpaired spin density in some niobocene complexes with σ -donor, π -acceptors (A. Antiñolo, M. Fajardo, E. De Jesús, Y. Mugnier and A. Otero), 127

Pauson-Khand reaction

A convenient synthesis of hexacarbonyldicobalt complexes of chiral (non-racemic) terminal alkoxyacetylenes (V. Bernardes, X. Verdager, A. Moyano, M.A. Pericàs, A. Riera and A.E. Greene), C12

Pentacoordinate silicon

Pentacoordinate silicon compounds: stereochemical non-rigidity of chelates formed by intramolecular ring-closure. Crystal structure of 8-dimethylamino-1-trifluorosilylnaphthalene (F. Carré, R.J.P. Corriu, A. Kpoton, M. Poirier, G. Royo, J.C. Young and C. Belin), 43

Pentafluorophenyl

Unusual stabilization of cationic "M(η^3 -allyl)⁺ⁿ (M = Pt, Pd) units by a dianionic *cis*-{Pt(C₆F₅)₂(C≡CSiMe₃)₂}²⁻ fragment (J.R. Berenguer, J. Forniés, E. Lalinde and F. Martínez), C15

Phosphaalkene

Übergangsmetall-substituierte Acylphosphane und Phosphaalkene. XX. Dipolare [3 + 2]-Cycloadditionen eines Acetylendicarbonsäureesters an die Metallophosphaalkene (η^5 -C₅Me₅)(CO)₂Fe-P=C(R)SiMe₃ (R = Ph, SiMe₃) (L. Weber und A. Rühlicke), C1

Phosphazane

Formation and structure of [Co{Ph₂PN(ⁱBu)PPh₂-P,P'}₂(CO)]Co(CO)₄ – a cage molecule-pair or an ion-pair complex? (Z.-Z. Zhang, A. Yu, H.-P. Xi, R.-J. Wang and H.-G. Wang), 223

Phosphide

Synthesis and derivative chemistry of [Ru₂(μ -PPh₂)(μ -OH)₂(η^6 -*p*-cymene)₂]⁺ (J.A. Cabeza, F. Mulla and V. Riera), 173

Phosphine

Molecular mechanics (MM2) calculations and cone angles of phosphine ligands (M. Chin, G.L. Durrst, S.R. Head, P.L. Bock and J.A. Mosbo), 73

The electrochemical reduction of Cp₂TiCl₂ in the presence of trimethylphosphine (I.M.M. Fusing, D. Pletcher and R.J. Whitby), 119

Phosphorus

Analogues of DIOP. Synthesis, reactivity, and NMR behaviour (A. Börner, R. Kadyrov, M. Michalik and D. Heller), 237

Determination of the free activation energy for the breaking of the ³¹P–¹⁰⁷(¹⁰⁹)Ag bond in [(η^6 -*p*-cymene)Ru(μ -pz)₃Ag(PPh₃)

(D. Carmona, J. Ferrer, M.P. Lamata, L.A. Oro, H.-H. Limbach, G. Scherer, J. Elguero and M.L. Jimeno), 271

Ein Beitrag zur Darstellung und Reaktivität der η^3 -Allyliridium-Komplexe [Ir(η^3 -2-RC₃H₄)(PⁱPr₃)₂] (M. Schulz und H. Werner), 243

Syntheses and structures of dimesitylbismuth(III) bromide, Mes₂BiBr, and bis(diphenyldithiophosphinato)mesitylbismuth(III), MesBi(S₂PPh₂)₂ (K.H. Ebert, R.E. Schulz, H.J. Breunig, C. Silvestru and I. Haiduc), 93

Photochemistry

Thermally and photochemically induced vinyl-hydrogen activation of [η^4 -1,2 : 3,4-*trans*-1,3,5-hexatriene)(η^5 -cyclopentadienyl)cobalt: regio- and stereospecific hydrogen migrations (J.A. King Jr. and K.P.C. Vollhardt), 207

Platinum

Molecular structure and electrochemical properties of the Pt^{II} complex of 1,1'-bis(methylthio)ferrocene and the Pd^{II} complexes of 1,4,7-trithia[7]- and 1,5,9-trithia[9] (1,1')ferrocenophanes (M. Sato, K. Suzuki, H. Asano, M. Sekino, Y. Kawata, Y. Habata and S. Akabori), 263

Unusual stabilization of cationic "M(η^3 -allyl)⁺ⁿ (M = Pt, Pd) units by a dianionic *cis*-{Pt(C₆F₅)₂(C≡CSiMe₃)₂}²⁻ fragment (J.R. Berenguer, J. Forniés, E. Lalinde and F. Martínez), C15

Polymer

Synthesis and characterisation of monomeric, dimeric and polymeric ferrocenylacetylides. Crystal structure of 1,1'-bis(phenylethynyl)ferrocene (S.L. Ingham, M.S. Khan, J. Lewis, N.J. Long and P.R. Raithby), 153

Praseodymium

Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organo-holmium hydroxide complexes (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99

Pseudo-rotation

Pentacoordinate silicon compounds: stereochemical non-rigidity of chelates formed by intramolecular ring-closure. Crystal structure of 8-dimethylamino-1-trifluorosilylnaphthalene (F. Carré, R.J.P. Corriu, A. Kpoton, M. Poirier, G. Royo, J.C. Young and C. Belin), 43

Pyrrole

Direct synthesis of (η^6 -arene)- tricarbonylchromium complexes by arene displacement from tricarbonyl(η^5 -1-methylpyrrole)chromium(0) (A. Goti and M.F. Semmelhack), C4

Quinoline

Direct synthesis of (η^6 -arene)- tricarbonylchromium complexes by arene displacement from tricarbonyl(η^5 -1-methylpyrrole)chromium(0) (A. Goti and M.F. Semmelhack), C4

Reduction

Reduction with lithium dialkylamides (M. Majewski and D.M. Gleave), 1

The electrochemical reduction of Cp₂TiCl₂ in the presence of trimethylphosphine (I.M.M. Fusing, D. Pletcher and R.J. Whitby), 119

The synthesis and electrochemistry of CpTiCl₂(OR) (R = alkyl, aryl) complexes (I.M.M. Fusing, D. Pletcher and R.J. Whitby), 109

Rhodium

Analogues of DIOP. Synthesis, reactivity, and NMR behaviour (A. Börner, R. Kadyrov, M. Michalik and D. Heller), 237

Ruthenium

η^5 -Pentamethylcyclopentadienylruthenium(II) complexes contain-

- ing η^6 -coordinated α -amino acids (W.S. Sheldrick and A. Gleichmann), 183
- Determination of the free activation energy for the breaking of the $^{31}\text{P}-^{107}\text{Ag}$ bond in $[(\eta^6\text{-}p\text{-cymene})\text{Ru}(\mu\text{-pz})_3\text{Ag}(\text{PPh}_3)]$ (D. Carmona, J. Ferrer, M.P. Lamata, L.A. Oro, H.-H. Limbach, G. Scherer, J. Elguero and M.L. Jimeno), 271
- Molecular orbital calculations on $\text{Fe}_4(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ (S.N. Datta, R.-K. Kondru and P. Mathur), 169
- NMR conformational studies of (1,2-bis(diarylphosphino)ethane)(diorgano sulfide)ruthenium complexes (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 179
- Relative stability of η^5 -cyclopentadienyl(olefin)ruthenium cations: the first estimation of the nature of the Ru^{II} -olefin bond (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 189
- Synthesis and derivative chemistry of $[\text{Ru}_2(\mu\text{-PPh}_2)(\mu\text{-OH})_2(\eta^6\text{-}p\text{-cymene})_2]^+$ (J.A. Cabeza, F. Mulla and V. Riera), 173
- Synthesis of aromatic acids via catalyzed methyl formate-chloroarene reactions (G. Jenner and A. Ben Taleb), 257
- The reaction of dihydridotetrakis(triphenylphosphane)ruthenium(II) with methyl acrylate. Crystal structure of bis(methylacrylate)bis(triphenylphosphane)(aqua)ruthenium(0) (R. Sustmann, B. Patzke and R. Boese), 191
- Selenium**
- Molecular orbital calculations on $\text{Fe}_4(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ and $\text{Fe}_3\text{Ru}(\text{CO})_{10}(\mu\text{-CO})(\mu_4\text{-Se})_2$ (S.N. Datta, R.-K. Kondru and P. Mathur), 169
- Silicon**
- Cyclotetrasilazane als Precursor für Cyclotrisilazane und Bis(cyclodisilazan-1-yl)silane (E. Werner und U. Klingebiel), 35
- Pentacoordinate silicon compounds: stereochemical non-rigidity of chelates formed by intramolecular ring-closure. Crystal structure of 8-dimethylamino-1-trifluorosilylnaphthalene (F. Carré, R.J.P. Corriu, A. Kpoton, M. Poirier, G. Royo, J.C. Young and C. Belin), 43
- Zur Struktur des Diethylgermanium-diphenylsiliciumoxids und zur Konformation von Achtringen mit Silicium und Germanium (M. Akkurt, T.R. Kök, P. Faleschini, L. Randaccio, H. Puff und W. Schuh), 59
- Silver**
- Determination of the free activation energy for the breaking of the $^{31}\text{P}-^{107}\text{Ag}$ bond in $[(\eta^6\text{-}p\text{-cymene})\text{Ru}(\mu\text{-pz})_3\text{Ag}(\text{PPh}_3)]$ (D. Carmona, J. Ferrer, M.P. Lamata, L.A. Oro, H.-H. Limbach, G. Scherer, J. Elguero and M.L. Jimeno), 271
- Sodium**
- Reaction of sodium amalgam with (μ -enyne)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum(I)) and (μ - η^2, η^3 -allenyl)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum) tetrafluoroborate complexes. Crystal structure of $[[\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-HC}\equiv\text{CCH}(\text{CH}_2\text{CH}_3))_2]$ (J.F. Capon, S. Cornen, N. Le Berre-Cosquer, R. Pichon, R. Kergoat and P. L'Haridon), 137
- Substitution**
- Cyclotetrasilazane als Precursor für Cyclotrisilazane und Bis(cyclodisilazan-1-yl)silane (E. Werner und U. Klingebiel), 35
- Sulfide**
- NMR conformational studies of (1,2-bis(diarylphosphino)ethane)(diorgano sulfide)ruthenium complexes (K. Ohkita, H. Kurosawa, T. Hirao and I. Ikeda), 179
- Sulfur inversion**
- Nuclear magnetic resonance studies of sulfur inversion in bis(cyclopentadienyl)-molybdenum and -tungsten complexes with dithioethers (J.R. Ascenso, M. De Deus Carvalho, A.R. Dias, C.C. Romão, M.J. Calhorda and L.F. Veiros), 147
- Synthesis**
- The synthesis and electrochemistry of $\text{CpTiCl}_2(\text{OR})$ (R = alkyl, aryl) complexes (I.M.M. Fussing, D. Pletcher and R.J. Whitby), 109
- Tetrabenzocyclo tetrayne**
- Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17, 18:21,22-hexabenzocyclo-3,7,11,15,19,23-hexadehydro[24]annulene (HBC), 1,2:5,6:9,10:13,14-tetrabenzocyclo-3,7,11,15-tetradehydro[16]annulene (QBC) and a tetracobalt complex of QBC. The first example of a transition metal complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231
- Tetrafluoroborate**
- Reaction of sodium amalgam with (μ -enyne)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum(I)) and (μ - η^2, η^3 -allenyl)bis(dicarbonyl- η^5 -cyclopentadienylmolybdenum) tetrafluoroborate complexes. Crystal structure of $[[\{\text{Mo}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-HC}\equiv\text{CCH}(\text{CH}_2\text{CH}_3))_2]$ (J.F. Capon, S. Cornen, N. Le Berre-Cosquer, R. Pichon, R. Kergoat and P. L'Haridon), 137
- Thermochemistry**
- Thermally and photochemically induced vinyl-hydrogen activation of $[\eta^4\text{-1,2:3,4-(trans-1,3,5-hexatriene)}](\eta^5\text{-cyclopentadienyl})$ cobalt: regio- and stereospecific hydrogen migrations (J.A. King Jr. and K.P.C. Vollhardt), 207
- Tin**
- Synthesis and characterisation of monomeric, dimeric and polymeric ferrocenylacetylides. Crystal structure of 1,1'-bis(phenylethynyl)ferrocene (S.L. Ingham, M.S. Khan, J. Lewis, N.J. Long and P.R. Raithby), 153
- Synthesis of bimetallic $[\text{Sn}^{\text{IV}}, \text{Al}^{\text{III}}]\text{-}\mu\text{-oxoisopropoxyacetate}$ and $\text{-}\mu\text{-oxoisopropoxide}$ and their acetylacetone and benzoylacetone derivatives (Sonika, A.K. Narula, O.P. Vermani, H.K. Sharma and A.S. Sarpal), 67
- Titanium**
- The electrochemical reduction of Cp_2TiCl_2 in the presence of trimethylphosphine (I.M.M. Fussing, D. Pletcher and R.J. Whitby), 119
- The synthesis and electrochemistry of $\text{CpTiCl}_2(\text{OR})$ (R = alkyl, aryl) complexes (I.M.M. Fussing, D. Pletcher and R.J. Whitby), 109
- Tribenzocyclotriene**
- Syntheses and crystal structures of 1,2:5,6:9,10:13,14:17, 18:21,22-hexabenzocyclo-3,7,11,15,19,23-hexadehydro[24]annulene (HBC), 1,2:5,6:9,10:13,14-tetrabenzocyclo-3,7,11,15-tetradehydro[16]annulene (QBC) and a tetracobalt complex of QBC. The first example of a transition metal complex of QBC (D. Solooki, J.D. Bradshaw, C.A. Tessier, W.J. Youngs, R.F. See, M. Churchill and J.D. Ferrara), 231
- Trimethylsilyl**
- Übergangsmetall-substituierte Acylphosphane und Phosphoalkene. XX. Dipolare [3 + 2]-Cycloadditionen eines Acetylendicarbonylsäureesters an die Metallophosphoalkene ($\eta^5\text{-C}_5\text{Me}_5(\text{CO})_2\text{Fe-P-C(R)SiMe}_3$ (R = Ph, SiMe₃)) (L. Weber und A. Rühlicke), C1
- Tungsten**
- Nuclear magnetic resonance studies of sulfur inversion in bis(cyclopentadienyl)-molybdenum and -tungsten complexes with dithioethers (J.R. Ascenso, M. De Deus Carvalho, A.R. Dias, C.C. Romão, M.J. Calhorda and L.F. Veiros), 147

- Urea**
 Reactivity of mono- and di-methoxycarbonyl complexes of Pd^{II} towards amines and copper amine complexes: role of copper in the catalyzed palladium–copper oxidative carbonylation of amines (P. Giannoccaro), 249
- Vinyl**
 Ein Beitrag zur Darstellung und Reaktivität der η^3 -Allyliridium-Komplexe $[\text{Ir}(\eta^3\text{-2-RC}_3\text{H}_4)(\text{P}^i\text{Pr}_3)_2]$ (M. Schulz und H. Werner), 243
- Voltammetry**
 The electrochemical reduction of Cp₂TiCl₂ in the presence of trimethylphosphine (I.M.M. Fussing, D. Pletcher and R.J. Whitby), 119
- X-ray diffraction**
 Molecular structure and electrochemical properties of the Pt^{II} complex of 1,1'-bis(methylthio)ferrocene and the Pd^{II} complexes of 1,4,7-trithia[7]- and 1,5,9-trithia[9] (1,1')ferrocenophanes (M. Sato, K. Suzuki, H. Asano, M. Sekino, Y. Kawata, Y. Habata and S. Akabori), 263
- Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organoholmium hydroxide complexes** (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99
- The reaction of dihydridotetrakis(triphenylphosphane)ruthenium(II) with methyl acrylate. Crystal structure of bis(methylacrylate)bis(triphenylphosphane)(aqua)ruthenium(0)** (R. Sustmann, B. Patzke and R. Boese), 191
- X-ray structure**
 Unusual stabilization of cationic " $\text{M}(\eta^3\text{-allyl})^{+n}$ (M = Pt, Pd) units by a dianionic $\text{cis-}[\text{Pt}(\text{C}_6\text{F}_5)_2(\text{C}\equiv\text{CSiMe}_3)_2]^{2-}$ fragment (J.R. Berenguer, J. Forniés, E. Lalinde and F. Martínez), C15
- Yttrium**
 Studies on organolanthanide complexes. LXIII. Synthesis, spectroscopic and X-ray crystallographic characterization of new early organolanthanide, organoyttrium hydride and organoholmium hydroxide complexes (D. Deng, Y. Jiang, C. Qian, G. Wu and P. Zheng), 99