

Subject Index of Volume 465

Ab initio

Molecular mechanical studies on the olefin metathesis reaction.
I. Development and evaluation of tungsten carbene parameters: METMOD1 (L. Bencze and R. Szilagyi), 211

Alkene

Palladium-catalyzed cross-coupling of aryl and alkenyl boronic acids with alkenes via oxidative addition of a carbon–boron bond to palladium(0) (C.S. Cho and S. Uemura), 85

Alkenyl

Palladium-catalyzed cross-coupling of aryl and alkenyl boronic acids with alkenes via oxidative addition of a carbon–boron bond to palladium(0) (C.S. Cho and S. Uemura), 85

Alkyne

A route to η^2 -alkyne complexes of vanadium(I): crystal structure of *trans*-[IV(CO)₂(PMe₂Ph)₂(2-butyne)] (H. Gailus, H. Maelger and D. Rehder), 181

Aluminium

Reactions of trimethylaluminium with *cis*-2-buten-1,4-diol (W. Ziemkowska, S. Pasynkiewicz and E. Kalbarczyk), 93

Amide

Darstellung und Struktur monomerer N-borylierter Gallium-amide (G. Linti), 79

Arene

Photoabsorption study of electronic excited states of (η^5 -cyclopentadienyl)(η^6 -arene)manganese (S.Yu. Ketkov), 225

Aryl

Palladium-catalyzed cross-coupling of aryl and alkenyl boronic acids with alkenes via oxidative addition of a carbon–boron bond to palladium(0) (C.S. Cho and S. Uemura), 85

Azides

Chemie der schweren Carben-Analogen R₂M (M = Si, Ge, Sn). XIX. Reaktionen von stabilem Diaminogermylen und -stannylen Me₂Si(^tBuN)₂M (M = Ge, Sn) mit organischen Aziden (B. Klein and W.P. Neumann), 119

Boron

Darstellung und Struktur monomerer N-borylierter Gallium-amide (G. Linti), 79

Boronic acid

Palladium-catalyzed cross-coupling of aryl and alkenyl boronic acids with alkenes via oxidative addition of a carbon–boron bond to palladium(0) (C.S. Cho and S. Uemura), 85

But-2-yne

Reactions of the cationic but-2-yne complex [WI(CO)(NCMe)-(Ph₂P(CH₂)PPh₂)(η^2 -MeC₂Me)][BF₄] with bidentate nitrogen donor ligands (P.K. Baker and K.R. Flower), 221

Cadmium

Some complexes of neopentylcadmium species with dithio- and di-selenocarbamates: the synthesis, characterization and single crystal X-ray structure of a mixed neopentyl/diethyldiselenocarbamate of cadmium: [(CH₃)₃CCH₂Cd-Se₂CNET₂]₂ (I. Abrahams, A. Malik, M. Mottevalli and P. O'Brien), 73

Carbene

Chemie der schweren Carben-Analogen R₂M (M = Si, Ge, Sn). XIX. Reaktionen von stabilem Diaminogermylen und -stannylen Me₂Si(^tBuN)₂M (M = Ge, Sn) mit organischen Aziden (B. Klein and W.P. Neumann), 119

Molecular mechanical studies on the olefin metathesis reaction. I. Development and evaluation of tungsten carbene parameters: METMOD1 (L. Bencze and R. Szilagyi), 211

Carbonyl

Dinuclear molybdenum carbonyls bridged by hydride ligands (J.T. Lin, Y.C. Lee, H.M. Kao, T.-Y. Dong and Y.S. Wen), 199

Catalysis

Electronic effects of bis(2-aryl-4,5,6,7-tetrahydroindenyl)titano-cene dichlorides on the catalytic epoxidation of *trans*-3-hexene (R.L. Halterman and T.M. Ramsey), 175

Charge transfer

Synthesis and magnetic studies of the metallocarborene sandwich salts [Fe(C₅Me₅)₂][M(C₂B₉H₁₂)₂] (M = Cr, Fe, Ni) (J.M. Forward, D.M.P. Mingos and A.V. Powell), 251

Cobalt

Heterometallic sulfido clusters by metal exchange: syntheses and X-ray crystal structures of (η^5 -C₅H₄CH₃)MoFeCo(CO)₈(μ_3 -S) and (η^5 -C₅H₄CH₃)₂Mo₂Fe(CO)₇(μ_3 -S) (W.-H. Sun, S.-Y. Yang, H.-Q. Wang, Q.-F. Zhou and K.-B. Yu), 263

Stabilisierung niederkoordinierter M(CO)-Bausteine: Synthese und Reaktivität von (η^5 -C₅H₄SiMe₃)₂Hf(C≡CPh)₂)M(CO) (M = Ni, Co) (H. Lang, M. Herres and W. Imhof), 283

Crystal structure

A route to η^2 -alkyne complexes of vanadium(I): crystal structure of *trans*-[IV(CO)₂(PMe₂Ph)₂(2-butyne)] (H. Gailus, H. Maelger and D. Rehder), 181

Darstellung und Struktur monomerer N-borylierter Gallium-amide (G. Linti), 79

Heterometallic sulfido clusters by metal exchange: syntheses and X-ray crystal structures of (η^5 -C₅H₄CH₃)MoFeCo(CO)₈(μ_3 -S) and (η^5 -C₅H₄CH₃)₂Mo₂Fe(CO)₇(μ_3 -S) (W.-H. Sun, S.-Y. Yang, H.-Q. Wang, Q.-F. Zhou and K.-B. Yu), 263

Magnesium compounds: classification and analysis of crystallographic and structural data (C.E. Holloway and M. Melnik), 1

Organotin-substituted organosiloxanes: synthesis and characterisation, including the crystal structure of 1,3-bis(triphenylstanoxo)-1,1,3,3-tetraphenyl-1,3-disiloxane (B.J. Bridson, M.F. Mahon, K.C. Molloy and P.J. Schofield), 145

Some complexes of neopentylcadmium species with dithio- and di-selenocarbamates: the synthesis, characterization and single crystal X-ray structure of a mixed neopentyl/diethyldiselenocarbamate of cadmium: [(CH₃)₃CCH₂Cd-Se₂CNET₂]₂ (I. Abrahams, A. Malik, M. Mottevalli and P. O'Brien), 73

Synthesis and magnetic studies of the metallocarborene sandwich salts [Fe(C₅Me₅)₂][M(C₂B₉H₁₂)₂] (M = Cr, Fe, Ni) (J.M. Forward, D.M.P. Mingos and A.V. Powell), 251

Cycloaddition

1,3-Dipolar cycloaddition to a Fe–N=C fragment. XII. Reactivity of Fe(2,6-xylyl-isocyanide)₃(ⁱPr- α -diimine) towards heteroal-

- lenic C=S bonds (P.P.M. De Lange, E. Alberts, M. Van Wijndooper, H.-W. Frühauf, K. Vrieze, H. Kooijman and A.L. Spek), 241
- Cyclopentadienyl**
- Etude électrochimique des cations (η^5 -cyclopentadiényl fer η^6 -nitrobenzène ou η^6 -nitrotoluènes) $^+$. Electrosynthèse des dérivés nitrosés correspondants (N. Guennec and C. Moinet), 233
- Komplexchemie perhalogenierter Cyclopentadiene und Alkine. XVI. Darstellung von π -Cyclopentadienylkomplexen mit zwei und drei Chalkogenen-substituenten am Ring, $[C_5Cl_{4-m}(ER)_m(ER')_m]ML_n$ ($ML_n = Mn(CO)_3, Fe(C_5H_5)$; $m = 1$ oder 2 ; $E = S, Se; R, R' = Me, Ph$). Struktur von $[C_5Cl_3(SMe)_2]IC_5H_5Fe$ (K. Sünkel, U. Birk, A. Blum and W. Kempinger), 167
- Photoabsorption study of electronic excited states of (η^5 -cyclopentadiényl)(η^6 -arene)manganese (S.Yu. Ketkov), 225
- Syntheses and electrochemical behaviour of novel dithiolatoniobium-(IV) and -(V) complexes with two η^5 -cyclopentadienyl rings (F. Guyon, J. Amaudrut, M.-F. Mercier and K. Shimizu), 187
- Difluoromethyl**
- Synthesis and properties of trifluoromethylated stannanes. The formation and NMR spectra of the difluoromethylstannanes $(CF_3)_n(CF_2H)_mSn(CH_3)_{4-n-m}$ (R. Eujen, N. Jahn and U. Thurmann), 153
- Diimine**
- 1,3-Dipolar cycloaddition to a Fe-N=C fragment. XII. Reactivity of $Fe(2,6\text{-xylyl-isocyanide})_3(^iPr\text{-}\alpha\text{-diimine})$ towards heteroalenic C=S bonds (P.P.M. De Lange, E. Alberts, M. Van Wijndooper, H.-W. Frühauf, K. Vrieze, H. Kooijman and A.L. Spek), 241
- Dimethyltin**
- Coordination complexes of dimethyltin dichloride with C-nitroso compounds (M. Cameron, B.G. Gowenlock, R.V. Parish and G. Vasapollo), 161
- Diphosphine**
- Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275
- Diphosphinomethanide**
- Substituentenabhängigkeit der Lithiumkoordination von Phosphinomethaniden und Molekülstrukturen von (TMEDA)Li-[C(PPH₂)₂(SiMe₃)] und ((THF)Li[CH(PMe₂)₂]₂] (H.H. Karsch, G. Grauvogl, P. Mikulcik, P. Bissinger and G. Müller), 65
- Diselenocarbamate**
- Some complexes of neopentylcadmium species with dithio- and di-selenocarbamates: the synthesis, characterization and single crystal X-ray structure of a mixed neopentyl/diethyldiselenocarbamate of cadmium: $[(CH_3)_3CCH_2-CdSe_2CNEt_2]_2$ (I. Abrahams, A. Malik, M. Mottevalli and P. O'Brien), 73
- Dithiocarbamate**
- Some complexes of neopentylcadmium species with dithio- and di-selenocarbamates: the synthesis, characterization and single crystal X-ray structure of a mixed neopentyl/diethyldiselenocarbamate of cadmium: $[(CH_3)_3CCH_2-CdSe_2CNEt_2]_2$ (I. Abrahams, A. Malik, M. Mottevalli and P. O'Brien), 73
- Electrochemistry**
- Etude électrochimique des cations (η^5 -cyclopentadiényl fer η^6 -nitrobenzène ou η^6 -nitrotoluènes) $^+$. Electrosynthèse des dérivés nitrosés correspondants (N. Guennec and C. Moinet), 233
- Syntheses and electrochemical behaviour of novel dithiolatoniobium-(IV) and -(V) complexes with two η^5 -cyclopentadienyl rings (F. Guyon, J. Amaudrut, M.-F. Mercier and K. Shimizu), 187
- Electronic**
- Photoabsorption study of electronic excited states of (η^5 -cyclopentadiényl)(η^6 -arene)manganese (S.Yu. Ketkov), 225
- Epoxidation**
- Electronic effects of bis(2-aryl-4,5,6,7-tetrahydroindenyl)titano-cene dichlorides on the catalytic epoxidation of *trans*-3-hexene (R.L. Halterman and T.M. Ramsey), 175
- Ferrocene**
- Synthesis and magnetic studies of the metallocarborane sandwich salts $[Fe(C_5Me_5)_2IM(C_2B_9H_{12})_2]$ ($M = Cr, Fe, Ni$) (J.M. Forward, D.M.P. Mingos and A.V. Powell), 251
- Ferromagnetism**
- Synthesis and magnetic studies of the metallocarborane sandwich salts $[Fe(C_5Me_5)_2IM(C_2B_9H_{12})_2]$ ($M = Cr, Fe, Ni$) (J.M. Forward, D.M.P. Mingos and A.V. Powell), 251
- Fluorine**
- Synthesis and properties of trifluoromethylated stannanes. The formation and NMR spectra of the difluoromethylstannanes $(CF_3)_n(CF_2H)_mSn(CH_3)_{4-n-m}$ (R. Eujen, N. Jahn and U. Thurmann), 153
- Gallium**
- Darstellung und Struktur monomerer N-borylierter Gallium-amide (G. Linti), 79
- Gas phase**
- Gas phase reactivity of aromatic silanes. The reaction of $Ph(CH_2)_xSiMe_3$ ($x = 0$ or 1) with cationic electrophiles (M.E. Crestoni, S. Fornarini and G. Occhiucci), 109
- Germanium**
- Chemie der schweren Carben-Analogen R_2M ($M = Si, Ge, Sn$). XIX. Reaktionen von stabilem Diaminogermylen und -stannylen $Me_2Si(^iBuN)_2M$ ($M = Ge, Sn$) mit organischen Aziden (B. Klein and W.P. Neumann), 119
- Germany**
- Photochemical reactions of 2-(pentamethyldisilanyl)furan and 2-(pentamethylgermyl)furan. Formation of a radical pair (K. Mochida, K. Kimijima, M. Wakasa and H. Hayashi), 101
- Group 6**
- Formation and *fac-mer* isomerization of cationic phosphonium transition metal complexes formulated as $[M(bpy)(CO)_3(PN(Me)CH_2CH_2NMe)]^+$ ($M = Cr, Mo, W$) and $[M(dppe)(CO)_3(PN(Me)CH_2CH_2NMe)]^+$ ($M = Mo, W$) (H. Nakazawa, Y. Yamaguchi and K. Miyoshi), 193
- Group 8**
- Hydrogen-bonded complexes involving the metal atom and protonation of metallocenes of the iron subgroup (E.S. Shubina, A.N. Krylov, A.Z. Kreindlin, M.I. Rybinskaya and L.M. Epstein), 259
- Group 14**
- Chemie der schweren Carben-Analogen R_2M ($M = Si, Ge, Sn$). XIX. Reaktionen von stabilem Diaminogermylen und -stannylen $Me_2Si(^iBuN)_2M$ ($M = Ge, Sn$) mit organischen Aziden (B. Klein and W.P. Neumann), 119
- Synthesis and properties of trifluoromethylated stannanes. The formation and NMR spectra of the difluoromethylstannanes $(CF_3)_n(CF_2H)_mSn(CH_3)_{4-n-m}$ (R. Eujen, N. Jahn and U. Thurmann), 153

Hafnium

Stabilisierung niederkoordinierter M(CO)-Bausteine: Synthese und Reaktivität von $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Hf}(\text{C}\equiv\text{CPh})_2\text{M}(\text{CO})$ ($\text{M} = \text{Ni}, \text{Co}$) (H. Lang, M. Herres and W. Imhof), 283

Heteroallenes

1,3-Dipolar cycloaddition to a Fe-N=C fragment. XII. Reactivity of $\text{Fe}(2,6\text{-xylyl-isocyanide})_3(\text{iPr-}\alpha\text{-diimine})$ towards heteroallic C=S bonds (P.P.M. De Lange, E. Alberts, M. Van Wijnkoop, H.-W. Frühauf, K. Vrieze, H. Kooijman and A.L. Spek), 241

Heterometallics

Heterometallic sulfido clusters by metal exchange: syntheses and X-ray crystal structures of $(\eta^5\text{-C}_5\text{H}_4\text{CH}_3)_2\text{MoFeCo}(\text{CO})_8(\mu_3\text{-S})$ and $(\eta^5\text{-C}_5\text{H}_4\text{CH}_3)_2\text{Mo}_2\text{Fe}(\text{CO})_7(\mu_3\text{-S})$ (W.-H. Sun, S.-Y. Yang, H.-Q. Wang, Q.-F. Zhou and K.-B. Yu), 263

Hindered (bulky) ligands

Phosphine-nitrile ligands: the molecular structure of *cis*-2-di-phenylphosphino-1-cyanocyclopentane and studies of the donor/acceptor properties of this and related ligands (E.J. Fredericks, M.J. Gindling, L.C. Kroll and B.N. Storhoff), 289

Hydride

Dinuclear molybdenum carbonyls bridged by hydride ligands (J.T. Lin, Y.C. Lee, H.M. Kao, T.-Y. Dong and Y.S. Wen), 199

Hydrogen bonding

Hydrogen-bonded complexes involving the metal atom and protonation of metallocenes of the iron subgroup (E.S. Shubina, A.N. Krylov, A.Z. Kreindlin, M.I. Rybinskaya and L.M. Epstein), 259

Imidazole

N-substituted imidazole derivatives of rhodium(I) and iridium(I). Crystal structure of the head-to-head dinuclear compound $[\text{Ir}_2(\mu\text{-N-benzylimidazolato-}\text{N}^3,\text{C}^2)_2(\text{CO})_4]$ (F. Bonati, L.A. Oro, M.T. Pinillos, C. Tejel and B. Bovio), 267

Infrared spectroscopy

Schwingungsspektren, Normalkoordinatenanalysen und Synthesen der t-Butylsilane $^t\text{BuSiX}_3$, $\text{X} = \text{H}, \text{D}, \text{F}, \text{Cl}, \text{Br}, \text{I}$ (K. Hassler and M. Weidenbruch), 127

Schwingungsspektren und Normalkoordinatenanalysen der Di-t-butylsilane $^t\text{Bu}_2\text{SiX}_2$, $\text{X} = \text{H}, \text{D}, \text{F}, \text{Cl}, \text{Br}, \text{I}$ (K. Hassler and M. Weidenbruch), 137

Ion cyclotron resonance

Gas phase reactivity of aromatic silanes. The reaction of $\text{Ph}(\text{CH}_2)_x\text{SiMe}_3$ ($x = 0$ or 1) with cationic electrophiles (M.E. Crestoni, S. Fornarini and G. Occhiucci), 109

Iridium

N-substituted imidazole derivatives of rhodium(I) and iridium(I). Crystal structure of the head-to-head dinuclear compound $[\text{Ir}_2(\mu\text{-N-benzylimidazolato-}\text{N}^3,\text{C}^2)_2(\text{CO})_4]$ (F. Bonati, L.A. Oro, M.T. Pinillos, C. Tejel and B. Bovio), 267

Iron

1,3-Dipolar cycloaddition to a Fe-N=C fragment. XII. Reactivity of $\text{Fe}(2,6\text{-xylyl-isocyanide})_3(\text{iPr-}\alpha\text{-diimine})$ towards heteroallic C=S bonds (P.P.M. De Lange, E. Alberts, M. Van Wijnkoop, H.-W. Frühauf, K. Vrieze, H. Kooijman and A.L. Spek), 241

Etude electrochimique des cations $(\eta^5\text{-cyclopentadiényl fer-}\eta^6\text{-nitrobenzène ou }\eta^6\text{-nitrotoluènes})^+$. Electrosynthèse des dérivés nitrosés correspondants (N. Guennec and C. Moinet), 233

Heterometallic sulfido clusters by metal exchange: syntheses and X-ray crystal structures of $(\eta^5\text{-C}_5\text{H}_4\text{CH}_3)_2\text{MoFeCo}(\text{CO})_8(\mu_3\text{-S})$ and $(\eta^5\text{-C}_5\text{H}_4\text{CH}_3)_2\text{Mo}_2\text{Fe}(\text{CO})_7(\mu_3\text{-S})$ (W.-H. Sun, S.-Y. Yang, H.-Q. Wang, Q.-F. Zhou and K.-B. Yu), 263

Komplexchemie perhalogenierter Cyclopentadiene und Alkine.

XVI. Darstellung von π -Cyclopentadienylkomplexen mit zwei und drei Chalkogensubstituenten am Ring, $[\text{C}_5\text{Cl}_{4-m}\text{-}(\text{ER})_m(\text{ER}')]\text{ML}_n$ ($\text{ML}_n = \text{Mn}(\text{CO})_3, \text{Fe}(\text{C}_5\text{H}_5)$; $m = 1$ oder 2 ; $\text{E} = \text{S}, \text{Se}; \text{R}, \text{R}' = \text{Me}, \text{Ph}$). Struktur von $[\text{C}_5\text{Cl}_3(\text{SMe})_2]\text{IC}_5\text{H}_5\text{Fe}$ (K. Sünkel, U. Birk, A. Blum and W. Kempinger), 167

Stabilisierung niederkoordinierter M(CO)-Bausteine: Synthese und Reaktivität von $(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Hf}(\text{C}\equiv\text{CPh})_2\text{M}(\text{CO})$ ($\text{M} = \text{Ni}, \text{Co}$) (H. Lang, M. Herres and W. Imhof), 283

Isocyanide

1,3-Dipolar cycloaddition to a Fe-N=C fragment. XII. Reactivity of $\text{Fe}(2,6\text{-xylyl-isocyanide})_3(\text{iPr-}\alpha\text{-diimine})$ towards heteroallic C=S bonds (P.P.M. De Lange, E. Alberts, M. Van Wijnkoop, H.-W. Frühauf, K. Vrieze, H. Kooijman and A.L. Spek), 241

Isomerism

Formation and *fac-mer* isomerization of cationic phosphonium transition metal complexes formulated as $[\text{M}(\text{bpy})(\text{CO})_3(\text{PN}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe})]^+$ ($\text{M} = \text{Cr}, \text{Mo}, \text{W}$) and $[\text{M}(\text{dppe})(\text{CO})_3(\text{PN}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe})]^+$ ($\text{M} = \text{Mo}, \text{W}$) (H. Nakazawa, Y. Yamaguchi and K. Miyoshi), 193

Lithium

Substituentenabhängigkeit der Lithiumkoordination von Phosphinomethaniden und Molekülstrukturen von $(\text{TMEDA})\text{Li-C}(\text{PPh}_2)_2(\text{SiMe}_3)$ und $((\text{THF})\text{Li}[\text{CH}(\text{PMe}_2)_2])_2$ (H.H. Karsch, G. Grauvogl, P. Mikulcik, P. Bissinger and G. Müller), 65

Magnesium

Magnesium compounds: classification and analysis of crystallographic and structural data (C.E. Holloway and M. Melnik), 1

Manganese

Komplexchemie perhalogenierter Cyclopentadiene und Alkine. XVI. Darstellung von π -Cyclopentadienylkomplexen mit zwei und drei Chalkogensubstituenten am Ring, $[\text{C}_5\text{Cl}_{4-m}\text{-}(\text{ER})_m(\text{ER}')]\text{ML}_n$ ($\text{ML}_n = \text{Mn}(\text{CO})_3, \text{Fe}(\text{C}_5\text{H}_5)$; $m = 1$ oder 2 ; $\text{E} = \text{S}, \text{Se}; \text{R}, \text{R}' = \text{Me}, \text{Ph}$). Struktur von $[\text{C}_5\text{Cl}_3(\text{SMe})_2]\text{IC}_5\text{H}_5\text{Fe}$ (K. Sünkel, U. Birk, A. Blum and W. Kempinger), 167

Photoabsorption study of electronic excited states of $(\eta^5\text{-cyclopentadienyl})(\eta^6\text{-arene})\text{manganese}$ (S.Yu. Ketkov), 225

Metallocarborane

Synthesis and magnetic studies of the metallocarborane sandwich salts $[\text{Fe}(\text{C}_5\text{Me}_5)_2\text{IM}(\text{C}_2\text{B}_9\text{H}_{12})_2]$ ($\text{M} = \text{Cr}, \text{Fe}, \text{Ni}$) (J.M. Forward, D.M.P. Mingos and A.V. Powell), 251

Metallocenes

Electronic effects of bis(2-aryl-4,5,6,7-tetrahydroindenyl)titano-cene dichlorides on the catalytic epoxidation of *trans*-3-hexene (R.L. Halterman and T.M. Ramsey), 175

Hydrogen-bonded complexes involving the metal atom and protonation of metallocenes of the iron subgroup (E.S. Shubina, A.N. Krylov, A.Z. Kreindlin, M.I. Rybinskaya and L.M. Epstein), 259

Komplexchemie perhalogenierter Cyclopentadiene und Alkine. XVI. Darstellung von π -Cyclopentadienylkomplexen mit zwei und drei Chalkogensubstituenten am Ring, $[\text{C}_5\text{Cl}_{4-m}\text{-}(\text{ER})_m(\text{ER}')]\text{ML}_n$ ($\text{ML}_n = \text{Mn}(\text{CO})_3, \text{Fe}(\text{C}_5\text{H}_5)$; $m = 1$ oder 2 ; $\text{E} = \text{S}, \text{Se}; \text{R}, \text{R}' = \text{Me}, \text{Ph}$). Struktur von $[\text{C}_5\text{Cl}_3(\text{SMe})_2]\text{IC}_5\text{H}_5\text{Fe}$ (K. Sünkel, U. Birk, A. Blum and W. Kempinger), 167

Molecular mechanics

Molecular mechanical studies on the olefin metathesis reaction.
I. Development and evaluation of tungsten carbene parameters: METMOD1 (L. Bencze and R. Szilagyi), 211

Phosphine-nitrile ligands: the molecular structure of *cis*-2-di-phenylphosphino-1-cyanocyclopentane and studies of the donor/acceptor properties of this and related ligands (E.J. Fredericks, M.J. Gindling, L.C. Kroll and B.N. Storhoff), 289

Molybdenum

Dinuclear molybdenum carbonyls bridged by hydride ligands (J.T. Lin, Y.C. Lee, H.M. Kao, T.-Y. Dong and Y.S. Wen), 199

Heterometallic sulfido clusters by metal exchange: syntheses and X-ray crystal structures of ($\eta^5\text{-C}_5\text{H}_4\text{CH}_3$) $\text{MoFeCo}(\text{CO})_8(\mu_3\text{-S})$ and ($\eta^5\text{-C}_5\text{H}_4\text{CH}_3$) $_2\text{Mo}_2\text{Fe}(\text{CO})_7(\mu_3\text{-S})$ (W.-H. Sun, S.-Y. Yang, H.-Q. Wang, Q.-F. Zhou and K.-B. Yu), 263

Mössbauer spectroscopy

Coordination complexes of dimethyltin dichloride with C-nitroso compounds (M. Cameron, B.G. Gowenlock, R.V. Parish and G. Vasapollo), 161

Organotin-substituted organosiloxanes: synthesis and characterisation, including the crystal structure of 1,3-bis(triphenylstannoxy)-1,1,3,3-tetraphenyl-1,3-disiloxane (B.J. Brisdon, M.F. Mahon, K.C. Molloy and P.J. Schofield), 145

Neopentyl

Some complexes of neopentylcadmium species with dithio- and di-selenocarbamates: the synthesis, characterization and single crystal X-ray structure of a mixed neopentyl/diethyldiselenocarbamate of cadmium: $[(\text{CH}_3)_3\text{CCH}_2\text{CdSe}_2\text{CNEt}_2]_2$ (I. Abrahams, A. Malik, M. Motavalli and P. O'Brien), 73

Nickel

Phosphine-nitrile ligands: the molecular structure of *cis*-2-di-phenylphosphino-1-cyanocyclopentane and studies of the donor/acceptor properties of this and related ligands (E.J. Fredericks, M.J. Gindling, L.C. Kroll and B.N. Storhoff), 289

Stabilisierung niederkоordinierter $\text{M}(\text{CO})$ -Bausteine: Synthese und Reaktivitat von ($\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$) $_2\text{Hf}(\text{C}\equiv\text{CPh})_2\text{M}(\text{CO})$ ($\text{M} = \text{Ni, Co}$) (H. Lang, M. Herres and W. Imhof), 283

Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275

Niobium

Syntheses and electrochemical behaviour of novel dithiolatonio-bium-(IV) and -(V) complexes with two η^5 -cyclopentadienyl rings (F. Guyon, J. Amaudrut, M.-F. Mercier and K. Shimizu), 187

Nitrobenzene

Etude lectrochimique des cations (η^5 -cyclopentadienyl fer η^6 -nitrobenzene ou η^6 -nitrotoluenes) $^+$. Electrosynth se des d riv s nitros s correspondants (N. Guennec and C. Moinet), 233

Nitrosobenzene

Etude lectrochimique des cations (η^5 -cyclopentadienyl fer η^6 -nitrobenzene ou η^6 -nitrotoluenes) $^+$. Electrosynth se des d riv s nitros s correspondants (N. Guennec and C. Moinet), 233

Nuclear magnetic resonance

Coordination complexes of dimethyltin dichloride with C-nitroso compounds (M. Cameron, B.G. Gowenlock, R.V. Parish and G. Vasapollo), 161

Olefin metathesis

Molecular mechanical studies on the olefin metathesis reaction.
I. Development and evaluation of tungsten carbene parameters: METMOD1 (L. Bencze and R. Szilagyi), 211

Oxidative addition

Palladium-catalyzed cross-coupling of aryl and alkenyl boronic acids with alkenes via oxidative addition of a carbon–boron bond to palladium(0) (C.S. Cho and S. Uemura), 85

Palladium

On the regioselectivity of palladium catalyzed cross-coupling reactions of alkenylsilanes: participation of β -cationic organosilicate-palladium species during the transmetalation (Y. Hatanaka, K. Goda and T. Hiyama), 97

Palladium-catalyzed cross-coupling of aryl and alkenyl boronic acids with alkenes via oxidative addition of a carbon–boron bond to palladium(0) (C.S. Cho and S. Uemura), 85

Phosphine-nitrile ligands: the molecular structure of *cis*-2-di-phenylphosphino-1-cyanocyclopentane and studies of the donor/acceptor properties of this and related ligands (E.J. Fredericks, M.J. Gindling, L.C. Kroll and B.N. Storhoff), 289

Reactions of unsaturated electrophiles with *trans*-(PMe_3) $_2\text{Pd}(\text{Ph})(\text{NHPH})$ (J.M. Boncella and L.A. Villanueva), 297

Phosphonium

Formation and *fac-mer* isomerization of cationic phosphonium transition metal complexes formulated as $[\text{M}(\text{bpy})\text{(CO)}_3\{\text{PN}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}\}]^+$ ($\text{M} = \text{Cr, Mo, W}$) and $[\text{M}(\text{dppe})\text{(CO)}_3\{\text{PN}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}\}]^+$ ($\text{M} = \text{Mo, W}$) (H. Nakazawa, Y. Yamaguchi and K. Miyoshi), 193

Phosphine-nitrile ligands

Phosphine-nitrile ligands: the molecular structure of *cis*-2-di-phenylphosphino-1-cyanocyclopentane and studies of the donor/acceptor properties of this and related ligands (E.J. Fredericks, M.J. Gindling, L.C. Kroll and B.N. Storhoff), 289

Phosphate

Formation and *fac-mer* isomerization of cationic phosphonium transition metal complexes formulated as $[\text{M}(\text{bpy})\text{(CO)}_3\{\text{PN}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}\}]^+$ ($\text{M} = \text{Cr, Mo, W}$) and $[\text{M}(\text{dppe})\text{(CO)}_3\{\text{PN}(\text{Me})\text{CH}_2\text{CH}_2\text{NMe}\}]^+$ ($\text{M} = \text{Mo, W}$) (H. Nakazawa, Y. Yamaguchi and K. Miyoshi), 193

Photoabsorption

Photoabsorption study of electronic excited states of (η^5 -cyclopentadienyl)(η^6 -arene)manganese (S.Yu. Ketkov), 225

Photochemistry

Photochemical reactions of 2-(pentamethyldisilanyl)furan and 2-(pentamethyldigermanyl)furan. Formation of a radical pair (K. Mochida, K. Kimijima, M. Wakasa and H. Hayashi), 101

Platinum

Phosphine-nitrile ligands: the molecular structure of *cis*-2-di-phenylphosphino-1-cyanocyclopentane and studies of the donor/acceptor properties of this and related ligands (E.J. Fredericks, M.J. Gindling, L.C. Kroll and B.N. Storhoff), 289

Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275

Polymer

Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275

Polymerization

Molecular mechanical studies on the olefin metathesis reaction.
I. Development and evaluation of tungsten carbene parameters: METMOD1 (L. Bencze and R. Szilagyi), 211

Radical

Photochemical reactions of 2-(pentamethyldisilanyl)furan and 2-(pentamethyldigermanyl)furan. Formation of a radical pair (K. Mochida, K. Kimijima, M. Wakasa and H. Hayashi), 101

Radiolysis

Gas phase reactivity of aromatic silanes. The reaction of $\text{Ph}(\text{CH}_2)_x\text{SiMe}_3$ ($x = 0$ or 1) with cationic electrophiles (M.E. Crestoni, S. Fornarini and G. Occhiucci), 109

Raman spectroscopy

Schwingungsspektren, Normalkoordinatenanalysen und Synthesen der t-Butylsilane ' BuSiX_3 , $X = \text{H}, \text{D}, \text{F}, \text{Cl}, \text{Br}, \text{I}$ (K. Hassler and M. Weidenbruch), 127

Raman spectroscopy

Schwingungsspektren und Normalkoordinatenanalysen der Di-t-butylsilane ' Bu_2SiX_2 , $X = \text{H}, \text{D}, \text{F}, \text{Cl}, \text{Br}, \text{I}$ (K. Hassler and M. Weidenbruch), 137

Regioselectivity

On the regioselectivity of palladium catalyzed cross-coupling reactions of alkenylsilanes: participation of β -cationic organosilicate-palladium species during the transmetallation (Y. Hatanaka, K. Goda and T. Hiyama), 97

Rhodium

N-substituted imidazole derivatives of rhodium(I) and iridium(I).

Crystal structure of the head-to-head dinuclear compound [$\text{Ir}_2(\mu\text{-N-benzylimidazolato-N}^3,\text{C}^2)_2(\text{CO})_4$] (F. Bonati, L.A. Oro, M.T. Pinillos, C. Tejel and B. Bovio), 267

Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275

Ruthenium

Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275

Selenium

Komplexchemie perhalogenierter Cyclopentadiene und Alkine.

XVI. Darstellung von π -Cyclopentadienylkomplexen mit zwei und drei Chalkogensubstituenten am Ring, $[\text{C}_5\text{Cl}_{4-m}-(\text{ER})_m(\text{ER}')]\text{ML}_n$ ($\text{ML}_n = \text{Mn}(\text{CO})_3, \text{Fe}(\text{C}_5\text{H}_5)$; $m = 1$ oder 2 ; $E = \text{S}, \text{Se}$; $R, R' = \text{Me}, \text{Ph}$). Struktur von $[\text{C}_5\text{Cl}_3-(\text{SMe})_2]\text{C}_5\text{H}_5\text{Fe}$ (K. Sünkel, U. Birk, A. Blum and W. Kempinger), 167

Silane

Gas phase reactivity of aromatic silanes. The reaction of $\text{Ph}(\text{CH}_2)_x\text{SiMe}_3$ ($x = 0$ or 1) with cationic electrophiles (M.E. Crestoni, S. Fornarini and G. Occhiucci), 109

On the regioselectivity of palladium catalyzed cross-coupling reactions of alkenylsilanes: participation of β -cationic organosilicate-palladium species during the transmetallation (Y. Hatanaka, K. Goda and T. Hiyama), 97

Schwingungsspektren, Normalkoordinatenanalysen und Synthesen der t-Butylsilane ' BuSiX_3 , $X = \text{H}, \text{D}, \text{F}, \text{Cl}, \text{Br}, \text{I}$ (K. Hassler and M. Weidenbruch), 127

Schwingungsspektren und Normalkoordinatenanalysen der Di-t-butylsilane ' Bu_2SiX_2 , $X = \text{H}, \text{D}, \text{F}, \text{Cl}, \text{Br}, \text{I}$ (K. Hassler and M. Weidenbruch), 137

Silanyl

Photochemical reactions of 2-(pentamethylidisilanyl)furan and 2-(pentamethyldigermanyl)furan. Formation of a radical pair (K. Mochida, K. Kimijima, M. Wakasa and H. Hayashi), 101

Silicon

Stabilisierung niederkordinierter $\text{M}(\text{CO})$ -Bausteine: Synthese und Reaktivität von $\{(\eta^5\text{C}_5\text{H}_4\text{SiMe}_3)_2\text{Hf}(\text{C}\equiv\text{CPh})_2\}\text{M}(\text{CO})$ ($\text{M} = \text{Ni}, \text{Co}$) (H. Lang, M. Herres and W. Imhof), 283

Siloxanes

Organotin-substituted organosiloxanes: synthesis and characterization, including the crystal structure of 1,3-bis(triphenylstannoxy)-1,1,3,3-tetraphenyl-1,3-disiloxane (B.J. Brisdon, M.F. Mahon, K.C. Molloy and P.J. Schofield), 145

Silyl

Substituentenabhängigkeit der Lithiumkoordination von Phosphinomethaniden und Molekülstrukturen von $(\text{TMEDA})\text{Li}-[\text{C}(\text{PPH}_2)_2(\text{SiMe}_3)]$ und $\{(\text{THF})\text{Li}[\text{CH}(\text{PMe}_2)_2]\}_2$ (H.H. Karsch, G. Grauvogl, P. Mikulcik, P. Bissinger and G. Müller), 65

Stannox

Organotin-substituted organosiloxanes: synthesis and characterization, including the crystal structure of 1,3-bis(triphenylstannoxy)-1,1,3,3-tetraphenyl-1,3-disiloxane (B.J. Brisdon, M.F. Mahon, K.C. Molloy and P.J. Schofield), 145

Sulfido

Heterometallic sulfido clusters by metal exchange: syntheses and X-ray crystal structures of $(\eta^5\text{C}_5\text{H}_4\text{CH}_3)\text{MoFeCo}(\text{CO})_8(\mu_3-\text{S})$ and $(\eta^5\text{C}_5\text{H}_4\text{CH}_3)_2\text{Mo}_2\text{Fe}(\text{CO})_7(\mu_3-\text{S})$ (W.-H. Sun, S.-Y. Yang, H.-Q. Wang, Q.-F. Zhou and K.-B. Yu), 263

Sulfur

Komplexchemie perhalogenierter Cyclopentadiene und Alkine. XVI. Darstellung von π -Cyclopentadienylkomplexen mit zwei und drei Chalkogensubstituenten am Ring, $[\text{C}_5\text{Cl}_{4-m}-(\text{ER})_m(\text{ER}')]\text{ML}_n$ ($\text{ML}_n = \text{Mn}(\text{CO})_3, \text{Fe}(\text{C}_5\text{H}_5)$; $m = 1$ oder 2 ; $E = \text{S}, \text{Se}$; $R, R' = \text{Me}, \text{Ph}$). Struktur von $[\text{C}_5\text{Cl}_3-(\text{SMe})_2]\text{C}_5\text{H}_5\text{Fe}$ (K. Sünkel, U. Birk, A. Blum and W. Kempinger), 167

Supported catalyst

Synthesis of a polymer-supported *o*-phenylene(diphosphine) and characterization of some metal complexes (N.R. Champness, W. Levason, R.D. Oldroyd and D.J. Gulliver), 275

Thiolate

Syntheses and electrochemical behaviour of novel dithiolatotinibium-(IV) and -(V) complexes with two η^5 -cyclopentadienyl rings (F. Guyon, J. Amaudrut, M.-F. Mercier and K. Shimizu), 187

Tin

Chemie der schweren Carben-Analogen R_2M ($\text{M} = \text{Si}, \text{Ge}, \text{Sn}$). XIX. Reaktionen von stabilem Diaminogermanyl und -stannylen $\text{Me}_2\text{Si}(\text{BuN})_2\text{M}$ ($\text{M} = \text{Ge}, \text{Sn}$) mit organischen Aziden (B. Klein and W.P. Neumann), 119

Synthesis and properties of trifluoromethylated stannanes. The formation and NMR spectra of the difluoromethylstannanes $(\text{CF}_3)_n(\text{CF}_2\text{H})_m\text{Sn}(\text{CH}_3)_{4-n-m}$ (R. Eujen, N. Jahn and U. Thurmann), 153

Titanium

Electronic effects of bis(2-aryl-4,5,6,7-tetrahydroindenyl)titano-cene dichlorides on the catalytic epoxidation of *trans*-3-hexene (R.L. Halterman and T.M. Ramsey), 175

Stabilisierung niederkordinierter $\text{M}(\text{CO})$ -Bausteine: Synthese und Reaktivität von $\{(\eta^5\text{C}_5\text{H}_4\text{SiMe}_3)_2\text{Hf}(\text{C}\equiv\text{CPh})_2\}\text{M}(\text{CO})$ ($\text{M} = \text{Ni}, \text{Co}$) (H. Lang, M. Herres and W. Imhof), 283

Transmetallation

On the regioselectivity of palladium catalyzed cross-coupling reactions of alkenylsilanes: participation of β -cationic organosilicate-palladium species during the transmetallation (Y. Hatanaka, K. Goda and T. Hiyama), 97

Tungsten

Molecular mechanical studies on the olefin metathesis reaction. I. Development and evaluation of tungsten carbene parameters: METMOD1 (L. Bencze and R. Szilagyi), 211

Reactions of the cationic but-2-yne complex $[\text{W}(\text{CO})(\text{NCMe})-(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPH}_2)(\eta^2\text{-MeC}_2\text{Me})]\text{BF}_4^-$ with bidentate nitrogen donor ligands (P.K. Baker and K.R. Flower), 221

Unsaturated electrophiles

Reactions of unsaturated electrophiles with *trans*-(PMe₃)₂-Pd(Ph)(NHPH) (J.M. Boncella and L.A. Villanueva), 297

Vanadium

A route to η^2 -alkyne complexes of vanadium(I): crystal structure of *trans*-[IV(CO)₂(PMe₂Ph)₂(2-butyne)] (H. Gailus, H. Maelger and D. Rehder), 181

X-ray diffraction

Magnesium compounds: classification and analysis of crystallographic and structural data (C.E. Holloway and M. Melnik), 1
Synthesis and magnetic studies of the metallocarborane sandwich salts [Fe(C₅Me₅)₂]M(C₂B₉H₁₂)₂] (M = Cr, Fe, Ni) (J.M. Forward, D.M.P. Mingos and A.V. Powell), 251