

Subject Index of Volume 485

Acetato complexes

Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π -Alkin-, Alkynyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

Acetylide(alkynyl)

Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-C}\equiv\text{C-Fp})\text{Ru}_3(\mu\text{-H})(\text{CO})_9]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-C}_2\text{H}_x)\text{M}_y$ cluster compounds ($x = 0, 1; y = 7, 8$), $\text{CpFeRu}_6(\mu_5\text{-C}_2)(\mu_5\text{-C}_2\text{H})(\text{CO})_{16}$, $\text{Cp}_2\text{Fe}_2\text{Ru}_5(\mu_5\text{-C}_2)_2(\text{CO})_{17}$ and $\text{Cp}_2\text{Fe}_2\text{Ru}_6(\mu_6\text{-C}_2)_2(\text{CO})_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14

Alcohol

Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β -keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55

Alkynyl(hydrido) complexes

Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π -Alkin-, Alkynyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

Alkoxides

Synthesis and characterization of new alkoxide and aryloxy derivatives of titanium and zirconium. X-ray molecular structure of $[\text{ZrCp}_2(\text{OC}_6\text{F}_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Alkyl halide

First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19

Alkylidenes

Synthesis of a tris(pyrazolyl)borate-stabilized molybdenum alkylidene and its hydrolysis products. Crystal structures of $\text{TpMo}(\text{CH}_2\text{C}(\text{Me})_2\text{Ph})(\text{NAr})(\text{O})$ and $[\text{TpMo}(\text{NAr})(\text{O})]_2\text{O}$ (W.M. Vaughan, K.A. Abboud and J.M. Boncella), 37

Alkyne complexes

Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π -Alkin-, Alkynyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

Allyl

Severe distortion of π -allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π -cinnamyl)-dicar-

bonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1

Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β -keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55

Amine

Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β -keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55

Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127

Ansa-bridging

Ansa-[(2,2-bis- η -cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18

Arene

Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^1H , ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109

Aryl

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Aryl bismuth compounds

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Aurated clusters

Aurated clusters derived from $(\mu_3\text{-ketenylidene})\text{-Fe}_3$ and $-\text{Fe}_2\text{Co}$ clusters (C. Thöne and H. Vahrenkamp), 185

Azides

Reaction of $(\text{CO})_5\text{W}(\text{THF})$ with triphenylmethyl azide and triptycyl azide (S.T. Massey, B. Mansour and L. McElwee-White), 123

Binuclear complexes

Mononuclear(η^6 -arene)ruthenium(II) and (η^5 -pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)-platinum(II) and ruthenium(II)-rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

2,2-bis- η -cyclopentadienyl(propane)

Ansa-[(2,2-bis- η -cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18

Bismuth

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, ($^1\text{h-C}_5\text{H}_5$)₃Bi" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Borate

Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127

Boron

He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215

Organotin(IV) polypyrazolylborates. VII. Hydridotris(3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of K(HB(3,4,5-Me₃Pz)₃) and [HB(3,4,5-Me₃Pz)₃ SnMeCl₂] (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45

Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127

Bridging ligand

Mononuclear(η^6 -arene)ruthenium(II) and (η^5 -pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)-platinum(II) and ruthenium(II)-rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

Bulky ligands

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Cage

Bimetallic siloxane cluster of higher valent transition metals: Na[(η^6 -cyclo-(PhSiO₂)₆)₂Co₂Ni₄(μ_6 -Cl)] (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257

Carbanions

Synthesis and structure of triphenylmethyl derivatives of ytterbium [Yb(THF)₆]²⁺[Ph₃C]₃⁻ and [(THF)₄Yb(μ -Cl)₂Yb(THF)₄]²⁺[Ph₃Cl]₂⁻ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

Carbonyl

Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ¹H, ¹³C and ¹⁷O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109

Carboxylic acid synthesis

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. XI. Synthese und Charakterisierung von [Li(TMEDA)]₂Ni(CH₂CH₂CH₂CH₂C(O)O)₂ und seine Umsetzung mit CH₃I und einem C₂₂-Steroidiodid (R. Wyrwa, H.-O. Fröhlich und R. Fischer), 227

Catalyst

Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethylamine]bis[μ -(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethylamine]bis[μ -(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209

Catalyst

Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β -keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55

C–C-bond formation

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. XI. Synthese und Charakterisierung von [Li(TMEDA)]₂Ni(CH₂CH₂CH₂CH₂C(O)O)₂ und seine Umsetzung mit CH₃I und einem C₂₂-Steroidiodid (R. Wyrwa, H.-O. Fröhlich und R. Fischer), 227

Charge-transfer

Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165

Chromium

Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ¹H, ¹³C and ¹⁷O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109

Cluster

Bimetallic siloxane cluster of higher valent transition metals: Na[(η^6 -cyclo-(PhSiO₂)₆)₂Co₂Ni₄(μ_6 -Cl)] (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257

Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Structure expansion of the mixed-metal dicarbide cluster core in [(μ_3 -C≡C-Fp)Ru₃(μ -HXCO)₉] via direct radical and thermal coupling reactions leading to higher-nuclearity (μ -C₂H_x)_y cluster compounds ($x = 0, 1; y = 7, 8$), CpFeRu₆(μ_5 -C₂)(μ_5 -C₂H)(CO)₁₆, Cp₂Fe₂Ru₅(μ_5 -C₂)₂(CO)₁₇ and Cp₂Fe₂Ru₆(μ_6 -C₂)₂(CO)₁₇ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14

Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10

Cobalt

Aurated clusters derived from (μ_3 -ketenylidene)-Fe₃ and -Fe₂Co clusters (C. Thöne and H. Vahrenkamp), 185

Bimetallic siloxane cluster of higher valent transition metals: Na[(η^6 -cyclo-(PhSiO₂)₆)₂Co₂Ni₄(μ_6 -Cl)] (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257

Formation constants of some phosphine and phosphite adducts of (η^5 -cyclopentadienyl) (substituted 1,2-ethylenedicalcogenolato)cobalt(III) complexes and their ¹H, ¹³C and ³¹P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31

Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ¹H, ¹³C and ¹⁷O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109

Reactions of diacetylene ligands with trinuclear clusters. I. Reactions of 2,4-hexadiyne-1,6-diol and its dicobalthexacarbonyl derivatives with H₂Os₃(CO)₁₀ (M.G. Karpov, S.P. Tunik, V.R.

- Denisov, G.L. Starova, A.B. Nikol'skii, F.M. Dolgushin, A.I. Yanovsky and Yu.T. Struchkov), 219
- Cone angles**
The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[\text{Ag}_2\text{Ru}_4(\mu_3\text{-H})_2(\text{CO})_{12} \{\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3\}_2]$. An investigation of the steric properties of the $\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191
- Coupling**
Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-C}\equiv\text{C-Fp})\text{Ru}_3(\mu\text{-H})(\text{CO})_9]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-C}_2\text{H}_x)\text{M}_y$ cluster compounds ($x = 0, 1; y = 7, 8$), $\text{CpFeRu}_6(\mu_5\text{-C}_2)(\mu_5\text{-C}_2\text{H})(\text{CO})_{16}$, $\text{Cp}_2\text{Fe}_2\text{Ru}_5(\mu_5\text{-C}_2)_2(\text{CO})_{17}$ and $\text{Cp}_2\text{Fe}_2\text{Ru}_6(\mu_6\text{-C}_2)_2(\text{CO})_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14
- Crystal structure**
Crystal and molecular structure of [1-(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161
Polysulfonamide. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di- μ -hydroxo-bis[(dimesylamido-*N*)dimethylzinn(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179
Severe distortion of π -allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π -cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1
Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6
Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165
The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[\text{Ag}_2\text{Ru}_4(\mu_3\text{-H})_2(\text{CO})_{12} \{\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3\}_2]$. An investigation of the steric properties of the $\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191
Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10
- Cyclohexane**
Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β -keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55
- Cyclometallated derivative**
Crystal and molecular structure of [1-(4-chlorophenyl)imino]ethyl] ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161
- Cyclopentadienyl**
Formation constants of some phosphine and phosphite adducts of (η^5 -cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H , ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31
Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[\text{ZrCp}_2(\text{OC}_6\text{F}_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153
Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylobismuth, ($^1\text{h-C}_5\text{H}_5$) $_3\text{Bi}$ " (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149
Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6
- Cyclopolyphosphine**
Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10
- Cyclosiloxane**
Bimetallic siloxane cluster of higher valent transition metals: $\text{Na}\{[\eta^6\text{-cyclo-(PhSiO}_2)_6]_2\text{Co}_2\text{Ni}_4(\mu_6\text{-Cl})\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257
- Dicarbide**
Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-C}\equiv\text{C-Fp})\text{Ru}_3(\mu\text{-H})(\text{CO})_9]$ via direct radical and thermal coupling reactions leading to higher-nuclearity $(\mu\text{-C}_2\text{H}_x)\text{M}_y$ cluster compounds ($x = 0, 1; y = 7, 8$), $\text{CpFeRu}_6(\mu_5\text{-C}_2)(\mu_5\text{-C}_2\text{H})(\text{CO})_{16}$, $\text{Cp}_2\text{Fe}_2\text{Ru}_5(\mu_5\text{-C}_2)_2(\text{CO})_{17}$ and $\text{Cp}_2\text{Fe}_2\text{Ru}_6(\mu_6\text{-C}_2)_2(\text{CO})_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14
- Dichlorosilanes**
Reactivity of dichlorosilanes with lithium ester enolates: synthesis of 3,3-disubstituted 3-silaglutarates (A. Djerourou and L. Blanco), 63
- Diketones**
Ferrocenyl salts as synthons: new ferrocenyl-1,3-diketones (C.M. Zakaria, C.A. Morrison, D. McAndrew, W. Bell and C. Glidewell), 201
- Dimesylamide**
Polysulfonamide. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di- μ -hydroxo-bis[(dimesylamido-*N*)dimethylzinn(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179
- 2,6-Dimethoxyphenyl**
Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127
- Diselenides**
First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19
- Disulfide ligand**
Comparative studies in the $\text{Cp}_2\text{M}(\eta^2\text{-S}_2)\text{H}$ series (Cp = *t*-BuC₅H₄, C₅Me₄Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237
- Ferrocene**
Ferrocenyl salts as synthons: new ferrocenyl-1,3-diketones (C.M. Zakaria, C.A. Morrison, D. McAndrew, W. Bell and C. Glidewell), 201

Ferrocenes

Synthesis and characterization of Fe(1-Me₃Si-3-CH₃C₅H₅)₂, an open ferrocene derived from an unsymmetric pentadienyl ligand (W. Trakarnpruk, A.M. Arif and R.D. Ernst), 25

Ferrocenylimine

Crystal and molecular structure of [1-(4-chlorophenyl)imino]ethyl ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161

Functionalised polysilanes

Synthese und Eigenschaften funktioneller Polysilane: Die Tetrasilane MeSi(SiMe₂X)₃ und Hexasilane (Me₂XSi)₂MeSiSiMe(SiMe₂X)₂, X = Me, H, F, Cl, Br, I (G. Kollegger und K. Hassler), 233

Group 4

Synthesis and characterization of new alkoxide and aryloxy derivatives of titanium and zirconium. X-ray molecular structure of [ZrCp₂(OC₆F₅)₂] (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Group 11

The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster [Ag₂Ru₄(μ₃-H)₂(CO)₁₂{P(C₆H₄Me-2)₃}₂]. An investigation of the steric properties of the P(C₆H₄Me-2)₃ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191

Group 15

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, (1^h-C₅H₅)₃Bi" (J. Lorbërth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Hafnium

He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215

Heterobimetallics

Bimetallic siloxane cluster of higher valent transition metals: Na[η⁶-cyclo-(PhSiO₂)₆]₂Co₂Ni₄(μ₆-Cl) (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257

Holmium

Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6

Hunsdiecker reaction

Surprisingly facile decomposition of the dication [(C₅Me₅)Ir(MeO-C₆H₄-CH₂CO₂NC₂H₄CO)]²⁺: a metal-mediated Hunsdiecker reaction of a succinimidyl ester? (H. El Amouri, M. Gruselle, J. Vaissermann, M.J. McGlinchey and G. Jaouen), 79

Hydride ligand

Comparative studies in the Cp₂M(η²-S₂)H series (Cp = t-BuC₅H₄, C₅Me₄Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237

Hydrogenation

Dicarbonylbis[2-(diphenylphosphino)-N,N-dimethylethylamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-N,N,N-trimethylethylamine]bis[μ-(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209

Hydrogen bonding

Molekül- und Kristallstruktur von μ-Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von [(μ-OH)Rh(COD-1.5)]₂ mit [η⁵-C₅(CH₃)₅]₂Ti(IV)(CH₃)₂ und (η⁵-C₅H₅)₂Ti(II)[P(CH₃)₃]₂ (D. Selent und M. Ramm), 135

Hydrolysis

Polysulfonamide. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-N)dimethylzinn(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179

Hydroxide

Polysulfonamide. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di-μ-hydroxo-bis[(dimesylamido-N)dimethylzinn(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179

Hydroxylamine

Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Iridium

Surprisingly facile decomposition of the dication [(C₅Me₅)Ir(MeO-C₆H₄-CH₂CO₂NC₂H₄CO)]²⁺: a metal-mediated Hunsdiecker reaction of a succinimidyl ester? (H. El Amouri, M. Gruselle, J. Vaissermann, M.J. McGlinchey and G. Jaouen), 79

Iron

Aurated clusters derived from (μ₃-ketenylidene)-Fe₃ and -Fe₂Co clusters (C. Thöne and H. Vahrenkamp), 185

Crystal and molecular structure of [1-(4-chlorophenyl)imino]ethyl ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161

Ferrocenyl salts as synthons: new ferrocenyl-1,3-diketones (C.M. Zakaria, C.A. Morrison, D. McAndrew, W. Bell and C. Glidewell), 201

Structure expansion of the mixed-metal dicarbide cluster core in [(μ₃-C≡C-Fp)Ru₃(μ-H)(CO)₉] via direct radical and thermal coupling reactions leading to higher-nuclearity (μ-C₂H₂)_{M_y} cluster compounds (x = 0, 1; y = 7, 8), CpFeRu₆(μ₅-C₂H)(μ₅-C₂H)(CO)₁₆, Cp₂Fe₂Ru₅(μ₅-C₂)(CO)₁₇ and Cp₂Fe₂Ru₆(μ₆-C₂)(CO)₁₇ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14

Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165

Isomerization

Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoacetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π-Alkin-, Alkynyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

Ketenylidene-Fe₃-cluster

Aurated clusters derived from (μ₃-ketenylidene)-Fe₃ and -Fe₂Co clusters (C. Thöne and H. Vahrenkamp), 185

Ketenylidene-Fe₂Co-cluster

Aurated clusters derived from (μ₃-ketenylidene)-Fe₃ and -Fe₂Co clusters (C. Thöne and H. Vahrenkamp), 185

Lanthanide complexes

Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6

Synthesis and structure of triphenylmethyl derivatives of ytterbium $[\text{Yb}(\text{THF})_6]^{2+}[\text{Ph}_3\text{C}]_2^-$ and $[(\text{THF})_4\text{Yb}(\mu\text{-Cl})_2\text{Yb}(\text{THF})_4]^{2+}[\text{Ph}_3\text{C}]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

Lithium enolates

Reactivity of dichlorosilanes with lithium ester enolates: synthesis of 3,3-disubstituted 3-silaglutarates (A. Djerourou and L. Blanco), 63

Magnetic susceptibility

Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165

Mercury

Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl]ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161

Metallocenes

Comparative studies in the $\text{Cp}_2\text{M}(\eta^2\text{-S}_2)\text{H}$ series ($\text{Cp} = \text{t-BuC}_5\text{H}_4$, $\text{C}_5\text{Me}_4\text{Et}$; $\text{M} = \text{Nb}$, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237

Synthesis and characterization of new one-carbon-bridged titanocene and zirconocene derivatives (R. Fierro, M.D. Rausch, G.S. Herrman and H.G. Alt), 11

Molybdenum

Ansa-[(2,2-bis- η -cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18

Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Severe distortion of π -allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π -cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1

Synthesis of a tris(pyrazolyl)borate-stabilized molybdenum alkylidene and its hydrolysis products. Crystal structures of $\text{TpMo}(\text{CH}_2\text{C}(\text{Me})_2\text{Ph})(\text{NAr})(\text{O})$ and $[\text{TpMo}(\text{NAr})(\text{O})]_2\text{O}$ (W.M. Vaughan, K.A. Abboud and J.M. Boncella), 37

2-Naphthoyltrifluoroacetato

Synthesis and structural investigation of lanthanide organometallics involving cyclopentadienyl and 2-naphthoyltrifluoroacetato chelate ligands (F. Shen, J. Hu, M. Xie, S. Wang and X. Huang), C6

Nickel

Bimetallic siloxane cluster of higher valent transition metals: $\text{Na}\{\{\eta^6\text{-cyclo}(\text{PhSiO}_2)_6\}_2\text{Co}_2\text{Ni}_4(\mu_6\text{-Cl})\}$ (G. Gavioli, R. Battistuzzi, P. Santi, C. Zucchi, G. Pályi, R. Ugo, A. Vizi-Orosz, O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman and A.A. Zhdanov), 257

Synthesis of highly crowded cyclopentadienes via reactions of Cp_2Ni and $\text{CpSn}(\text{CH}_3)_3$ with triarylmethylhalogenides. Molecular structure of 1-(CH_3)₃Si-(3- Ph_3C)- C_5H_4 (K.A. Rufanov, N.B. Kazennova, A.V. Churakov, D.A. Lemenovskii and L.G. Kuz'mina), 173

Nickelaoxacycloheptanones

Beiträge zur Chemie organometallischer metallacyclischer Nebengruppenmetallverbindungen. XI. Synthese und Charakterisierung von $[\text{Li}(\text{TMEDA})]_2\text{Ni}(\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{O})_2$ und seine Umsetzung mit CH_3I und einem C_{22} -Steroidiodid (R. Wyrwa, H.-O. Fröhlich und R. Fischer), 227

Nitrenes

Reaction of $(\text{CO})_5\text{W}(\text{THF})$ with triphenylmethyl azide and triptycyl azide (S.T. Massey, B. Mansour and L. McElwee-White), 123

¹⁷O-NMR spectroscopy

Mehrfachbindungen zwischen Übergangsmetallen und Hauptgruppenelementen. CXXXIV. ¹⁷O-NMR-Spektroskopie an Organorhenium(VII)-oxiden (W.A. Herrmann, F.E. Kühn und P.W. Roesky), 243

Nuclear magnetic resonance

Formation constants of some phosphine and phosphite adducts of (η^5 -cyclopentadienyl) (substituted 1,2-ethylenedichalogenolato)cobalt(III) complexes and their ¹H, ¹³C and ³¹P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31

Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ¹H, ¹³C and ¹⁷O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109

Severe distortion of π -allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π -cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1

Open ferrocenes

Synthesis and characterization of $\text{Fe}(1\text{-Me}_3\text{Si-3-CH}_3\text{C}_5\text{H}_5)_2$, an open ferrocene derived from an unsymmetric pentadienyl ligand (W. Trakarnpruk, A.M. Arif and R.D. Ernst), 25

Organorhenium(VII) oxides

Mehrfachbindungen zwischen Übergangsmetallen und Hauptgruppenelementen. CXXXIV. ¹⁷O-NMR-Spektroskopie an Organorhenium(VII)-oxiden (W.A. Herrmann, F.E. Kühn und P.W. Roesky), 243

Osmium

Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69

Reactions of diacetylene ligands with trinuclear clusters. I. Reactions of 2,4-hexadiyne-1,6-diol and its dicobalthexacarbonyl derivatives with $\text{H}_2\text{Os}_3(\text{CO})_{10}$ (M.G. Karpov, S.P. Tunik, V.R. Denisov, G.L. Starova, A.B. Nikol'skii, F.M. Dolgushin, A.I. Yanovsky and Yu.T. Struchkov), 219

Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10

Palladium

Crystal and molecular structure of [1-[(4-chlorophenyl)imino]ethyl]ferrocene and a structural comparison with its cyclometallated derivatives (Y.J. Wu, S.Q. Huo and Y. Zhu), 161

Pentadienyl complexes

Synthesis and characterization of $\text{Fe}(1\text{-Me}_3\text{Si-3-CH}_3\text{C}_5\text{H}_5)_2$, an open ferrocene derived from an unsymmetric pentadienyl ligand (W. Trakarnpruk, A.M. Arif and R.D. Ernst), 25

Phosphine

Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethylamine]bis[μ -(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethylaminium]bis[μ -(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209

Formation constants of some phosphine and phosphite adducts of (η^5 -cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H , ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31

Phosphine ligands

The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[\text{Ag}_2\text{Ru}_4(\mu_3\text{-H})_2(\text{CO})_{12} \{\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3\}_2]$. An investigation of the steric properties of the $\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191

Phosphonium

Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127

Phosphorus

Triosmium carbonyl cluster derivatives containing pentameric phenyl cyclopolyphosphine (H.-G. Ang, S.-G. Ang, W.-L. Kwik and Q. Zhang), C10

Photoelectron spectroscopy

He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zánthy), 215

Platinum

Mononuclear(η^6 -arene)ruthenium(II) and (η^5 -pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)-platinum(II) and ruthenium(II)-rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

Properties

Synthese und Eigenschaften funktioneller Polysilane: Die Tetrasilane $\text{MeSi}(\text{SiMe}_2\text{X})_3$ und Hexasilane $(\text{Me}_2\text{XSi})_2\text{MeSiSiMe}(\text{SiMe}_2\text{X})_2$, X = Me, H, F, Cl, Br, I (G. Kollegger und K. Hassler), 233

Pyrazole

Severe distortion of π -allyl orientation in a Molybdenum complex containing a sterically demanding ligand: Crystal structure of Hydrotris(3,5-dimethylpyrazolyl)borato-(π -cinnamyl)-dicarbonyl molybdenum(II) (V.S. Joshi, K.M. Sathe, M. Nandi, P. Chakrabarti and A. Sarkar), C1

Pyrazolylborate

Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), I65

Pyrazolylborates

Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of $\text{K}[\text{HB}(\text{3,4,5-Me}_3\text{Pz})_3]$ and $[\text{HB}(\text{3,4,5-Me}_3\text{Pz})_3 \text{SnMeCl}_2]$ (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45

Reactivity

Comparative studies in the $\text{Cp}_2\text{M}(\eta^2\text{-S}_2)\text{H}$ series (Cp = *t*- BuC_5H_4 , $\text{C}_5\text{Me}_4\text{Et}$; M = Nb, Ta) on the reactivity of hydride

and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237

Red modification

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, ($^1\text{h-C}_5\text{H}_5)_3\text{Bi}$ " (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Rhodium

Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethylamine]bis[μ -(2-methyl)-2-propanethiolato]dirhodium and dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethylaminium]bis[μ -(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209

Molekül- und Kristallstruktur von μ -Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von $[(\mu\text{-OH})\text{Rh}(\text{COD-1.5})_2]$ mit $[\eta^5\text{-C}_5(\text{CH}_3)_5]_2\text{Ti}(\text{IV})(\text{CH}_3)_2$ und $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{II})[\text{P}(\text{CH}_3)_3]_2$ (D. Selent und M. Ramm), 135

Mononuclear(η^6 -arene)ruthenium(II) and (η^5 -pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)-platinum(II) and ruthenium(II)-rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π -Alkin-, Alkynyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85

Ruthenium

Mono-, bi- and polynuclear complexes of diphenylmethane with Cr, Co and Ru. Synthesis and investigation by ^1H , ^{13}C and ^{17}O NMR (E. Kolehmainen, K. Laihia, J. Korvola, V.S. Kaganovich, M.I. Rybinskaya and Z.A. Kerzina), 109

Mononuclear(η^6 -arene)ruthenium(II) and (η^5 -pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)-platinum(II) and ruthenium(II)-rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

Reactions of dodecacarbonyltriruthenium with enones: η^3 -coordinated dihydropyranil rings (L.V. Rybin, S.V. Osintseva, M.I. Rybinskaya, F.M. Dolgushin, A.I. Yanovsky and Yu.T. Struchkov), 253

Structure expansion of the mixed-metal dicarbide cluster core in $[(\mu_3\text{-C}\equiv\text{C-Fp})\text{Ru}_3(\mu\text{-H})(\text{CO})_9]$ via direct radical and thermal coupling reactions leading to higher-nuclearity ($\mu\text{-C}_2\text{H}_x$) M_y cluster compounds ($x = 0, 1$; $y = 7, 8$), $\text{CpFeRu}_6(\mu_5\text{-C}_2)(\mu_5\text{-C}_2\text{H})(\text{CO})_{16}$, $\text{Cp}_2\text{Fe}_2\text{Ru}_5(\mu_5\text{-C}_2)_2(\text{CO})_{17}$ and $\text{Cp}_2\text{Fe}_2\text{Ru}_6(\mu_6\text{-C}_2)_2(\text{CO})_{17}$ (M. Akita, H. Hirakawa, M. Tanaka and Y. Moro-oka), C14

Syntheses of cyclohexenone derivatives by the ruthenium complex-catalysed reactions of allylic compounds with β -keto esters (S.-W. Zhang, T. Mitsudo, T. Kondo and Y. Watanabe), 55

The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[\text{Ag}_2\text{Ru}_4(\mu_3\text{-H})_2(\text{CO})_{12} \{\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3\}_2]$. An investigation of the steric properties of the $\text{P}(\text{C}_6\text{H}_4\text{Me-2})_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191

Selenium

First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl dis-

- lenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19
- Selenoamides**
First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19
- Selenourea**
First example of selenium transfer reaction of primary selenoamides and selenourea. Novel synthesis of dialkyl diselenides from alkyl halides (M.-D. Ruan, H.-R. Zhao, W.-Q. Fan and X.-J. Zhou), 19
- 3-Silaglutarates**
Reactivity of dichlorosilanes with lithium ester enolates: synthesis of 3,3-disubstituted 3-silaglutarates (A. Djerourou and L. Blanco), 63
- Silicon**
Synthesis of highly crowded cyclopentadienes via reactions of Cp_2Ni and $\text{CpSn}(\text{CH}_3)_3$ with triarylmethylhalogenides. Molecular structure of 1-(CH_3)₃Si-(3- Ph_3C)- C_5H_4 (K.A. Rufanov, N.B. Kazennova, A.V. Churakov, D.A. Lemenovskii and L.G. Kuz'mina), 173
- Silver**
The heteronuclear cluster chemistry of the Group IB metals. Part XVII. Synthesis and X-ray crystal structure of the hexanuclear mixed-metal cluster $[\text{Ag}_2\text{Ru}_4(\mu_3\text{-H})_2(\text{CO})_{12} \{\text{P}(\text{C}_6\text{H}_4\text{Me}_2)_3\}_2]$. An investigation of the steric properties of the $\text{P}(\text{C}_6\text{H}_4\text{Me}_2)_3$ ligand (P.J. McCarthy, I.D. Salter and T. Adatia), 191
- Solubility**
Tris(2,6-dimethoxyphenyl)borane and its adducts (M. Wada, M. Kanzaki, H. Ogura, S. Hayase and T. Erabi), 127
- Sulfur**
Formation constants of some phosphine and phosphite adducts of (η^5 -cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H , ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31
- Synthesis**
Synthese und Eigenschaften funktioneller Polysilane: Die Tetrasilane $\text{MeSi}(\text{SiMe}_2\text{X})_3$ und Hexasilane $(\text{Me}_2\text{XSi})_2\text{MeSiSiMe}(\text{SiMe}_2\text{X})_2$, X = Me, H, F, Cl, Br, I (G. Kollegger und K. Hassler), 233
- Tantalum**
He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zanathy), 215
- TCNQ**
Synthesis, crystal structures and magnetic properties of salts containing bis[hydrotris(3,5-dimethyl-1-pyrazolyl)borate]iron(III) (S.J. Mason, C.M. Hill, V.J. Murphy, D. O'Hare and D.J. Watkin), 165
- Thiolate**
Formation constants of some phosphine and phosphite adducts of (η^5 -cyclopentadienyl) (substituted 1,2-ethylenedichalcogenolato)cobalt(III) complexes and their ^1H , ^{13}C and ^{31}P NMR spectra (M. Kajitani, A. Igarashi, H. Hatano, T. Akiyama, A. Sugimori, S. Matsumoto, Y. Iguchi, H. Boennemann, K. Shimizu and G.P. Satô), 31
- Tin**
Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of $\text{K}(\text{HB}(3,4,5\text{-Me}_3\text{Pz})_3)$ and $[\text{HB}(3,4,5\text{-Me}_3\text{Pz})_3 \text{SnMeCl}_2]$ (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45
- Polysulfonilamine. LX. Ein dimeres Diorganozinnhydroxid mit einer Zinn-Stickstoff-Bindung: Bildung und Röntgenstrukturanalyse von Di- μ -hydroxo-bis[(dimesylamido-*N*)dimethylzinn(IV)] (I. Lange, P.G. Jones und A. Blaschette), 179
- Synthesis of highly crowded cyclopentadienes via reactions of Cp_2Ni and $\text{CpSn}(\text{CH}_3)_3$ with triarylmethylhalogenides. Molecular structure of 1-(CH_3)₃Si-(3- Ph_3C)- C_5H_4 (K.A. Rufanov, N.B. Kazennova, A.V. Churakov, D.A. Lemenovskii and L.G. Kuz'mina), 173
- Titanium**
Molekül- und Kristallstruktur von μ -Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von $[(\mu\text{-OH})\text{Rh}(\text{COD}\text{-}1,5)]_2$ mit $\{\eta^5\text{-C}_5(\text{CH}_3)_5\}_2\text{Ti}(\text{IV})(\text{CH}_3)_2$ und $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{II})[\text{P}(\text{CH}_3)_3]_2$ (D. Selent und M. Ramm), 135
- Synthesis and characterization of new alkoxide and aryloxy derivatives of titanium and zirconium. X-ray molecular structure of $[\text{ZrCp}_2(\text{OC}_6\text{F}_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153
- Synthesis and characterization of new one-carbon-bridged titanocene and zirconocene derivatives (R. Fierro, M.D. Rausch, G.S. Herrman and H.G. Alt), 11
- Triphenylmethyl derivatives**
Synthesis and structure of triphenylmethyl derivatives of ytterbium $[\text{Yb}(\text{THF})_6]^{2+} [\text{Ph}_3\text{C}]^-$ and $[(\text{THF})_4\text{Yb}(\mu\text{-Cl})_2\text{Yb}(\text{THF})_4]^{2+} [\text{Ph}_3\text{C}]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101
- 2,4,6-Triphenylphenyl**
Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141
- Tris-cyclopentadienyl bismuth**
Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, ($^1\text{h-C}_5\text{H}_5$)₃Bi" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149
- Trispyrazolylborates**
Synthesis of a tris(pyrazolyl)borate-stabilized molybdenum alkylidene and its hydrolysis products. Crystal structures of $\text{TpMo}(\text{CH}_2\text{C}(\text{Me})_2\text{Ph})(\text{NAr})(\text{O})$ and $[\text{TpMo}(\text{NAr})(\text{O})]_2\text{O}$ (W.M. Vaughan, K.A. Abboud and J.M. Boncella), 37
- Tungsten**
Ansa-(2,2-bis- η -cyclopentadienyl)propane]-molybdenum and -tungsten chemistry and related studies (L. Labella, A. Chernega and M.L.H. Green), C18
- Hydroxylamines as new decarbonylating reagents in the synthesis of metal carbonyl derivatives (V.A. Maksakov and V.A. Ershova), 69
- Reaction of $(\text{CO})_5\text{W}(\text{THF})$ with triphenylmethyl azide and triptycyl azide (S.T. Massey, B. Mansour and L. McElwee-White), 123
- Vinylidene complexes**
Vinyliden-Übergangsmetallkomplexe. XXX. Reaktionen von Acetato- und Trifluoracetato-Rhodium(I)-Komplexen mit 1-Alkinen: Ein Beispiel für die Balance zwischen isomeren π -Alkin-, Alkynyl(hydrido)- und Vinyliden-Metallverbindungen (M. Schäfer, J. Wolf und H. Werner), 85
- X-ray**
Organotin(IV) polypyrazolylborates. VII. Hydridotris (3,4,5-trimethyl-1*H*-pyrazol-1-yl)borates. X-ray crystal structures of

K(HB(3,4,5-Me₃Pz)₃) and [HB(3,4,5-Me₃Pz)₃ SnMeCl₂] (G. Gioia Lobbia, P. Cecchi, R. Spagna, M. Colapietro, A. Pifferi and C. Pettinari), 45

X-ray analysis

Molekül- und Kristallstruktur von μ -Hydroxy-cyclooctadien(1,5)-rhodium(I)-Dimer bei 173 K. Zur Reaktion von $[(\mu\text{-OH})\text{Rh}(\text{COD-1,5})_2]_2$ mit $[\eta^5\text{-C}_5(\text{CH}_3)_5\text{Ti}(\text{IV})(\text{CH}_3)_2]$ und $(\eta^5\text{-C}_5\text{H}_5)_2\text{Ti}(\text{II})[\text{P}(\text{CH}_3)_3]_2$ (D. Selent und M. Ramm), 135

X-ray diffraction

Dicarbonylbis[2-(diphenylphosphino)-*N,N*-dimethylethylamine]bis[μ -(2-methyl)-2-propanethiolato]dirhodium und dicarbonylbis[2-(diphenylphosphino)-*N,N,N*-trimethylethylamine]bis[μ -(2-methyl)-2-propanethiolato]dirhodium tetraphenylborate. Their syntheses, characterization and application as hydrogenation catalyst (H. Schumann, H. Hemling, N. Goren and J. Blum), 209

Synthesis and characterization of aryl bismuth compounds using 2,4,6-triphenylphenyl as a bulky ligand (X.-W. Li, J. Lorberth, W. Massa and S. Wocadlo), 141

Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[\text{ZrCp}_2(\text{OC}_6\text{F}_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Synthesis and crystal structure of E.O. Fischer's "red crystalline modification of tris-cyclopentadienylbismuth, ($^1\text{h-C}_5\text{H}_5$)₃Bi" (J. Lorberth, W. Massa, S. Wocadlo, I. Sarraje, S.-H. Shin and X.-W. Li), 149

Synthesis and structure of triphenylmethyl derivatives of ytterbium $[\text{Yb}(\text{THF})_6]^{2+}[\text{Ph}_3\text{C}]_2^-$ and $[(\text{THF})_4\text{Yb}(\mu\text{-Cl})_2\text{Yb}(\text{THF})_4]^{2+}[\text{Ph}_3\text{Cl}]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

X-ray structure

Aurated clusters derived from (μ_3 -ketenylidene)-Fe₃ and -Fe₂Co clusters (C. Thöne and H. Vahrenkamp), 185

Comparative studies in the Cp₂M(η^2 -S₂)H series (Cp = t-BuC₅H₄, C₅Me₄Et; M = Nb, Ta) on the reactivity of hydride and disulfide ligands (J.-C. Leblanc, C. Moise, F. Volpato, H. Brunner, G. Gehart, J. Wachter and B. Nuber), 237

Mononuclear(η^6 -arene)ruthenium(II) and (η^5 -pentamethylcyclopentadienyl)rhodium(III) and binuclear ruthenium(II)-platinum(II) and ruthenium(II)-rhodium(I) complexes containing 2-(diphenylphosphino)pyridine (D. Drommi, C.G. Arena, F. Nicolò, G. Bruno and F. Faraone), 115

Surprisingly facile decomposition of the dication $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{MeO-C}_6\text{H}_4\text{-CH}_2\text{CO}_2\text{NCOCH}_2\text{CH}_2\text{CO})]^{2+}$: a metal-mediated Hunsdiecker reaction of a succinimidyl ester? (H. El Amouri, M. Gruselle, J. Vaissermann, M.J. McGlinchey and G. Jaouen), 79

Ytterbium

Synthesis and structure of triphenylmethyl derivatives of ytterbium $[\text{Yb}(\text{THF})_6]^{2+}[\text{Ph}_3\text{C}]_2^-$ and $[(\text{THF})_4\text{Yb}(\mu\text{-Cl})_2\text{Yb}(\text{THF})_4]^{2+}[\text{Ph}_3\text{Cl}]_2^-$ (L.N. Bochkarev, N.E. Molosnova, L.N. Zakharov, G.K. Fukin, A.I. Yanovsky and Y.T. Struchkov), 101

Zirconium

He(I) photoelectron spectroscopic study of half- and bent sandwich tetrahydroborates (A. Nagy, L. Szepes, G. Vass and L. Zánthy), 215

Synthesis and characterization of new one-carbon-bridged titanocene and zirconocene derivatives (R. Fierro, M.D. Rausch, G.S. Herrman and H.G. Alt), 11

Synthesis and characterization of new alkoxide and aryloxide derivatives of titanium and zirconium. X-ray molecular structure of $[\text{ZrCp}_2(\text{OC}_6\text{F}_5)_2]$ (J.I. Amor, N.C. Burton, T. Cuenca, P. Gómez-Sal and P. Royo), 153

Book Reviews

Catalysis of Organic Reactions, John R. Kosak and Thomas A. Johnson (eds.) (J.R. Hanson), C22

Leaving No Stone Unturned: Pathways in Organometallic Chemistry F. Gordon A. Stone (*Profiles, Pathway, and Dreams: Autobiographies of Eminent Chemists*, Jeffrey I. Seeman, Series Editor) (M.F. Lappert), C22

Molybdenum Enzymes, Cofactors, and Model Systems, E.I. Stiefel, D. Coucouvanis and W.E. Newton (Eds.) (P.G.J. Leigh), C22