

Subject Index of Volume 495

Acetylacetonate- C^{γ} complexes

Synthesis and reactivity of acetylacetonato- C^{γ} complexes of M^{II} ($M = Pd$ or Pt): X-ray crystal structure of $[Pd(C_6F_5)(OOCPh)(bipy)] \cdot CHCl_3$ (J. Forniés, F. Martínez, R. Navarro and E.P. Urriolabeitia), 185

Alanes

Darstellung und Molekülstruktur von Bis(3-dimethylaminopropylalan) (U. Dümichen, K.-H. Thiele, T. Gelbrich und J. Sieler), 71

Alkylrhenium(VII)-oxide

Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXLVIII. Alkylrhenium(VII)-oxide—Synthese, Eigenschaften und Abbaureaktionen (W.A. Herrmann, F.E. Kühn und C.C. Romão), 209

Alkyl

Synthesis and characterization of titanium complexes containing the 1-(3-butenyl)-2,3,4,5-tetramethylcyclopentadienyl ligand (J. Okuda, K.E. Du Plooy and P.J. Toscano), 195

Alkyne

Lewis-Basen Addukte monomerer Bis(η^2 -alkin) Ag^I X-Verbindungen ($X = BF_4$ oder OSO_2CF_3) (H. Lang, K. Köhler und B. Schiemenz), 135

Alkyne ligands

Cluster chemistry. Part 97. Some unusual complexes obtained from reactions of $Ru_3(\mu-dppm)(CO)_{10}$ and C_7 or C_8 hydrocarbons (M.I. Bruce, J.R. Hinchliffe, B.W. Skelton and A.H. White), 141

Alkynes

Matrix-isolation studies of alkyne complexes in the photochemical reactions of $Mo(CO)_6$ with alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 149

The photochemical behaviour of matrix-isolated $W(CO)_6$ in the presence of alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 163

Aluminium

Darstellung und Molekülstruktur von Bis(3-dimethylaminopropylalan) (U. Dümichen, K.-H. Thiele, T. Gelbrich und J. Sieler), 71

Amide

Chemie polyfunktioneller Moleküle. CXVII. Synthese und Molekülstruktur von $[Co^{+1}(CO)_2\{Ph_2P \equiv N \equiv P(O)Ph_2^{\ominus}-P\}-(Ph_2P-NH-PPh_2)] \cdot 2 CH_3OH$ (D. Pohl, J. Ellermann, F.A. Knoch and M. Moll), C6

Amine

Chemie polyfunktioneller Moleküle. CXVII. Synthese und Molekülstruktur von $[Co^{+1}(CO)_2\{Ph_2P \equiv N \equiv P(O)Ph_2^{\ominus}-P\}-(Ph_2P-NH-PPh_2)] \cdot 2 CH_3OH$ (D. Pohl, J. Ellermann, F.A. Knoch and M. Moll), C6

Arsenic ligands

Solid angles. III. The role of conformers in solid angle calculations (D. White, B.C. Taverner, N.J. Coville and P.W. Wade), 41

Benzoferrylphosphine complex

Synthesis of the tertiaryphosphine derivatives of iron carbonyl phosphines: preparation of tetracarbonyl[tris(2-furyl)phosphine]iron(0), pentacarbonylbis[μ -bis(2-furyl)phosphido]-[tris(2-furyl)phosphine]diiron(0), tetracarbonyl[tris(2-benzofuryl)phosphine]iron(0) and pentacarbonylbis[μ -bis(2-benzofuryl)phosphido]-[tris(2-benzofuryl)phosphine]diiron(0) (C. Santelli-Rouvier, C. Coin, L. Toupet and M. Santelli), 91

Bifunctional ligands

Chemie polyfunktioneller Moleküle. CXVII. Synthese und Molekülstruktur von $[Co^{+1}(CO)_2\{Ph_2P \equiv N \equiv P(O)Ph_2^{\ominus}-P\}-(Ph_2P-NH-PPh_2)] \cdot 2 CH_3OH$ (D. Pohl, J. Ellermann, F.A. Knoch and M. Moll), C6

Bis(bis(tert.-butyl)fluorophosphine)-deca(carbonyl)-triosmium(0)

1Bu_2PF as a ligand in tri-osmium clusters (L. Heuer and D. Schomburg), 53

Bis(diphenylphosphino)methane

On the synthesis and $^{31}P\{^1H\}$ NMR spectrum of $(CO)_5W(\mu-PPh_2CH_2PPh_2)W(CO)_5$ (J.W. Benson, E.A. Keiter and R.L. Keiter), 77

Bis(tert.-butyl)fluorophosphine

1Bu_2PF as a ligand in tri-osmium clusters (L. Heuer and D. Schomburg), 53

Carbon monoxide

Substitution of SO_2 for CO in triruthenium carbonyl anions (G.B. Karet, C.L. Stern, J.A. Cody, S.J. Lange, M.A. Pell, C. Sleboznick and D.F. Shriver), 33

Carbonyl

Cluster chemistry. Part 97. Some unusual complexes obtained from reactions of $Ru_3(\mu-dppm)(CO)_{10}$ and C_7 or C_8 hydrocarbons (M.I. Bruce, J.R. Hinchliffe, B.W. Skelton and A.H. White), 141

Koordinationschemie funktioneller Phosphine. II. Carbonyl-(nitrosyl)wolfram-Komplexe mit 2-Diphenylphosphinoanisol sowie 2-Diphenylphosphinoanilid, -benzoat und -phenolat als Liganden (J. Pietsch, L. Dahlenburg, A. Wolski, H. Berke und I.L. Eremenko), 113

Matrix-isolation studies of alkyne complexes in the photochemical reactions of $Mo(CO)_6$ with alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 149

On the synthesis and $^{31}P\{^1H\}$ NMR spectrum of $(CO)_5W(\mu-PPh_2CH_2PPh_2)W(CO)_5$ (J.W. Benson, E.A. Keiter and R.L. Keiter), 77

Photochemical reaction of (η^5 -cyclopentadienyl) dicarbonyliron iodide with 1-substituted uracils in the presence of diisopropylamine: crystal structure of the (η^5 - C_5H_5) $Fe(CO)_2$ complex of deprotonated 5-fluoro-1-(tetrahydro-2-furyl)uracil (Ftorafur) (J. Zakrzewski, A. Tosik and M. Bukowska-Strzyżewska), 83

- Substitution of SO₂ for CO in triruthenium carbonyl anions (G.B. Karet, C.L. Stern, J.A. Cody, S.J. Lange, M.A. Pell, C. Slebočnick and D.F. Shriver), 33
- The photochemical behaviour of matrix-isolated W(CO)₆ in the presence of alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 163
- Carbonyls**
- Synthesis of the tertiaryphosphine derivatives of iron carbonyl phosphines: preparation of tetracarbonyl[tris(2-furyl)phosphine]iron(0), pentacarbonylbis[μ-bis(2-furyl)phosphido]-[tris(2-furyl)phosphine]diiron(0), tetracarbonyl[tris(2-benzofuryl)phosphine]iron(0) and pentacarbonylbis[μ-bis(2-benzofuryl)phosphido]-[tris(2-benzofuryl)phosphine]diiron(0) (C. Santelli-Rouvier, C. Coin, L. Toupet and M. Santelli), 91
- Carboxylate**
- Synthesis and reactivity of acetylacetonato-C^γ complexes of M^{II} (M = Pd or Pt): X-ray crystal structure of [Pd(C₆F₅)(OOCPh)(bipy)] · CHCl₃ (J. Forniés, F. Martínez, R. Navarro and E.P. Urriolabeitia), 185
- Catalysis**
- Chiral tetraphenylcyclopentadienyl complexes of rhodium: synthesis and crystal structure of [Rh(η⁵-C₅Ph₄CH(Me)Ph)(cod)] (J.A. Ramsden, D. Milner, H. Adams, N.A. Bailey and C. White), 215
- Chirality**
- Chiral tetraphenylcyclopentadienyl complexes of rhodium: synthesis and crystal structure of [Rh(η⁵-C₅Ph₄CH(Me)Ph)(cod)] (J.A. Ramsden, D. Milner, H. Adams, N.A. Bailey and C. White), 215
- Cobalt**
- Chemie polyfunktioneller Moleküle. CXVII. Synthese und Molekülstruktur von [Co⁺¹(CO)₂(Ph₂P=N=P(O)Ph₂^o-P)(Ph₂P-NH-PPh₂)] · 2 CH₃OH (D. Pohl, J. Ellermann, F.A. Knoch and M. Moll), C6
- Low-valent complexes of cobalt and phosphorus donor ligands as mediators in Reformatsky-type reactions (F. Orsini, M. Pulici and L.M. Vallarino), C1
- Reactions of molybdenum-cobalt complex with phenylacetylene: X-ray crystal structure of [MoCo(CO)₄(CPhCHCHCPh)(η⁵-C₅H₅)] (F.-E. Hong, I.-R. Lue, S.-C. Lo and C.-C. Lin), 97
- Cone angles**
- Solid angles. III. The role of conformers in solid angle calculations (D. White, B.C. Taverner, N.J. Coville and P.W. Wade), 41
- Coupling constants**
- On the synthesis and ³¹P{¹H} NMR spectrum of (CO)₅W(μ-PPh₂CH₂PPh₂)W(CO)₅ (J.W. Benson, E.A. Keiter and R.L. Keiter), 77
- Crystal structure**
- Chemie polyfunktioneller Moleküle. CXVII. Synthese und Molekülstruktur von [Co⁺¹(CO)₂(Ph₂P=N=P(O)Ph₂^o-P)(Ph₂P-NH-PPh₂)] · 2 CH₃OH (D. Pohl, J. Ellermann, F.A. Knoch and M. Moll), C6
- Chiral tetraphenylcyclopentadienyl complexes of rhodium: synthesis and crystal structure of [Rh(η⁵-C₅Ph₄CH(Me)Ph)(cod)] (J.A. Ramsden, D. Milner, H. Adams, N.A. Bailey and C. White), 215
- Cluster chemistry. Part 97. Some unusual complexes obtained from reactions of Ru₃(μ-dppm)(CO)₁₀ and C₇ or C₈ hydrocarbons (M.I. Bruce, J.R. Hinchliffe, B.W. Skelton and A.H. White), 141
- Metallorganische Verbindungen der Lanthanoide. XCVIII. Synthese und Strukturklärung von Dichloro(η⁵-2,5-di-tert.-butylpyrrolyl)bis(tetrahydrofuran)ytterbium(III) (H. Schumann, E.C.E. Rosenthal, J. Winterfeld and G. Kociok-Köhn), C12
- Photochemical reaction of (η⁵-cyclopentadienyl) dicarbonyliron iodide with 1-substituted uracils in the presence of diisopropylamine: crystal structure of the (η⁵-C₅H₅)Fe(CO)₂ complex of deprotonated 5-fluoro-1-(tetrahydro-2-furyl)uracil (Ftorafur) (J. Zakrzewski, A. Tosik and M. Bukowska-Strzyżewska), 83
- Synthesis and crystal structure of [(η⁵-η⁵-C₅H₄PPh₂)Ti]_n and the divalent ytterbium derivative [(η⁵-C₅H₄PPh₂)₂Yb(DME)] (G. Lin and W.-T. Wong), 203
- Cycloheptatriene**
- Cluster chemistry. Part 97. Some unusual complexes obtained from reactions of Ru₃(μ-dppm)(CO)₁₀ and C₇ or C₈ hydrocarbons (M.I. Bruce, J.R. Hinchliffe, B.W. Skelton and A.H. White), 141
- Cyclopentadienyl**
- Photochemical reaction of (η⁵-cyclopentadienyl) dicarbonyliron iodide with 1-substituted uracils in the presence of diisopropylamine: crystal structure of the (η⁵-C₅H₅)Fe(CO)₂ complex of deprotonated 5-fluoro-1-(tetrahydro-2-furyl)uracil (Ftorafur) (J. Zakrzewski, A. Tosik and M. Bukowska-Strzyżewska), 83
- Synthesis and characterization of titanium complexes containing the 1-(3-butenyl)-2,3,4,5-tetramethylcyclopentadienyl ligand (J. Okuda, K.E. Du Plooy and P.J. Toscano), 195
- Synthesis and crystal structure of [(η⁵-η⁵-C₅H₄PPh₂)Ti]_n and the divalent ytterbium derivative [(η⁵-C₅H₄PPh₂)₂Yb(DME)] (G. Lin and W.-T. Wong), 203
- Cyclopentadienyl derivatives**
- Synthesis and reactivity investigation of iridium maleonitriledithiolate complexes. Redox studies and extended Hückel molecular orbital calculations on Cp*IrL(mnt) (where L = PMe₃, PPh₃, CN-t-Bu) (K. Yang, M.-J. Don, D.K. Sharma, S.G. Bott and M.G. Richmond), 61
- Decomposition**
- Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXLVIII. Alkylirrhodium(VII)-oxide—Synthese, Eigenschaften und Abbaureaktionen (W.A. Herrmann, F.E. Kühn und C.C. Romão), 209
- Diazomethane**
- Übergangsmetall-substituierte Diphosphene. XXXVI. Cyclopropanierung der P=P-Bindung in den [3 + 2]-Cycloaddukten aus Metallodiphosphenen und Hexafluoracetone, (η⁵-C₅R₅)(CO)M-P(=PMes*)OC(CF₃)₂C(O)(M = Fe; R₅ = Me₄Et) (M = Ru; R₅ = Me₅) (L. Weber und H. Misiak), 127
- Dibutyltin(V)**
- Studies on the diorganotin(IV) complexes of some dihydroxamic acids (M.K. Das and S. De), 177
- Dihydroxamic acids**
- Studies on the diorganotin(IV) complexes of some dihydroxamic acids (M.K. Das and S. De), 177
- Dinuclear**
- Synthesis of the tertiaryphosphine derivatives of iron carbonyl phosphines: preparation of tetracarbonyl[tris(2-furyl)phosphine]iron(0), pentacarbonylbis[μ-bis(2-furyl)phosphido]-[tris(2-furyl)phosphine]diiron(0), tetracarbonyl[tris(2-benzofuryl)phosphine]iron(0) and pentacarbonylbis[μ-bis(2-benzofuryl)phosphido]-[tris(2-benzofuryl)phosphine]diiron(0) (C. Santelli-Rouvier, C. Coin, L. Toupet and M. Santelli), 91
- Diphenyltin(IV)**
- Studies on the diorganotin(IV) complexes of some dihydroxamic acids (M.K. Das and S. De), 177
- Ditertiary phosphines**
- On the synthesis and ³¹P{¹H} NMR spectrum of (CO)₅W(μ-PPh₂CH₂PPh₂)W(CO)₅ (J.W. Benson, E.A. Keiter and R.L. Keiter), 77
- Electrochemistry**
- Synthesis and reactivity investigation of iridium maleonitriledithiolate complexes. Redox studies and extended Hückel molecular orbital calculations on Cp*IrL(mnt) (where L = PMe₃, PPh₃, CN-t-Bu) (K. Yang, M.-J. Don, D.K. Sharma, S.G. Bott and M.G. Richmond), 61

Group 4

Synthesis and characterization of titanium complexes containing the 1-(3-butenyl)-2,3,4,5-tetramethylcyclopentadienyl ligand (J. Okuda, K.E. Du Plooy and P.J. Toscano), 195

Hexafluoroacetone adducts

Übergangsmetall-substituierte Diphosphene. XXXVI. Cyclopropanierung der P=P-Bindung in den [3 + 2]-Cycloaddukten aus Metallo(diphosphenen) und Hexafluoroacetone, $(\eta^5\text{-C}_5\text{R}_5\text{XCO})\text{M}^-\text{P}(=\text{PMes}^*)\text{OC}(\text{CF}_3)_2\text{C}(\text{O})\text{X}$ (M = Fe; R₅ = Me₄Et) (M = Ru; R₅ = Me₅) (L. Weber und H. Misiak), 127

Hückel molecular orbital calculations

Synthesis and reactivity investigation of iridium maleonitriledithiolate complexes. Redox studies and extended Hückel molecular orbital calculations on Cp*IrL(mnt) (where L = PMe₃, PPh₃, CN-t-Bu) (K. Yang, M.-J. Don, D.K. Sharma, S.G. Bott and M.G. Richmond), 61

Hydrogenation

Chiral tetraphenylcyclopentadienyl complexes of rhodium: synthesis and crystal structure of [Rh(η⁵-C₅Ph₄CH(Me)Ph)(cod)] (J.A. Ramsden, D. Milner, H. Adams, N.A. Bailey and C. White), 215

Hydroxyethylphosphines

Synthesis of monohydroxy -methyl- and -ethyl-phosphines PPh₂CHROH (G. Muller and D. Sainz), 103

Hydroxymethylphosphines

Synthesis of monohydroxy -methyl- and -ethyl-phosphines PPh₂CHROH (G. Muller and D. Sainz), 103

Infrared spectroscopy

Studies on the diorganotin(IV) complexes of some dihydroxamic acids (M.K. Das and S. De), 177

The photochemical behaviour of matrix-isolated W(CO)₆ in the presence of alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 163

Iridium

Synthesis and reactivity investigation of iridium maleonitriledithiolate complexes. Redox studies and extended Hückel molecular orbital calculations on Cp*IrL(mnt) (where L = PMe₃, PPh₃, CN-t-Bu) (K. Yang, M.-J. Don, D.K. Sharma, S.G. Bott and M.G. Richmond), 61

Iron

Photochemical reaction of (η⁵-cyclopentadienyl) dicarbonyliron iodide with 1-substituted uracils in the presence of diisopropylamine: crystal structure of the (η⁵-C₅H₅)Fe(CO)₂ complex of deprotonated 5-fluoro-1-(tetrahydro-2-furyl)uracil (Ftorafur) (J. Zakrzewski, A. Tosik and M. Bukowska-Strzyżewska), 83

Synthesis of the tertiaryphosphine derivatives of iron carbonyl phosphines: preparation of tetracarbonyl[tris(2-furyl)phosphine]iron(0), pentacarbonylbis[μ-bis(2-furyl)phosphido]-[tris(2-furyl)phosphine]diiron(0), tetracarbonyl[tris(2-benzofuryl)phosphine]iron(0) and pentacarbonylbis[μ-bis(2-benzofuryl)phosphido]-[tris(2-benzofuryl)phosphine]diiron(0) (C. Santelli-Rouvier, C. Coin, L. Toupet and M. Santelli), 91

Lewis base adducts

Lewis-Basen Addukte monomerer Bis(η²-alkin)Ag^IX-Verbindungen (X = BF₄ oder OSO₂CF₃) (H. Lang, K. Köhler und B. Schiemenz), 135

Magnesium

Reaction of magnesium with benzyl halides in an optically active solvent (A.M. Egorov and A.V. Anisimov), 131

Maleonitriledithiolate complexes

Synthesis and reactivity investigation of iridium maleonitriledithiolate complexes. Redox studies and extended Hückel molecular orbital

calculations on Cp*IrL(mnt) (where L = PMe₃, PPh₃, CN-t-Bu) (K. Yang, M.-J. Don, D.K. Sharma, S.G. Bott and M.G. Richmond), 61

Matrix isolation

Matrix-isolation studies of alkyne complexes in the photochemical reactions of Mo(CO)₆ with alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 149

Metal clusters

Substitution of SO₂ for CO in triruthenium carbonyl anions (G.B. Karet, C.L. Stern, J.A. Cody, S.J. Lange, M.A. Pell, C. Sleboznick and D.F. Shriver), 33

Metallocenes

Synthesis and characterization of titanium complexes containing the 1-(3-butenyl)-2,3,4,5-tetramethylcyclopentadienyl ligand (J. Okuda, K.E. Du Plooy and P.J. Toscano), 195

Metallo(diphosphene)

Übergangsmetall-substituierte Diphosphene. XXXVI. Cyclopropanierung der P=P-Bindung in den [3 + 2]-Cycloaddukten aus Metallo(diphosphenen) und Hexafluoroacetone, $(\eta^5\text{-C}_5\text{R}_5\text{XCO})\text{M}^-\text{P}(=\text{PMes}^*)\text{OC}(\text{CF}_3)_2\text{C}(\text{O})\text{X}$ (M = Fe; R₅ = Me₄Et) (M = Ru; R₅ = Me₅) (L. Weber und H. Misiak), 127

Molecular mechanics calculation

Solid angles. III. The role of conformers in solid angle calculations (D. White, B.C. Taverner, N.J. Coville and P.W. Wade), 41

Molybdenum

Matrix-isolation studies of alkyne complexes in the photochemical reactions of Mo(CO)₆ with alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 149

Reactions of molybdenum-cobalt complex with phenylacetylene: X-ray crystal structure of [MoCo(CO)₄(CPhCHCHCPh)(η⁵-C₅H₅)] (F.-E. Hong, I.-R. Lue, S.-C. Lo and C.-C. Lin), 97

Nitrosyl

Koordinationschemie funktioneller Phosphine. II. Carbonyl(nitrosyl)-wolfram-Komplexe mit 2-Diphenylphosphinoanisol sowie 2-Diphenylphosphinoanilid, -benzoat und -phenolat als Liganden (J. Pietsch, L. Dahlenburg, A. Wolski, H. Berke und I.L. Eremenko), 113

P-31 NMR

On the synthesis and ³¹P{¹H} NMR spectrum of (CO)₅W(μ-PPh₂CH₂PPh₂)W(CO)₅ (J.W. Benson, E.A. Keiter and R.L. Keiter), 77

NMR spectroscopy

Darstellung und Molekülstruktur von Bis(3-dimethylaminopropylalan) (U. Dümichen, K.-H. Thiele, T. Gelbrich und J. Sieler), 71

Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXLVIII. Alkynylrhenium(VII)-oxide—Synthese, Eigenschaften und Abbaureaktionen (W.A. Herrmann, F.E. Kühn und C.C. Romão), 209

Studies on the diorganotin(IV) complexes of some dihydroxamic acids (M.K. Das and S. De), 177

Optical activity

Reaction of magnesium with benzyl halides in an optically active solvent (A.M. Egorov and A.V. Anisimov), 131

Palladium

Synthesis and reactivity of acetylacetonato-C^γ complexes of M^{II} (M = Pd or Pt): X-ray crystal structure of [Pd(C₆F₅)(OOCPh)(bipy)]·CHCl₃ (J. Fornies, F. Martínez, R. Navarro and E.P. Urriolabeitia), 185

Pentachlorophenyl

Synthesis and reactivity of acetylacetonato-C^γ complexes of M^{II} (M = Pd or Pt): X-ray crystal structure of [Pd(C₆F₅)(OOCPh)-

- (bipy) · CHCl₃ (J. Forniés, F. Martínez, R. Navarro and E.P. Urriolabeitia), 185
- Pentafluorophenyl**
 Synthesis and reactivity of acetylacetonato-*C*^γ complexes of M^{II} (M = Pd or Pt): X-ray crystal structure of [Pd(C₆F₅)(OOCPh)(bipy)] · CHCl₃ (J. Forniés, F. Martínez, R. Navarro and E.P. Urriolabeitia), 185
- Phosphine**
 Chemie polyfunktioneller Moleküle. CXVII. Synthese und Molekülstruktur von [Co⁺¹(CO)₂{Ph₂P=N=N=P(O)Ph₂-P}(Ph₂P-NH-PPh₂)} · 2 CH₃OH (D. Pohl, J. Ellermann, F.A. Knoch and M. Moll), C6
 Koordinationschemie funktioneller Phosphine. II. Carbonyl(nitrosyl)-wolfram-Komplexe mit 2-Diphenylphosphinoanisol sowie 2-Diphenylphosphinoanilid, -benzoat und -phenolat als Liganden (J. Pietsch, L. Dahlenburg, A. Wolski, H. Berke und I.L. Eremenko), 113
- Phosphine complexes**
 Low-valent complexes of cobalt and phosphorus donor ligands as mediators in Reformatsky-type reactions (F. Orsini, M. Pulici und L.M. Vallarino), C1
- Phosphorus**
 Synthesis of monohydroxy -methyl- and -ethyl-phosphines PPh₂CHROH (G. Muller and D. Sainz), 103
- Phosphorus ligands**
 Solid angles. III. The role of conformers in solid angle calculations (D. White, B.C. Taverner, N.J. Coville and P.W. Wade), 41
- Photolysis**
 Matrix-isolation studies of alkyne complexes in the photochemical reactions of Mo(CO)₆ with alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 149
- Platinum**
 Synthesis and reactivity of acetylacetonato-*C*^γ complexes of M^{II} (M = Pd or Pt): X-ray crystal structure of [Pd(C₆F₅)(OOCPh)(bipy)] · CHCl₃ (J. Forniés, F. Martínez, R. Navarro and E.P. Urriolabeitia), 185
- Polymerization of alkynes**
 The photochemical behaviour of matrix-isolated W(CO)₆ in the presence of alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 163
- Preparative methods**
 Synthesis of monohydroxy -methyl- and -ethyl-phosphines PPh₂CHROH (G. Muller and D. Sainz), 103
- Raman spectroscopy**
 Studies on the diorganotin(IV) complexes of some dihydroxamic acids (M.K. Das and S. De), 177
- Reformatsky reaction**
 Low-valent complexes of cobalt and phosphorus donor ligands as mediators in Reformatsky-type reactions (F. Orsini, M. Pulici und L.M. Vallarino), C1
- Rhenium**
 Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXLVIII. Alkynylrhenium(VII)-oxide—Synthese, Eigenschaften und Abbaureaktionen (W.A. Herrmann, F.E. Kühn und C.C. Romão), 209
- Rhodium**
 Chiral tetraphenylcyclopentadienyl complexes of rhodium: synthesis and crystal structure of [Rh(η⁵-C₅Ph₄CH(Me)Ph)(cod)] (J.A. Ramsden, D. Milner, H. Adams, N.A. Bailey and C. White), 215
- Ruthenium**
 Cluster chemistry. Part 97. Some unusual complexes obtained from reactions of Ru₃(μ-dppm)(CO)₁₀ and C₇ or C₈ hydrocarbons (M.I. Bruce, J.R. Hinchliffe, B.W. Skelton and A.H. White), 141
- Substitution of SO₂ for CO in triruthenium carbonyl anions (G.B. Karet, C.L. Stern, J.A. Cody, S.J. Lange, M.A. Pell, C. Slebochnick and D.F. Shriver), 33
- Silver**
 Lewis-Basen Addukte monomerer Bis(η²-alkin)Ag^IX-Verbindungen (X = BF₄ oder OSO₂CF₃) (H. Lang, K. Köhler und B. Schiemenz), 135
- Solid angles**
 Solid angles. III. The role of conformers in solid angle calculations (D. White, B.C. Taverner, N.J. Coville and P.W. Wade), 41
- Steric size**
 Solid angles. III. The role of conformers in solid angle calculations (D. White, B.C. Taverner, N.J. Coville and P.W. Wade), 41
- Sulfur dioxide**
 Substitution of SO₂ for CO in triruthenium carbonyl anions (G.B. Karet, C.L. Stern, J.A. Cody, S.J. Lange, M.A. Pell, C. Slebochnick and D.F. Shriver), 33
- Synthesis**
 Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen. CXLVIII. Alkynylrhenium(VII)-oxide—Synthese, Eigenschaften und Abbaureaktionen (W.A. Herrmann, F.E. Kühn und C.C. Romão), 209
- Tetraphenylcyclopentadienyl**
 Chiral tetraphenylcyclopentadienyl complexes of rhodium: synthesis and crystal structure of [Rh(η⁵-C₅Ph₄CH(Me)Ph)(cod)] (J.A. Ramsden, D. Milner, H. Adams, N.A. Bailey and C. White), 215
- Thallium**
 Synthesis and crystal structure of [(η⁵-η⁵-C₅H₄PPh₂)Tl]₂ and the divalent ytterbium derivative [(η⁵-C₅H₄PPh₂)₂Yb(DME)] (G. Lin and W.-T. Wong), 203
- Titanium**
 Synthesis and characterization of titanium complexes containing the 1-(3-butenyl)-2,3,4,5-tetramethylcyclopentadienyl ligand (J. Okuda, K.E. Du Plooy and P.J. Toscano), 195
- Triflate**
 Synthesis and characterization of titanium complexes containing the 1-(3-butenyl)-2,3,4,5-tetramethylcyclopentadienyl ligand (J. Okuda, K.E. Du Plooy and P.J. Toscano), 195
- Tungsten**
 Koordinationschemie funktioneller Phosphine. II. Carbonyl(nitrosyl)-wolfram-Komplexe mit 2-Diphenylphosphinoanisol sowie 2-Diphenylphosphinoanilid, -benzoat und -phenolat als Liganden (J. Pietsch, L. Dahlenburg, A. Wolski, H. Berke und I.L. Eremenko), 113
 On the synthesis and ³¹P{¹H} NMR spectrum of (CO)₅W(μ-PPh₂CH₂PPh₂)W(CO)₅ (J.W. Benson, E.A. Keiter and R.L. Keiter), 77
 The photochemical behaviour of matrix-isolated W(CO)₆ in the presence of alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 163
- Uracils**
 Photochemical reaction of (η⁵-cyclopentadienyl) dicarbonyliron iodide with 1-substituted uracils in the presence of diisopropylamine: crystal structure of the (η⁵-C₅H₅)Fe(CO)₂ complex of deprotonated 5-fluoro-1-(tetrahydro-2-furyl)uracil (Ftorafur) (J. Zakrzewski, A. Tosik and M. Bukowska-Strzyżewska), 83
- Vinylidene**
 The photochemical behaviour of matrix-isolated W(CO)₆ in the presence of alkynes (T. Szymańska-Buzar, A.J. Downs, T.M. Greene and A.S. Marshall), 163

X-ray diffraction

Darstellung und Molekülstruktur von Bis(3-dimethylaminopropylalan) (U. Dümichen, K.-H. Thiele, T. Gelbrich und J. Sieler), 71

Koordinationschemie funktioneller Phosphine. II. Carbonyl(nitrosyl)-wolfram-Komplexe mit 2-Diphenylphosphinoanisol sowie 2-Diphenylphosphinoanilid, -benzoat und -phenolat als Liganden (J. Pietsch, L. Dahlenburg, A. Wolski, H. Berke und I.L. Eremenko), 113

X-ray structure

Lewis-Basen Addukte monomerer Bis(η^2 -alkin)Ag¹X-Verbindungen (X = BF₄ oder OSO₂CF₃) (H. Lang, K. Köhler und B. Schiemenz), 135

Substitution of SO₂ for CO in triruthenium carbonyl anions (G.B. Karet, C.L. Stern, J.A. Cody, S.J. Lange, M.A. Pell, C. Slebodnick and D.F. Shriver), 33

Synthesis of the tertiaryphosphine derivatives of iron carbonyl phosphines: preparation of tetracarbonyl[tris(2-furyl)phosphine]iron(0), pentacarbonylbis[μ -bis(2-furyl)phosphido]-[tris(2-furyl)phosphine]diiron(0), tetracarbonyl[tris(2-benzofuryl)phosphine]iron(0) and pentacarbonylbis[μ -bis(2-benzofuryl)phosphido]-[tris(2-benzofuryl)phosphine]diiron(0) (C. Santelli-Rouvier, C. Coin, L. Toupet and M. Santelli), 91

Ytterbium

Metallorganische Verbindungen der Lanthanoide. XCVIII. Synthese und Strukturaufklärung von Dichloro(η^5 -2,5-di-tert.-butylpyrrolyl)bis(tetrahydrofuran)ytterbium(III) (H. Schumann, E.C.E. Rosenthal, J. Winterfeld und G. Kociok-Köhn), C12

Synthesis and crystal structure of [(η^5 : η^5 -C₅H₄PPh₂)Tl]₂ and the divalent ytterbium derivative [(η^5 -C₅H₄PPh₂)₂Yb(DME)] (G. Lin and W.-T. Wong), 203