



ELSEVIER

www.elsevier.nl/locate/jorgchem

Journal of Organometallic Chemistry 597 (2000) 209–212

Journal
of Organo
metallic
Chemistry

Subject Index of Volume 597

Acetone azine

Reactions of methyl-substituted titanocene–bis(trimethylsilyl)acetylene complexes with acetone azine: crystal structures of $(\eta^5:\eta^1\text{C}_5\text{HMe}_3\text{CH}_2\text{CMe}_2\text{NH})_2\text{Ti}$ and $(\text{C}_5\text{Me}_5)_2\text{Ti}(\text{N}=\text{CMe}_2)$, 146

Acetylene π -complex

Synthesis and molecular structures of $\text{Co}_2(\text{CO})_{6-n}(\text{PPh}_3)_n[\mu\text{C}_2(\text{CH}_2\text{R})_2]$ ($\text{R} = \text{CH}_2\text{OH}$ or CH_2SCMe_3) derivatives containing chelated $[\text{CpFe}(\text{CO})_2]_2\text{Sn}$ or $\text{W}(\text{CO})_4$ fragments, 196

Acyl complexes

Synthesis, characterization and decomposition behavior of novel acyl(hydrido)platinum(IV) complexes, 10

Alkylalanes

Syntheses of some bulky alkylalanes and alkyltrihydroaluminates: crystal structures of $[\text{Li}(\text{THF})_2\text{AlH}_3\text{C}(\text{SiMe}_3)_2(\text{SiMe}_2\text{NMe}_2)]_2$, $\text{Li}(\text{THF})_2\text{Al}_2\text{H}_5\{\text{C}(\text{SiMe}_3)_3\}_2$, $(\text{Me}_3\text{Si})_2\text{CAIH}_2\cdot\text{THF}$ and the pyrazolato derivative $[\text{LiAlH}(\text{C}_3\text{H}_3\text{N}_2)_2\text{C}(\text{SiMe}_3)_3]_2$ ($\text{THF} = \text{tetrahydrofuran}$), 3

Alkynes

1-Azadienes as bridging ligands in nickel(0) complexes of the type $[(1\text{-aza-1.3-diene})\text{Ni}(\text{L})_2]$ ($\text{L} = 1\text{-aza-1.3-diene, alkyne, triphenylphosphine}$), 116

Allyl

Komplexe LVI: Zur Kation-Anion-Wechselwirkung in den $\text{C}_{12}\text{-Allylnickel(II)-Komplexen } [\text{Ni}(\eta^3,\eta^2,\eta^2\text{-C}_{12}\text{H}_{19})]\text{X}$ als Steuergröße für die Katalyse der stereospezifischen Butadienpolymerisation. Synthese des Tetrakis(pentafluorophenyl)borats ($\text{X} = \text{B}(\text{C}_6\text{F}_5)_4$) und Charakterisierung des Tetrafluoroborats sowie des Trifluor-methansulfonats ($\text{X} = \text{BF}_4$, CF_3SO_3) durch Röntgenkristallstrukturanalyse und Leitfähigkeitsmessungen, 92

Aluminium

Syntheses of some bulky alkylalanes and alkyltrihydroaluminates: crystal structures of $[\text{Li}(\text{THF})_2\text{AlH}_3\text{C}(\text{SiMe}_3)_2(\text{SiMe}_2\text{NMe}_2)]_2$, $\text{Li}(\text{THF})_2\text{Al}_2\text{H}_5\{\text{C}(\text{SiMe}_3)_3\}_2$, $(\text{Me}_3\text{Si})_2\text{CAIH}_2\cdot\text{THF}$ and the pyrazolato derivative $[\text{LiAlH}(\text{C}_3\text{H}_3\text{N}_2)_2\text{C}(\text{SiMe}_3)_3]_2$ ($\text{THF} = \text{tetrahydrofuran}$), 3

Aluminum

Sterically crowded aryloxides of aluminum: intramolecular coordination of bidentate ligands, 29

Synthesis and structural characterization of polymeric dimethylaluminum phenylthiolate, 105

Aryoxide

Sterically crowded aryloxides of aluminum: intramolecular coordination of bidentate ligands, 29

1-Aza-1.3-dienes

1-Azadienes as bridging ligands in nickel(0) complexes of the type $[(1\text{-aza-1.3-diene})\text{Ni}(\text{L})_2]$ ($\text{L} = 1\text{-aza-1.3-diene, alkyne, triphenylphosphine}$), 116

Barbiturate

$(\eta^5\text{-Cyclopentadienyl})\text{Fe}(\text{CO})_2$ complexes of barbiturate anions, 17 bis-Cyclopentadienyl zirconium and titanium complexes Activation of CH_4 and H_2 by zirconium(IV) and titanium(IV) cationic complexes. Theoretical DFT study, 182

Branched silanes

Reactions of the Si–H bonds in tetrakis(dimethylsilyl)methane and silane, 111

Butadiene

Komplexe LVI: Zur Kation-Anion-Wechselwirkung in den $\text{C}_{12}\text{-Allylnickel(II)-Komplexen } [\text{Ni}(\eta^3,\eta^2,\eta^2\text{-C}_{12}\text{H}_{19})]\text{X}$ als Steuergröße für die Katalyse der stereospezifischen Butadienpolymerisation. Synthese des Tetrakis(pentafluorophenyl)borats ($\text{X} = \text{B}(\text{C}_6\text{F}_5)_4$) und Charakterisierung des Tetrafluoroborats sowie des Trifluor-methansulfonats ($\text{X} = \text{BF}_4$, CF_3SO_3) durch Röntgenkristallstrukturanalyse und Leitfähigkeitsmessungen, 92

Carbon monoxide

Molecular structure and properties of the rhodium(II) complexes with chemilabile ether–phosphine ligands $\text{P}(2\text{-MeOC}_6\text{H}_4)_3$ and $\text{P}(2,6\text{-}(\text{MeO})_2\text{C}_6\text{H}_3)_3$, 20

Carbonylmetallates

Halophilic reactions of pentafluorohalobenzenes with transition-metal carbonyl anions, 77

Catalysis

Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO_2 , 139

Catalyzed reaction of triethylborane with pyrazole, 190

Komplexe LVI: Zur Kation-Anion-Wechselwirkung in den $\text{C}_{12}\text{-Allylnickel(II)-Komplexen } [\text{Ni}(\eta^3,\eta^2,\eta^2\text{-C}_{12}\text{H}_{19})]\text{X}$ als Steuergröße für die Katalyse der stereospezifischen Butadienpolymerisation. Synthese des Tetrakis(pentafluorophenyl)borats ($\text{X} = \text{B}(\text{C}_6\text{F}_5)_4$) und Charakterisierung des Tetrafluoroborats sowie des Trifluor-methansulfonats ($\text{X} = \text{BF}_4$, CF_3SO_3) durch Röntgenkristallstrukturanalyse und Leitfähigkeitsmessungen, 92

C–H activation

Oxo and imido/imido exchange and C–H activation reactions based on pentamethylcyclopentadienyl imido tantalum complexes, 61

Chelate

Syntheses and molecular structures of $\text{Co}_2(\text{CO})_{6-n}(\text{PPh}_3)_n[\mu\text{C}_2(\text{CH}_2\text{R})_2]$ ($\text{R} = \text{CH}_2\text{OH}$ or CH_2SCMe_3) derivatives containing chelated $[\text{CpFe}(\text{CO})_2]_2\text{Sn}$ or $\text{W}(\text{CO})_4$ fragments, 196

Chiral HPLC

Intramolecular versus intermolecular mechanism of metallotropic rearrangements of chiral chromium tricarbonyl complexes of substituted naphthalenes and other polycyclic aromatic compounds investigated by chiral HPLC of its racemic mixtures, 42

Chiral ligands

Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO_2 , 139

- Chromium tri(carbonyl) aromatic
Intramolecular versus intermolecular mechanism of metallotropic rearrangements of chiral chromium tricarbonyl complexes of substituted naphthalenes and other polyaromatic compounds investigated by chiral HPLC of its racemic mixtures, 42
- closo*-Rhodacarborane
Mercury derivatives of *exo-nido*-ruthenacarborane, 48
- Clusters
Analogies between Group 13 metal clusters in organometallic and intermetallic structures, 54
- Reactions of nickelocene with sodium in the presence of terminal alkenes, 133
- CO₂
Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139
- Cobalt
Synthesis and molecular structures of Co₂(CO)_{6-n}(PPh₃)_n[μ-C₂(CH₂R)₂] (R = CH₃OH or CH₂SCMe₃) derivatives containing chelated [CpFe(CO)₂]₂Sn or W(CO)₄ fragments, 196
- CPMAS
Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139
- Crystal structure
Biscyclopentadienylzirconium(IV) η³-propargyl/allenyl complexes, 125
- Reactions of methyl-substituted titanocene–bis(trimethylsilyl)acetylene complexes with acetone azine: crystal structures of (η⁵:η¹-C₅HMe₃CH₂CMe₂NH)₂Ti and (C₅Me₅)₂Ti(N=CMe₂), 146
- Reactions of nickelocene with sodium in the presence of terminal alkenes, 133
- Syntheses and reactions of some new C-pentafluorophenyl and tetrafluorophenylene carborane systems, 157
- Cyanide bridging
Cyanide-bridged tetranuclear complexes with T-shaped Pt(CN–M)₃ arrangements, 38
- Cyclopentadienyl
(η⁵-Cyclopentadienyl)Fe(CO)₂ complexes of barbiturate anions, 17
- Reactions of nickelocene with sodium in the presence of terminal alkenes, 133
- Cyclopentadienyl imido tantalum complexes
Oxo and imido/imido exchange and C–H activation reactions based on pentamethylcyclopentadienyl imido tantalum complexes, 61
- Dehydrogenative (coupling) silylation
Catalysis of hydrosilylation: Part XXXIV. High catalytic efficiency of the nickel equivalent of Karstedt catalyst [{Ni(η-CH₂=CHSiMe₂)₂O}₂{μ-(η-CH₂=CHSiMe₂)₂O}], 175
- Deltahedra
Analogies between Group 13 metal clusters in organometallic and intermetallic structures, 54
- Density functional calculations
Activation of CH₄ and H₂ by zirconium(IV) and titanium(IV) cationic complexes. Theoretical DFT study, 182
- Molecular structure and conformational composition of de-
- camethyl-*n*-tetrasilane, Si₄Me₁₀, by gas electron diffraction and density functional theory calculations, 87
- Dirhodium complexes
Molecular structure and properties of the rhodium(II) complexes with chemilabile ether–phosphine ligands P(2-MeOC₆H₄)₃ and P(2,6-(MeO)₂C₆H₃)₃, 20
- Effect of methyl substituents
Reactions of methyl-substituted titanocene–bis(trimethylsilyl)acetylene complexes with acetone azine: crystal structures of (η⁵:η¹-C₅HMe₃CH₂CMe₂NH)₂Ti and (C₅Me₅)₂Ti(N=CMe₂), 146
- Electrochemistry
Cyanide-bridged tetranuclear complexes with T-shaped Pt(CN–M)₃ arrangements, 38
- Electron diffraction
Molecular structure and conformational composition of de-
- camethyl-*n*-tetrasilane, Si₄Me₁₀, by gas electron diffraction and density functional theory calculations, 87
- Ether-phosphines
Molecular structure and properties of the rhodium(II) complexes with chemilabile ether–phosphine ligands P(2-MeOC₆H₄)₃ and P(2,6-(MeO)₂C₆H₃)₃, 20
- exo-nido*-Ruthenacarborane
Mercury derivatives of *exo-nido*-ruthenacarborane, 48
- Ferrocenes
Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139
- Fluxional behavior
Biscyclopentadienylzirconium(IV) η³-propargyl/allenyl complexes, 125
- Group 13
Analogies between Group 13 metal clusters in organometallic and intermetallic structures, 54
- Halogen–metal exchange
Halophilic reactions of pentafluorohalobenzenes with transition-metal carbonyl anions, 77
- Hammett correlations
Benzhydroximic acids — NMR study of trimethylsilyl derivatives, 200
- Hexafluorobenzene
Syntheses and reactions of some new C-pentafluorophenyl and tetrafluorophenylene carborane systems, 157
- H–H bond activation
Activation of CH₄ and H₂ by zirconium(IV) and titanium(IV) cationic complexes. Theoretical DFT study, 182
- Hybrid liquid crystals
Reactions of the Si–H bonds in tetrakis(dimethylsilyl)methane and silane, 111
- Hydrido complexes
Synthesis, characterization and decomposition behavior of novel acyl(hydrido)platinum(IV) complexes, 10
- Hydrogenation
New insight into role of *ortho*-metallation in rhodium triphenylphosphite complexes. Hydrogen mobility in hydrogenation and isomerization of unsaturated substrates, 69
- Hydrosilylation
Catalysis of hydrosilylation: Part XXXIV. High catalytic efficiency of the nickel equivalent of Karstedt catalyst [{Ni(η-CH₂=CHSiMe₂)₂O}₂{μ-(η-CH₂=CHSiMe₂)₂O}], 175

- Reactions of the Si–H bonds in tetrakis(dimethylsilyl)methane and silane, 111
- Intermetallic**
- Analogies between Group 13 metal clusters in organometallic and intermetallic structures, 54
- Iron**
- (η^5 -Cyclopentadienyl)Fe(CO)₂ complexes of barbiturate anions, 17
 - Synthesis and molecular structures of Co₂(CO)_{6-n}(PPh₃)_n[μ -C₂(CH₂R)₂] (R = CH₂OH or CH₂SCMe₃) derivatives containing chelated [CpFe(CO)₂]Sn or W(CO)₄ fragments, 196
- Isomerization**
- New insight into role of *ortho*-metallation in rhodium triphenylphosphite complexes. Hydrogen mobility in hydrogenation and isomerization of unsaturated substrates, 69
- Lactone**
- Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139
- Lithium**
- Syntheses of some bulky alkylalanes and alkyltrihydroaluminates: crystal structures of [Li(THF)₂AlH₃C(SiMe₃)₂(SiMe₂-NMe₂)₂], Li(THF)₂Al₂H₅{C(SiMe₃)₃}₂, (Me₃Si)₃CAIH₂·THF and the pyrazolato derivative [LiAlH(C₃H₃N)₂]₂(C(SiMe₃)₃]₂ (THF = tetrahydrofuran), 3
- Mercuration**
- Mercury derivatives of *exo-nido*-ruthenacarborane, 48
- meta Carborane**
- Syntheses and reactions of some new C-pentafluorophenyl and tetrafluorophenylene carborane systems, 157
- Metal acyl anions**
- Halophilic reactions of pentafluorohalobenzenes with transition-metal carbonyl anions, 77
- Metallacarboranes**
- Mercury derivatives of *exo-nido*-ruthenacarborane, 48
- Metallotropic**
- Intramolecular versus intermolecular mechanism of metallotropic rearrangements of chiral chromium tricarbonyl complexes of substituted naphthalenes and other polyaromatic compounds investigated by chiral HPLC of its racemic mixtures, 42
- Metal–metal charge transfer**
- Cyanide-bridged tetranuclear complexes with T-shaped Pt(CN–M)₃ arrangements, 38
- Methane C–H bond activation**
- Activation of CH₄ and H₂ by zirconium(IV) and titanium(IV) cationic complexes. Theoretical DFT study, 182
- Mixed-metal cluster**
- Synthesis and molecular structures of Co₂(CO)_{6-n}(PPh₃)_n[μ -C₂(CH₂R)₂] (R = CH₂OH or CH₂SCMe₃) derivatives containing chelated [CpFe(CO)₂]Sn or W(CO)₄ fragments, 196
- Molecular dynamics**
- X-ray structure and fluxional behaviour of *N,N'*-di-*p*-fluorophenyltriazenido complex of nickel(II), 164
- Molecular orbital**
- Molecular structure and conformational composition of deamethyl-*n*-tetrasilane, Si₄Me₁₀, by gas electron diffraction and density functional theory calculations, 87
- M multinuclear NMR**
- Syntheses and reactions of some new C-pentafluorophenyl and tetrafluorophenylene carborane systems, 157
- Multiple bonding**
- Analogies between Group 13 metal clusters in organometallic and intermetallic structures, 54
- Nickel**
- Komplexkatalyse LVI: Zur Kation-Anion-Wechselwirkung in den C₁₂-Allylnickel(II)-Komplexen [Ni(η^3 , η^2 , η^2 -C₁₂H₁₉)X] als Steuergröße für die Katalyse der stereospezifischen Butadienpolymerisation. Synthese des Tetrakis(pentafluorophenyl)borats (X = B(C₆F₅)₄) und Charakterisierung des Tetrafluoroborats sowie des Trifluor-methansulfonats (X = BF₄, CF₃SO₃) durch Röntgenkristallstrukturanalyse und Leitfähigkeitsmessungen, 92
 - Reactions of nickelocene with sodium in the presence of terminal alkenes, 133
- Nickel(0)**
- 1-Azadienes as bridging ligands in nickel(0) complexes of the type [(1-aza-1,3-diene)Ni(L)]₂ (L = 1-aza-1,3-diene, alkyne, triphenylphosphine), 116
- Nickel complexes**
- Catalysis of hydrosilylation: Part XXXIV. High catalytic efficiency of the nickel equivalent of Karstedt catalyst [{Ni(η -CH₂-CHSiMe₂)₂O}₂{ μ -(η -CH₂=CHSiMe₂)₂O}], 175
 - X-ray structure and fluxional behaviour of *N,N'*-di-*p*-fluorophenyltriazenido complex of nickel(II), 164
- NMR**
- Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139
- NMR chemical shifts**
- Benzhydroximic acids — NMR study of trimethylsilyl derivatives, 200
- Nucleophilic aromatic substitution**
- Halophilic reactions of pentafluorohalobenzenes with transition-metal carbonyl anions, 77
- ortho Carborane**
- Syntheses and reactions of some new C-pentafluorophenyl and tetrafluorophenylene carborane systems, 157
- ortho-Metallation**
- New insight into role of *ortho*-metallation in rhodium triphenylphosphite complexes. Hydrogen mobility in hydrogenation and isomerization of unsaturated substrates, 69
- Oxo/imido imido/imido exchange**
- Oxo and imido/imido exchange and C–H activation reactions based on pentamethylcyclopentadienyl imido tantalum complexes, 61
- para Carborane**
- Syntheses and reactions of some new C-pentafluorophenyl and tetrafluorophenylene carborane systems, 157
- Phenylthiolate**
- Synthesis and structural characterization of polymeric dimethylaluminum phenylthiolate, 105
- Phosphines**
- Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139
- Photolysis**
- (η^5 -Cyclopentadienyl)Fe(CO)₂ complexes of barbiturate anions, 17

Platinum

Synthesis, characterization and decomposition behavior of novel acyl(hydrido)platinum(IV) complexes, 10

Polyfluorinated aryl halides

Halophilic reactions of pentafluorohalobenzenes with transition-metal carbonyl anions, 77

Polymer

Synthesis and structural characterization of polymeric dimethylaluminum phenylthiolate, 105

Polymerization

Komplexbekatalyse LVI: Zur Kation-Anion-Wechselwirkung in den C₁₂-Allylnickel(II)-Komplexen [Ni(η³,η²,η²-C₁₂H₁₉)X]X als Steuergröße für die Katalyse der stereospezifischen Butadieneopolymerisation. Synthese des Tetrakis(pentafluorophenyl)borats (X = B(C₆F₅)₄) und Charakterisierung des Tetrafluoroborats sowie des Trifluor-methansulfonats (X = BF₄⁻, CF₃SO₃⁻) durch Röntgenkristallstrukturanalyse und Leitfähigkeitsmessungen, 92

Propargyl/allenyl complexes

Biscyclopentadienylzirconium(IV) η³-propargyl/allenyl complexes, 125

Pyrazolato

Syntheses of some bulky alkylalanes and alkyltrihydroaluminates: crystal structures of [Li(THF)₂AlH₃C(SiMe₃)₂(SiMe₂-NMe₂)₂], Li(THF)₂Al₂H₅{C(SiMe₃)₃}₂, (Me₃Si)₂CAIH₂·THF and the pyrazolato derivative [LiAlH(C₃H₃N₂)₂-C(SiMe₃)₃]₂ (THF = tetrahydrofuran), 3

Pyrazole

Catalyzed reaction of triethylborane with pyrazole, 190

Reaction mechanisms

Halophilic reactions of pentafluorohalobenzenes with transition-metal carbonyl anions, 77

Reducing agents

Reactions of the Si-H bonds in tetrakis(dimethylsilyl)methane and silane, 111

Rhodium complexes

New insight into role of *ortho*-metallation in rhodium triphenylphosphite complexes. Hydrogen mobility in hydrogenation and isomerization of unsaturated substrates, 69

Silanes

Catalysis of hydrosilylation: Part XXXIV. High catalytic efficiency of the nickel equivalent of Karstedt catalyst [{Ni(η-CH₂=CHSiMe₂)₂O}₂{μ-(η-CH₂=CHSiMe₂)₂O}], 175

Molecular structure and conformational composition of decamethyl-*n*-tetrasilane, Si₄Me₁₀, by gas electron diffraction and density functional theory calculations, 87

Silicon

Catalysis of hydrosilylation: Part XXXIV. High catalytic efficiency of the nickel equivalent of Karstedt catalyst [{Ni(η-CH₂=CHSiMe₂)₂O}₂{μ-(η-CH₂=CHSiMe₂)₂O}], 175

Sodium

Reactions of nickelocene with sodium in the presence of terminal alkenes, 133

Sterically crowded

Sterically crowded aryloxides of aluminum: intramolecular coordination of bidentate ligands, 29

Substituted benzhydrolimic acids

Benzhydrolimic acids — NMR study of trimethylsilyl derivatives, 200

Tetranuclear complexes

Cyanide-bridged tetranuclear complexes with T-shaped Pt(CN-M)₃ arrangements, 38

Tin

Synthesis and molecular structures of Co₂(CO)_{6-n}(PPh₃)_n[μ-C₂(CH₂R)₂] (R = CH₂OH or CH₂SCMe₃) derivatives containing chelated [CpFe(CO)₂]₂Sn or W(CO)₄ fragments, 196

Titanium

Reactions of methyl-substituted titanocene-bis(trimethylsilyl)-acetylene complexes with acetone azine: crystal structures of (η⁵:η¹-C₅HMe₃CH₂CMe₂NH)₂Ti and (C₅Me₅)₂Ti-(N=CMe₂), 146

Titanocene-amide complexes

Reactions of methyl-substituted titanocene-bis(trimethylsilyl)-acetylene complexes with acetone azine: crystal structures of (η⁵:η¹-C₅HMe₃CH₂CMe₂NH)₂Ti and (C₅Me₅)₂Ti-(N=CMe₂), 146

Titanocene-bis(trimethylsilyl)acetylene complexes

Reactions of methyl-substituted titanocene-bis(trimethylsilyl)-acetylene complexes with acetone azine: crystal structures of (η⁵:η¹-C₅HMe₃CH₂CMe₂NH)₂Ti and (C₅Me₅)₂Ti-(N=CMe₂), 146

Triarylphosphite

New insight into role of *ortho*-metallation in rhodium triphenylphosphite complexes. Hydrogen mobility in hydrogenation and isomerization of unsaturated substrates, 69

Triazenido complexes

X-ray structure and fluxional behaviour of *N,N'*-di-*p*-fluorophenyltriazenido complex of nickel(II), 164

Triethylborane

Catalyzed reaction of triethylborane with pyrazole, 190

Trimethylsilyl derivatives

Benzhydroximic acids — NMR study of trimethylsilyl derivatives, 200

Tungsten

Synthesis and molecular structures of Co₂(CO)_{6-n}(PPh₃)_n[μ-C₂(CH₂R)₂] (R = CH₂OH or CH₂SCMe₃) derivatives containing chelated [CpFe(CO)₂]₂Sn or W(CO)₄ fragments, 196

X-ray analysis

Synthesis and molecular structures of Co₂(CO)_{6-n}(PPh₃)_n[μ-C₂(CH₂R)₂] (R = CH₂OH or CH₂SCMe₃) derivatives containing chelated [CpFe(CO)₂]₂Sn or W(CO)₄ fragments, 196

X-ray diffraction

Synthesis, characterization and decomposition behavior of novel acyl(hydrido)platinum(IV) complexes, 10

X-ray structures

1-Azadienes as bridging ligands in nickel(0) complexes of the type [(1-aza-1,3-diene)Ni(L)]₂ (L = 1-aza-1,3-diene, alkyne, triphenylphosphine), 116

Bidentate ferrocenylphosphines and their palladium(II)dichloride complexes — X-ray structural and NMR spectroscopic investigations and first results of their characteristics in the Pd-catalysed cooligomerisation of 1,3-butadiene with CO₂, 139

X-ray structure and fluxional behaviour of *N,N'*-di-*p*-fluorophenyltriazenido complex of nickel(II), 164

Zirconium

Biscyclopentadienylzirconium(IV) η³-propargyl/allenyl complexes, 125