



ELSEVIER

Subject index of Volume 564

Acetamidinate

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzamidinate supporting ligands, 209

Acid reactivity

Reactions of protic reagents with a tungsten phosphonium ion complex, 13

Acyl

Synthesis, characterization and chemistry of some long chain alkoxy carbonyl manganese pentacarbonyl complexes $[\text{Mn}(\text{CO})_5\{\text{C}(\text{O})\text{OR}\}]$ (where R = $n\text{-C}_7\text{H}_{15}$ to $n\text{-C}_{10}\text{H}_{21}$, $n\text{-C}_{12}\text{H}_{25}$, $n\text{-C}_{14}\text{H}_{29}$, and $n\text{-C}_{16}\text{H}_{33}$), 267

Alcohol

Synthesis, characterization and chemistry of some long chain alkoxy carbonyl manganese pentacarbonyl complexes $[\text{Mn}(\text{CO})_5\{\text{C}(\text{O})\text{OR}\}]$ (where R = $n\text{-C}_7\text{H}_{15}$ to $n\text{-C}_{10}\text{H}_{21}$, $n\text{-C}_{12}\text{H}_{25}$, $n\text{-C}_{14}\text{H}_{29}$, and $n\text{-C}_{16}\text{H}_{33}$), 267

Alkoxy carbonyl

Synthesis, characterization and chemistry of some long chain alkoxy carbonyl manganese pentacarbonyl complexes $[\text{Mn}(\text{CO})_5\{\text{C}(\text{O})\text{OR}\}]$ (where R = $n\text{-C}_7\text{H}_{15}$ to $n\text{-C}_{10}\text{H}_{21}$, $n\text{-C}_{12}\text{H}_{25}$, $n\text{-C}_{14}\text{H}_{29}$, and $n\text{-C}_{16}\text{H}_{33}$), 267

Alkyl

Synthesis and reactivity of cyclopentadienyl chloro, imido and alkylidene tungsten (VI) complexes, 277

Synthesis, characterization and chemistry of some long chain alkoxy carbonyl manganese pentacarbonyl complexes $[\text{Mn}(\text{CO})_5\{\text{C}(\text{O})\text{OR}\}]$ (where R = $n\text{-C}_7\text{H}_{15}$ to $n\text{-C}_{10}\text{H}_{21}$, $n\text{-C}_{12}\text{H}_{25}$, $n\text{-C}_{14}\text{H}_{29}$, and $n\text{-C}_{16}\text{H}_{33}$), 267

Alkylidene

Synthesis and reactivity of cyclopentadienyl chloro, imido and alkylidene tungsten (VI) complexes, 277

Alkyldenep

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of $\text{MoCl}_3(\text{NAr})\text{L}_2$ (Ar = 2,6-diisopropyl phenyl; L = N-2',6'-diisopropyl phenyl-2,5-dimethyl benzamide), 85

Alkyls

Synthesis and structural characterisation of nickel(II) alkyl complexes NiR_2 , $[\text{Ni}(\text{Cp})\text{R}]_2$ and a 'head-to-tail' coupled alkyl intermediate compound $[\text{NiR}(\text{Cl})(\text{R}-\text{R})]$ [R = $\text{C}(\text{SiMe}_3)_2\text{-C}_5\text{H}_4\text{N-2}$, $\text{Cp}(\text{SiMe}_3)\text{C}_5\text{H}_4\text{N-2}$ or $\text{CH}(\text{SiMe}_3)\text{C}_5\text{H}_4\text{N-2}$; $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$], 193

Alkyltitanium

Reactivity of chlorodimethylsilyl- η^5 -cyclopentadienyltrichlorotitanium with nitrogen based donors. X-ray molecular structure of $[\text{Ti}\{\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\|\text{N}(2,6\text{-Me}_2\text{C}_6\text{H}_3)\}\text{Cl}_2]$, 93

Alkyne complexes

Synthesis, structure and reactivity of novel W–Ge chlorocarbonyl compounds. X-ray crystal structure of $[\text{WCl}(\text{GeCl}_3)(\text{CO})_3(\text{NCEt})_2]$, 143

Amido-silylcyclopentadienyl

Reactivity of chlorodimethylsilyl- η^5 -cyclopentadienyltrichlorotitanium with nitrogen based donors. X-ray molecular structure of $[\text{Ti}\{\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\|\text{N}(2,6\text{-Me}_2\text{C}_6\text{H}_3)\}\text{Cl}_2]$, 93

Amidotitanium

Reactivity of chlorodimethylsilyl- η^5 -cyclopentadienyltrichlorotitanium with nitrogen based donors. X-ray molecular structure of $[\text{Ti}\{\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\|\text{N}(2,6\text{-Me}_2\text{C}_6\text{H}_3)\}\text{Cl}_2]$, 93

Amino silyl cyclopentadienyl

Synthesis, characterization and polymerization potential of *ansa*-metallocene dichloride complexes of titanium, zirconium and hafnium containing a Si–N–Si bridging unit, 109

Anils

Transmetallation reaction of cyclomercurated anils of benzoylferrocene with metallic tin. X-ray crystal structure of $[\text{SnCl}_2\{\eta^5\text{-C}_5\text{H}_5\}\text{Fe}(\eta^5\text{-C}_5\text{H}_3\text{CPh=NC}_6\text{H}_4\text{-4-CH}_3)\}_2$, 233

Ansa-Metallocene dichloride

Synthesis, characterization and polymerization potential of *ansa*-metallocene dichloride complexes of titanium, zirconium and hafnium containing a Si–N–Si bridging unit, 109

Ansa-metallocenes

Synthesis and structural study of the ate-complex $\text{rac-}(CH_3)_2C(C_5H_3\text{-3-Si(CH}_3)_3)_2Yb(\mu_2\text{-Cl})_2Li(OEt_2)_2$ and the polymeric compound $\{\text{rac-}(CH_3)_2C(C_5H_3\text{-3-Si(CH}_3)_3)_2Yb[(\mu_2\text{-H})_2B(\mu_2\text{-H})_2]_2Li(\text{THF})_2\}_\infty$, 5

Au···Au interaction

The synthesis of $[\text{AuCo}(\text{CO})_4]_2(\mu\text{-dppfe})$ (dppfe = 1,1'-bis(diphenyl phosphino)ferrocene) and the P–Au–Co–CO(axial) skeletal pair formation between neighboring molecules in the crystal, 249

Au-transition metal bond

The synthesis of $[\text{AuCo}(\text{CO})_4]_2(\mu\text{-dppfe})$ (dppfe = 1,1'-bis(diphenyl phosphino)ferrocene) and the P–Au–Co–CO(axial) skeletal pair formation between neighboring molecules in the crystal, 249

Azavinylidene

Reactivity of the 1-azavinylidene cluster $[\text{Ru}_3(\mu\text{-H})(\mu\text{-N=CPH}_2)(\text{CO})_{10}]$ with hydrogen, tertiary silanes and tertiary stannanes, 201

Basicity studies

Relationships between basicity, redox behaviour of ferrocenylamines and their reactivity with Pt[II] compounds, 125

Benzamidinate

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzamidinate supporting ligands, 209

Bidentate

Trans-4,5-bis(2-pyridyl)norbornane: a bidentate nitrogen ligand with a potentially large bite angle, 29

Bis methylene hetero(Si, Ge, P)cyclopentanes

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Bite angle

Trans-4,5-bis(2-pyridyl)norbornane: a bidentate nitrogen ligand with a potentially large bite angle, 29

Borohydrides

Synthesis and structural study of the ate-complex $\text{rac-}(CH_3)_2C(C_5H_3\text{-3-Si(CH}_3)_3)_2Yb(\mu_2\text{-Cl})_2Li(OEt_2)_2$ and the polymeric compound $\{\text{rac-}(CH_3)_2C(C_5H_3\text{-3-Si(CH}_3)_3)_2Yb[(\mu_2\text{-H})_2B(\mu_2\text{-H})_2]_2Li(\text{THF})_2\}_\infty$, 5

Carbonyl

Synthesis and reactivity of tetramethylthioureatriosmium clusters: crystal structures of $[(\mu\text{-H})\text{Os}_3(\text{CO})_8(\mu_3\text{-S})(\mu\text{-OMeCO})\{\eta^1\text{-SC}(\text{NMe}_2)_2\}]$ and $[\text{Os}_3(\text{CO})_9(\mu\text{-OH})(\mu\text{-OMeCO})(\text{PPh}_3)]$, 133

Catalysis

Chirale Bisphosphane VIII. Die enantioselektive Hydrierung von α -Acetamidozimtsäure durch kationische Rhodium(I)-Komplexe mit optisch reinen (1S,2S)-Cyclopantan-1,2-diyl-bis(phosphonigsäurediester)- und (1S,2S)-Cyclopantan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden, 227

Unusual Pt-catalyzed stereo and regioselective intramolecular hydrosilylation of propargyl alcohols to (E,E)-1,6-dioxa-2,7-disila-3,8-ecidienes, 57

C–H activation

Chirale Bisphosphane VI. Chelatkomplexe von Platin(II) mit racemischen und optisch reinen trans-Cyclopantan-1,2-diyl-bis(phosphonigsäurediester)- und trans-Cyclopantan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden: Synthesen, Strukturen und C–H-Aktivierungsreaktionen, 155

C–H activation

C–H activation of methyl vinyl ketone in $\text{Ir}(\text{acac})\{\eta^2\text{-CH}_2=\text{CHC}(\text{O})\text{CH}_3\}(\text{PCy}_3)$, 241

Chirality

Chirale Bisphosphane VI. Chelatkomplexe von Platin(II) mit racemischen und optisch reinen trans-Cyclopantan-1,2-diyl-bis(phosphonigsäurediester)- und trans-Cyclopantan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden: Synthesen, Strukturen und C–H-Aktivierungsreaktionen, 155

Chirale Bisphosphane VIII. Die enantioselektive Hydrierung von α -Acetamidozimtsäure durch kationische Rhodium(I)-Komplexe mit optisch reinen (1S,2S)-Cyclopantan-1,2-diyl-bis(phosphonigsäurediester)- und (1S,2S)-Cyclopantan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden, 227

Cluster

Reactivity of the 1-azavinylidene cluster $[\text{Ru}_3(\mu\text{-H})(\mu\text{-N}=\text{CPh}_2)(\text{CO})_{10}]$ with hydrogen, tertiary silanes and tertiary stananes, 201

Clusters

Synthesis and reactivity of tetramethylthioureatriosmium clusters: crystal structures of $[(\mu\text{-H})\text{Os}_3(\text{CO})_8(\mu_3\text{-S})(\mu\text{-OMeCO})\{\eta^1\text{-SC}(\text{NMe}_2)_2\}]$ and $[\text{Os}_3(\text{CO})_9(\mu\text{-OH})(\mu\text{-OMeCO})(\text{PPh}_3)]$, 133

Conformation

Base-free tris(indenyl)lanthanoid(III) complexes ($\text{Ln} = \text{La, Pr, Nd, Sm}$): solid-state structure and solution NMR/NIR–vis spectroscopy, 167

Conjugated pyridines

Polynuclear metal complexes linked by conjugated pyridines, 257

Crystal structure

Base-free tris(indenyl)lanthanoid(III) complexes ($\text{Ln} = \text{La, Pr, Nd, Sm}$): solid-state structure and solution NMR/NIR–vis spectroscopy, 167

Halbsandwich-Komplexe von (S)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Organotin compounds containing a bulky $(\text{Me}_3\text{Si})_3\text{C}$ or related ligand. Crystal structures of $\{(\text{Me}_3\text{Si})_3\text{CMe}(\text{O}_2\text{NO})\text{Sn}\}_2\text{O}$, $(\text{PhMe}_2\text{Si})_3\text{CSnMeCl}_2$ and $(\text{PhMe}_2\text{Si})_3\text{CSnCl}_3$, 215

Synthesis and structural study of the ate-complex $rac\text{-}(\text{CH}_3)_2\text{C}(\text{C}_5\text{H}_3\text{-3-Si}(\text{CH}_3)_3)_2\text{Yb}(\mu_2\text{-Cl})_2\text{Li}(\text{OEt}_2)_2$ and the polymeric compound $\{rac\text{-}(\text{CH}_3)_2\text{C}(\text{C}_5\text{H}_3\text{-3-Si}(\text{CH}_3)_3)_2\text{-Yb}[(\mu_2\text{-H})_2\text{B}(\mu_2\text{-H})_2]\}_2\text{Li}(\text{THF})_2\}_{\infty}$, 5

Synthesis, structure and reactivity of novel W–Ge chlorocarbonyl compounds. X-ray crystal structure of $[\text{WCl}(\text{GeCl}_3)(\text{CO})_3\text{(NCEt)}_2]$, 143

Crystal structures

A synthetic and X-ray crystallographic study of the indenyl-phosphine complexes 1,3-($\text{Ph}_2\text{P}=\text{X})_2\text{C}_9\text{H}_6$, ($\text{X} = \text{O, S}$) and $(\eta^5\text{-C}_9\text{H}_5(\text{Ph}_2\text{P}=\text{S})_2)[\text{Mn}(\text{CO})_3]$: versatile ligands for the preparation of heteropolymetallic complexes, 101

Synthesis and reactivity of tetramethylthioureatriosmium clusters: crystal structures of $[(\mu\text{-H})\text{Os}_3(\text{CO})_8(\mu_3\text{-S})(\mu\text{-OMeCO})\{\eta^1\text{-SC}(\text{NMe}_2)_2\}]$ and $[\text{Os}_3(\text{CO})_9(\mu\text{-OH})(\mu\text{-OMeCO})(\text{PPh}_3)]$, 133

Cycloaddition

[3 + 2] Cycloaddition reactions of the phospha-alkyne, $\text{Bu}'\text{CP}$, with a transition metal trisulphide containing terminal thio ligands: syntheses, crystal and molecular structures of $[\text{W}(\eta^5\text{-C}_5\text{Me}_5)(\text{S})(\text{S}_2\text{PC}'\text{Bu})]_2$ and $[\text{W}(\eta^5\text{-C}_5\text{Me}_5)(\text{S})(\text{S}_2\text{C}_2\text{Ph}_2)]_2$, 189

Cyclopentadienyl

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzimidinate supporting ligands, 209

Synthesis and reactivity of cyclopentadienyl chloro, imido and alkylidene tungsten (VI) complexes, 277

Synthesis and structural characterisation of nickel(II) alkyl complexes NiR_2 , $[\text{Ni}(\text{Cp})\text{R}]_2$ and a ‘head-to-tail’ coupled alkyl intermediate compound $[\text{NiR}(\text{Cl})(\text{R}-\text{R})]$ [$\text{R} = \text{C}(\text{SiMe}_3)_2\text{C}_5\text{H}_4\text{N-2}$, $\text{CPh}(\text{SiMe}_3)\text{C}_5\text{H}_4\text{N-2}$ or $\text{CH}(\text{SiMe}_3)\text{C}_5\text{H}_4\text{N-2}$; $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$], 193

Cyclopentadienylnickel(II)

Reversible sulfur dioxide reactions with cyclopentadienylnickel(II) organochalcogenide complexes, 37

Cyclopropene

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of $\text{MoCl}_3(\text{NAr})\text{L}_2$ ($\text{Ar} = 2,6\text{-diisopropyl phenyl}$; $\text{L} = \text{N-2',6'-diisopropyl phenyl-2,5-dimethyl benzamide}$), 85

Cyclozirconation

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Dehydrocoupling

Zirconocene-catalyzed dehydrogenative coupling of phenylgermane and properties of the resulting partially network polyphenylgermanes, 81

Dihydrogen complexes

New osmium hydrides containing the bulky phosphine 1,2-bis(diisopropylphosphino)ethane (dippe), 21

Dioxygen complexes

New osmium hydrides containing the bulky phosphine 1,2-bis(diisopropylphosphino)ethane (dippe), 21

Diphosphines

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Ethylene

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands, 115

Ethylene polymerization

Synthesis, characterization and polymerization potential of *ansa*-metallocene dichloride complexes of titanium, zirconium and hafnium containing a Si–N–Si bridging unit, 109

Ferrocene

Transmetallation reaction of cyclomercurated anils of benzoylferrocene with metallic tin. X-ray crystal structure of $[\text{SnCl}_2\{(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_3\text{CPh=NC}_6\text{H}_4\text{-4-CH}_3)\}_2]$, 233

Ferrocenylamines

Relationships between basicity, redox behaviour of ferrocenylamines and their reactivity with Pt[II] compounds, 125

Fluorenyl

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands, 115

German

Synthesis, structure and reactivity of novel W–Ge chlorocarbonyl compounds. X-ray crystal structure of $[WCl(GeCl_3)(CO)_3(NCEt)_2]$, 143

Germyl selenide

Palladium(0)-catalyzed regio- and stereoselective addition of heteroatom compounds bearing Si–Se, Ge–Se, and Si–Ge bonds to phenylacetylene, 1

Group IV metal

Synthesis, characterization and polymerization potential of *ansa*-metallocene dichloride complexes of titanium, zirconium and hafnium containing a Si–N–Si bridging unit, 109

Half-sandwich complexes

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands, 115

Hetero(Si, Ge, P)cyclopentanes

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Hydride complexes

New osmium hydrides containing the bulky phosphine 1,2-bis(diisopropylphosphino)ethane (dippe), 21

Hydroido complexes

C–H activation of methyl vinyl ketone in Ir(acac){ η^2 -CH₂=CHC(O)CH₃} (PCy₃), 241

Hydrogenation

Chirale Bisphosphane VIII. Die enantioselektive Hydrierung von α -Acetamidozimtsäure durch kationische Rhodium(I)-Komplexe mit optisch reinen (1*S*,2*S*)-Cyclopantan-1,2-diyl-bis(phosphonigsäurediester)- und (1*S*,2*S*)-Cyclopantan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden, 227

Reactivity of the 1-azavinylidene cluster $[Ru_3(\mu\text{-H})(\mu\text{-N=CPh}_2)(CO)]_{10}$ with hydrogen, tertiary silanes and tertiary stannanes, 201

Imide

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzamidinate supporting ligands, 209

Imido

Synthesis and reactivity of cyclopentadienyl chloro, imido and alkylidene tungsten (VI) complexes, 277

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of MoCl₃(NAr)L₂ (Ar = 2,6-diisopropyl phenyl; L = N-2',6'-diisopropyl phenyl-2,5-dimethyl benzamide), 85

Indenyl

Base-free tris(indenyl)lanthanoid(III) complexes (Ln = La, Pr, Nd, Sm): solid-state structure and solution NMR/NIR-vis spectroscopy, 167

Synthesis, characterization and polymerization potential of *ansa*-metallocene dichloride complexes of titanium, zirconium and hafnium containing a Si–N–Si bridging unit, 109

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands, 115

Indenyl-phosphine complexes

A synthetic and X-ray crystallographic study of the indenyl-phosphine complexes 1,3-(Ph₂P=)₂(C₉H₆), (X = O, S) and (η^5 -C₉H₅(Ph₂P=S)₂)[Mn(CO)₃]: versatile ligands for the preparation of heteropolymetallic complexes, 101

Intramolecular hydrosilylation

Unusual Pt-catalyzed stereo and regioselective intramolecular hydrosilylation of propargyl alcohols to (*E,E*)-1,6-dioxa-2,7-disila-3,8-ecidienes, 57

IR(ν (CO)) splitting

The synthesis of [AuCo(CO)₄]₂(μ -dppfe) (dppfe = 1,1'-bis(diphenyl phosphino)ferrocene) and the P–Au–Co–CO(axial) skeletal pair formation between neighboring molecules in the crystal, 249

Iridium

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Iridium complexes

C–H activation of methyl vinyl ketone in Ir(acac){ η^2 -CH₂=CHC(O)CH₃} (PCy₃), 241

Lanthanoid

Base-free tris(indenyl)lanthanoid(III) complexes (Ln = La, Pr, Nd, Sm): solid-state structure and solution NMR/NIR-vis spectroscopy, 167

Manganese

Synthesis, characterization and chemistry of some long chain alkoxy carbonyl manganese pentacarbonyl complexes [Mn(CO)₅{C(O)OR}] (where R = n-C₇H₁₅ to n-C₁₀H₂₁, n-C₁₂H₂₅, n-C₁₄H₂₉, and n-C₁₆H₃₃), 267

Mercury

Transmetallation reaction of cyclomercurated anils of benzoylferrocene with metallic tin. X-ray crystal structure of [SnCl₂{(η^5 -C₅H₅)Fe(η^5 -C₅H₃CPh=NC₆H₄-4-CH₃)}₂], 233

Metal complexes

Polynuclear metal complexes linked by conjugated pyridines, 257

Methyl vinyl ketone complexes

C–H activation of methyl vinyl ketone in Ir(acac){ η^2 -CH₂=CHC(O)CH₃} (PCy₃), 241

Molybdenum

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of MoCl₃(NAr)L₂ (Ar = 2,6-diisopropyl phenyl; L = N-2',6'-diisopropyl phenyl-2,5-dimethyl benzamide), 85

Nickel(II)

Synthesis and structural characterisation of nickel(II) alkyl complexes NiR₂, [Ni(Cp)R]₂ and a ‘head-to-tail’ coupled alkyl intermediate compound [NiR(Cl)(R–R)][R = C(SiMe₃)₂C₅H₄N-2, CPh(SiMe₃)C₅H₄N-2 or CH(SiMe₃)C₅H₄N-2; Cp = η^5 -C₅H₅], 193

Niobocene

Syntheses and properties of tetrathio- and tetraseleno metalates [(C₅Me₄R)₂NbE₂]₂M (E = S, Se; M = Cr, Mo; R = Me, Et) with peripheric niobocene ligands, 71

Oligophenylene

Polymeric organosilicon systems. XXIX. Thermal properties of poly[(disilanylene)oligophenylenes], 47

Organochalcogenide

Reversible sulfur dioxide reactions with cyclopentadienylnickel(II) organochalcogenide complexes, 37

Organosilicon polymer

Polymeric organosilicon systems. XXIX. Thermal properties of poly[(disilanylene)oligophenylenes], 47

Osmium

New osmium hydrides containing the bulky phosphine 1,2-bis-diisopropylphosphinoethane (dippe), 21

Synthesis and reactivity of tetramethylthioureatetrosmium clusters: crystal structures of $[(\mu\text{-H})\text{Os}_3(\text{CO})_8(\mu_3\text{-S})(\mu\text{-OMeCO})\{\eta^1\text{-SC}(\text{NMe}_2)_2\}]$ and $[\text{Os}_3(\text{CO})_9(\mu\text{-OH})(\mu\text{-OMeCO})(\text{PPh}_3)]$, 133

Oxo

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of $\text{MoCl}_3(\text{NAr})\text{L}_2$ ($\text{Ar} = 2,6\text{-diisopropyl phenyl}$; $\text{L} = \text{N}-2',6'\text{-diisopropyl phenyl-2,5-dimethyl benzamide}$), 85

Palladium

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Palladium catalyst

Palladium(0)-catalyzed regio- and stereoselective addition of heteroatom compounds bearing Si–Se, Ge–Se, and Si–Ge bonds to phenylacetylene, 1

Phosphenium ion

Reactions of protic reagents with a tungsten phosphenium ion complex, 13

Phosphine

New osmium hydrides containing the bulky phosphine 1,2-bis-diisopropylphosphinoethane (dippe), 21

Phosphines

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Phosphorus

Reactions of protic reagents with a tungsten phosphenium ion complex, 13

Platinum

Chirale Bisphosphane VI. Chelatkoplexe von Platin(II) mit racemischen und optisch reinen *trans*-Cyclopentan-1,2-diyli-bis(phosphonigsäurediester)- und *trans*-Cyclopentan-1,2-diyli-bis(phosphonigsäurediamid)-Liganden: Synthesen, Strukturen und C–H-Aktivierungsreaktionen, 155

Platinum atoms

Unusual Pt-catalyzed stereo and regioselective intramolecular hydrosilylation of propargyl alcohols to (*E,E*)-1,6-dioxa-2,7-disila-3,8-ecidienes, 57

Platinum compounds

Relationships between basicity, redox behaviour of ferrocenylamines and their reactivity with Pt[II] compounds, 125

P-ligands

Chirale Bisphosphane VI. Chelatkoplexe von Platin(II) mit racemischen und optisch reinen *trans*-Cyclopentan-1,2-diyli-bis(phosphonigsäurediester)- und *trans*-Cyclopentan-1,2-diyli-bis(phosphonigsäurediamid)-Liganden: Synthesen, Strukturen und C–H-Aktivierungsreaktionen, 155

Chirale Bisphosphane VIII. Die enantioselektive Hydrierung von α -Acetamidozimtsäure durch kationische Rhodium(I)-Komplexe mit optisch reinen (1*S,2S*)-Cyclopentan-1,2-diyli-bis(phosphonigsäurediester)- und (1*S,2S*)-Cyclopentan-1,2-diyli-bis(phosphonigsäurediamid)-Liganden, 227

Polygermane

Zirconocene-catalyzed dehydrogenative coupling of phenylgermane and properties of the resulting partially network polyphenylgermanes, 81

Polymerization

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands, 115

Synthesis, structure and reactivity of novel W–Ge chlorocarbonyl compounds. X-ray crystal structure of $[\text{WCl}(\text{GeCl}_3)(\text{CO})_3(\text{NCEt})_2]$, 143

Polynuclear complexes

Syntheses and properties of tetrathio- and tetraseleno metalates $[(\text{C}_5\text{Me}_4\text{R})_2\text{NbE}_2]_2\text{M}$ ($\text{E} = \text{S}, \text{Se}$; $\text{M} = \text{Cr}, \text{Mo}$; $\text{R} = \text{Me}, \text{Et}$) with peripheric niobocene ligands, 71

Polynuclear

Polynuclear metal complexes linked by conjugated pyridines, 257

Propargyl alcohols

Unusual Pt-catalyzed stereo and regioselective intramolecular hydrosilylation of propargyl alcohols to (*E,E*)-1,6-dioxa-2,7-disila-3,8-ecidienes, 57

Pyridine substituted alkyls

Synthesis and structural characterisation of nickel(II) alkyl complexes NiR_2 , $[\text{Ni}(\text{Cp})\text{R}]_2$ and a ‘head-to-tail’ coupled alkyl intermediate compound $[\text{NiR}(\text{Cl})(\text{R–R})]$ [$\text{R} = \text{C}(\text{SiMe}_3)_2\text{C}_5\text{H}_4\text{N-2}$, $\text{CPh}(\text{SiMe}_3)\text{C}_5\text{H}_4\text{N-2}$ or $\text{CH}(\text{SiMe}_3)\text{C}_5\text{H}_4\text{N-2}$; $\text{Cp} = \eta^5\text{-C}_5\text{H}_5$], 193

Pyrolysis

Polymeric organosilicon systems. XXIX. Thermal properties of poly[(disilanylene)oligophenylenes], 47

Redox studies

Relationships between basicity, redox behaviour of ferrocenylamines and their reactivity with Pt[II] compounds, 125

Regio- and stereoselective addition

Palladium(0)-catalyzed regio- and stereoselective addition of heteroatom compounds bearing Si–Se, Ge–Se, and Si–Ge bonds to phenylacetylene, 1

Reversible

Reversible sulfur dioxide reactions with cyclopentadienylnickel(II) organochalcogenide complexes, 37

Rhodium

Chirale Bisphosphane VIII. Die enantioselektive Hydrierung von α -Acetamidozimtsäure durch kationische Rhodium(I)-Komplexe mit optisch reinen (1*S,2S*)-Cyclopentan-1,2-diyli-bis(phosphonigsäurediester)- und (1*S,2S*)-Cyclopentan-1,2-diyli-bis(phosphonigsäurediamid)-Liganden, 227

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Ruthenium

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Reactivity of the 1-azavinylidene cluster $[\text{Ru}_3(\mu\text{-H})(\mu\text{-N=CPh}_2)(\text{CO})_{10}]$ with hydrogen, tertiary silanes and tertiary stannanes, 201

SAMP

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Schiff base complexes

Halbsandwich-Komplexe von (*S*)-1-Amino-2-(methoxymethyl)-pyrrolidin (SAMP), 179

Selectivity

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Selenium

Syntheses and properties of tetrathio- and tetraseleno metalates $[(\text{C}_5\text{Me}_4\text{R})_2\text{NbE}_2]_2\text{M}$ ($\text{E} = \text{S}, \text{Se}$; $\text{M} = \text{Cr}, \text{Mo}$; $\text{R} = \text{Me}, \text{Et}$) with peripheric niobocene ligands, 71

Silicon

Organotin compounds containing a bulky $(\text{Me}_3\text{Si})_3\text{C}$ or related ligand. Crystal structures of $\{(\text{Me}_3\text{Si})_3\text{CMe}(\text{O}_2\text{NO})\text{Sn}\}_2\text{O}$, $(\text{PhMe}_2\text{Si})_3\text{CSnMeCl}_2$ and $(\text{PhMe}_2\text{Si})_3\text{CSnCl}_3$, 215

Unusual Pt-catalyzed stereo and regioselective intramolecular hydrosilylation of propargyl alcohols to (*E,E*)-1,6-dioxa-2,7-disila-3,8-ecidienes, 57

Silylgermane

Palladium(0)-catalyzed regio- and stereoselective addition of heteroatom compounds bearing Si–Se, Ge–Se, and Si–Ge bonds to phenylacetylene, 1

Silyl selenide

Palladium(0)-catalyzed regio- and stereoselective addition of heteroatom compounds bearing Si–Se, Ge–Se, and Si–Ge bonds to phenylacetylene, 1

Single crystal X-ray analysis

The synthesis of $[\text{AuCo}(\text{CO})_4]_2(\mu\text{-dppfe})$ (dppfe = 1,1'-bis(diphenyl phosphino)ferrocene) and the P–Au–Co–CO(axial) skeletal pair formation between neighboring molecules in the crystal, 249

Sulfur

Syntheses and properties of tetrathio- and tetraseleno metalates $[(\text{C}_5\text{Me}_4\text{R})_2\text{NbE}_2]_2\text{M}$ ($\text{E} = \text{S}, \text{Se}; \text{M} = \text{Cr}, \text{Mo}; \text{R} = \text{Me}, \text{Et}$) with peripheric niobocene ligands, 71

Sulfur dioxide

Reversible sulfur dioxide reactions with cyclopentadienylnickel(II) organochalcogenide complexes, 37

Tertiary stannane

Reactivity of the 1-azavinylidene cluster $[\text{Ru}_3(\mu\text{-H})(\mu\text{-N}=\text{CPh}_2)(\text{CO})_{10}]$ with hydrogen, tertiary silanes and tertiary stannanes, 201

Tetramethylthiourea

Synthesis and reactivity of tetramethylthioureatriosmium clusters: crystal structures of $[(\mu\text{-H})\text{Os}_3(\text{CO})_8(\mu_3\text{-S})(\mu\text{-OMeCO})\{\eta^1\text{-SC}(\text{NMe}_2)_2\}]$ and $[\text{Os}_3(\text{CO})_9(\mu\text{-OH})(\mu\text{-OMeCO})(\text{PPh}_3)]$, 133

Thio ligands

[3 + 2] Cycloaddition reactions of the phospha-alkyne, Bu'CP, with a transition metal trisulphide containing terminal thio ligands: syntheses, crystal and molecular structures of $[\text{W}(\eta^5\text{-C}_5\text{Me}_5)(\text{S})(\text{S}_2\text{PC}'\text{Bu})]_2$ and $[\text{W}(\eta^5\text{-C}_5\text{Me}_5)(\text{S})(\text{S}_2\text{C}_2\text{Ph}_2)]_2$, 189

Tin

Organotin compounds containing a bulky $(\text{Me}_3\text{Si})_3\text{C}$ or related ligand. Crystal structures of $\{(\text{Me}_3\text{Si})_3\text{CMe}(\text{O}_2\text{NO})\text{Sn}\}_2\text{O}$, $(\text{PhMe}_2\text{Si})_3\text{CSnMeCl}_2$ and $(\text{PhMe}_2\text{Si})_3\text{CSnCl}_3$, 215

Transmetallation reaction of cyclomercurated anils of benzoyl-ferrocene with metallic tin. X-ray crystal structure of $[\text{SnCl}_2\{\eta^5\text{-C}_5\text{H}_5\}\text{Fe}(\eta^5\text{-C}_5\text{H}_3\text{CPh}=\text{NC}_6\text{H}_4\text{-4-CH}_3)\}_2$, 233

Titanium

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzimidinate supporting ligands, 209

Reactivity of chlorodimethylsilyl- η^5 -cyclopentadienyltrichlorotitanium with nitrogen based donors. X-ray molecular structure of $[\text{Ti}\{\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2[\eta^1\text{-N}(2,6\text{-Me}_2\text{C}_6\text{H}_3)]\}\text{Cl}_2]_2$, 93

Trans-4,5-bis(2-pyridyl)norbornane

Trans-4,5-bis(2-pyridyl)norbornane: a bidentate nitrogen ligand with a potentially large bite angle, 29

Transmetallation

Transmetallation reaction of cyclomercurated anils of benzoyl-ferrocene with metallic tin. X-ray crystal structure of $[\text{SnCl}_2\{\eta^5\text{-C}_5\text{H}_5\}\text{Fe}(\eta^5\text{-C}_5\text{H}_3\text{CPh}=\text{NC}_6\text{H}_4\text{-4-CH}_3)\}_2$, 233

Tris(dimethylphenylsilyl)methyl

Organotin compounds containing a bulky $(\text{Me}_3\text{Si})_3\text{C}$ or related ligand. Crystal structures of $\{(\text{Me}_3\text{Si})_3\text{CMe}(\text{O}_2\text{NO})\text{Sn}\}_2\text{O}$, $(\text{PhMe}_2\text{Si})_3\text{CSnMeCl}_2$ and $(\text{PhMe}_2\text{Si})_3\text{CSnCl}_3$, 215

Tris(trimethylsilyl)methyl

Organotin compounds containing a bulky $(\text{Me}_3\text{Si})_3\text{C}$ or related ligand. Crystal structures of $\{(\text{Me}_3\text{Si})_3\text{CMe}(\text{O}_2\text{NO})\text{Sn}\}_2\text{O}$, $(\text{PhMe}_2\text{Si})_3\text{CSnMeCl}_2$ and $(\text{PhMe}_2\text{Si})_3\text{CSnCl}_3$, 215

Trisulfide

[3 + 2] Cycloaddition reactions of the phospha-alkyne, Bu'CP, with a transition metal trisulphide containing terminal thio ligands: syntheses, crystal and molecular structures of $[\text{W}(\eta^5\text{-C}_5\text{Me}_5)(\text{S})(\text{S}_2\text{PC}'\text{Bu})]_2$ and $[\text{W}(\eta^5\text{-C}_5\text{Me}_5)(\text{S})(\text{S}_2\text{C}_2\text{Ph}_2)]_2$, 189

Trisyl

Organotin compounds containing a bulky $(\text{Me}_3\text{Si})_3\text{C}$ or related ligand. Crystal structures of $\{(\text{Me}_3\text{Si})_3\text{CMe}(\text{O}_2\text{NO})\text{Sn}\}_2\text{O}$, $(\text{PhMe}_2\text{Si})_3\text{CSnMeCl}_2$ and $(\text{PhMe}_2\text{Si})_3\text{CSnCl}_3$, 215

Tungsten

Reactions of protic reagents with a tungsten phosphonium ion complex, 13

Synthesis and reactivity of cyclopentadienyl chloro, imido and alkylidene tungsten (VI) complexes, 277

Synthesis, structure and reactivity of molybdenum imido and oxo complexes. X-ray structure of $\text{MoCl}_3(\text{NAr})\text{L}_2$ ($\text{Ar} = 2,6\text{-diisopropyl phenyl}; \text{L} = \text{N}-2',6'\text{-diisopropyl phenyl-2,5-dimethyl benzamide}$), 85

Tungsten(II)

Synthesis, structure and reactivity of novel W–Ge chlorocarbonyl compounds. X-ray crystal structure of $[\text{WCl}(\text{GeCl}_3)(\text{CO})_3(\text{NCEt})_2]$, 143

Versatile ligands

A synthetic and X-ray crystallographic study of the indenyl-phosphine complexes $1,3-(\text{Ph}_2\text{P}=\text{X})_2(\text{C}_9\text{H}_6)$, ($\text{X} = \text{O}, \text{S}$) and $(\eta^5\text{-C}_9\text{H}_5(\text{Ph}_2\text{P}=\text{S})_2)[\text{Mn}(\text{CO})_3]$: versatile ligands for the preparation of heteropolymetallic complexes, 101

X-ray diffraction

New binuclear alkyl and half-sandwich cyclopentadienyl imido titanium complexes containing acetamidinate and benzimidinate supporting ligands, 209

X-ray structure

Reactivity of the 1-azavinylidene cluster $[\text{Ru}_3(\mu\text{-H})(\mu\text{-N}=\text{CPh}_2)(\text{CO})_{10}]$ with hydrogen, tertiary silanes and tertiary stannanes, 201

X-ray structure analysis

Chirale Bisphosphane VI. Chelatkomplexe von Platin(II) mit racemischen und optisch reinen *trans*-Cyclopentan-1,2-diyl-bis(phosphonigsäurediester)- und *trans*-Cyclopentan-1,2-diyl-bis(phosphonigsäurediamid)-Liganden: Synthesen, Strukturen und C–H-Aktivierungsreaktionen, 155

Ytterbium (III)

Synthesis and structural study of the ate-complex *rac*-($\text{CH}_3)_2\text{C}(\text{C}_5\text{H}_3\text{-3-Si}(\text{CH}_3)_3)_2\text{Yb}(\mu_2\text{-Cl})_2\text{Li}(\text{OEt}_2)_2$ and the polymeric compound $\{\text{rac-}(\text{CH}_3)_2\text{C}(\text{C}_5\text{H}_3\text{-3-Si}(\text{CH}_3)_3)_2\text{Yb}[(\mu_2\text{-H})_2\text{B}(\mu_2\text{-H})_2]\}_2\text{Li}(\text{THF})_2\}_{\infty}$, 5

Zirconacomplexes

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61

Zirconium

Synthesis, characterization and polymerization potential of unbridged and bridged half-sandwich complexes of zirconium containing *N*-donor ligands, 115

Zirconocene-catalyzed dehydrogenative coupling of phenylgermane and properties of the resulting partially network polyphenylgermanes, 81

Zirconocene

Stereoselective synthesis of new hetero(P, Si, Ge, Sn)cyclic derivatives from zirconium diyne and diene complexes, 61