

Subject Index of Volume 580

Ab initio

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, 320

Acetylides

Solid-state aggregation of mercury bis-acetylides, $\text{Hg}(\text{C}\equiv\text{CR})_2$, $\text{R} = \text{Ph}, \text{SiMe}_3$, 363

4-Acyl-5-pyrazolones

Tin(IV) and organotin(IV) derivatives of novel β -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), 344

Aggregation

Solid-state aggregation of mercury bis-acetylides, $\text{Hg}(\text{C}\equiv\text{CR})_2$, $\text{R} = \text{Ph}, \text{SiMe}_3$, 363

Alkenes

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

Alkoxide

Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, 108

Alkyl

Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, 110

Alkyl amido imido cyclopentadienyl complexes

Half-sandwich dichloro, alkyl chloro, dialkyl, alkyl methyl and amido methyl imido cyclopentadienyl niobium and tantalum(V) complexes. Dynamic behaviour of amido imido tantalum derivatives, 161

Alkylation

Reactivity of AlMe_3 with titanium(IV) Schiff base complexes: X-ray structure of $[\text{Ti}\{(\mu\text{-Br})(\text{AlMe}_2)\}\{(\mu\text{-Br})(\text{AlMe}_2\text{X})\}(\text{salen})] \cdot \text{C}_7\text{H}_8$ ($\text{X} = \text{Me}$ or Br) and reactivity studies of mono-alkylated $[\text{Ti}(\text{Me})\text{X}(\text{L})]$ complexes, 304

Alkylidene

Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, 110

Alkyl lithium

NMR study of the dynamic behaviour of the system 2-ethylhexylsodium-2-ethylhexyllithium in heptane, 205

Alkylsodium

NMR study of the dynamic behaviour of the system 2-ethylhexylsodium-2-ethylhexyllithium in heptane, 205

Alkynes

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L-}P,P')(\eta^2\text{-RC}_2\text{R})]$ $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$. X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L-}P,P')(\eta^2\text{-PhC}_2\text{Ph})]$ ($\text{X} = \text{Br}$ and I) and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPPh}_2\text{-}P,P',O\}]$, 265

Allyl

Syntheses and X-ray structures of complexes with $(\eta^3\text{-allyl})\text{Mo}$ units in oxygen-rich coordination spheres, 214

1,1-Allylboration

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

1,2-Allylboration

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

Amine

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPPh}_3)$, 82

Aromatic compounds

Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, 290

Arylation

Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, 191

Assembling ligands

Synthesis of Co_2Pt , Co_2Pd and MoPd_2 mixed-metal clusters with the P–N–P assembling ligands $(\text{Ph}_2\text{P})_2\text{NH}$ (dppa) and $(\text{Ph}_2\text{P})_2\text{NMe}$ (dppaMe). Crystal structure of $[\text{Co}_2\text{Pt}(\mu_3\text{-CO})(\text{CO})_6(\mu\text{-dppa})]$, 257

Aza-allyl

Erratum to “The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry”: [J. Organometallic Chem. 529 (1997) 243–255], 385

Benzene

Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$, $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bistrimethylsilylcyclopentadienone})_2$ and 2,5-bis(trimethylsilyl) cyclopentadiene-2,4-dien-1-one, 98

Benzimidazole

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26

Benzylamine

Cyclopalladation of mono-, di- and tribenzylamine by palladium(II) acetate. Influence of bulkiness around the nitrogen atom of benzylamine upon internal metallation, 273

Bimetallic

Synthesis of Co_2Pt , Co_2Pd and MoPd_2 mixed-metal clusters with the P–N–P assembling ligands $(\text{Ph}_2\text{P})_2\text{NH}$ (dppa) and $(\text{Ph}_2\text{P})_2\text{NMe}$ (dppaMe). Crystal structure of $[\text{Co}_2\text{Pt}(\mu_3\text{-CO})(\text{CO})_6(\mu\text{-dppa})]$, 257

Bis(indenylidene) isomers

Synthesis and molecular structure of *cisoid* and *transoid* isomers of $[\text{Re}_2(\text{CO})_6(1,1'\text{-bis(indenylidene)})]$ derived from the reaction of $[\text{Re}_2(\text{CO})_8(\text{MeCN})_2]$ and diazoindene, 339

Bond strength

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282

Borane adducts

Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, 320

Borole ligand

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PET}_3)_2]$, 66

Boron

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

Bridging-sulfur ligand

Synthesis of $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$ and their reactivity toward electrophiles, 201

Carbene

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPh}_3)$, 82

Carbon dioxide

Bis(pentamethylcyclopentadienyl) uranium(IV) thiolate compounds. Synthesis and reactions with CO_2 and CS_2 , 137

Carbon disulfide

Bis(pentamethylcyclopentadienyl) uranium(IV) thiolate compounds. Synthesis and reactions with CO_2 and CS_2 , 137

Carbon–hydrogen and carbon–sulfur bond activation

Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, 370

Carbon monoxide

Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, 290

Carbonyls

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the η^1 -complexes $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}_2^5)]$ ($\text{M} = \text{Cr}, \text{W}$) and $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}_2^5)]$, 156

Site-selective carbonyl substitution in the mixed-metal cluster anion $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, 225

Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L-}P, P')(\eta^2\text{-RC}_2\text{R})]$ $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$. X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L-}P, P')(\eta^2\text{-PhC}_2\text{Ph})]$ ($\text{X} = \text{Br}$ and I) and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P, P', O\}]$, 265

Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$, $[\text{Os}_3\text{Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$ and $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})(\text{CO})_9(\eta^5\text{-Cp}^*)]$, 48

Carbonyl activation

Synthesis, reactions and X-ray crystal structures of metallocrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, 328

Carbonylmanganese cations

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, 295

Carboxylate

Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, 108

Carboxylation

Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, 290

Catalysis

C_2 -bridged metallocene dichloride complexes of the types $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_9\text{H}_6\text{-}n\text{R}'_n)\text{ZrCl}_2$ and $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_{13}\text{H}_8)\text{MCl}_2$ ($n = 0, 1$; $\text{R} = \text{H}$, alkenyl; $\text{R}' =$ alkenyl, benzyl; $\text{M} = \text{Zr}, \text{Hf}$) as self-immobilizing catalyst precursors for ethylene polymerization, 1

C–H activation

Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, 22

Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, 191

Chiral cyclodextrin sorbents

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26

Chiral ligands

A C_2 -symmetric analogue of dpfp: synthesis and structures of the indenyl-diphosphine complex $1,3\text{-}(\text{Ph}_2\text{PSe})_2(\text{C}_9\text{H}_6)$ and the racemic and *meso* isomers of the heterobimetallic complex $\text{Mo}(\text{CO})_4(1\text{-PPh}_2\text{-}\eta^5\text{-C}_9\text{H}_6)_2\text{Fe}$, 245

Chromium

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the η^1 -complexes $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}_2^5)]$ ($\text{M} = \text{Cr}, \text{W}$) and $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}_2^5)]$, 156

Clusters

Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$, $[\text{Os}_3\text{Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$ and $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})(\text{CO})_9(\eta^5\text{-Cp}^*)]$, 48

Synthesis of Co_2Pt , Co_2Pd and MoPd_2 mixed-metal clusters with the P–N–P assembling ligands $(\text{Ph}_2\text{P})_2\text{NH}$ (dppa) and $(\text{Ph}_2\text{P})_2\text{NMe}$ (dppaMe). Crystal structure of $[\text{Co}_2\text{Pt}(\mu_3\text{-CO})(\text{CO})_6(\mu\text{-dppa})]$, 257

Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, 370

Complexes

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the η^1 -complexes $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}_2^5)]$ ($\text{M} = \text{Cr}, \text{W}$) and $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}_2^5)]$, 156

Synthesis of $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$ and their reactivity toward electrophiles, 201

Coordination polymer

Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, 22

 Cp_2TiCl_2 catalyst

Cp_2TiCl_2 catalyzed one-pot synthesis of *n*- Bu_3GeH from GeCl_4 , 378

Crystal structure

Erratum to “The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry”: [J. Organometallic Chem. 529 (1997) 243–255], 385

Ring-opening reactions of thiomorpholine and thiazolidine by $[\text{Ru}_3(\text{CO})_{12}]$: crystal structures of $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{CH}_2\text{NH}_2)(\text{CO})_9]$ and $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$, 60

Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L-}P, P')(\eta^2\text{-RC}_2\text{R})]$ $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$. X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L-}P, P')(\eta^2\text{-PhC}_2\text{Ph})]$ ($\text{X} = \text{Br}$ and I) and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P, P', O\}]$, 265

Synthesis and molecular structure of the tetranuclear μ_4 -oxo-centered chromium complex $[\text{Cr}_4(\text{CH}_2\text{CH}_2\text{CH}_2\text{NMe}_2)_4(\mu_4\text{-O})(\mu\text{-Cl})_6]$, 56

- Synthesis and structural characterization of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$ and $[(\text{dppm})(\text{CO})_2\text{Mo}(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2\text{Mo}(\text{O})]_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-O})$ and electrochemical property of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$, 313
- Tin(IV) and organotin(IV) derivatives of novel β -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), 344
- Cyanide ligand
Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PEt}_3)_2]$, 66
- Cyclocarbaphosphazene
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41
- Cyclopalladation
Cyclopalladation of mono-, di- and tribenzylamine by palladium(II) acetate. Influence of bulkiness around the nitrogen atom of benzylamine upon internal metallation, 273
- Cyclopentadienone
Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$, $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bistrimethylsilylcyclopentadienone})_2$ and $2,5\text{-bis}(\text{trimethylsilyl})\text{cyclopenta-}2,4\text{-dien-}1\text{-one}$, 98
- Cyclopentadienyl
Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, 169
- Cyclopentadienyl–cyclohexadienyl bond
A directly ring-to-ring linked ferrocene–pseudotitanocene complex, 210
- Cyotoxicity
Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, 17
- Dealkylation reaction
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41
- Dialkylaminomethylferrocenes
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41
- Diazoindene
Synthesis and molecular structure of *cisoid* and *transoid* isomers of $[\text{Re}_2(\text{CO})_6(1,1'\text{-bis}(\text{indenylidene}))]$ derived from the reaction of $[\text{Re}_2(\text{CO})_8(\text{MeCN})_2]$ and diazoindene, 339
- Di-cobalt complexes
Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$, $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bistrimethylsilylcyclopentadienone})_2$ and $2,5\text{-bis}(\text{trimethylsilyl})\text{cyclopenta-}2,4\text{-dien-}1\text{-one}$, 98
- Diels–Alder reaction
Syntheses with organoboranes. IX. Vinyl- and 1-alkenyldichloroboranes as ethylene and 1-alkene equivalents for the Diels–Alder reaction, 354
- 2,6-Dimethoxyphenyl
Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282
- Dinuclear complexes
Syntheses of polymethylene bridged dinuclear zirconocenes and investigation of their polymerisation activities, 90
- 1,3-Diphenyldisiloxane
 CO_2 laser photosensitised decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, 188
- Diphosphines
Synthesis of Co_2Pt , Co_2Pd and MoPd_2 mixed-metal clusters with the P–N–P assembling ligands $(\text{Ph}_2\text{P})_2\text{NH}$ (dppa) and $(\text{Ph}_2\text{P})_2\text{NMe}$ (dppaMe). Crystal structure of $[\text{Co}_2\text{Pt}(\mu_3\text{-CO})(\text{CO})_6(\mu\text{-dppa})]$, 257
- Distannylated derivatives
Syntheses of polymethylene bridged dinuclear zirconocenes and investigation of their polymerisation activities, 90
- DNMR
NMR study of the dynamic behaviour of the system 2-ethylhexylsodium-2-ethylhexyllithium in heptane, 205
- Electrochemistry
Synthesis and structural characterization of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$ and $[(\text{dppm})(\text{CO})_2\text{Mo}(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2\text{Mo}(\text{O})]_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-O})$ and electrochemical property of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$, 313
- Electron transfer
Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, 295
- Electro-oxidation
Electrochemical and optical properties of poly[(disilanyl)oligophenylenes], peculiar behavior in the solid state, 77
- Exchange
Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, 320
- Face-capping ligand
A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, 239
- Ferrocene
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41
- Ferrocene derivatives
Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26
- Ferrocene–pseudotitanocene complex
A directly ring-to-ring linked ferrocene–pseudotitanocene complex, 210
- Ferrocenylacetylene
A directly ring-to-ring linked ferrocene–pseudotitanocene complex, 210
- α -Ferrocenylalkylation
Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26
- Flyover bridge
Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$, $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bistrimethylsilylcyclopentadienone})_2$ and $2,5\text{-bis}(\text{trimethylsilyl})\text{cyclopenta-}2,4\text{-dien-}1\text{-one}$, 98
- Formate
Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, 108
- Fungitoxicity
Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, 17
- GeCl_4
 Cp_2TiCl_2 catalyzed one-pot synthesis of *n*- Bu_3GeH from GeCl_4 , 378

Geometry constrained catalysts

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, 145

Germanium

New optically active organogermane compounds containing the adamantyl radical for heterogeneous bimetallic catalysis. Part II, 128

Halide

Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L}-P,P')(\eta^2\text{-RC}_2\text{R})]$ $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$. X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L}-P,P')(\eta^2\text{-PhC}_2\text{Ph})]$ ($\text{X} = \text{Br}$ and I) and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$, 265

Heterobimetallic

A C_2 -symmetric analogue of dpfp: synthesis and structures of the indenyl-diphosphine complex $1,3\text{-}(\text{Ph}_2\text{PSe})_2(\text{C}_9\text{H}_6)$ and the racemic and *meso* isomers of the heterobimetallic complex $\text{Mo}(\text{CO})_4(1\text{-PPH}_2\text{-}\eta^5\text{-C}_9\text{H}_6)_2\text{Fe}$, 245

Heterogeneous bimetallic catalysts

New optically active organogermane compounds containing the adamantyl radical for heterogeneous bimetallic catalysis. Part II, 128

Heterogeneous Ziegler–Natta polymerization

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, 145

Heterometallic complexes

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PEt}_3)_2]$, 66

Homodinuclear complexes

Preparations and characterizations of homodinuclear complexes containing flyover bridge or cyclopentadienone ligands; X-ray crystal structures of $\text{Co}_2(\text{CO})_3(\text{PPh}_3)\{\mu\text{-C}(\text{SiMe}_3)=\text{CH}-\text{CH}=\text{C}(\text{SiMe}_3)-\text{CH}=\text{C}(\text{SiMe}_3)\}$, $\text{Co}_2(\text{CO})_2(\mu\text{-CO})_2(\eta^4\text{-}2,5\text{-bistrimethylsilylcyclopentadienone})_2$ and $2,5\text{-bis}(\text{trimethylsilyl})\text{cyclopenta-}2,4\text{-dien-}1\text{-one}$, 98

Homogeneous Ziegler–Natta polymerization

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, 145

HPLC

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26

Hydride

Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, 169

Hydrido

Site-selective carbonyl substitution in the mixed-metal cluster anion $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, 225

Imido

Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, 110

 μ -Indenylidene

Synthesis and molecular structure of *cisoid* and *transoid* isomers of $[\text{Re}_2(\text{CO})_6(1,1'\text{-bis}(\text{indenylidene}))]$ derived from the reaction of $[\text{Re}_2(\text{CO})_8(\text{MeCN})_2]$ and diazoindene, 339

Indenyl-phosphine complexes

A C_2 -symmetric analogue of dpfp: synthesis and structures of the indenyl-diphosphine complex $1,3\text{-}(\text{Ph}_2\text{PSe})_2(\text{C}_9\text{H}_6)$ and the racemic and *meso* isomers of the heterobimetallic complex $\text{Mo}(\text{CO})_4(1\text{-PPH}_2\text{-}\eta^5\text{-C}_9\text{H}_6)_2\text{Fe}$, 245

Indeyl

A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, 239

Insertion

Exchange and insertion reactions involving borane adducts of

phosphirane and phosphirene: a G2(MP2) ab initio investigation, 320

Insertion reactions

Bis-pentamethylcyclopentadienyl uranium(IV) thiolate compounds. Synthesis and reactions with CO_2 and CS_2 , 137

Ion-pair annihilation

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, 295

IR

Tin(IV) and organotin(IV) derivatives of novel β -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), 344

Iridium

Site-selective carbonyl substitution in the mixed-metal cluster anion $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, 225

Iron

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PEt}_3)_2]$, 66

Synthesis of $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$ and their reactivity toward electrophiles, 201

IR spectroscopy

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, 36

Isocyanide

Methylcyclopentadienyl-vanadium(III) complexes, 72

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPH}_3)$, 82

Laser-induced decomposition

CO_2 laser photosensitized decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, 188

Linkage isomerism

Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of $[\{\eta^5\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PEt}_3)_2]$, 66

Lithium

Erratum to “The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry”: [J. Organometallic Chem. 529 (1997) 243–255], 385

Mercuriophilic interactions

Solid-state aggregation of mercury bis-acetylides, $\text{Hg}(\text{C}=\text{CR})_2$, $\text{R} = \text{Ph}, \text{SiMe}_3$, 363

Mercury

Solid-state aggregation of mercury bis-acetylides, $\text{Hg}(\text{C}=\text{CR})_2$, $\text{R} = \text{Ph}, \text{SiMe}_3$, 363

Metal carbonyls

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, 36

Metallacycles

IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, 36

Metallacrown ethers

Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, 328

Metallocene complexes

C_2 -bridged metallocene dichloride complexes of the types $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_9\text{H}_6\text{-}_n\text{R}'_n)\text{ZrCl}_2$ and $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_{13}\text{H}_8)\text{MCl}_2$ ($n = 0, 1$; $\text{R} = \text{H}, \text{alkenyl}$; $\text{R}' = \text{alkenyl}, \text{benzyl}$; $\text{M} = \text{Zr}, \text{Hf}$) as self-immobilizing catalyst precursors for ethylene polymerization, 1

- Methylcyclopentadienyl
Methylcyclopentadienyl-vanadium(III) complexes, 72
- Mixed-metal clusters
Site-selective carbonyl substitution in the mixed-metal cluster anion $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, 225
- Mixed-ring indenyl and bis-indenyl halides complexes
Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, 169
- Molybdenum
Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L}-P,P')(\eta^2\text{-RC}_2\text{R})]$ $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$. X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L}-P,P')(\eta^2\text{-PhC}_2\text{Ph})]$ $(\text{X} = \text{Br} \text{ and } \text{I})$ and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$, 265
Synthesis and structural characterization of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$ and $[(\text{dppm})(\text{CO})_2\text{Mo}(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2\text{Mo}(\text{O})_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-O})]$ and electrochemical property of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$, 313
- Molybdenum complexes
Syntheses and X-ray structures of complexes with $(\eta^3\text{-allyl})\text{Mo}$ -units in oxygen-rich coordination spheres, 214
- Mo_2S_2 unit
Synthesis and structural characterization of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$ and $[(\text{dppm})(\text{CO})_2\text{Mo}(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2\text{Mo}(\text{O})_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-O})]$ and electrochemical property of $\text{Mo}_2(\mu\text{-SC}_6\text{H}_4\text{-Cl-}p)_2(\mu\text{-dppm})(\text{CO})_6$, 313
- Niobium
Half-sandwich dichloro, alkyl chloro, dialkyl, alkyl methyl and amido methyl imido cyclopentadienyl niobium and tantalum(V) complexes. Dynamic behaviour of amido imido tantalum derivatives, 161
- NMR
 Cp_2TiCl_2 catalyzed one-pot synthesis of $n\text{-Bu}_3\text{GeH}$ from GeCl_4 , 378
Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234
Tin(IV) and organotin(IV) derivatives of novel β -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), 344
- NMR spectra
A directly ring-to-ring linked ferrocene-pseudotitanocene complex, 210
- N*-Trimethylsilylimines
Synthèse régiosélective par voie organozincique de 1,4-diamines primaires propargyliques et applications, 178
- Nucleophilic attack
Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, 295
- α -Olefin polymerization
Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, 145
- Oligophenylene polymer
Electrochemical and optical properties of poly[(disilanyl)ene]oligophenylenes, peculiar behavior in the solid state, 77
- One-pot synthesis
 Cp_2TiCl_2 catalyzed one-pot synthesis of $n\text{-Bu}_3\text{GeH}$ from GeCl_4 , 378
- Onium
Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282
- Optically active organogermane compounds
New optically active organogermane compounds containing the adamantly radical for heterogeneous bimetallic catalysis. Part II, 128
- Organoborate anions
Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, 295
- Organosilicon polymer
Electrochemical and optical properties of poly[(disilanyl)ene]oligophenylenes, peculiar behavior in the solid state, 77
- Organozincs
Synthèse régiosélective par voie organozincique de 1,4-diamines primaires propargyliques et applications, 178
- Osmium
Synthesis, characterization and crystal structures of osmium-rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$, $[\text{Os}_3\text{Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$ and $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})(\text{CO})_9(\eta^5\text{-Cp}^*)]$, 48
Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, 370
- Osmium carbonyl clusters synthesis
High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of $[\text{M}(\text{CO})_3\text{Cl}_2]_2$ or MCl_3 ($\text{M} = \text{Ru}, \text{Os}$) in the presence of Na_2CO_3 or K_2CO_3 , 117
- Osmium trichloride reductive carbonylation
High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of $[\text{M}(\text{CO})_3\text{Cl}_2]_2$ or MCl_3 ($\text{M} = \text{Ru}, \text{Os}$) in the presence of Na_2CO_3 or K_2CO_3 , 117
- Oxide complex
Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L}-P,P')(\eta^2\text{-RC}_2\text{R})]$ $\{\text{M} = \text{Mo}, \text{W}; \text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}; \text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2; \text{R} = \text{Me}, \text{Ph}\}$. X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L}-P,P')(\eta^2\text{-PhC}_2\text{Ph})]$ $(\text{X} = \text{Br} \text{ and } \text{I})$ and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$, 265
- Oxygen
Syntheses and X-ray structures of complexes with $(\eta^3\text{-allyl})\text{Mo}$ -units in oxygen-rich coordination spheres, 214
- Palladacycles
Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, 191
- Palladium
Cyclopalladation of mono-, di- and tribenzylamine by palladium(II) acetate. Influence of bulkiness around the nitrogen atom of benzylamine upon internal metallation, 273
Palladium(II)-catalyzed carboxylation of benzene and other aromatic compounds with carbon monoxide under very mild conditions, 290
- Pentamethylcyclopentadienyl
Synthesis, characterization and crystal structures of osmium-rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$, $[\text{Os}_3\text{Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$ and $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})(\text{CO})_9(\eta^5\text{-Cp}^*)]$, 48
- Phenylsilanone
 CO_2 laser photosensitised decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, 188
- Phosphazene
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41

- Phosphetidine**
Erratum to "The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry": [J. Organometallic Chem. 529 (1997) 243–255], 385
- Phosphine**
Site-selective carbonyl substitution in the mixed-metal cluster anion $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, 225
- Phosphinite**
Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, 328
- Phosphirane**
Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, 320
- Phosphirene**
Exchange and insertion reactions involving borane adducts of phosphirane and phosphirene: a G2(MP2) ab initio investigation, 320
- Phosphite**
Synthesis, reactions and X-ray crystal structures of metallacrown ethers with unsymmetrical bis(phosphinite) and bis(phosphite) ligands derived from 2-hydroxy-2'-(1,4-bisoxo-6-hexanol)-1,1'-biphenyl, 328
- Phosphonium salt**
Erratum to "The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry": [J. Organometallic Chem. 529 (1997) 243–255], 385
- Phosphorus**
Erratum to "The role of lithium 1,3-bis(trimethylsilyl)-1-aza-allyls in phosphorus chemistry": [J. Organometallic Chem. 529 (1997) 243–255], 385
- Platinum**
Cyano-bridged Fe–CN–Pt–NC–Fe chain complexes: structural and spectroscopic characterization of four isomers of $[\{\eta^2\text{-}(1\text{-phenylborole})\text{Fe}(\text{CO})_2(\mu\text{-CN})\}_2\text{Pt}(\text{PEt}_3)_2]$, 66
- Polyaromatics**
Multiple arylations of aromatics via removable palladacycle scaffolds: polycyclic compounds and their way of formation, 191
- Polyfluoro-benzimidazoles**
Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26
- Polymerization**
 C_2 -bridged metallocene dichloride complexes of the types $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_9\text{H}_6\text{-}_n\text{R}'_n)\text{ZrCl}_2$ and $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_{13}\text{H}_8)\text{MCl}_2$ ($n = 0, 1$; $\text{R} = \text{H}$, alkenyl; $\text{R}' =$ alkenyl, benzyl; $\text{M} = \text{Zr}$, Hf) as self-immobilizing catalyst precursors for ethylene polymerization, 1
- Polymethylene bridged dinuclear zirconocenes**
Syntheses of polymethylene bridged dinuclear zirconocenes and investigation of their polymerisation activities, 90
- Poly(phenylsiloxanes)**
 CO_2 laser photosensitised decomposition of 1,3-diphenyldisiloxane in the liquid phase: formation of poly(phenylsiloxanes) via extrusion/insertion of phenylsilanone, 188
- Porphyrin**
Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, 22
- Primary 1,4-diamines**
Synthèse régiosélective par voie organozincique de 1,4-diamines primaires propargyliques et applications, 178
- Reaction**
Synthesis of $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$ and their reactivity toward electrophiles, 201
- Reaction rate**
Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282
- Regiospecificity**
Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41
- Rhenium**
Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPH}_3)$, 82
- Rhenium carbonyls**
Synthesis and molecular structure of *cisoid* and *transoid* isomers of $[\text{Re}_2(\text{CO})_6(1,1'\text{-bis(indenylidene)})]$ derived from the reaction of $[\text{Re}_2(\text{CO})_8(\text{MeCN})_2]$ and diazoindene, 339
- Rhenium complexes**
IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, 36
- Rhodium**
Intermolecular C–H activation to a novel Rh nitrile bridged porphyrin coordination polymer, 22
Synthesis, characterization and crystal structures of osmium–rhodium mixed-metal clusters containing pentamethylcyclopentadienyl ligand: $[\text{Os}_3\text{Rh}(\mu\text{-CO})_2(\text{CO})_9(\eta^5\text{-Cp}^*)]$, $[\text{Os}_3\text{Rh}_2(\mu\text{-H})(\mu\text{-CO})_2(\text{CO})_8(\eta^5\text{-Cp}^*)(\mu_2\text{-}\eta^5, \eta^1\text{-CH}_2\text{C}_5\text{Me}_4)]$ and $[\text{Os}_3\text{Rh}(\mu_3\text{-H})(\mu\text{-Cl})(\mu\text{-CO})(\text{CO})_9(\eta^5\text{-Cp}^*)]$, 48
- Ring-opening**
Ring-opening reactions of thiomorpholine and thiazolidine by $[\text{Ru}_3(\text{CO})_{12}]$: crystal structures of $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{-CH}_2\text{NH}_2)(\text{CO})_9]$ and $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$, 60
- Rotational isomerism**
IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, 36
- Ruthenium**
A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, 239
Reactions of dinuclear carboxylate ruthenium (I) complexes with alcohols. The unexpected oxidation of primary alcohols to carboxylic ligands, 108
Ring-opening reactions of thiomorpholine and thiazolidine by $[\text{Ru}_3(\text{CO})_{12}]$: crystal structures of $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{-CH}_2\text{NH}_2)(\text{CO})_9]$ and $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$, 60
Site-selective carbonyl substitution in the mixed-metal cluster anion $[\text{H}_2\text{Ru}_3\text{Ir}(\text{CO})_{12}]^-$: synthesis and characterization of phosphine, phosphite, arsine and stibine derivatives, 225
- Ruthenium carbonyl clusters synthesis**
High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of $[\text{M}(\text{CO})_5\text{Cl}_2]_2$ or MCl_3 ($\text{M} = \text{Ru}$, Os) in the presence of Na_2CO_3 or K_2CO_3 , 117
- Ruthenium complexes**
IR study of rotational isomerism in dirhenium and diruthenium complexes of the ferrole type, 36
- Ruthenium trichloride reductive carbonylation**
High-yield syntheses, under mild conditions, of various neutral and anionic ruthenium and osmium carbonyl clusters by controlled reduction in ethylene glycol of $[\text{M}(\text{CO})_5\text{Cl}_2]_2$ or MCl_3 ($\text{M} = \text{Ru}$, Os) in the presence of Na_2CO_3 or K_2CO_3 , 117
- Schiff base**
Reactivity of AlMe_3 with titanium(IV) Schiff base complexes: X-ray structure of $[\text{Ti}\{\mu\text{-Br}(\text{AlMe}_2)\}\{\mu\text{-Br}(\text{AlMe}_2\text{X})\}\text{(salen)}] \cdot \text{C}_7\text{H}_8$ ($\text{X} = \text{Me}$ or Br) and reactivity studies of mono-alkylated $[\text{Ti}(\text{Me})\text{X}(\text{L})]$ complexes, 304
- Selenadiphosphole**
First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the η^1 -complexes $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}_2^*)]$ ($\text{M} = \text{Cr}$, W) and $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}_2^*)]$, 156
- Selenium**
Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282

Separation of enantiomers

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26

Silicon

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

Structure

Synthesis, structure, and reactivity of rhenium *N*-isocyanide complexes $\text{ReBr}(\text{CO})_3(\text{CNR})(\text{CNNPPPh}_3)$, 82

Sulfur

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282

Synthesis

Synthesis and molecular structure of the tetranuclear μ_4 -oxo-centered chromium complex $[\text{Cr}_4(\text{CH}_2\text{CH}_2\text{CH}_2\text{NMe}_2)_4(\mu_4\text{-O})(\mu\text{-Cl})_6]$, 56

Synthesis of $[(\mu\text{-RS})(\mu\text{-S})\{\text{Fe}_2(\text{CO})_6\}_2(\mu_4\text{-S})]^-$ and their reactivity toward electrophiles, 201

Tantalum

Half-sandwich dichloro, alkyl chloro, dialkyl, alkyl methyl and amido methyl imido cyclopentadienyl niobium and tantalum(V) complexes. Dynamic behaviour of amido imido tantalum derivatives, 161

Telluradiphosphole

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the η^1 -complexes $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}'_2)]$ ($\text{M} = \text{Cr}, \text{W}$) and $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}'_2)]$, 156

Tellurium

Bis(2,6-dimethoxyphenyl) sulfide, selenide and telluride, and their derivatives, 282

Tetrachlorodicarbaphosphatriazine

Dealkylation reactions of dialkylaminomethylferrocenes with cyclic chlorocarbaphosphazene, $(\text{ClCN})_2(\text{Cl}_2\text{PN})$, 41

Tetrachromium complexes

Synthesis and molecular structure of the tetranuclear μ_4 -oxo-centered chromium complex $[\text{Cr}_4(\text{CH}_2\text{CH}_2\text{CH}_2\text{NMe}_2)_4(\mu_4\text{-O})(\mu\text{-Cl})_6]$, 56

Tetramethylborate

Electron transfer versus nucleophilic pathways in the ion-pair annihilation of organoborate anions by carbonylmanganese(I) cations, 295

Tetranuclear cluster

A tetraruthenium carbonyl cluster bearing indenyl-type ligand as the facial bonding mode, 239

Thiazolidine

Ring-opening reactions of thiomorpholine and thiazolidine by $[\text{Ru}_3(\text{CO})_{12}]$: crystal structures of $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{CH}_2\text{NH}_2)(\text{CO})_9]$ and $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$, 60

Thiocarbonate compounds

Bispendamethylcyclopentadienyl uranium(IV) thiolate compounds. Synthesis and reactions with CO_2 and CS_2 , 137

Thiomorpholine

Ring-opening reactions of thiomorpholine and thiazolidine by $[\text{Ru}_3(\text{CO})_{12}]$: crystal structures of $[\text{Ru}_3(\mu\text{-H})(\mu\text{-}\eta^2\text{-SCH}_2\text{CH}_2\text{NH}_2)(\text{CO})_9]$ and $[\text{Ru}_2(\mu\text{-}\eta^3\text{-SCH}_2\text{CH}_2\text{NHCH}_2)(\text{CO})_6]$, 60

Thiophene

Thermal and photochemical transformations of 2-methylthiophene in triosmium clusters: photoinduced migrations of MeS from carbon to metal and back again, 370

Thiosemicarbazone

Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, 17

Tin

Synthesis, crystal structure and biological activity of thiophene-2-carboxaldehyde thiosemicarbazone and its tin complexes, 17

Titanium

Reactivity of AlMe_3 with titanium(IV) Schiff base complexes: X-ray structure of $[\text{Ti}\{\mu\text{-Br}(\text{AlMe}_2)\}\{\mu\text{-Br}(\text{AlMe}_2\text{X})\}(\text{salen})] \cdot \text{C}_7\text{H}_8$ ($\text{X} = \text{Me}$ or Br) and reactivity studies of mono-alkylated $[\text{Ti}(\text{Me})\text{X}(\text{L})]$ complexes, 304

Titanium complexes

Synthesis and olefin polymerization using supported and non-supported geometry constrained titanium complexes, 145

Titanium(II)

A directly ring-to-ring linked ferrocene-pseudotitanocene complex, 210

Triallylborane

Reaction of 1-alkynylsilanes with triallylborane—competition between 1,1- and 1,2-allylboration, 234

tri-*n*-Butylgermane

Cp_2TiCl_2 catalyzed one-pot synthesis of *n*- Bu_3GeH from GeCl_4 , 378

Triorganotin(IV) complex

Tin(IV) and organotin(IV) derivatives of novel β -diketones: Part IV. Triorganotin(IV) complexes of fluorinated 4-acyl-5-pyrazolones. Crystal structure of (1-(4-trifluoromethylphenyl)-3-methyl-4-acetylpyrazolon-5-ato)triphenyltin(IV), 344

Triphos

Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L-}P,P')(\eta^2\text{-RC}_2\text{R})]$ ($\text{M} = \text{Mo}, \text{W}$; $\text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}$; $\text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$; $\text{R} = \text{Me}, \text{Ph}$). X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L-}P,P')(\eta^2\text{-PhC}_2\text{Ph})]$ ($\text{X} = \text{Br}$ and I) and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$, 265

Tungsten

Alkyl and alkylidene imido cyclopentadienyl tungsten complexes, 110

First structural characterisation of 1,2,4-selenadiphosphole and 1,2,4-telluradiphosphole ring systems. Crystal and molecular structures of the η^1 -complexes $[\text{M}(\text{CO})_5(\text{P}_2\text{SeC}_2\text{Bu}'_2)]$ ($\text{M} = \text{Cr}, \text{W}$) and $[\text{W}(\text{CO})_5(\text{P}_2\text{TeC}_2\text{Bu}'_2)]$, 156

Synthesis and characterisation of the bidentately attached linear triphos complexes $[\text{MXY}(\text{CO})(\text{L-}P,P')(\eta^2\text{-RC}_2\text{R})]$ ($\text{M} = \text{Mo}, \text{W}$; $\text{X}, \text{Y} = \text{Cl}, \text{Br}, \text{I}$; $\text{L} = \text{PhP}(\text{CH}_2\text{CH}_2\text{PPh}_2)_2$; $\text{R} = \text{Me}, \text{Ph}$). X-ray crystal structures of $[\text{WX}_2(\text{CO})(\text{L-}P,P')(\eta^2\text{-PhC}_2\text{Ph})]$ ($\text{X} = \text{Br}$ and I) and $[\text{MoI}_2(\text{O})\{\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}(\text{CH}_2)_2\text{POPh}_2\text{-}P,P',O\}]$, 265

Tungstenocene

Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, 169

Uranium(IV) thiolate compounds

Bispendamethylcyclopentadienyl uranium(IV) thiolate compounds. Synthesis and reactions with CO_2 and CS_2 , 137

Vanadium

Methylcyclopentadienyl-vanadium(III) complexes, 72

Vinylic organoborane dienophiles

Syntheses with organoboranes. IX. Vinyl- and 1-alkenyldichloroboranes as ethylene and 1-alkene equivalents for the Diels–Alder reaction, 354

X-ray crystal structure

Synthesis and structure of biologically active ferrocenylalkyl polyfluoro benzimidazoles, 26

X-ray diffraction

Synthesis of bis-cyclopentadienyl, bis-indenyl and mixed-ring indenyl halides of tungsten, 169

Zirconium

C_2 -bridged metallocene dichloride complexes of the types $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_9\text{H}_6\text{-}_n\text{R}'_n)\text{ZrCl}_2$ and $(\text{C}_{13}\text{H}_8\text{-CH}_2\text{CHR-C}_{13}\text{H}_8)\text{MCl}_2$ ($n = 0, 1$; $\text{R} = \text{H}, \text{alkenyl}$; $\text{R}' = \text{alkenyl}, \text{benzyl}$; $\text{M} = \text{Zr}, \text{Hf}$) as self-immobilizing catalyst precursors for ethylene polymerization, 1