

Subject Index of Volume 591

- Alkoxide
C–H activation of pendant alkoxides by tungsten imide complexes, 104
- Alkyl
Cationic Group 4 metal alkyl compounds containing aryloxide ligation: synthesis, structure, reactivity and polymerization studies, 148
Imido complexes of manganese, 36
The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, 8
The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, 71
- Aluminum
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand [P₂N₂] ([P₂N₂] = [PhP(CH₂-SiMe₂NSiMe₂CH₂)₂PPh]), 63
- Amide
Imido complexes of manganese, 36
New titanium imido complexes containing piperazine-based diamido–diamine ligands, 114
- ansa*-Metallocene
Synthesis and structural characterization of uranium *ansa*-metallocene complexes containing organoimido functional groups; electronic effects of ancillary ligands, 14
- Aryloxide
Ancillary aryloxide ligands in ethylene polymerization catalyst precursors, 185
- Atom abstraction
(SiOx)₃Nb(η²-N,C-4-picoline) as a source of (siOx)₃Nb: structure of [(siOx)₃Nb]₂(μ-η²,η²-C₆H₆) (siOx = 'Bu₃SiO), 194
- Benzene
(SiOx)₃Nb(η²-N,C-4-picoline) as a source of (siOx)₃Nb: structure of [(siOx)₃Nb]₂(μ-η²,η²-C₆H₆) (siOx = 'Bu₃SiO), 194
- Benzyl cation
Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, 127
- N-tert*-Butylanilide
Dimolybdenum–μ-cyanide complexes supported by *N-tert*-butylanilide ligation: in pursuit of cyanide reductive cleavage, 24
- Cationic complexes
Stoichiometric olefin insertion into the Ti–C bond of four-coordinate cationic bis(phenolate) titanium aryl and benzyl complexes, 88
- C–H
Synthesis of a highly strained uranacycle: molecular structures of organometallic products arising from reduction, oxidation and protonolysis, 174
- C–H activation
C–H activation of pendant alkoxides by tungsten imide complexes, 104
Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, 2
- Cyano
Dimolybdenum–μ-cyanide complexes supported by *N-tert*-butylanilide ligation: in pursuit of cyanide reductive cleavage, 24
- Cyclopentadienyl
Ancillary aryloxide ligands in ethylene polymerization catalyst precursors, 185
- Density functional theory
The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, 204
- Diamido
Synthesis of Group 4 [(RN-*o*-C₆H₄)₂O]²⁻ complexes where R is SiMe₃ or 0.5 Me₂SiCH₂CH₂SiMe₂, 163
- Diazaphosphametallidines
P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn-synthesis and structural investigations, 138
- Dinuclear complexes
Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, 2
- Electrochemistry
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Ethene polymerization
Ancillary aryloxide ligands in ethylene polymerization catalyst precursors, 185
- Ethylene polymerization
Transition metal imido catalysts for ethylene polymerisation, 78
Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, 127
The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, 204
- Gallium
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand [P₂N₂] ([P₂N₂] = [PhP(CH₂SiMe₂NSiMe₂CH₂)₂PPh]), 63
- Hafnium
Synthesis of Group 4 [(RN-*o*-C₆H₄)₂O]²⁻ complexes where R is SiMe₃ or 0.5 Me₂SiCH₂CH₂SiMe₂, 163

- The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, 71
- Heterocubane**
The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, 71
- Hydride**
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand $[P_2N_2]$ ($[P_2N_2] = [PhP(CH_2SiMe_2NSiMe_2CH_2)_2PPh]$), 63
The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, 8
- Hydrogenolysis**
The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, 8
- Hydrotris(pyrazolyl)borate ligands**
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Imide**
C–H activation of pendant alkoxides by tungsten imide complexes, 104
New titanium imido complexes containing piperazine-based diamido–diamine ligands, 114
- Imido**
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
Imido complexes of manganese, 36
The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, 8
- Imido complexes**
Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, 2
Transition metal imido catalysts for ethylene polymerisation, 78
- Imines**
Reversible C–H activation of an *N*-methyl group in a bis-Zr(IV) diazametallacycle: generation of an end-on, side-on coordinated bridging alkylideneamido ligand, 2
- Indium**
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand $[P_2N_2]$ ($[P_2N_2] = [PhP(CH_2SiMe_2NSiMe_2CH_2)_2PPh]$), 63
- Iron carbene**
The metal–carbon multiple bond in iron(I)– and iron(II)–dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, 45
- Iron–carbon monoxide**
The metal–carbon multiple bond in iron(I)– and iron(II)–dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, 45
- Iron macrocycle redox**
The metal–carbon multiple bond in iron(I)– and iron(II)–dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, 45
- Linked amido–cyclopentadienyl ligand**
Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, 127
- Macrocyclic**
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand $[P_2N_2]$ ($[P_2N_2] = [PhP(CH_2SiMe_2NSiMe_2CH_2)_2PPh]$), 63
- Manganese**
Imido complexes of manganese, 36
- Metallacycle**
Synthesis of a highly strained uranacycle: molecular structures of organometallic products arising from reduction, oxidation and protonolysis, 174
- Metalla-macrocyclic**
The metal–carbon multiple bond in iron(I)– and iron(II)–dibenzotetramethyltetra[14]azaannulene: carbene, carbonyl, and isocyanide derivatives, 45
- Metallaspirobicycloheptanes**
P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn–synthesis and structural investigations, 138
- Metal–metal bonds**
Synthesis and characterization of Group 13 hydrides and metal–metal bonded dimers stabilized by the macrocyclic bis(amidophosphine) ligand $[P_2N_2]$ ($[P_2N_2] = [PhP(CH_2SiMe_2NSiMe_2CH_2)_2PPh]$), 63
- Molecular mechanism**
The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, 204
- Molybdenum**
Dimolybdenum– μ -cyanide complexes supported by *N*-*tert*-butylanilide ligation: in pursuit of cyanide reductive cleavage, 24
- Nickel**
The influence of electronic and steric factors on chain branching in ethylene polymerization by Brookhart-type Ni(II) diimine catalysts: a combined density functional theory and molecular mechanics study, 204
- Niobium**
(Silox)₃Nb(η^2 -N,C-4-picoline) as a source of (silox)₃Nb: structure of [(silox)₃Nb]₂(μ - η^2 , η^2 -C₆H₆) (silox = 'Bu₃SiO), 194
- Nitrido**
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Olefin insertion**
Stoichiometric olefin insertion into the Ti–C bond of four-coordinate cationic bis(phenolate) titanium aryl and benzyl complexes, 88
- Organoimido**
Synthesis and structural characterization of uranium *ansa*-metallocene complexes containing organoimido functional groups; electronic effects of ancillary ligands, 14
- Osmium**
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Oxo**
Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Phenoxides**
Cationic Group 4 metal alkyl compounds containing aryloxide ligation: synthesis, structure, reactivity and polymerization studies, 148

- Picoline**
 (Silox)₃Nb(η²-N,C-4-picoline) as a source of (silox)₃Nb: structure of [(silox)₃Nb]₂(μ-η²,η²-C₆H₆) (silox = 'Bu₃SiO), 194
- Polydentate ligand**
 New titanium imido complexes containing piperazine-based di-amido–diamine ligands, 114
- Polymerisation of hexene**
 Synthesis of Group 4 [(RN-*o*-C₆H₄)₂O]²⁻ complexes where R is SiMe₃ or 0.5 Me₂SiCH₂CH₂SiMe₂, 163
- Protonolysis**
 Synthesis of a highly strained uranacycle: molecular structures of organometallic products arising from reduction, oxidation and protonolysis, 174
- Redox reactions**
 Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Reduction**
 (Silox)₃Nb(η²-N,C-4-picoline) as a source of (silox)₃Nb: structure of [(silox)₃Nb]₂(μ-η²,η²-C₆H₆) (silox = 'Bu₃SiO), 194
- Rhenium**
 Electrochemical and reactivity comparisons among isoelectronic oxo, imido, and nitrido complexes of rhenium and osmium, 96
- Tantalum**
 (Silox)₃Nb(η²-N,C-4-picoline) as a source of (silox)₃Nb: structure of [(silox)₃Nb]₂(μ-η²,η²-C₆H₆) (silox = 'Bu₃SiO), 194
- Thioether ligand**
 Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, 127
- Thiolate**
 Imido complexes of manganese, 36
- Tin**
 P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn-synthesis and structural investigations, 138
- Titanium**
 Ancillary aryloxide ligands in ethylene polymerization catalyst precursors, 185
 Cationic Group 4 metal alkyl compounds containing aryloxide ligation: synthesis, structure, reactivity and polymerization studies, 148
 Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, 127
 New titanium imido complexes containing piperazine-based di-amido–diamine ligands, 114
- P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn-synthesis and structural investigations, 138
 Synthesis of Group 4 [(RN-*o*-C₆H₄)₂O]²⁻ complexes where R is SiMe₃ or 0.5 Me₂SiCH₂CH₂SiMe₂, 163
- Titanium alkyls**
 Stoichiometric olefin insertion into the Ti–C bond of four-coordinate cationic bis(phenolate) titanium aryl and benzyl complexes, 88
- Transition metals**
 Transition metal imido catalysts for ethylene polymerisation, 78
- Tripodal amide**
 The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, 71
- Tungsten**
 C–H activation of pendant alkoxides by tungsten imide complexes, 104
 The synthesis of tungsten imido hydride complexes by the hydrogenolysis of dialkyl complexes, 8
- Uranium**
 Synthesis and structural characterization of uranium *ansa*-metallocene complexes containing organoimido functional groups; electronic effects of ancillary ligands, 14
- X-ray**
 P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn-synthesis and structural investigations, 138
- X-ray diffraction**
 New titanium imido complexes containing piperazine-based di-amido–diamine ligands, 114
- Zirconium**
 Cationic Group 4 metal alkyl compounds containing aryloxide ligation: synthesis, structure, reactivity and polymerization studies, 148
 Complexes of titanium and zirconium containing a tridentate linked amido–cyclopentadienyl ligand with a soft donor group: synthesis, structure, and ethylene polymerization catalysis, 127
 P-functionally substituted aminoiminophosphoranate chelates of Ti, Zr, and Sn-synthesis and structural investigations, 138
 Synthesis of Group 4 [(RN-*o*-C₆H₄)₂O]²⁻ complexes where R is SiMe₃ or 0.5 Me₂SiCH₂CH₂SiMe₂, 163
 The tendency of tripodal amidozirconium and hafnium complexes to form hexacoordinate structures: alkali metal halide cages versus solvent adducts, 71
- Zwitterionic polymerization**
 Cationic Group 4 metal alkyl compounds containing aryloxide ligation: synthesis, structure, reactivity and polymerization studies, 148