

Subject Index of Volume 595

Alkenyl ligand

Reactions of $[\{\mu\text{-}\eta^2\text{-C(R}^1\text{)=CHR}^2\}(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ with $(\mu\text{-S}_2)\text{-Fe}_2(\text{CO})_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-C(R}^1\text{)=CHR}^2](\mu\text{-RS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})$ (R = alkyl, phenyl or acyl), 158

Alkoxides

Monomeric organoaluminium complexes $\text{RAl(OR}^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21

Alkoxysilanes

Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Alkylidene

Binuclear hydrido platinum(II) complexes: syntheses of $\{\{\text{Pt(P-P)}\}_2(\mu\text{-CHCH}_2\text{R})(\mu\text{-H})[\text{BF}_4]\}$ (R = C_6H_5 , H) and easy cleavage of a P–C bond in a chelating 1,4-bis(diphenyl)phosphinobutane (P–P), 224

Alkyne

Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2 -alkyne carbonyl complexes, 70

Alkynyl complexes

Selectivity in alkynylation: the reaction between $\text{Ru}_2(\text{LL})_4\text{Cl}$ and $\text{Me}_3\text{SnC}\equiv\text{CR}$ (LL = 2-anilinopyridine, 2-chloro- and 2-bromo-oxypyridine), 20

Alpha-lithiated alkoxysilanes

Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Alpha-lithiation

Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Alpha-silyl carbanions

Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Aluminium

Monomeric organoaluminium complexes $\text{RAl(OR}^*)_2$ and $\text{R}_2\text{-AlOR}^*$ with an optically active amino alkoxide ligand, 21

Amines

Monomeric organoaluminium complexes $\text{RAl(OR}^*)_2$ and $\text{R}_2\text{-AlOR}^*$ with an optically active amino alkoxide ligand, 21

Aminocarbene complexes

Synthese und Struktur von $\{\{N\text{-[Bis(diisopropylamino)phosphanyl]amino(phenyl)carben}\} \text{dicarbonyl}(\eta^5\text{-methylcyclopentadienyl)mangan(0)}\}$ und versuchte Darstellung von $\{\{2\text{-Bis(trimethylsilyl)methyl-3-phenyl-2H-azaphosphiren-}\kappa\text{P}\} \text{dicarbonyl}(\eta^5\text{-methylcyclopentadienyl)mangan(0)}\}$, 12

Aminophosphines

Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdocyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166

Aromatic bridge

Convenient routes to mono- and dinuclear $(\text{C}_5\text{Me}_5)\text{iron(II)}$ complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81

Aromatic ditellurides

The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232

Aryloxy-zirconocene complex

Dehydrocoupling polymerization of arylsilanes with chloro(aryloxy)bis(cyclopentadienyl)zirconium complex catalysts, 1

Arylsilanes

Dehydrocoupling polymerization of arylsilanes with chloro(aryloxy)bis(cyclopentadienyl)zirconium complex catalysts, 1

Binuclear complexes

Binuclear hydrido platinum(II) complexes: syntheses of $\{\{\text{Pt(P-P)}\}_2(\mu\text{-CHCH}_2\text{R})(\mu\text{-H})[\text{BF}_4]\}$ (R = C_6H_5 , H) and easy cleavage of a P–C bond in a chelating 1,4-bis(diphenyl)phosphinobutane (P–P), 224

Bis(trimesitylgermyl)-carbodiimide

Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153

Bis(trimesitylgermylcarbodiimido)germylene

Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153

Borane adduct

P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[\text{Cp}_2\text{Zr(Me)(cyclo-1-O-2-PH(Tipp)-C}_6\text{H}_{10})]$ (Tipp = 2,4,6- $\text{Pr}_i^3\text{C}_6\text{H}_2$), $[\text{Cp}_2\text{Zr(Me)(cyclo-1-O-2-PH(Tipp)\{Mo(CO)}_3\text{-C}_6\text{H}_{10})}]$ and $[\text{Cp}_2\text{Zr(BH}_4\text{)(cyclo-1-O-2-PH(Mes)(BH}_3\text{-C}_6\text{H}_{10})}]$ (Mes = 2,4,6- $\text{Me}_3\text{C}_6\text{H}_2$), and molecular structure of $(\text{R}_C, \text{R}_C, \text{R}_P/\text{S}_C, \text{S}_P)\text{-[Cp}_2\text{Zr(Me)(cyclo-1-O-2-PH(Tipp)-C}_6\text{H}_{10})]$, 126

Borasiloxanes

cyclo-Boratrilsiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-BrC}_6\text{H}_4\text{BO})_2(\text{Ph}_2\text{SiO})_2$, 241

Boron

cyclo-Boratrilsiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-BrC}_6\text{H}_4\text{BO})_2(\text{Ph}_2\text{SiO})_2$, 241

Synthesis of dioxazaborocenes from *N,N'*-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215

Bridging ligand

Synthesis and spectroscopic characterisation of compounds with formula $\{\{\text{Fe}_2\text{S}_2(\text{CO})_5\}\text{Ph}_2\text{P(CH}_2\text{)}_n\text{PPh}_2\{\text{Fe(CO)}_4\}\}$ and $\{\{\text{Fe}_2\text{S}_2(\text{CO})_5\}_2\text{Ph}_2\text{P(CH}_2\text{)}_n\text{PPh}_2\}$ ($n = 3, 4$ and 6), 191

Bridging-sulfur ligand

Reactions of $[\{\mu\text{-}\eta^2\text{-C(R}^1\text{)=CHR}^2\}(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ with $(\mu\text{-S}_2)\text{Fe}_2(\text{CO})_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-C(R}^1\text{)=CHR}^2](\mu\text{-RS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})$ (R = alkyl, phenyl or acyl), 158

Carbazole

Electron-transfer processes in the reaction of K^- , K^+ (15-crown-5)₂ with carbazole: unexpected formation of metallic potassium, 66

Carbene

Convenient routes to mono- and dinuclear $(\text{C}_5\text{Me}_5)\text{iron(II)}$ complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81

Carbon-carbon bond formation

N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186

- Carbon disulfide
Reactivity of $[\text{Fe}_2(\text{CO})_6(\mu\text{-CO})(\mu\text{-dppm})]$ towards unsaturated sulfur-containing compounds, SO_2 , CS_2 and ArNCS : facile carbon–sulfur bond cleavage at the diiron centre, 134
- Carbon–fluorine bond activation
Regioselective C–F and C–H bond activation: synthesis and structure of an asymmetric rhodium complex comprising a $\eta^2:\eta^1:\eta^1$ -cyclopentadienyl-bis(phosphine) ligand, 292
- Carbon–hydrogen bond activation
Regioselective C–F and C–H bond activation: synthesis and structure of an asymmetric rhodium complex comprising a $\eta^2:\eta^1:\eta^1$ -cyclopentadienyl-bis(phosphine) ligand, 292
- Carbonylation
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
- Chirality
Monomeric organoaluminium complexes $\text{RAl}(\text{OR}^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21
- Cobalt
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2 -alkyne carbonyl complexes, 70
- Complex
Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}_2(\mu\text{-S})(\mu\text{-Te})(\mu\text{-SPh})_2]$, 140
- Complexes
Reactions of $[\{\mu\text{-}\eta^2\text{-C}(\text{R}^1)=\text{CHR}^2\}(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ with $(\mu\text{-S}_2)\text{Fe}_2(\text{CO})_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-C}(\text{R}^1)=\text{CHR}^2](\mu\text{-RS})[\text{Fe}_2(\text{CO})_6](\mu_4\text{-S})$ (R = alkyl, phenyl or acyl), 158
- Coupling reaction
Selectivity in alkynylation: the reaction between $\text{Ru}_2(\text{LL})_4\text{Cl}$ and $\text{Me}_3\text{SnC}\equiv\text{CR}$ (LL = 2-anilino-pyridine, 2-chloro- and 2-bromo-oxy-pyridine), 300
- Cross coupling
N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186
- Cryogenic reactions
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
- Crystal structure
Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[\text{Ph}_2\text{Sn}\{2\text{-}(\text{Ph}_2\text{P})\text{C}_6\text{H}_4\text{S}\}_2]$ and $[\text{Me}_2\text{Sn}\{2\text{-}(\text{Ph}_2\text{PO})\text{-6-(Me}_3\text{Si})\text{C}_6\text{H}_3\text{S}\}_2]$, 59
Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[\text{PtMe}_3\text{-}(\text{bpy})(\text{PPh}_3)][\text{O}_3\text{SCF}_3]$ (bpy = 2,2'-bipyridine), 296
- Crystal structures
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $\text{Tp}^{\text{Ph,Me}}\text{RhL}_1\text{L}_2$, 178
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2 -alkyne carbonyl complexes, 70
- [2 + 2]-Cycloaddition
[2 + 2]-Cycloaddukte PH-funktioneller Phosphonium-Komplexe mit Alkylisothiocyanaten: Darstellung von $\text{C}_5\text{R}_5(\text{OC})_2\text{C}=\text{O}$ type = "tsqb" $\text{M-P}(\text{H})(\text{t-Bu})\text{-C}(\text{=NR}')\text{-S}$ (R = H, Me; M = Mo, W; R' = Me, Et, *t*-Bu) und Reaktion unter Beanspruchung der PH-Funktion: Phosphonium-Übergangsmetallkomplexe, 37, 285
- Cyclometallated
New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[\text{Pt}\{4\text{-MeC}_6\text{H}_3\text{C}(\text{Me})=\text{NN}=\text{C}(\text{S})\text{NH}_2\}(\text{PPh}_3)]$, 199
- Cyclometallation
Cyclometallation of anthracen-9-ylmethylene-phenyl-amine by palladium(II) compounds, 54
- Dehydrocoupling polymerization
Dehydrocoupling polymerization of arylsilanes with chloro(aryloxy)bis(cyclopentadienyl)zirconium complex catalysts, 1
- Density functional calculations
Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248
- Dimeric carbonyl
Reactivity of $[\text{Fe}_2(\text{CO})_6(\mu\text{-CO})(\mu\text{-dppm})]$ towards unsaturated sulfur-containing compounds, SO_2 , CS_2 and ArNCS : facile carbon–sulfur bond cleavage at the diiron centre, 134
- Dimolybdacyclophanes
Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166
- Dinuclear
Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81
- Dioxazaborocine
Synthesis of dioxazaborocines from *N,N'*-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215
- Diphosphenes
Synthese und Struktur von $\{\{N\text{-}[\text{Bis}(\text{diisopropylamino})\text{-phosphanyl}]\text{amino}(\text{phenyl})\text{carben}\}\text{dicarbonyl}(\eta^5\text{-methylcyclopentadienyl})\text{mangan(0)}\}$ und versuchte Darstellung von $\{[2\text{-Bis}(\text{trimethylsilyl})\text{methyl-3-phenyl-2H-azaphosphiren-}\kappa\text{P}]\text{dicarbonyl}(\eta^5\text{-methylcyclopentadienyl})\text{mangan(0)}\}$, 12
- Diphosphine
Synthesis and spectroscopic characterisation of compounds with formula $[\{\text{Fe}_2\text{S}_2(\text{CO})_5\}\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\{\text{Fe}(\text{CO})_4\}]$ and $[\{\text{Fe}_2\text{S}_2(\text{CO})_5\}_2\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2]$ (n = 3, 4 and 6), 191
- Diruthenium
Selectivity in alkynylation: the reaction between $\text{Ru}_2(\text{LL})_4\text{Cl}$ and $\text{Me}_3\text{SnC}\equiv\text{CR}$ (LL = 2-anilino-pyridine, 2-chloro- and 2-bromo-oxy-pyridine), 300
- Disiloxane-bridged metallocene
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediy bridges, 147
- Doubly bridged metallocene
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanediy bridges, 147
- Electrochemistry
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
- Electron transfer
Electron-transfer processes in the reaction of K^- , K^+ (15-crown-5) $_2$ with carbazole: unexpected formation of metallic potassium, 66
- Ferrocene
Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93
- Ferrocenyl
Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2 -alkyne carbonyl complexes, 70

Fluorene

Synthesis, electrochemistry and structural characterization of new dimeric cobalt η^2 -alkyne carbonyl complexes, 70

Fragmentation

Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Hafnium

Synthesis and characterization of Group 4 metallocene complexes with two disiloxanedyl bridges, 147

2*H*-azaphosphirene complexes

Synthese und Struktur von $\{\{N$ -[Bis(diisopropylamino)-phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 -methylcyclopentadienyl)mangan(0)) und versuchte Darstellung von $\{[2$ -Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren- κP]dicarbonyl(η^5 -methylcyclopentadienyl)mangan(0))\}, 12

Heterodinuclear molybdenum phosphine complex

P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[Cp_2Zr(Me)(cyclo-1-O-2-PH(Tipp)-C_6H_{10})]$ (Tipp = 2,4,6- $Pr^i_3C_6H_2$), $[Cp_2Zr(Me)(cyclo-1-O-2-PH(Tipp)\{Mo(CO)_3\}-C_6H_{10})]$ and $[Cp_2Zr(BH_4)(cyclo-1-O-2-PH(Mes)(BH_3)-C_6H_{10})]$ (Mes = 2,4,6- $Me_3C_6H_2$), and molecular structure of $(R_C, R_C, R_P/S_C, S_C, S_P)$ - $[Cp_2Zr(Me)(cyclo-1-O-2-PH(Tipp)-C_6H_{10})]$, 126

Heterometallic

Reactions of $[Pt_2(\mu-S)_2(PPh_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[Pt_2Re(\mu_3-S)_2(PPh_3)_4(CO)_3]^+$, 276

High dilution

Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166

2*H*-1,3-Naphthoxazine

Synthesis of dioxazaborocines from *N,N'*-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215

Hydride

Binuclear hydrido platinum(II) complexes: syntheses of $\{[Pt(P-P)]_2(\mu-CHCH_2R)(\mu-H)]BF_4\}$ (R = C_6H_5 , H) and easy cleavage of a P–C bond in a chelating 1,4-bis(diphenyl)phosphinobutane (P–P), 224

Hydroformylation

Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93

Imido η^2 -iminoacyl, η^2 -acyl cyclopentadienyl complexes

Chemical behaviour of alkyl imido cyclopentadienyl niobium and tantalum(V) complexes in insertion processes. X-ray crystal structures of $[MCpCl(NAr)\{\eta^2-C(Me)=NAr\}]$ (Ar = 2,6- $Me_2C_6H_3$; M = Nb, Cp = $\eta^5-C_5H_4SiMe_3$; M = Ta, Cp = $\eta^5-C_5Me_5$) and $[Ta(\eta^5-C_5Me_5)Me(NAr)\{\eta^2-C(CH_2CMe_2Ph)=O\}]$ (Ar = 2,6- $Me_2C_6H_3$), 36

Infrared spectroscopy

Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248

Intramolecular proton transfer

Electron-transfer processes in the reaction of K^+ , K^+ (15-crown-5)₂ with carbazole: unexpected formation of metallic potassium, 66

Iron

Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81

Reactions of $[\{\mu-\eta^2-C(R^1)=CHR^2\}(\mu-CO)Fe_2(CO)_6]^-$ with $(\mu-S_2)Fe_2(CO)_6$: synthesis and structures of $[\mu-\eta^2-C(R^1)=CHR^2](\mu-RS)[Fe_2(CO)_6]_2(\mu_4-S)$ (R = alkyl, phenyl or acyl), 158

Reactivity of $[Fe_2(CO)_6(\mu-CO)(\mu-dppm)]$ towards unsaturated sulfur-containing compounds, SO_2 , CS_2 and $ArNCS$: facile carbon–sulfur bond cleavage at the diiron centre, 134

Synthesis and spectroscopic characterisation of compounds with formula $\{[Fe_2S_2(CO)_5]Ph_2P(CH_2)_nPPH_2\{Fe(CO)_4\}]$ and $\{[Fe_2S_2(CO)_5]_2Ph_2P(CH_2)_nPPH_2\}$ ($n = 3, 4$ and 6), 191

The thermal rearrangement of $-[Me_2Si-\eta^5-C_5H_4(CO)Fe(CO)_2Fe(CO)-\eta^5-C_5H_4SiMe_2]-$ in the presence of phosphorus ligands, 268

Isocyanide complexes

Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114

Isomer

Synthesis and spectroscopic characterisation of compounds with formula $\{[Fe_2S_2(CO)_5]Ph_2P(CH_2)_nPPH_2\{Fe(CO)_4\}]$ and $\{[Fe_2S_2(CO)_5]_2Ph_2P(CH_2)_nPPH_2\}$ ($n = 3, 4$ and 6), 191

Lithium

Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

Macrocyclization

Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166

Manganese

Synthese und Struktur von $\{\{N$ -[Bis(diisopropylamino)-phosphanyl]amino(phenyl)carben}dicarbonyl(η^5 -methylcyclopentadienyl)mangan(0)) und versuchte Darstellung von $\{[2$ -Bis(trimethylsilyl)methyl-3-phenyl-2*H*-azaphosphiren- κP]dicarbonyl(η^5 -methylcyclopentadienyl)mangan(0))\}, 12

Matrix isolation

Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248

Mechanism

Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Metallocene complexes

Synthesis and characterization of Group 4 metallocene complexes with two disiloxanedyl bridges, 147

Metalloligand

Reactions of $[Pt_2(\mu-S)_2(PPh_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[Pt_2Re(\mu_3-S)_2(PPh_3)_4(CO)_3]^+$, 276

Metathesis

The thermal rearrangement of $-[Me_2Si-\eta^5-C_5H_4(CO)Fe(CO)_2Fe(CO)-\eta^5-C_5H_4SiMe_2]-$ in the presence of phosphorus ligands, 268

Molybdenum

Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5-C_5H_5)_2Mo_2(\mu-S)(\mu-Te)(\mu-SPh)_2]$, 140

Molybdenum complexes

Preparation, properties and reactions of metal-containing heterocycles: Part C: tetraazatetraphosphadimolybdacyclophanes: synthesis, isolation, characterization, and X-ray crystal structures, 166

Mössbauer

Synthesis and spectroscopic characterisation of compounds with formula $[\{\text{Fe}_2\text{S}_2(\text{CO})_5\}\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\{\text{Fe}(\text{CO})_4\}]$ and $[\{\text{Fe}_2\text{S}_2(\text{CO})_5\}_2\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2]$ ($n = 3, 4$ and 6), 191

N-arylsulfonyl aldimines

Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31

N-Heterocyclic carbenes

N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186

Niobium

Chemical behaviour of alkyl imido cyclopentadienyl niobium and tantalum(V) complexes in insertion processes. X-ray crystal structures of $[\text{MCpCl}(\text{NAr})\{\eta^2\text{-C}(\text{Me})=\text{NAr}\}]$ ($\text{Ar} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$; $\text{M} = \text{Nb}$, $\text{Cp} = \eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$; $\text{M} = \text{Ta}$, $\text{Cp} = \eta^5\text{-C}_5\text{Me}_5$) and $[\text{Ta}(\eta^5\text{-C}_5\text{Me}_5)\text{Me}(\text{NAr})\{\eta^2\text{-C}(\text{CH}_2\text{CMe}_2\text{Ph})=\text{O}\}]$ ($\text{Ar} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), 36

NMR spectroscopy

Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $\text{Tp}^{\text{Ph,Me}}\text{RhL}_2$, 178

The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232

¹⁵N-NMR

Synthesis and structural analysis of palladium(II) pyridinylpyrazole complexes by ¹H-, ¹³C-, ¹⁵N-NMR and X-ray diffraction. Comparison of binuclear methylpalladium, chloromethylpalladium, and dichloropalladium complexes by ¹⁵N-NMR, 208

Nuclear magnetic resonance

Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93

Nucleophilicity

Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Organotin complexes

Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[\text{Ph}_2\text{Sn}\{2\text{-(Ph}_2\text{P)C}_6\text{H}_4\text{S}\}_2]$ and $[\text{Me}_2\text{Sn}\{2\text{-(Ph}_2\text{PO)-6-(Me}_3\text{Si)C}_6\text{H}_3\text{S}\}_2]$, 59

Oxypyridine

Selectivity in alkylation: the reaction between $\text{Ru}_2(\text{LL})_4\text{Cl}$ and $\text{Me}_3\text{SnC}\equiv\text{CR}$ ($\text{LL} = 2\text{-anilinopyridine}$, 2-chloro- and $2\text{-bromo-oxypyridine}$), 300

Palladium

Cyclometallation of anthracen-9-ylmethylene-phenyl-amine by palladium(II) compounds, 54

N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186

Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[\text{PtMe}_3\text{(bpy)(PPh}_3)][\text{O}_3\text{SCF}_3]$ ($\text{bpy} = 2,2'\text{-bipyridine}$), 296

Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93

Synthesis and structural analysis of palladium(II) pyridinylpyrazole complexes by ¹H-, ¹³C-, ¹⁵N-NMR and X-ray diffraction. Comparison of binuclear methylpalladium, chloromethylpalladium, and dichloropalladium complexes by ¹⁵N-NMR, 208

Palladium complex

The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232

P–C cleavage

Binuclear hydrido platinum(II) complexes: syntheses of $\{[\text{Pt}(\text{P-P})]_2(\mu\text{-CHCH}_2\text{R})(\mu\text{-H})[\text{BF}_4]\}$ ($\text{R} = \text{C}_6\text{H}_5$, H) and easy cleavage of a P–C bond in a chelating 1,4-bis(diphenyl)phosphinobutane (P–P), 224

Pentacoordinated intermediate

Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phosphabicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

P–H functionalised phosphinoalkoxide

P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[\text{Cp}_2\text{Zr}(\text{Me})(\text{cyclo-1-O-2-PH}(\text{Tipp})\text{-C}_6\text{H}_{10})]$ ($\text{Tipp} = 2,4,6\text{-Pr}_3\text{C}_6\text{H}_2$), $[\text{Cp}_2\text{Zr}(\text{Me})(\text{cyclo-1-O-2-PH}(\text{Tipp})\{\text{Mo}(\text{CO})_5\}\text{-C}_6\text{H}_{10})]$ and $[\text{Cp}_2\text{Zr}(\text{BH}_4)(\text{cyclo-1-O-2-PH}(\text{Mes})(\text{BH}_3)\text{-C}_6\text{H}_{10})]$ ($\text{Mes} = 2,4,6\text{-Me}_3\text{C}_6\text{H}_2$), and molecular structure of $(\text{R}_{\text{C}}\text{R}_{\text{C}}\text{R}_{\text{P}}/\text{S}_{\text{C}}\text{S}_{\text{C}}\text{S}_{\text{P}})\text{-}[\text{Cp}_2\text{Zr}(\text{Me})(\text{cyclo-1-O-2-PH}(\text{Tipp})\text{-C}_6\text{H}_{10})]$, 126

Phosphametallacycles

$[2 + 2]$ -Cycloaddukte PH-funktioneller Phosphenium-Komplexe mit Alkylisothiocyanaten: Darstellung von $\text{C}_5\text{R}_5(\text{OC})_2 < \text{ovl type} = \text{“tsqb”} > \text{M-P}(\text{H})(t\text{-Bu})\text{-C}(\text{=NR})\text{-S}$ ($\text{R} = \text{H}$, Me ; $\text{M} = \text{Mo}$, W ; $\text{R}' = \text{Me}$, Et , $t\text{-Bu}$) und Reaktion unter Beanspruchung der PH-Funktion: Phosphenium-Übergangsmetallkomplexe, 37, 285

Phosphenium complexes

$[2 + 2]$ -Cycloaddukte PH-funktioneller Phosphenium-Komplexe mit Alkylisothiocyanaten: Darstellung von $\text{C}_5\text{R}_5(\text{OC})_2 < \text{ovl type} = \text{“tsqb”} > \text{M-P}(\text{H})(t\text{-Bu})\text{-C}(\text{=NR})\text{-S}$ ($\text{R} = \text{H}$, Me ; $\text{M} = \text{Mo}$, W ; $\text{R}' = \text{Me}$, Et , $t\text{-Bu}$) und Reaktion unter Beanspruchung der PH-Funktion: Phosphenium-Übergangsmetallkomplexe, 37, 285

Phosphide

Binuclear hydrido platinum(II) complexes: syntheses of $\{[\text{Pt}(\text{P-P})]_2(\mu\text{-CHCH}_2\text{R})(\mu\text{-H})[\text{BF}_4]\}$ ($\text{R} = \text{C}_6\text{H}_5$, H) and easy cleavage of a P–C bond in a chelating 1,4-bis(diphenyl)phosphinobutane (P–P), 224

Phosphine

Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93

Phosphine complexes

Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[\text{PtMe}_3(\text{bpy})(\text{PPh}_3)][\text{O}_3\text{SCF}_3]$ ($\text{bpy} = 2,2'\text{-bipyridine}$), 296

Phosphinothiolate complexes

Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[\text{Ph}_2\text{Sn}\{2\text{-(Ph}_2\text{P)C}_6\text{H}_4\text{S}\}_2]$ and $[\text{Me}_2\text{Sn}\{2\text{-(Ph}_2\text{PO)-6-(Me}_3\text{Si)C}_6\text{H}_3\text{S}\}_2]$, 59

Phospholes

Convenient route for the preparation of unsymmetrical phospholes via zirconacyclopentadienes, 261

Phosphorus

The thermal rearrangement of $-\text{[Me}_2\text{Si-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{-Fe}(\text{CO})\text{-}\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{-}]$ in the presence of phosphorus ligands, 268

Phosphorus heterocycle

Convenient route for the preparation of unsymmetrical phospholes via zirconacyclopentadienes, 261

Photolysis

Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phospha-bicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Reactivity of $[\text{Fe}_2(\text{CO})_6(\mu\text{-CO})(\mu\text{-dppm})]$ towards unsaturated sulfur-containing compounds, SO_2 , CS_2 and ArNCS : facile carbon–sulfur bond cleavage at the diiron centre, 134

Platinum

Binuclear hydrido platinum(II) complexes: syntheses of $[\{\text{Pt}(\text{P}-\text{P})\}_2(\mu\text{-CHCH}_2\text{R})(\mu\text{-H})][\text{BF}_4]$ ($\text{R} = \text{C}_6\text{H}_5$, H) and easy cleavage of a P–C bond in a chelating 1,4-bis(diphenyl)phosphinobutane (P–P), 224

New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[\text{Pt}\{4\text{-MeC}_6\text{H}_3\text{C}(\text{Me})=\text{NN}=\text{C}(\text{S})\text{NH}_2\}(\text{PPh}_3)]$, 199

Organopalladium(IV) complexes containing phosphine ligands, and the structure of the platinum(IV) complex $[\text{PtMe}_3(\text{bpy})(\text{PPh}_3)][\text{O}_3\text{SCF}_3]$ ($\text{bpy} = 2,2'$ -bipyridine), 296

Platinum complexes of heteroannularly bridged heterobidentate ferrocenyl diphosphine ligands: their molecular structure and their use in catalytic carbonylation reactions, 93

Reactions of $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[\text{Pt}_2\text{Re}(\mu_3\text{-S})_2(\text{PPh}_3)_4(\text{CO})_3]^+$, 276

Platinum complex

The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)-platinum or -palladium, 232

Polycarbodiimido-germylene

Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermyl-cyanamide and trimesitylgermylcarbodiimide, 153

Porphyrins

Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114

Potassium anions

Electron-transfer processes in the reaction of K^- , K^+ (15-crown-5)₂ with carbazole: unexpected formation of metallic potassium, 66

Pyrazolylborate complexes

Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $\text{Tp}^{\text{Ph,Me}}\text{RhL}_1\text{L}_2$, 178

Pyridinylpyrazole

Synthesis and structural analysis of palladium(II) pyridinylpyrazole complexes by ^1H -, ^{13}C -, ^{15}N -NMR and X-ray diffraction. Comparison of binuclear methylpalladium, chloromethylpalladium, and dichloropalladium complexes by ^{15}N -NMR, 208

Rearrangement

The thermal rearrangement of $-\text{[Me}_2\text{Si-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{Fe}(\text{CO})-\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{-]}$ in the presence of phosphorus ligands, 268

Reversibility

Evidence for the involvement of pentacoordinate P-intermediates during the UV light-mediated fragmentation of a 2-phospha-bicyclo[2.2.2]octene 2-oxide in the presence of O-nucleophiles, 109

Rhenium

Reactions of $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[\text{Pt}_2\text{Re}(\mu_3\text{-S})_2(\text{PPh}_3)_4(\text{CO})_3]^+$, 276

Rhodium

Regioselective C–F and C–H bond activation: synthesis and structure of an asymmetric rhodium complex comprising a $\eta^5:\eta^1:\eta^1$ -cyclopentadienyl-bis(phosphine) ligand, 292

Synthesis and structure of Rh(I) tris(pyrazolyl)borate complexes of the type $\text{Tp}^{\text{Ph,Me}}\text{RhL}_1\text{L}_2$, 178

Rhodium-catalyzed phenylation

Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31

Ring compounds

cyclo-Boratrissiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-BrC}_6\text{H}_4\text{BO})_2(\text{Ph}_2\text{SiO})_2$, 241

Ruthenium

Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114

Schiff bases

New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[\text{Pt}\{4\text{-MeC}_6\text{H}_3\text{C}(\text{Me})=\text{NN}=\text{C}(\text{S})\text{NH}_2\}(\text{PPh}_3)]$, 199

Silanones

Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248

Silicon

cyclo-Boratrissiloxane and *cyclo*-diboratetrasiloxane derivatives and their reactions with amines: crystal and molecular structure of $(p\text{-BrC}_6\text{H}_4\text{BO})_2(\text{Ph}_2\text{SiO})_2$, 241

Regiospecific synthesis of alpha-lithiated alkoxysilanes, 87

The thermal rearrangement of $-\text{[Me}_2\text{Si-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{Fe}(\text{CO})-\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{-]}$ in the presence of phosphorus ligands, 268

Silicon atoms

Matrix isolation infrared and density functional theoretical studies of cryogenic reactions of silicon atoms with dimethyl ether and methanol: new route to generation and stabilization of transient silanones, 248

Sodium tetraphenylborate

Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31

Stable germylene

Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermyl-cyanamide and trimesitylgermylcarbodiimide, 153

Structure determinations

Synthese und Struktur von $\{\{N\text{-[Bis(diisopropylamino)phosphanyl]amino(phenyl)carben}\} \text{dicarbonyl}(\eta^5\text{-methylcyclopentadienyl)mangan(0)}\}$ und versuchte Darstellung von $\{\{2\text{-Bis(trimethylsilyl)methyl-3-phenyl-2H-azaphosphiren-}\kappa\text{P}\} \text{dicarbonyl}(\eta^5\text{-methylcyclopentadienyl)mangan(0)}\}$, 12

Structures

Reactions of $[\{\mu\text{-}\eta^2\text{-C}(\text{R}^1)=\text{CHR}^2\}(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ with $(\mu\text{-S}_2)\text{Fe}_2(\text{CO})_6$: synthesis and structures of $[\{\mu\text{-}\eta^2\text{-C}(\text{R}^1)=\text{CHR}^2\}(\mu\text{-RS})\text{[Fe}_2(\text{CO})_6\text{]}_2(\mu_4\text{-S})]$ ($\text{R} = \text{alkyl, phenyl or acyl}$), 158

Substitution

The thermal rearrangement of $-\text{[Me}_2\text{Si-}\eta^5\text{-C}_5\text{H}_4(\text{CO})\text{Fe}(\text{CO})_2\text{Fe}(\text{CO})-\eta^5\text{-C}_5\text{H}_4\text{SiMe}_2\text{-]}$ in the presence of phosphorus ligands, 268

Sulfide

Reactions of $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[\text{Pt}_2\text{Re}(\mu_3\text{-S})_2(\text{PPh}_3)_4(\text{CO})_3]^+$, 276

Sulfur

Synthesis and spectroscopic characterisation of compounds with formula $\{\{\text{Fe}_2\text{S}_2(\text{CO})_5\}\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\{\text{Fe}(\text{CO})_4\}\}$ and $\{\{\text{Fe}_2\text{S}_2(\text{CO})_5\}\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}$ ($n = 3, 4$ and 6), 191

- Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}_2(\mu\text{-S})(\mu\text{-Te})(\mu\text{-SPH})_2]$, 140
- Sulfur dioxide
Reactivity of $[\text{Fe}_2(\text{CO})_6(\mu\text{-CO})(\mu\text{-dppm})]$ towards unsaturated sulfur-containing compounds, SO_2 , CS_2 and ArNCS : facile carbon–sulfur bond cleavage at the diiron centre, 134
- Suzuki reaction
N-Heterocyclic carbenes: Part 26. *N*-Heterocyclic carbene complexes of palladium(0): synthesis and application in the Suzuki cross-coupling reaction, 186
- Synthesis
Reactions of $[\{\mu\text{-}\eta^2\text{-C}(\text{R}^1)=\text{CHR}^2\}(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ with $(\mu\text{-S}_2)\text{Fe}_2(\text{CO})_6$: synthesis and structures of $[\mu\text{-}\eta^2\text{-C}(\text{R}^1)=\text{CHR}^2](\mu\text{-RS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})$ (R = alkyl, phenyl or acyl), 158
- Tantalum
Chemical behaviour of alkyl imido cyclopentadienyl niobium and tantalum(V) complexes in insertion processes. X-ray crystal structures of $[\text{MCpCl}(\text{NAr})\{\eta^2\text{-C}(\text{Me})=\text{NAr}\}]$ (Ar = 2,6- $\text{Me}_2\text{C}_6\text{H}_3$; M = Nb, Cp = $\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3$; M = Ta, Cp = $\eta^5\text{-C}_5\text{Me}_5$) and $[\text{Ta}(\eta^5\text{-C}_5\text{Me}_5)\text{Me}(\text{NAr})\{\eta^2\text{-C}(\text{CH}_2\text{CMe}_2\text{Ph})=\text{O}\}]$ (Ar = 2,6- $\text{Me}_2\text{C}_6\text{H}_3$), 36
- Tellurium
Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}_2(\mu\text{-S})(\mu\text{-Te})(\mu\text{-SPH})_2]$, 140
- Tertiary amine
Isocyanide ligation at ruthenium(II) complexes with chelating tertiary amine and porphyrin ligands. Structural and electrochemical studies, 114
- Thienylcopper
Tetrameric thienylcopper and pentanuclear thienylcuprate, 102
- Thienylcuprate
Tetrameric thienylcopper and pentanuclear thienylcuprate, 102
- Thiolato
Synthesis, optical nonlinearity and crystal structure of a novel mixed-chalcogen, thiolato-bridged complex $[(\eta^5\text{-C}_5\text{H}_5)_2\text{Mo}_2(\mu\text{-S})(\mu\text{-Te})(\mu\text{-SPH})_2]$, 140
- Thiosemicarbazone
New cyclometallated platinum(II) compounds with thiosemicarbazones: crystal and molecular structure of $[\text{Pt}\{4\text{-MeC}_6\text{H}_3\text{C}(\text{Me})=\text{NN}=\text{C}(\text{S})\text{NH}_2\}(\text{PPh}_3)]$, 199
- Tin complexes
Diorganotin(IV) derivatives of arenephosphinothiol ligands. The crystal structure of $[\text{Ph}_2\text{Sn}\{2\text{-(Ph}_2\text{P)C}_6\text{H}_4\text{S}\}_2]$ and $[\text{Me}_2\text{Sn}\{2\text{-(Ph}_2\text{PO)-6-(Me}_3\text{Si)C}_6\text{H}_3\text{S}\}_2]$, 59
- Trien
Synthesis of dioxazaborocines from *N,N'*-alkylbridged-bis(bis(2-hydroxybenzyl)aminomethyl)amines, 215
- Trimesitylgermylcarbodiimide
Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153
- Trimesitylgermylcyanamide
Bis(trimesitylgermylcarbodiimido)germylene, trimesitylgermylcyanamide and trimesitylgermylcarbodiimide, 153
- Trimethyl(phenyl)stannane
Rhodium-catalyzed phenylation of *N*-arylsulfonyl aldimines with sodium tetraphenylborate or trimethyl(phenyl)stannane, 31
- Tungsten
Reactions of $[\text{Pt}_2(\mu\text{-S})_2(\text{PPh}_3)_4]$ with Group 6 and 7 metal carbonyls; crystal structure of the apparently unsaturated heterometallic complex $[\text{Pt}_2\text{Re}(\mu_3\text{-S})_2(\text{PPh}_3)_4(\text{CO})_3]^+$, 276
- Unsaturated ligand
Convenient routes to mono- and dinuclear (C_5Me_5)iron(II) complexes bearing acyl, alkynyl, aryl and thienyl ligands, 81
- X-ray crystallography
The X-ray crystallographic study of the reaction of bis(2-thienyl)ditelluride with tetrakis(triphenylphosphine)platinum or -palladium, 232
- X-ray crystal structure determination
Tetrameric thienylcopper and pentanuclear thienylcuprate, 102
- X-ray diffraction
Monomeric organoaluminium complexes $\text{RAl}(\text{OR}^*)_2$ and R_2AlOR^* with an optically active amino alkoxide ligand, 21
- Zirconacyclopentadienes
Convenient route for the preparation of unsymmetrical phosphoranes via zirconacyclopentadienes, 261
- Zirconium
Synthesis and characterization of Group 4 metallocene complexes with two disiloxanedyl bridges, 147
- Zirconocene complex
P–H functionalised phosphinoalkoxides as ligands: synthesis and catalytic properties of $[\text{Cp}_2\text{Zr}(\text{Me})(\text{cyclo-1-O-2-PH}(\text{Tipp})\text{-C}_6\text{H}_{10})]$ (Tipp = 2,4,6- $\text{Pr}_3\text{C}_6\text{H}_2$), $[\text{Cp}_2\text{Zr}(\text{Me})(\text{cyclo-1-O-2-PH}(\text{Tipp})\{\text{Mo}(\text{CO})_5\}\text{-C}_6\text{H}_{10})]$ and $[\text{Cp}_2\text{Zr}(\text{BH}_4)(\text{cyclo-1-O-2-PH}(\text{Mes})(\text{BH}_3)\text{-C}_6\text{H}_{10})]$ (Mes = 2,4,6- $\text{Me}_3\text{C}_6\text{H}_2$), and molecular structure of $(\text{R}_\text{C},\text{R}_\text{C},\text{R}_\text{P}/\text{S}_\text{C},\text{S}_\text{C},\text{S}_\text{P})\text{-}[\text{Cp}_2\text{Zr}(\text{Me})(\text{cyclo-1-O-2-PH}(\text{Tipp})\text{-C}_6\text{H}_{10})]$, 126